

Table 2.4
All Cancer Sites (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|--------|---------|--------|--------|---------|--------|--------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 400.38 | 466.77 | 365.83 | 402.13 | 468.60 | 369.56 | 426.21 | 525.13 | 356.43 |
| 1976 | 407.39 | 481.16 | 367.39 | 409.73 | 481.86 | 373.20 | 427.38 | 553.02 | 340.44 |
| 1977 | 407.73 | 486.49 | 363.69 | 409.28 | 487.10 | 367.86 | 442.62 | 577.65 | 352.01 |
| 1978 | 407.31 | 488.43 | 362.17 | 409.80 | 489.83 | 367.91 | 436.33 | 570.52 | 343.90 |
| 1979 | 412.47 | 496.49 | 365.46 | 414.09 | 497.62 | 369.66 | 451.54 | 589.90 | 358.94 |
| 1980 | 417.94 | 505.70 | 368.00 | 419.42 | 505.16 | 373.29 | 467.16 | 614.68 | 366.32 |
| 1981 | 425.23 | 510.87 | 377.04 | 427.68 | 510.90 | 383.85 | 472.65 | 633.89 | 362.85 |
| 1982 | 424.45 | 511.07 | 376.13 | 427.75 | 512.48 | 383.39 | 465.24 | 621.86 | 359.54 |
| 1983 | 431.13 | 519.92 | 380.75 | 434.40 | 520.93 | 388.01 | 486.67 | 645.50 | 381.75 |
| 1984 | 439.88 | 525.37 | 392.34 | 443.51 | 527.51 | 399.47 | 487.89 | 643.89 | 385.60 |
| 1985 | 448.73 | 531.08 | 403.06 | 453.36 | 533.39 | 411.46 | 488.36 | 642.77 | 387.63 |
| 1986 | 451.27 | 538.15 | 402.16 | 455.89 | 542.83 | 408.98 | 493.67 | 643.90 | 396.23 |
| 1987 | 467.93 | 563.07 | 413.18 | 474.83 | 568.67 | 423.31 | 500.86 | 665.97 | 393.93 |
| 1988 | 463.70 | 558.30 | 409.05 | 470.12 | 564.95 | 417.67 | 503.74 | 657.37 | 404.16 |
| 1989 | 467.52 | 566.79 | 408.09 | 474.87 | 573.24 | 418.31 | 492.25 | 660.66 | 384.43 |
| 1990 | 481.65 | 592.31 | 415.26 | 488.57 | 598.84 | 424.61 | 520.53 | 698.47 | 407.60 |
| 1991 | 502.69 | 637.14 | 418.54 | 510.01 | 644.02 | 428.04 | 548.68 | 764.00 | 412.53 |
| 1992 | 510.22 | 657.61 | 414.60 | 516.46 | 662.70 | 423.39 | 567.59 | 807.11 | 412.60 |
| 1993 | 492.99 | 621.48 | 408.04 | 496.58 | 619.62 | 416.90 | 567.33 | 816.59 | 403.48 |
| 1994 | 482.92 | 589.01 | 413.00 | 486.81 | 586.74 | 422.57 | 556.45 | 781.26 | 408.05 |
| 1995 | 476.09 | 570.49 | 414.22 | 481.25 | 569.00 | 425.39 | 540.60 | 744.38 | 404.30 |
| 1996 | 478.28 | 571.34 | 417.39 | 483.85 | 572.40 | 427.49 | 538.33 | 724.41 | 415.06 |
| 1997 | 485.51 | 574.43 | 427.67 | 492.53 | 575.12 | 440.33 | 537.53 | 734.55 | 408.95 |
| 1998 | 486.93 | 571.03 | 432.87 | 495.45 | 574.48 | 446.04 | 531.57 | 716.49 | 412.59 |
| 1999 | 489.15 | 579.13 | 429.27 | 496.97 | 580.65 | 442.31 | 536.68 | 716.98 | 418.40 |
| 2000 | 484.03 | 576.80 | 420.65 | 494.13 | 580.18 | 436.40 | 522.60 | 705.44 | 400.05 |
| 2001 | 486.54 | 576.35 | 424.52 | 497.31 | 582.10 | 439.68 | 513.68 | 689.00 | 395.21 |
| 2002 | 482.96 | 568.14 | 424.40 | 492.61 | 574.07 | 437.14 | 526.46 | 687.19 | 418.40 |
| 2003 | 469.33 | 551.99 | 412.30 | 479.20 | 556.17 | 426.83 | 508.94 | 664.99 | 403.91 |
| 2004 | 467.70 | 546.61 | 412.98 | 477.13 | 551.69 | 425.77 | 506.12 | 656.99 | 405.70 |
| 2005 | 459.94 | 531.85 | 410.21 | 470.39 | 538.84 | 423.71 | 486.53 | 608.51 | 404.72 |
| 2006 | 456.20 | 527.31 | 406.01 | 464.31 | 530.15 | 418.49 | 477.70 | 598.85 | 394.60 |
| 1975-2006 | 461.24 | 551.98 | 404.12 | 466.62 | 554.34 | 413.30 | 504.87 | 672.06 | 394.00 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 2.5
All Cancer Sites (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|--------|---------|--------|--------|---------|--------|--------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 199.14 | 258.43 | 160.03 | 196.33 | 254.00 | 158.99 | 235.50 | 320.45 | 174.17 |
| 1976 | 202.30 | 262.23 | 162.70 | 199.38 | 257.50 | 161.69 | 241.76 | 328.45 | 179.10 |
| 1977 | 203.02 | 264.23 | 162.94 | 199.67 | 258.78 | 161.61 | 246.38 | 337.95 | 181.60 |
| 1978 | 204.42 | 267.07 | 163.76 | 201.14 | 261.65 | 162.54 | 247.24 | 340.91 | 181.53 |
| 1979 | 204.53 | 268.65 | 163.06 | 201.20 | 263.24 | 161.70 | 248.98 | 344.67 | 182.73 |
| 1980 | 206.96 | 271.08 | 165.75 | 203.27 | 265.03 | 164.17 | 256.18 | 353.91 | 189.25 |
| 1981 | 206.40 | 269.78 | 165.90 | 202.64 | 263.27 | 164.55 | 256.77 | 358.65 | 187.47 |
| 1982 | 208.32 | 272.89 | 167.45 | 204.54 | 266.28 | 166.11 | 260.36 | 364.86 | 190.17 |
| 1983 | 209.21 | 274.09 | 168.27 | 205.17 | 267.22 | 166.69 | 263.75 | 369.32 | 193.24 |
| 1984 | 210.91 | 275.08 | 170.55 | 206.66 | 267.55 | 169.02 | 268.15 | 377.93 | 195.67 |
| 1985 | 211.35 | 275.68 | 170.96 | 207.25 | 268.25 | 169.58 | 268.06 | 377.82 | 196.31 |
| 1986 | 211.79 | 276.04 | 171.59 | 207.73 | 268.83 | 170.01 | 269.45 | 378.92 | 199.05 |
| 1987 | 211.91 | 276.34 | 171.54 | 207.78 | 268.96 | 169.87 | 271.12 | 383.24 | 199.91 |
| 1988 | 212.62 | 276.89 | 172.53 | 208.48 | 269.46 | 170.94 | 271.59 | 384.17 | 200.36 |
| 1989 | 214.30 | 278.50 | 174.25 | 209.96 | 270.25 | 172.88 | 275.87 | 395.28 | 200.87 |
| 1990 | 214.95 | 279.82 | 174.65 | 210.41 | 271.48 | 172.89 | 279.30 | 399.14 | 205.39 |
| 1991 | 215.06 | 279.15 | 175.30 | 210.62 | 271.19 | 173.41 | 278.97 | 395.85 | 207.39 |
| 1992 | 213.46 | 276.51 | 174.39 | 209.24 | 268.91 | 172.58 | 274.93 | 389.20 | 205.80 |
| 1993 | 213.45 | 275.89 | 174.55 | 209.16 | 267.89 | 172.82 | 277.09 | 394.76 | 206.29 |
| 1994 | 211.74 | 272.12 | 174.11 | 207.83 | 264.73 | 172.62 | 271.96 | 385.20 | 204.06 |
| 1995 | 209.89 | 268.50 | 173.35 | 206.09 | 261.46 | 171.73 | 269.51 | 379.02 | 204.53 |
| 1996 | 207.00 | 263.75 | 171.21 | 203.40 | 256.83 | 169.85 | 265.28 | 373.70 | 200.94 |
| 1997 | 203.55 | 258.13 | 169.00 | 199.94 | 251.58 | 167.35 | 262.14 | 363.18 | 202.53 |
| 1998 | 200.82 | 253.63 | 166.93 | 197.56 | 247.62 | 165.38 | 255.66 | 353.13 | 198.38 |
| 1999 | 200.72 | 252.75 | 167.24 | 197.75 | 247.14 | 165.92 | 253.80 | 349.23 | 197.54 |
| 2000 | 198.69 | 248.21 | 166.64 | 196.28 | 243.14 | 165.92 | 248.05 | 340.92 | 193.19 |
| 2001 | 195.89 | 244.08 | 164.33 | 193.59 | 239.42 | 163.46 | 244.04 | 332.36 | 191.92 |
| 2002 | 193.71 | 240.22 | 162.87 | 191.74 | 236.33 | 162.04 | 240.27 | 322.97 | 191.05 |
| 2003 | 190.06 | 234.18 | 160.47 | 188.29 | 230.69 | 159.69 | 234.47 | 311.39 | 188.17 |
| 2004 | 185.92 | 228.61 | 157.10 | 184.29 | 225.12 | 156.54 | 228.41 | 304.01 | 182.97 |
| 2005 | 184.24 | 226.44 | 155.52 | 182.86 | 223.47 | 155.02 | 224.39 | 297.21 | 180.43 |
| 2006 | 181.07 | 221.11 | 153.66 | 180.10 | 218.74 | 153.44 | 218.79 | 287.75 | 176.92 |
| 1975-2006 | 203.72 | 260.43 | 167.11 | 200.61 | 254.69 | 165.92 | 255.30 | 353.38 | 193.65 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 3.4
Cancer of the Brain and Other Nervous System (Invasive)
Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 5.85 | 6.83 | 5.01 | 6.21 | 7.31 | 5.28 | 4.13 | 4.27 | 3.88 |
| 1976 | 5.82 | 7.14 | 4.68 | 6.19 | 7.51 | 5.05 | 3.32 | 5.37 | - |
| 1977 | 6.17 | 7.76 | 4.88 | 6.60 | 8.26 | 5.26 | 3.55 | 5.17 | 2.47 |
| 1978 | 5.76 | 6.79 | 4.91 | 6.10 | 7.20 | 5.19 | 3.86 | 4.34 | 3.51 |
| 1979 | 6.12 | 7.41 | 5.01 | 6.60 | 8.02 | 5.37 | 3.67 | 4.19 | 3.20 |
| 1980 | 6.29 | 7.57 | 5.24 | 6.80 | 8.18 | 5.65 | 3.14 | 3.35 | 2.92 |
| 1981 | 6.50 | 8.07 | 5.18 | 6.89 | 8.45 | 5.60 | 5.02 | 7.24 | 3.16 |
| 1982 | 6.43 | 7.93 | 5.25 | 6.92 | 8.58 | 5.64 | 3.71 | 4.40 | 3.05 |
| 1983 | 6.31 | 7.60 | 5.19 | 6.88 | 8.20 | 5.74 | 2.75 | 3.79 | 1.84 |
| 1984 | 6.12 | 7.16 | 5.22 | 6.49 | 7.63 | 5.51 | 4.52 | 5.31 | 3.88 |
| 1985 | 6.94 | 8.53 | 5.63 | 7.43 | 9.11 | 6.04 | 4.88 | 5.62 | 4.18 |
| 1986 | 6.84 | 8.07 | 5.80 | 7.36 | 8.62 | 6.30 | 4.13 | 4.75 | 3.54 |
| 1987 | 6.99 | 8.12 | 6.03 | 7.59 | 8.71 | 6.68 | 4.18 | 5.49 | 3.13 |
| 1988 | 6.83 | 7.87 | 5.92 | 7.39 | 8.56 | 6.39 | 4.42 | 4.48 | 4.31 |
| 1989 | 6.85 | 8.49 | 5.46 | 7.42 | 9.20 | 5.91 | 4.90 | 5.93 | 4.12 |
| 1990 | 7.04 | 8.11 | 6.15 | 7.58 | 8.76 | 6.61 | 4.33 | 5.18 | 3.64 |
| 1991 | 6.94 | 8.55 | 5.63 | 7.60 | 9.28 | 6.22 | 3.66 | 4.51 | 3.09 |
| 1992 | 6.97 | 8.66 | 5.61 | 7.63 | 9.46 | 6.16 | 3.61 | 4.01 | 3.28 |
| 1993 | 6.77 | 8.08 | 5.67 | 7.39 | 8.89 | 6.14 | 3.93 | 3.97 | 3.80 |
| 1994 | 6.59 | 7.95 | 5.41 | 7.10 | 8.52 | 5.89 | 4.24 | 4.52 | 3.97 |
| 1995 | 6.47 | 7.91 | 5.31 | 7.03 | 8.56 | 5.78 | 3.97 | 5.02 | 3.15 |
| 1996 | 6.65 | 8.21 | 5.31 | 7.25 | 8.96 | 5.76 | 4.02 | 5.09 | 3.37 |
| 1997 | 6.73 | 8.08 | 5.58 | 7.39 | 8.87 | 6.11 | 4.03 | 5.30 | 3.12 |
| 1998 | 6.60 | 7.71 | 5.67 | 7.24 | 8.44 | 6.23 | 4.04 | 4.26 | 3.85 |
| 1999 | 6.89 | 8.36 | 5.64 | 7.67 | 9.23 | 6.34 | 3.70 | 5.12 | 2.56 |
| 2000 | 6.77 | 8.18 | 5.62 | 7.46 | 9.04 | 6.15 | 4.39 | 5.39 | 3.62 |
| 2001 | 6.55 | 7.90 | 5.44 | 7.19 | 8.57 | 6.02 | 4.10 | 5.27 | 3.31 |
| 2002 | 6.67 | 7.98 | 5.59 | 7.40 | 8.81 | 6.23 | 4.22 | 5.30 | 3.47 |
| 2003 | 6.56 | 7.51 | 5.77 | 7.14 | 8.21 | 6.23 | 4.60 | 4.37 | 4.65 |
| 2004 | 6.67 | 7.76 | 5.66 | 7.44 | 8.61 | 6.34 | 3.95 | 4.43 | 3.44 |
| 2005 | 6.60 | 7.63 | 5.72 | 7.31 | 8.52 | 6.25 | 4.43 | 4.88 | 4.07 |
| 2006 | 6.12 | 7.41 | 5.02 | 6.81 | 8.29 | 5.49 | 3.51 | 3.95 | 3.28 |
| 1975-2006 | 6.57 | 7.88 | 5.47 | 7.14 | 8.55 | 5.96 | 4.05 | 4.81 | 3.43 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 3.5
Cancer of the Brain and Other Nervous System (Invasive)
Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 4.11 | 4.96 | 3.40 | 4.32 | 5.20 | 3.60 | 2.41 | 3.20 | 1.75 |
| 1976 | 4.34 | 5.28 | 3.56 | 4.54 | 5.50 | 3.74 | 2.66 | 3.54 | 1.98 |
| 1977 | 4.40 | 5.39 | 3.59 | 4.63 | 5.69 | 3.77 | 2.54 | 2.99 | 2.18 |
| 1978 | 4.53 | 5.54 | 3.71 | 4.76 | 5.82 | 3.91 | 2.70 | 3.36 | 2.18 |
| 1979 | 4.26 | 5.25 | 3.45 | 4.48 | 5.50 | 3.64 | 2.63 | 3.30 | 2.11 |
| 1980 | 4.37 | 5.45 | 3.47 | 4.62 | 5.77 | 3.66 | 2.46 | 3.04 | 2.02 |
| 1981 | 4.36 | 5.33 | 3.59 | 4.60 | 5.58 | 3.80 | 2.55 | 3.31 | 1.98 |
| 1982 | 4.43 | 5.38 | 3.67 | 4.67 | 5.67 | 3.88 | 2.64 | 3.37 | 2.09 |
| 1983 | 4.39 | 5.31 | 3.64 | 4.66 | 5.64 | 3.85 | 2.49 | 2.91 | 2.18 |
| 1984 | 4.55 | 5.57 | 3.74 | 4.82 | 5.86 | 4.00 | 2.56 | 3.35 | 1.94 |
| 1985 | 4.57 | 5.60 | 3.75 | 4.85 | 5.92 | 3.99 | 2.54 | 3.22 | 2.02 |
| 1986 | 4.53 | 5.59 | 3.68 | 4.78 | 5.90 | 3.87 | 2.77 | 3.47 | 2.27 |
| 1987 | 4.71 | 5.68 | 3.93 | 4.99 | 6.00 | 4.17 | 2.89 | 3.55 | 2.36 |
| 1988 | 4.72 | 5.89 | 3.79 | 4.98 | 6.21 | 4.00 | 2.91 | 3.63 | 2.39 |
| 1989 | 4.73 | 5.82 | 3.88 | 5.01 | 6.15 | 4.12 | 2.73 | 3.47 | 2.23 |
| 1990 | 4.87 | 5.97 | 3.99 | 5.16 | 6.30 | 4.24 | 2.97 | 3.66 | 2.42 |
| 1991 | 4.95 | 6.01 | 4.11 | 5.27 | 6.37 | 4.37 | 2.84 | 3.42 | 2.41 |
| 1992 | 4.85 | 5.88 | 4.03 | 5.17 | 6.26 | 4.30 | 2.82 | 3.41 | 2.39 |
| 1993 | 4.79 | 5.87 | 3.91 | 5.09 | 6.23 | 4.17 | 2.83 | 3.59 | 2.27 |
| 1994 | 4.84 | 5.88 | 3.97 | 5.17 | 6.26 | 4.25 | 2.69 | 3.23 | 2.26 |
| 1995 | 4.67 | 5.67 | 3.86 | 4.97 | 6.02 | 4.11 | 2.77 | 3.40 | 2.33 |
| 1996 | 4.73 | 5.74 | 3.91 | 5.07 | 6.14 | 4.19 | 2.64 | 3.24 | 2.21 |
| 1997 | 4.68 | 5.66 | 3.90 | 4.99 | 6.03 | 4.13 | 2.87 | 3.30 | 2.55 |
| 1998 | 4.68 | 5.70 | 3.83 | 5.00 | 6.06 | 4.12 | 2.73 | 3.39 | 2.24 |
| 1999 | 4.64 | 5.66 | 3.81 | 4.98 | 6.04 | 4.10 | 2.73 | 3.28 | 2.29 |
| 2000 | 4.53 | 5.57 | 3.67 | 4.86 | 5.96 | 3.93 | 2.70 | 3.28 | 2.24 |
| 2001 | 4.45 | 5.49 | 3.59 | 4.78 | 5.87 | 3.86 | 2.58 | 3.13 | 2.19 |
| 2002 | 4.45 | 5.46 | 3.61 | 4.78 | 5.82 | 3.89 | 2.64 | 3.21 | 2.21 |
| 2003 | 4.40 | 5.36 | 3.62 | 4.73 | 5.72 | 3.91 | 2.67 | 3.39 | 2.11 |
| 2004 | 4.31 | 5.22 | 3.53 | 4.63 | 5.59 | 3.81 | 2.47 | 3.01 | 2.04 |
| 2005 | 4.34 | 5.32 | 3.53 | 4.68 | 5.68 | 3.83 | 2.54 | 3.27 | 2.00 |
| 2006 | 4.17 | 5.13 | 3.38 | 4.52 | 5.53 | 3.66 | 2.36 | 2.94 | 1.94 |
| 1975-2006 | 4.56 | 5.58 | 3.74 | 4.86 | 5.92 | 3.99 | 2.67 | 3.30 | 2.20 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 4.5
Cancer of the Breast (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 58.05 | 0.85 | 105.06 | 59.44 | 0.88 | 107.32 | 52.10 | - | 93.56 |
| 1976 | 56.44 | 1.10 | 101.90 | 58.12 | 1.12 | 104.80 | 48.03 | - | 85.34 |
| 1977 | 55.98 | 0.96 | 100.82 | 57.47 | 0.88 | 103.43 | 49.67 | - | 87.03 |
| 1978 | 55.84 | 0.90 | 100.61 | 57.54 | 0.89 | 103.61 | 48.27 | - | 86.09 |
| 1979 | 56.83 | 0.99 | 102.08 | 58.32 | 0.94 | 104.78 | 49.86 | - | 86.85 |
| 1980 | 56.88 | 0.90 | 102.19 | 58.53 | 0.95 | 105.09 | 50.52 | - | 89.35 |
| 1981 | 59.29 | 1.00 | 106.29 | 61.23 | 0.93 | 109.89 | 54.35 | - | 94.24 |
| 1982 | 59.22 | 0.98 | 106.45 | 61.08 | 0.94 | 109.97 | 53.98 | - | 93.66 |
| 1983 | 61.90 | 1.04 | 111.14 | 63.75 | 1.13 | 114.55 | 58.83 | - | 103.35 |
| 1984 | 64.46 | 0.92 | 115.96 | 66.59 | 0.93 | 119.93 | 58.32 | - | 102.29 |
| 1985 | 69.08 | 0.95 | 124.18 | 71.03 | 0.97 | 127.89 | 64.00 | - | 111.49 |
| 1986 | 70.44 | 0.94 | 126.79 | 72.31 | 0.95 | 130.42 | 66.21 | - | 114.95 |
| 1987 | 74.76 | 0.96 | 134.45 | 77.83 | 0.98 | 140.31 | 63.01 | - | 109.33 |
| 1988 | 73.06 | 1.23 | 131.27 | 75.64 | 1.17 | 136.36 | 68.95 | 2.76 | 118.57 |
| 1989 | 70.70 | 1.13 | 127.13 | 73.35 | 1.08 | 132.32 | 60.47 | - | 104.83 |
| 1990 | 73.10 | 1.13 | 131.72 | 75.32 | 1.15 | 136.20 | 68.25 | - | 118.09 |
| 1991 | 74.03 | 1.13 | 133.72 | 76.56 | 1.19 | 138.74 | 67.39 | - | 117.38 |
| 1992 | 72.97 | 1.04 | 131.94 | 74.75 | 1.02 | 135.78 | 71.32 | - | 123.52 |
| 1993 | 71.28 | 1.08 | 129.11 | 73.16 | 1.04 | 133.14 | 68.82 | - | 119.44 |
| 1994 | 72.01 | 1.13 | 130.83 | 74.25 | 1.08 | 135.62 | 70.41 | - | 122.13 |
| 1995 | 72.77 | 0.99 | 132.46 | 75.08 | 0.94 | 137.51 | 71.63 | - | 123.76 |
| 1996 | 73.25 | 1.32 | 133.49 | 75.28 | 1.33 | 138.02 | 71.22 | - | 123.51 |
| 1997 | 75.48 | 1.14 | 137.75 | 77.59 | 1.12 | 142.60 | 72.19 | - | 124.80 |
| 1998 | 77.18 | 1.07 | 141.12 | 79.65 | 1.13 | 146.62 | 70.50 | - | 122.84 |
| 1999 | 76.95 | 1.03 | 141.00 | 79.41 | 1.02 | 146.70 | 72.24 | 2.24 | 124.83 |
| 2000 | 74.18 | 1.19 | 136.13 | 77.14 | 1.16 | 142.79 | 67.41 | 2.94 | 116.00 |
| 2001 | 75.10 | 1.21 | 137.91 | 77.79 | 1.21 | 144.14 | 65.79 | 1.55 | 113.94 |
| 2002 | 73.01 | 1.16 | 134.57 | 75.30 | 1.13 | 140.12 | 69.92 | 2.06 | 121.07 |
| 2003 | 68.06 | 1.31 | 125.46 | 69.93 | 1.34 | 130.15 | 70.02 | 1.60 | 121.54 |
| 2004 | 68.13 | 1.21 | 125.89 | 69.60 | 1.25 | 129.85 | 69.28 | 1.57 | 120.34 |
| 2005 | 67.02 | 1.15 | 124.10 | 68.83 | 1.25 | 128.75 | 65.80 | - | 114.44 |
| 2006 | 66.41 | 1.27 | 123.04 | 67.62 | 1.34 | 126.61 | 68.51 | 1.48 | 119.22 |
| 1975-2006 | 68.58 | 1.09 | 124.60 | 70.51 | 1.09 | 128.61 | 64.76 | 1.60 | 112.63 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 4.6
Cancer of the Breast (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 17.83 | 0.37 | 31.45 | 18.03 | 0.38 | 31.79 | 16.82 | 0.41 | 29.49 |
| 1976 | 18.06 | 0.38 | 31.80 | 18.25 | 0.36 | 32.17 | 17.61 | 0.60 | 30.47 |
| 1977 | 18.47 | 0.36 | 32.48 | 18.56 | 0.35 | 32.67 | 18.90 | 0.51 | 32.80 |
| 1978 | 18.05 | 0.36 | 31.73 | 18.12 | 0.35 | 31.90 | 18.63 | 0.60 | 32.14 |
| 1979 | 17.74 | 0.28 | 31.21 | 17.87 | 0.26 | 31.48 | 17.87 | 0.54 | 30.82 |
| 1980 | 18.04 | 0.30 | 31.68 | 18.15 | 0.30 | 31.93 | 18.37 | 0.42 | 31.68 |
| 1981 | 18.20 | 0.32 | 31.92 | 18.28 | 0.31 | 32.12 | 18.95 | 0.46 | 32.55 |
| 1982 | 18.39 | 0.33 | 32.19 | 18.42 | 0.32 | 32.31 | 19.68 | 0.56 | 33.75 |
| 1983 | 18.33 | 0.31 | 32.07 | 18.36 | 0.29 | 32.20 | 19.57 | 0.50 | 33.53 |
| 1984 | 18.79 | 0.29 | 32.90 | 18.76 | 0.28 | 32.90 | 20.97 | 0.47 | 35.94 |
| 1985 | 18.84 | 0.34 | 32.98 | 18.88 | 0.34 | 33.11 | 20.35 | 0.43 | 34.85 |
| 1986 | 18.76 | 0.28 | 32.87 | 18.76 | 0.27 | 32.93 | 20.69 | 0.42 | 35.44 |
| 1987 | 18.66 | 0.34 | 32.66 | 18.57 | 0.34 | 32.57 | 21.53 | 0.47 | 36.73 |
| 1988 | 18.98 | 0.33 | 33.20 | 18.85 | 0.28 | 33.09 | 22.35 | 0.88 | 37.78 |
| 1989 | 19.00 | 0.33 | 33.23 | 18.95 | 0.33 | 33.23 | 21.56 | 0.45 | 36.61 |
| 1990 | 18.93 | 0.28 | 33.14 | 18.79 | 0.26 | 33.02 | 22.49 | 0.52 | 38.00 |
| 1991 | 18.66 | 0.28 | 32.69 | 18.46 | 0.27 | 32.45 | 22.62 | 0.44 | 38.28 |
| 1992 | 18.07 | 0.31 | 31.64 | 17.89 | 0.30 | 31.43 | 21.88 | 0.45 | 37.09 |
| 1993 | 17.98 | 0.36 | 31.39 | 17.74 | 0.34 | 31.09 | 22.62 | 0.66 | 38.04 |
| 1994 | 17.67 | 0.35 | 30.92 | 17.40 | 0.34 | 30.56 | 22.44 | 0.60 | 37.74 |
| 1995 | 17.44 | 0.35 | 30.55 | 17.11 | 0.32 | 30.09 | 22.70 | 0.77 | 38.18 |
| 1996 | 16.83 | 0.34 | 29.49 | 16.52 | 0.31 | 29.05 | 22.09 | 0.72 | 37.13 |
| 1997 | 16.07 | 0.32 | 28.21 | 15.67 | 0.30 | 27.62 | 22.25 | 0.63 | 37.43 |
| 1998 | 15.67 | 0.32 | 27.54 | 15.32 | 0.31 | 27.01 | 21.08 | 0.56 | 35.53 |
| 1999 | 15.18 | 0.34 | 26.61 | 14.77 | 0.32 | 25.98 | 20.93 | 0.69 | 35.21 |
| 2000 | 15.18 | 0.37 | 26.63 | 14.85 | 0.35 | 26.17 | 20.47 | 0.63 | 34.36 |
| 2001 | 14.74 | 0.35 | 25.98 | 14.34 | 0.33 | 25.39 | 20.50 | 0.65 | 34.52 |
| 2002 | 14.48 | 0.31 | 25.56 | 14.07 | 0.28 | 24.95 | 20.28 | 0.60 | 34.12 |
| 2003 | 14.23 | 0.31 | 25.19 | 13.81 | 0.30 | 24.56 | 20.19 | 0.46 | 34.06 |
| 2004 | 13.74 | 0.29 | 24.40 | 13.36 | 0.27 | 23.82 | 19.09 | 0.54 | 32.29 |
| 2005 | 13.52 | 0.29 | 24.03 | 13.08 | 0.28 | 23.36 | 19.37 | 0.50 | 32.83 |
| 2006 | 13.16 | 0.29 | 23.45 | 12.78 | 0.28 | 22.89 | 18.73 | 0.47 | 31.65 |
| 1975-2006 | 16.89 | 0.32 | 29.70 | 16.75 | 0.31 | 29.54 | 20.50 | 0.56 | 34.81 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 4.7
Cancer of the Female Breast (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

| | All Races, Females | | | White Females | | | Black Females | | |
|--------------------|--------------------|-------|--------|---------------|-------|--------|---------------|-------|--------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| Year of Diagnosis: | | | | | | | | | |
| 1975 | 105.06 | 40.61 | 273.82 | 107.32 | 40.71 | 281.77 | 93.56 | 42.59 | 227.04 |
| 1976 | 101.90 | 40.04 | 263.90 | 104.80 | 40.85 | 272.27 | 85.34 | 37.82 | 209.77 |
| 1977 | 100.82 | 39.12 | 262.40 | 103.43 | 39.40 | 271.08 | 87.03 | 37.62 | 216.41 |
| 1978 | 100.61 | 38.85 | 262.34 | 103.61 | 39.29 | 272.02 | 86.09 | 40.24 | 206.14 |
| 1979 | 102.08 | 37.97 | 269.95 | 104.78 | 38.63 | 278.00 | 86.85 | 37.78 | 215.34 |
| 1980 | 102.19 | 37.76 | 270.92 | 105.09 | 38.28 | 280.03 | 89.35 | 37.61 | 224.82 |
| 1981 | 106.29 | 39.19 | 282.02 | 109.89 | 40.12 | 292.60 | 94.24 | 38.86 | 239.26 |
| 1982 | 106.45 | 40.54 | 279.04 | 109.97 | 41.29 | 289.82 | 93.66 | 38.11 | 239.13 |
| 1983 | 111.14 | 40.24 | 296.79 | 114.55 | 40.92 | 307.37 | 103.35 | 41.24 | 266.00 |
| 1984 | 115.96 | 42.35 | 308.70 | 119.93 | 43.13 | 321.04 | 102.29 | 44.76 | 252.92 |
| 1985 | 124.18 | 44.35 | 333.24 | 127.89 | 44.56 | 346.11 | 111.49 | 45.45 | 284.41 |
| 1986 | 126.79 | 44.43 | 342.46 | 130.42 | 45.24 | 353.46 | 114.95 | 45.49 | 296.84 |
| 1987 | 134.45 | 45.49 | 367.41 | 140.31 | 46.55 | 385.82 | 109.33 | 43.13 | 282.66 |
| 1988 | 131.27 | 44.01 | 359.75 | 136.36 | 45.09 | 375.36 | 118.57 | 42.95 | 316.57 |
| 1989 | 127.13 | 43.36 | 346.48 | 132.32 | 44.46 | 362.42 | 104.83 | 42.16 | 268.93 |
| 1990 | 131.72 | 45.56 | 357.35 | 136.20 | 45.38 | 374.03 | 118.09 | 48.71 | 299.77 |
| 1991 | 133.72 | 47.16 | 360.39 | 138.74 | 47.13 | 378.63 | 117.38 | 50.93 | 291.39 |
| 1992 | 131.94 | 43.43 | 363.71 | 135.78 | 43.68 | 376.95 | 123.52 | 46.80 | 324.43 |
| 1993 | 129.11 | 42.79 | 355.14 | 133.14 | 43.26 | 368.48 | 119.44 | 45.31 | 313.55 |
| 1994 | 130.83 | 42.35 | 362.50 | 135.62 | 42.74 | 378.86 | 122.13 | 47.27 | 318.15 |
| 1995 | 132.46 | 43.15 | 366.34 | 137.51 | 43.76 | 382.99 | 123.76 | 45.36 | 329.03 |
| 1996 | 133.49 | 43.46 | 369.26 | 138.02 | 43.92 | 384.42 | 123.51 | 44.11 | 331.44 |
| 1997 | 137.75 | 43.43 | 384.75 | 142.60 | 44.05 | 400.68 | 124.80 | 44.18 | 335.91 |
| 1998 | 141.12 | 45.36 | 391.88 | 146.62 | 45.72 | 410.81 | 122.84 | 46.15 | 323.67 |
| 1999 | 141.00 | 43.60 | 396.05 | 146.70 | 44.93 | 413.21 | 124.83 | 43.36 | 338.18 |
| 2000 | 136.13 | 43.34 | 379.11 | 142.79 | 43.93 | 401.69 | 116.00 | 43.85 | 304.94 |
| 2001 | 137.91 | 43.53 | 385.08 | 144.14 | 44.44 | 405.22 | 113.94 | 39.14 | 309.80 |
| 2002 | 134.57 | 42.34 | 376.08 | 140.12 | 43.03 | 394.35 | 121.07 | 40.31 | 332.56 |
| 2003 | 125.46 | 42.68 | 342.22 | 130.15 | 43.50 | 357.07 | 121.54 | 42.33 | 328.97 |
| 2004 | 125.89 | 43.97 | 340.40 | 129.85 | 43.91 | 354.90 | 120.34 | 45.38 | 316.63 |
| 2005 | 124.10 | 42.97 | 336.52 | 128.75 | 43.29 | 352.56 | 114.44 | 43.75 | 299.54 |
| 2006 | 123.04 | 42.79 | 333.19 | 126.61 | 43.47 | 344.32 | 119.22 | 42.66 | 319.69 |
| 1975-2006 | 124.60 | 42.86 | 338.67 | 128.61 | 43.40 | 351.77 | 112.63 | 43.44 | 293.81 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 4.8
Cancer of the Female Breast (Invasive)
Age-adjusted U.S. Death^a Rates by Year, Race and Age

| Year of Death: | All Races, Females | | | White Females | | | Black Females | | |
|----------------|--------------------|------|-------|---------------|------|-------|---------------|-------|--------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| 1975 | 31.45 | 9.11 | 89.94 | 31.79 | 9.01 | 91.42 | 29.49 | 10.68 | 78.75 |
| 1976 | 31.80 | 8.74 | 92.21 | 32.17 | 8.64 | 93.80 | 30.47 | 10.22 | 83.50 |
| 1977 | 32.48 | 8.86 | 94.34 | 32.67 | 8.75 | 95.32 | 32.80 | 10.52 | 91.13 |
| 1978 | 31.73 | 8.69 | 92.08 | 31.90 | 8.46 | 93.29 | 32.14 | 11.37 | 86.53 |
| 1979 | 31.21 | 8.58 | 90.47 | 31.48 | 8.34 | 92.07 | 30.82 | 11.36 | 81.78 |
| 1980 | 31.68 | 8.57 | 92.22 | 31.93 | 8.39 | 93.56 | 31.68 | 10.82 | 86.31 |
| 1981 | 31.92 | 8.46 | 93.33 | 32.12 | 8.20 | 94.77 | 32.55 | 11.48 | 87.74 |
| 1982 | 32.19 | 8.42 | 94.43 | 32.31 | 8.19 | 95.46 | 33.75 | 11.19 | 92.83 |
| 1983 | 32.07 | 8.16 | 94.70 | 32.20 | 7.89 | 95.85 | 33.53 | 11.17 | 92.10 |
| 1984 | 32.90 | 8.58 | 96.57 | 32.90 | 8.22 | 97.55 | 35.94 | 12.55 | 97.21 |
| 1985 | 32.98 | 8.56 | 96.90 | 33.11 | 8.30 | 98.09 | 34.85 | 12.04 | 94.58 |
| 1986 | 32.87 | 8.64 | 96.32 | 32.93 | 8.26 | 97.53 | 35.44 | 12.86 | 94.58 |
| 1987 | 32.66 | 8.44 | 96.08 | 32.57 | 7.99 | 96.93 | 36.73 | 12.94 | 99.04 |
| 1988 | 33.20 | 8.39 | 98.18 | 33.09 | 8.01 | 98.78 | 37.78 | 12.71 | 103.42 |
| 1989 | 33.23 | 8.42 | 98.22 | 33.23 | 8.06 | 99.16 | 36.61 | 12.41 | 99.96 |
| 1990 | 33.14 | 8.32 | 98.14 | 33.02 | 7.99 | 98.55 | 38.00 | 11.99 | 106.13 |
| 1991 | 32.69 | 8.16 | 96.91 | 32.45 | 7.73 | 97.15 | 38.28 | 12.31 | 106.29 |
| 1992 | 31.64 | 7.77 | 94.15 | 31.43 | 7.37 | 94.45 | 37.09 | 12.03 | 102.73 |
| 1993 | 31.39 | 7.41 | 94.18 | 31.09 | 6.99 | 94.20 | 38.04 | 11.65 | 107.14 |
| 1994 | 30.92 | 7.37 | 92.58 | 30.56 | 6.92 | 92.47 | 37.74 | 11.70 | 105.91 |
| 1995 | 30.55 | 7.29 | 91.46 | 30.09 | 6.80 | 91.07 | 38.18 | 11.71 | 107.49 |
| 1996 | 29.49 | 6.95 | 88.53 | 29.05 | 6.40 | 88.35 | 37.13 | 11.74 | 103.61 |
| 1997 | 28.21 | 6.71 | 84.53 | 27.62 | 6.22 | 83.64 | 37.43 | 11.03 | 106.56 |
| 1998 | 27.54 | 6.34 | 83.05 | 27.01 | 5.82 | 82.50 | 35.53 | 10.65 | 100.68 |
| 1999 | 26.61 | 5.75 | 81.22 | 25.98 | 5.20 | 80.40 | 35.21 | 10.27 | 100.52 |
| 2000 | 26.63 | 5.89 | 80.96 | 26.17 | 5.37 | 80.61 | 34.36 | 9.94 | 98.30 |
| 2001 | 25.98 | 5.86 | 78.66 | 25.39 | 5.35 | 77.86 | 34.52 | 9.86 | 99.07 |
| 2002 | 25.56 | 5.63 | 77.75 | 24.95 | 5.05 | 77.05 | 34.12 | 10.04 | 97.17 |
| 2003 | 25.19 | 5.57 | 76.56 | 24.56 | 5.06 | 75.60 | 34.06 | 9.66 | 97.96 |
| 2004 | 24.40 | 5.20 | 74.67 | 23.82 | 4.65 | 74.03 | 32.29 | 9.30 | 92.49 |
| 2005 | 24.03 | 5.19 | 73.37 | 23.36 | 4.62 | 72.42 | 32.83 | 9.53 | 93.86 |
| 2006 | 23.45 | 5.04 | 71.64 | 22.89 | 4.55 | 70.90 | 31.65 | 8.96 | 91.08 |
| 1975-2006 | 29.70 | 7.20 | 88.60 | 29.54 | 6.83 | 89.00 | 34.81 | 10.98 | 97.22 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 4.9
Cancer of the Female Breast (*In Situ*)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

| Year of Diagnosis: | All Races, Females | | | White Females | | | Black Females | | |
|--------------------|--------------------|-------|-------|---------------|-------|-------|---------------|-------|-------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| 1975 | 5.83 | 3.78 | 11.19 | 5.89 | 3.81 | 11.32 | 3.94 | - | 8.31 |
| 1976 | 5.28 | 2.72 | 11.98 | 5.38 | 2.91 | 11.83 | 3.28 | - | 9.40 |
| 1977 | 4.26 | 2.56 | 8.71 | 4.33 | 2.59 | 8.89 | 3.03 | - | - |
| 1978 | 4.40 | 2.36 | 9.74 | 4.55 | 2.50 | 9.91 | 3.29 | - | - |
| 1979 | 4.58 | 2.81 | 9.22 | 4.67 | 2.87 | 9.38 | 4.78 | - | 9.55 |
| 1980 | 4.83 | 2.98 | 9.67 | 5.03 | 3.07 | 10.17 | 3.56 | - | - |
| 1981 | 4.96 | 2.64 | 11.05 | 4.95 | 2.54 | 11.26 | 4.08 | - | 8.72 |
| 1982 | 5.30 | 2.73 | 12.03 | 5.34 | 2.71 | 12.24 | 5.55 | 2.73 | 12.94 |
| 1983 | 6.09 | 3.03 | 14.09 | 6.22 | 3.03 | 14.57 | 5.41 | - | 13.20 |
| 1984 | 8.62 | 4.79 | 18.66 | 9.16 | 5.16 | 19.61 | 6.35 | 2.37 | 16.77 |
| 1985 | 11.53 | 5.93 | 26.19 | 12.10 | 6.27 | 27.38 | 8.21 | 3.49 | 20.57 |
| 1986 | 13.63 | 6.78 | 31.55 | 14.29 | 7.23 | 32.77 | 9.32 | 5.12 | 20.32 |
| 1987 | 17.38 | 7.83 | 42.39 | 18.21 | 8.24 | 44.33 | 12.30 | 5.64 | 29.73 |
| 1988 | 17.57 | 8.18 | 42.15 | 18.11 | 8.52 | 43.23 | 16.08 | 7.38 | 38.85 |
| 1989 | 16.52 | 7.02 | 41.41 | 17.08 | 7.27 | 42.76 | 14.53 | 5.52 | 38.14 |
| 1990 | 19.14 | 8.40 | 47.27 | 19.95 | 8.77 | 49.22 | 15.06 | 5.54 | 40.01 |
| 1991 | 19.78 | 8.77 | 48.61 | 20.45 | 8.87 | 50.80 | 15.31 | 7.29 | 36.29 |
| 1992 | 21.41 | 8.99 | 53.92 | 22.18 | 9.25 | 56.04 | 19.09 | 6.83 | 51.21 |
| 1993 | 20.70 | 8.36 | 53.03 | 21.35 | 8.78 | 54.26 | 16.51 | 6.06 | 43.87 |
| 1994 | 22.31 | 8.75 | 57.80 | 22.91 | 9.05 | 59.18 | 20.14 | 7.47 | 53.32 |
| 1995 | 24.69 | 9.63 | 64.14 | 25.06 | 9.98 | 64.55 | 22.64 | 6.39 | 65.19 |
| 1996 | 25.36 | 9.83 | 66.05 | 25.91 | 9.79 | 68.12 | 22.48 | 7.78 | 61.00 |
| 1997 | 28.31 | 11.05 | 73.51 | 28.97 | 11.25 | 75.39 | 28.08 | 11.40 | 71.77 |
| 1998 | 32.76 | 11.97 | 87.19 | 33.74 | 12.31 | 89.84 | 30.42 | 10.77 | 81.85 |
| 1999 | 32.74 | 11.97 | 87.11 | 33.17 | 12.22 | 88.00 | 29.58 | 10.39 | 79.83 |
| 2000 | 32.78 | 11.75 | 87.84 | 33.80 | 12.49 | 89.62 | 29.50 | 8.04 | 85.72 |
| 2001 | 33.58 | 11.52 | 91.34 | 34.10 | 11.95 | 92.11 | 31.57 | 8.94 | 90.82 |
| 2002 | 33.81 | 11.77 | 91.52 | 34.29 | 12.16 | 92.24 | 28.34 | 8.55 | 80.17 |
| 2003 | 32.14 | 11.84 | 85.31 | 33.07 | 12.34 | 87.34 | 29.52 | 9.16 | 82.83 |
| 2004 | 32.71 | 12.54 | 85.53 | 33.46 | 12.77 | 87.62 | 29.98 | 9.69 | 83.10 |
| 2005 | 32.12 | 12.18 | 84.33 | 33.00 | 12.67 | 86.23 | 31.03 | 9.70 | 86.91 |
| 2006 | 32.48 | 13.38 | 82.48 | 32.69 | 13.82 | 82.10 | 34.71 | 11.06 | 96.62 |
| 1975-2006 | 20.62 | 8.66 | 51.93 | 20.74 | 8.82 | 51.96 | 19.39 | 7.10 | 51.55 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 5.4
Cancer of the Cervix Uteri (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

| Year of Diagnosis: | All Races, Females | | | White Females | | | Black Females | | |
|--------------------|--------------------|------|-------|---------------|------|-------|---------------|-------|-------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| 1975 | 14.79 | 9.66 | 28.22 | 13.25 | 8.85 | 24.77 | 33.06 | 17.55 | 73.66 |
| 1976 | 14.26 | 9.49 | 26.74 | 12.90 | 8.76 | 23.74 | 29.25 | 15.84 | 64.38 |
| 1977 | 13.03 | 8.44 | 25.05 | 11.70 | 7.56 | 22.53 | 28.30 | 15.90 | 60.78 |
| 1978 | 12.54 | 8.31 | 23.63 | 11.39 | 7.77 | 20.87 | 24.14 | 12.35 | 55.00 |
| 1979 | 12.76 | 8.68 | 23.44 | 10.98 | 7.55 | 19.94 | 28.83 | 16.01 | 62.38 |
| 1980 | 12.23 | 8.11 | 23.03 | 10.98 | 7.30 | 20.61 | 23.43 | 13.50 | 49.44 |
| 1981 | 10.76 | 7.35 | 19.70 | 9.55 | 6.60 | 17.25 | 24.00 | 13.61 | 51.22 |
| 1982 | 10.62 | 7.12 | 19.77 | 9.40 | 6.68 | 16.51 | 21.82 | 10.15 | 52.38 |
| 1983 | 10.50 | 7.32 | 18.83 | 9.59 | 6.76 | 16.99 | 18.84 | 11.89 | 37.03 |
| 1984 | 11.04 | 8.00 | 19.00 | 10.00 | 7.41 | 16.79 | 21.08 | 12.59 | 43.32 |
| 1985 | 10.21 | 7.06 | 18.47 | 9.14 | 6.74 | 15.45 | 18.99 | 9.52 | 43.80 |
| 1986 | 10.82 | 7.87 | 18.57 | 9.71 | 7.31 | 15.98 | 18.39 | 10.72 | 38.46 |
| 1987 | 10.00 | 7.10 | 17.59 | 8.99 | 6.89 | 14.50 | 18.74 | 8.70 | 45.03 |
| 1988 | 10.58 | 7.86 | 17.70 | 9.51 | 7.43 | 14.97 | 18.98 | 10.45 | 41.33 |
| 1989 | 10.70 | 7.92 | 17.99 | 10.06 | 7.89 | 15.74 | 15.68 | 7.96 | 35.92 |
| 1990 | 10.64 | 8.04 | 17.46 | 9.94 | 7.73 | 15.74 | 16.72 | 9.65 | 35.23 |
| 1991 | 10.03 | 7.05 | 17.86 | 9.23 | 6.79 | 15.64 | 16.33 | 9.22 | 34.96 |
| 1992 | 9.93 | 7.46 | 16.41 | 9.42 | 7.21 | 15.22 | 14.07 | 9.90 | 24.98 |
| 1993 | 9.66 | 7.07 | 16.44 | 9.19 | 7.05 | 14.81 | 13.74 | 8.87 | 26.48 |
| 1994 | 9.45 | 6.88 | 16.18 | 8.71 | 6.79 | 13.73 | 14.03 | 6.94 | 32.58 |
| 1995 | 8.87 | 6.79 | 14.33 | 7.96 | 6.52 | 11.73 | 13.58 | 7.88 | 28.51 |
| 1996 | 9.63 | 7.38 | 15.52 | 8.74 | 7.14 | 12.94 | 14.37 | 8.16 | 30.62 |
| 1997 | 9.20 | 7.10 | 14.71 | 8.35 | 6.89 | 12.19 | 13.56 | 8.02 | 28.07 |
| 1998 | 9.10 | 7.08 | 14.39 | 8.42 | 6.85 | 12.53 | 13.09 | 8.97 | 23.90 |
| 1999 | 8.25 | 6.50 | 12.81 | 7.80 | 6.56 | 11.03 | 13.35 | 8.02 | 27.33 |
| 2000 | 7.70 | 5.84 | 12.57 | 7.42 | 5.90 | 11.40 | 10.22 | 6.42 | 20.18 |
| 2001 | 7.89 | 5.81 | 13.33 | 7.35 | 5.73 | 11.57 | 11.17 | 4.93 | 27.51 |
| 2002 | 7.37 | 5.64 | 11.90 | 6.98 | 5.66 | 10.43 | 10.29 | 6.48 | 20.29 |
| 2003 | 7.24 | 5.68 | 11.33 | 6.77 | 5.58 | 9.89 | 10.59 | 6.03 | 22.53 |
| 2004 | 7.05 | 5.43 | 11.28 | 6.77 | 5.56 | 9.95 | 10.39 | 6.40 | 20.83 |
| 2005 | 6.69 | 5.34 | 10.21 | 6.45 | 5.38 | 9.23 | 8.34 | 5.01 | 17.08 |
| 2006 | 6.72 | 5.40 | 10.18 | 6.58 | 5.50 | 9.39 | 8.30 | 4.95 | 17.07 |
| 1975-2006 | 9.71 | 7.05 | 16.68 | 8.98 | 6.80 | 14.70 | 15.73 | 8.83 | 33.81 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 5.5
Cancer of the Cervix Uteri (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Age

| Year of Death: | All Races, Females | | | White Females | | | Black Females | | |
|----------------|--------------------|------|-------|---------------|------|-------|---------------|------|-------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| 1975 | 5.55 | 2.38 | 13.87 | 4.75 | 1.95 | 12.08 | 13.35 | 5.73 | 33.29 |
| 1976 | 5.43 | 2.19 | 13.92 | 4.68 | 1.86 | 12.06 | 12.86 | 4.89 | 33.75 |
| 1977 | 4.96 | 2.00 | 12.71 | 4.22 | 1.62 | 11.02 | 12.19 | 4.97 | 31.10 |
| 1978 | 4.85 | 2.11 | 12.00 | 4.15 | 1.78 | 10.38 | 11.64 | 4.78 | 29.60 |
| 1979 | 4.58 | 1.91 | 11.58 | 3.95 | 1.58 | 10.15 | 10.61 | 4.48 | 26.67 |
| 1980 | 4.45 | 1.96 | 10.97 | 3.77 | 1.60 | 9.45 | 10.78 | 4.64 | 26.86 |
| 1981 | 4.32 | 1.83 | 10.86 | 3.74 | 1.57 | 9.39 | 9.89 | 3.74 | 26.00 |
| 1982 | 4.10 | 1.77 | 10.20 | 3.51 | 1.54 | 8.66 | 9.77 | 3.63 | 25.85 |
| 1983 | 4.04 | 1.80 | 9.92 | 3.43 | 1.53 | 8.40 | 9.76 | 3.87 | 25.20 |
| 1984 | 3.94 | 1.81 | 9.51 | 3.36 | 1.54 | 8.12 | 9.27 | 3.92 | 23.28 |
| 1985 | 3.82 | 1.71 | 9.36 | 3.27 | 1.49 | 7.92 | 8.98 | 3.36 | 23.70 |
| 1986 | 3.82 | 1.77 | 9.20 | 3.22 | 1.47 | 7.80 | 9.23 | 4.00 | 22.93 |
| 1987 | 3.64 | 1.74 | 8.61 | 3.13 | 1.51 | 7.40 | 8.22 | 3.60 | 20.31 |
| 1988 | 3.61 | 1.69 | 8.63 | 3.10 | 1.52 | 7.24 | 8.15 | 2.98 | 21.72 |
| 1989 | 3.59 | 1.70 | 8.54 | 3.05 | 1.50 | 7.10 | 8.60 | 3.26 | 22.57 |
| 1990 | 3.66 | 1.79 | 8.56 | 3.19 | 1.58 | 7.40 | 7.77 | 3.41 | 19.18 |
| 1991 | 3.49 | 1.70 | 8.20 | 3.01 | 1.51 | 6.97 | 7.65 | 3.15 | 19.41 |
| 1992 | 3.52 | 1.74 | 8.19 | 3.01 | 1.53 | 6.89 | 8.07 | 3.38 | 20.37 |
| 1993 | 3.41 | 1.67 | 7.99 | 2.95 | 1.48 | 6.78 | 7.69 | 3.14 | 19.63 |
| 1994 | 3.38 | 1.66 | 7.86 | 3.00 | 1.50 | 6.93 | 6.71 | 2.85 | 16.82 |
| 1995 | 3.24 | 1.63 | 7.47 | 2.82 | 1.45 | 6.39 | 6.98 | 2.97 | 17.50 |
| 1996 | 3.21 | 1.58 | 7.49 | 2.87 | 1.47 | 6.54 | 6.20 | 2.52 | 15.85 |
| 1997 | 3.14 | 1.61 | 7.16 | 2.76 | 1.45 | 6.20 | 6.47 | 2.84 | 15.98 |
| 1998 | 2.98 | 1.57 | 6.66 | 2.65 | 1.45 | 5.77 | 5.91 | 2.55 | 14.72 |
| 1999 | 2.83 | 1.37 | 6.66 | 2.57 | 1.33 | 5.82 | 5.41 | 1.88 | 14.64 |
| 2000 | 2.78 | 1.36 | 6.52 | 2.47 | 1.25 | 5.68 | 5.43 | 2.17 | 13.97 |
| 2001 | 2.67 | 1.37 | 6.09 | 2.43 | 1.31 | 5.36 | 4.82 | 1.92 | 12.44 |
| 2002 | 2.55 | 1.28 | 5.87 | 2.28 | 1.20 | 5.12 | 4.97 | 2.07 | 12.56 |
| 2003 | 2.48 | 1.24 | 5.74 | 2.23 | 1.15 | 5.05 | 4.66 | 1.93 | 11.80 |
| 2004 | 2.42 | 1.25 | 5.45 | 2.18 | 1.18 | 4.80 | 4.50 | 1.91 | 11.28 |
| 2005 | 2.42 | 1.24 | 5.50 | 2.21 | 1.16 | 4.94 | 4.37 | 1.96 | 10.68 |
| 2006 | 2.42 | 1.24 | 5.53 | 2.21 | 1.15 | 4.97 | 4.31 | 1.90 | 10.64 |
| 1975-2006 | 3.48 | 1.62 | 8.33 | 3.04 | 1.45 | 7.23 | 7.44 | 2.99 | 19.11 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 6.5
Cancer of the Colon and Rectum (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 59.51 | 68.40 | 53.64 | 60.17 | 69.72 | 54.04 | 56.85 | 59.33 | 54.66 |
| 1976 | 61.36 | 71.74 | 54.15 | 62.15 | 72.67 | 55.03 | 55.06 | 58.53 | 51.32 |
| 1977 | 62.40 | 73.50 | 54.95 | 63.19 | 74.32 | 55.76 | 60.72 | 75.75 | 51.47 |
| 1978 | 62.04 | 72.37 | 55.25 | 62.77 | 73.89 | 55.58 | 62.16 | 65.70 | 60.64 |
| 1979 | 62.37 | 74.11 | 54.57 | 62.95 | 75.20 | 54.94 | 58.57 | 61.64 | 56.29 |
| 1980 | 63.74 | 74.53 | 56.53 | 63.94 | 74.87 | 56.82 | 70.76 | 79.68 | 64.12 |
| 1981 | 64.24 | 75.84 | 56.49 | 64.89 | 76.72 | 57.14 | 62.78 | 72.81 | 55.94 |
| 1982 | 62.78 | 75.19 | 54.50 | 63.44 | 76.27 | 55.04 | 61.25 | 68.97 | 55.92 |
| 1983 | 63.67 | 75.58 | 55.62 | 64.28 | 76.56 | 56.07 | 66.61 | 75.39 | 61.14 |
| 1984 | 64.82 | 77.68 | 56.07 | 65.83 | 79.27 | 56.73 | 62.26 | 70.26 | 57.98 |
| 1985 | 66.31 | 79.14 | 57.37 | 67.22 | 80.18 | 58.43 | 63.87 | 74.12 | 56.27 |
| 1986 | 64.18 | 77.77 | 54.72 | 64.60 | 78.92 | 54.74 | 64.52 | 72.90 | 58.61 |
| 1987 | 62.71 | 77.47 | 52.67 | 62.88 | 78.20 | 52.62 | 66.21 | 77.48 | 59.01 |
| 1988 | 61.36 | 75.17 | 51.80 | 61.52 | 75.75 | 51.70 | 63.14 | 72.41 | 56.91 |
| 1989 | 61.67 | 74.94 | 52.41 | 61.95 | 75.42 | 52.67 | 64.38 | 79.28 | 54.60 |
| 1990 | 60.69 | 74.33 | 51.29 | 60.66 | 74.73 | 51.05 | 66.63 | 74.71 | 61.58 |
| 1991 | 59.47 | 72.89 | 50.05 | 59.28 | 73.08 | 49.67 | 65.59 | 76.29 | 58.16 |
| 1992 | 57.98 | 70.41 | 49.18 | 57.83 | 70.68 | 48.71 | 65.41 | 78.35 | 57.31 |
| 1993 | 56.80 | 68.42 | 48.32 | 56.34 | 68.08 | 47.82 | 64.33 | 77.85 | 55.14 |
| 1994 | 55.60 | 66.25 | 47.69 | 55.34 | 66.30 | 47.13 | 64.47 | 75.33 | 57.98 |
| 1995 | 54.02 | 63.77 | 46.93 | 53.67 | 63.37 | 46.58 | 61.97 | 71.10 | 56.15 |
| 1996 | 54.72 | 65.47 | 46.68 | 54.73 | 65.84 | 46.41 | 58.26 | 66.62 | 52.53 |
| 1997 | 56.40 | 68.09 | 47.94 | 56.41 | 67.93 | 48.06 | 64.23 | 77.07 | 55.88 |
| 1998 | 56.71 | 66.94 | 49.27 | 56.66 | 67.15 | 49.02 | 62.68 | 73.79 | 55.77 |
| 1999 | 55.44 | 65.62 | 47.85 | 55.24 | 65.73 | 47.20 | 63.07 | 71.71 | 58.04 |
| 2000 | 54.05 | 63.56 | 46.69 | 53.70 | 62.95 | 46.45 | 64.05 | 73.68 | 57.71 |
| 2001 | 53.32 | 62.59 | 46.04 | 52.69 | 62.16 | 45.17 | 63.73 | 74.08 | 57.04 |
| 2002 | 52.71 | 60.87 | 46.31 | 51.95 | 59.73 | 45.71 | 62.10 | 72.75 | 55.38 |
| 2003 | 50.26 | 58.78 | 43.47 | 49.78 | 58.01 | 43.07 | 61.05 | 73.35 | 53.30 |
| 2004 | 49.16 | 57.75 | 42.51 | 48.35 | 57.25 | 41.34 | 60.86 | 71.89 | 53.50 |
| 2005 | 46.70 | 53.78 | 41.06 | 46.35 | 53.75 | 40.34 | 55.52 | 61.16 | 52.24 |
| 2006 | 45.45 | 51.29 | 40.69 | 44.64 | 50.58 | 39.68 | 55.41 | 61.49 | 51.37 |
| 1975-2006 | 57.38 | 67.95 | 49.71 | 57.55 | 68.33 | 49.75 | 62.26 | 71.77 | 56.05 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 6.6
Cancer of the Colon (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 41.33 | 45.36 | 38.89 | 41.71 | 46.10 | 39.15 | 42.83 | 44.14 | 41.64 |
| 1976 | 43.06 | 47.76 | 39.96 | 43.45 | 48.19 | 40.41 | 43.11 | 43.60 | 41.95 |
| 1977 | 43.97 | 49.55 | 40.36 | 44.66 | 50.06 | 41.19 | 45.42 | 58.42 | 37.66 |
| 1978 | 43.85 | 49.13 | 40.59 | 44.11 | 49.87 | 40.61 | 47.77 | 48.81 | 47.88 |
| 1979 | 43.77 | 50.20 | 39.61 | 44.10 | 51.03 | 39.70 | 44.68 | 45.58 | 43.97 |
| 1980 | 45.56 | 50.67 | 42.33 | 45.58 | 50.88 | 42.32 | 55.12 | 58.32 | 52.74 |
| 1981 | 45.47 | 51.34 | 41.72 | 45.89 | 51.96 | 42.11 | 46.86 | 50.64 | 44.36 |
| 1982 | 45.75 | 53.45 | 40.66 | 46.07 | 54.09 | 40.84 | 48.27 | 52.48 | 45.19 |
| 1983 | 46.11 | 53.02 | 41.64 | 46.41 | 53.46 | 41.88 | 50.15 | 58.40 | 45.25 |
| 1984 | 46.51 | 54.61 | 41.31 | 47.04 | 55.57 | 41.58 | 48.56 | 54.02 | 45.83 |
| 1985 | 47.89 | 55.02 | 43.01 | 48.43 | 55.59 | 43.70 | 49.87 | 57.34 | 44.21 |
| 1986 | 46.79 | 54.36 | 41.58 | 47.07 | 55.21 | 41.51 | 49.20 | 55.04 | 45.46 |
| 1987 | 45.24 | 54.42 | 39.17 | 45.15 | 54.52 | 39.09 | 51.35 | 59.84 | 45.73 |
| 1988 | 44.30 | 52.59 | 38.66 | 44.25 | 52.85 | 38.39 | 48.93 | 54.02 | 45.61 |
| 1989 | 44.35 | 52.41 | 38.95 | 44.48 | 52.49 | 39.20 | 48.85 | 60.28 | 41.54 |
| 1990 | 44.19 | 51.96 | 38.93 | 44.06 | 51.98 | 38.76 | 52.02 | 57.69 | 48.57 |
| 1991 | 43.60 | 51.84 | 37.91 | 43.29 | 51.85 | 37.41 | 51.74 | 57.75 | 47.67 |
| 1992 | 42.28 | 49.65 | 37.20 | 42.05 | 49.77 | 36.71 | 50.79 | 60.21 | 45.17 |
| 1993 | 41.60 | 48.93 | 36.32 | 41.07 | 48.62 | 35.67 | 51.10 | 60.40 | 45.02 |
| 1994 | 40.82 | 46.93 | 36.31 | 40.61 | 47.03 | 35.79 | 50.11 | 57.03 | 46.30 |
| 1995 | 39.64 | 45.02 | 35.86 | 39.29 | 44.64 | 35.47 | 48.74 | 54.79 | 45.42 |
| 1996 | 39.34 | 45.31 | 34.88 | 39.30 | 45.75 | 34.46 | 44.45 | 48.40 | 41.84 |
| 1997 | 41.09 | 48.41 | 35.95 | 41.11 | 48.30 | 36.13 | 49.03 | 58.76 | 42.46 |
| 1998 | 41.00 | 46.76 | 37.01 | 40.92 | 47.04 | 36.60 | 47.41 | 54.38 | 43.71 |
| 1999 | 39.79 | 45.36 | 35.70 | 39.56 | 45.35 | 35.16 | 48.43 | 54.70 | 44.91 |
| 2000 | 39.30 | 44.34 | 35.44 | 38.92 | 43.82 | 35.13 | 50.85 | 57.16 | 46.89 |
| 2001 | 39.04 | 43.79 | 35.39 | 38.63 | 43.56 | 34.76 | 48.62 | 55.13 | 44.49 |
| 2002 | 38.37 | 42.86 | 34.87 | 37.68 | 41.83 | 34.37 | 48.72 | 56.09 | 44.03 |
| 2003 | 36.59 | 40.95 | 33.08 | 36.26 | 40.30 | 32.90 | 46.62 | 55.27 | 41.25 |
| 2004 | 35.80 | 40.80 | 32.05 | 35.24 | 40.54 | 31.21 | 46.39 | 53.21 | 41.91 |
| 2005 | 33.70 | 37.02 | 31.08 | 33.48 | 37.16 | 30.50 | 42.54 | 45.91 | 40.87 |
| 2006 | 32.95 | 35.93 | 30.51 | 32.42 | 35.46 | 29.87 | 42.76 | 46.52 | 40.16 |
| 1975-2006 | 41.43 | 47.32 | 37.25 | 41.46 | 47.51 | 37.18 | 48.04 | 54.25 | 44.13 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 6.7
Cancer of the Rectum (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 18.18 | 23.04 | 14.75 | 18.45 | 23.62 | 14.89 | 14.02 | 15.19 | 13.01 |
| 1976 | 18.30 | 23.98 | 14.19 | 18.70 | 24.48 | 14.63 | 11.96 | 14.93 | 9.37 |
| 1977 | 18.43 | 23.95 | 14.59 | 18.53 | 24.26 | 14.58 | 15.30 | 17.33 | 13.81 |
| 1978 | 18.19 | 23.25 | 14.66 | 18.67 | 24.02 | 14.97 | 14.39 | 16.89 | 12.76 |
| 1979 | 18.60 | 23.91 | 14.96 | 18.85 | 24.17 | 15.24 | 13.90 | 16.06 | 12.31 |
| 1980 | 18.18 | 23.86 | 14.20 | 18.36 | 23.99 | 14.51 | 15.65 | 21.36 | 11.38 |
| 1981 | 18.77 | 24.50 | 14.77 | 19.00 | 24.76 | 15.02 | 15.92 | 22.17 | 11.59 |
| 1982 | 17.03 | 21.74 | 13.84 | 17.38 | 22.18 | 14.20 | 12.98 | 16.49 | 10.73 |
| 1983 | 17.56 | 22.55 | 13.98 | 17.87 | 23.11 | 14.19 | 16.47 | 16.99 | 15.89 |
| 1984 | 18.30 | 23.07 | 14.76 | 18.79 | 23.70 | 15.15 | 13.71 | 16.24 | 12.15 |
| 1985 | 18.42 | 24.12 | 14.36 | 18.79 | 24.59 | 14.72 | 14.00 | 16.79 | 12.06 |
| 1986 | 17.39 | 23.41 | 13.13 | 17.53 | 23.71 | 13.23 | 15.32 | 17.86 | 13.15 |
| 1987 | 17.47 | 23.05 | 13.50 | 17.73 | 23.68 | 13.53 | 14.87 | 17.64 | 13.29 |
| 1988 | 17.06 | 22.58 | 13.14 | 17.27 | 22.90 | 13.31 | 14.21 | 18.38 | 11.30 |
| 1989 | 17.32 | 22.52 | 13.45 | 17.47 | 22.93 | 13.47 | 15.52 | 19.00 | 13.05 |
| 1990 | 16.51 | 22.37 | 12.36 | 16.61 | 22.75 | 12.29 | 14.61 | 17.02 | 13.01 |
| 1991 | 15.88 | 21.05 | 12.14 | 15.99 | 21.23 | 12.26 | 13.85 | 18.55 | 10.50 |
| 1992 | 15.71 | 20.76 | 11.98 | 15.77 | 20.90 | 11.99 | 14.62 | 18.14 | 12.14 |
| 1993 | 15.20 | 19.49 | 12.00 | 15.27 | 19.46 | 12.15 | 13.23 | 17.46 | 10.12 |
| 1994 | 14.78 | 19.32 | 11.38 | 14.73 | 19.27 | 11.34 | 14.36 | 18.30 | 11.68 |
| 1995 | 14.38 | 18.76 | 11.07 | 14.38 | 18.73 | 11.11 | 13.23 | 16.31 | 10.73 |
| 1996 | 15.39 | 20.16 | 11.80 | 15.43 | 20.09 | 11.95 | 13.81 | 18.22 | 10.69 |
| 1997 | 15.31 | 19.69 | 11.98 | 15.30 | 19.64 | 11.92 | 15.19 | 18.31 | 13.42 |
| 1998 | 15.70 | 20.19 | 12.26 | 15.74 | 20.11 | 12.41 | 15.27 | 19.41 | 12.07 |
| 1999 | 15.65 | 20.26 | 12.15 | 15.68 | 20.38 | 12.04 | 14.64 | 17.01 | 13.13 |
| 2000 | 14.75 | 19.22 | 11.25 | 14.78 | 19.14 | 11.32 | 13.19 | 16.53 | 10.82 |
| 2001 | 14.27 | 18.80 | 10.66 | 14.05 | 18.60 | 10.41 | 15.11 | 18.94 | 12.55 |
| 2002 | 14.34 | 18.01 | 11.43 | 14.27 | 17.90 | 11.35 | 13.38 | 16.65 | 11.35 |
| 2003 | 13.67 | 17.83 | 10.39 | 13.51 | 17.71 | 10.17 | 14.44 | 18.09 | 12.06 |
| 2004 | 13.36 | 16.95 | 10.46 | 13.10 | 16.72 | 10.13 | 14.47 | 18.68 | 11.59 |
| 2005 | 13.00 | 16.76 | 9.98 | 12.88 | 16.60 | 9.84 | 12.98 | 15.25 | 11.37 |
| 2006 | 12.50 | 15.37 | 10.18 | 12.21 | 15.12 | 9.81 | 12.65 | 14.97 | 11.21 |
| 1975-2006 | 15.95 | 20.63 | 12.46 | 16.09 | 20.82 | 12.57 | 14.22 | 17.52 | 11.92 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 6.8
Cancer of the Colon and Rectum (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 28.09 | 32.84 | 24.96 | 28.30 | 33.24 | 25.06 | 26.79 | 30.09 | 24.55 |
| 1976 | 28.58 | 33.34 | 25.44 | 28.74 | 33.76 | 25.45 | 28.23 | 31.16 | 26.26 |
| 1977 | 28.19 | 33.21 | 24.87 | 28.34 | 33.49 | 24.97 | 27.61 | 31.44 | 24.92 |
| 1978 | 28.54 | 33.75 | 25.13 | 28.70 | 34.12 | 25.17 | 27.91 | 31.20 | 25.66 |
| 1979 | 28.15 | 33.56 | 24.63 | 28.25 | 33.92 | 24.57 | 28.16 | 31.57 | 25.96 |
| 1980 | 28.05 | 33.74 | 24.35 | 28.12 | 33.99 | 24.32 | 28.81 | 33.18 | 25.89 |
| 1981 | 27.52 | 32.95 | 23.97 | 27.53 | 33.16 | 23.85 | 28.93 | 32.83 | 26.32 |
| 1982 | 27.24 | 32.96 | 23.55 | 27.21 | 33.11 | 23.42 | 29.11 | 33.71 | 26.12 |
| 1983 | 27.12 | 33.08 | 23.23 | 27.13 | 33.32 | 23.11 | 28.67 | 32.90 | 25.82 |
| 1984 | 27.35 | 33.02 | 23.61 | 27.30 | 33.12 | 23.49 | 29.58 | 34.65 | 26.22 |
| 1985 | 26.93 | 32.66 | 23.11 | 26.75 | 32.74 | 22.77 | 30.49 | 34.10 | 28.04 |
| 1986 | 26.16 | 32.05 | 22.31 | 26.00 | 32.01 | 22.07 | 29.82 | 35.36 | 26.35 |
| 1987 | 25.89 | 31.91 | 21.95 | 25.68 | 31.85 | 21.63 | 29.85 | 35.10 | 26.48 |
| 1988 | 25.28 | 31.08 | 21.46 | 25.08 | 31.04 | 21.16 | 29.18 | 34.29 | 25.88 |
| 1989 | 25.01 | 30.80 | 21.16 | 24.70 | 30.52 | 20.82 | 30.05 | 36.37 | 26.07 |
| 1990 | 24.65 | 30.77 | 20.64 | 24.25 | 30.43 | 20.18 | 30.77 | 37.64 | 26.59 |
| 1991 | 24.01 | 29.64 | 20.30 | 23.63 | 29.35 | 19.85 | 30.02 | 35.80 | 26.50 |
| 1992 | 23.62 | 29.32 | 19.85 | 23.25 | 28.99 | 19.42 | 29.47 | 36.23 | 25.43 |
| 1993 | 23.31 | 28.56 | 19.77 | 22.88 | 28.18 | 19.28 | 29.77 | 35.84 | 26.04 |
| 1994 | 22.92 | 28.28 | 19.28 | 22.50 | 27.89 | 18.84 | 29.04 | 35.66 | 24.88 |
| 1995 | 22.59 | 27.65 | 19.14 | 22.11 | 27.16 | 18.64 | 29.54 | 36.30 | 25.47 |
| 1996 | 21.86 | 26.71 | 18.51 | 21.39 | 26.29 | 17.98 | 28.59 | 34.52 | 24.86 |
| 1997 | 21.47 | 26.23 | 18.11 | 20.95 | 25.68 | 17.59 | 28.84 | 35.07 | 24.80 |
| 1998 | 21.19 | 25.62 | 18.03 | 20.68 | 25.13 | 17.48 | 28.31 | 33.90 | 24.78 |
| 1999 | 20.93 | 25.48 | 17.75 | 20.43 | 24.99 | 17.20 | 28.51 | 34.50 | 24.70 |
| 2000 | 20.65 | 25.05 | 17.48 | 20.14 | 24.43 | 17.01 | 28.14 | 34.94 | 23.87 |
| 2001 | 20.09 | 24.25 | 17.06 | 19.54 | 23.72 | 16.46 | 27.66 | 33.18 | 24.18 |
| 2002 | 19.65 | 23.83 | 16.52 | 19.13 | 23.23 | 16.03 | 26.92 | 33.37 | 22.82 |
| 2003 | 19.00 | 22.97 | 16.05 | 18.47 | 22.43 | 15.49 | 26.43 | 32.10 | 22.72 |
| 2004 | 17.93 | 21.60 | 15.17 | 17.46 | 21.09 | 14.69 | 24.80 | 30.29 | 21.19 |
| 2005 | 17.38 | 21.01 | 14.64 | 16.83 | 20.41 | 14.09 | 24.90 | 30.48 | 21.20 |
| 2006 | 17.10 | 20.51 | 14.54 | 16.60 | 19.87 | 14.08 | 24.26 | 30.84 | 20.12 |
| 1975-2006 | 23.37 | 28.23 | 20.03 | 23.12 | 28.02 | 19.73 | 28.31 | 33.70 | 24.85 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 7.4
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

| Year of Diagnosis: | All Races, Females | | | White Females | | | Black Females | | |
|--------------------|--------------------|------|--------|---------------|------|--------|---------------|------|-------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| 1975 | 35.47 | 6.48 | 111.37 | 37.34 | 6.83 | 117.22 | 19.91 | 3.64 | 62.51 |
| 1976 | 34.37 | 6.22 | 108.08 | 36.25 | 6.28 | 114.71 | 17.61 | 3.29 | 55.11 |
| 1977 | 31.66 | 5.73 | 99.58 | 33.13 | 5.91 | 104.41 | 20.06 | 3.52 | 63.37 |
| 1978 | 30.05 | 6.10 | 92.76 | 31.67 | 6.32 | 98.05 | 18.59 | 3.74 | 57.49 |
| 1979 | 27.89 | 5.30 | 87.07 | 29.23 | 5.33 | 91.83 | 16.96 | 3.94 | 51.06 |
| 1980 | 27.31 | 4.97 | 85.82 | 28.54 | 4.97 | 90.26 | 16.22 | - | 52.13 |
| 1981 | 27.26 | 5.58 | 84.02 | 28.35 | 5.45 | 88.33 | 17.42 | 2.86 | 55.52 |
| 1982 | 26.79 | 5.04 | 83.75 | 28.21 | 5.20 | 88.47 | 17.06 | 2.49 | 55.22 |
| 1983 | 26.53 | 4.86 | 83.27 | 27.77 | 4.89 | 87.69 | 18.81 | 3.37 | 59.25 |
| 1984 | 25.66 | 4.58 | 80.88 | 27.11 | 4.84 | 85.42 | 17.38 | - | 58.35 |
| 1985 | 25.22 | 4.60 | 79.21 | 26.53 | 4.80 | 83.42 | 17.51 | 2.23 | 57.52 |
| 1986 | 24.40 | 4.84 | 75.63 | 25.64 | 5.02 | 79.64 | 16.78 | 2.54 | 54.09 |
| 1987 | 24.72 | 4.99 | 76.38 | 26.15 | 5.28 | 80.80 | 15.77 | - | 51.79 |
| 1988 | 23.55 | 4.52 | 73.38 | 24.59 | 4.46 | 77.30 | 16.88 | 2.97 | 53.32 |
| 1989 | 24.42 | 4.49 | 76.61 | 25.49 | 4.78 | 79.72 | 19.36 | 1.78 | 65.40 |
| 1990 | 25.05 | 4.60 | 78.59 | 26.52 | 4.80 | 83.40 | 16.92 | 2.28 | 55.27 |
| 1991 | 24.63 | 4.39 | 77.64 | 26.00 | 4.49 | 82.31 | 17.48 | 3.12 | 55.09 |
| 1992 | 24.82 | 4.87 | 77.08 | 26.19 | 4.88 | 82.00 | 16.94 | 2.88 | 53.74 |
| 1993 | 24.28 | 5.18 | 74.27 | 25.67 | 5.14 | 79.42 | 17.06 | 3.48 | 52.63 |
| 1994 | 25.13 | 5.02 | 77.78 | 26.32 | 5.03 | 82.07 | 17.62 | 2.51 | 57.20 |
| 1995 | 25.31 | 4.69 | 79.31 | 26.46 | 4.62 | 83.65 | 18.86 | 2.65 | 61.29 |
| 1996 | 24.92 | 4.52 | 78.35 | 25.96 | 4.36 | 82.52 | 19.30 | 3.17 | 61.55 |
| 1997 | 25.87 | 5.04 | 80.40 | 27.35 | 5.04 | 85.75 | 16.47 | 2.46 | 53.15 |
| 1998 | 25.70 | 4.80 | 80.40 | 27.04 | 4.77 | 85.35 | 19.37 | 2.98 | 62.29 |
| 1999 | 25.31 | 5.18 | 78.02 | 26.76 | 5.06 | 83.56 | 17.47 | 2.79 | 55.91 |
| 2000 | 24.79 | 5.21 | 76.08 | 26.17 | 5.35 | 80.69 | 17.90 | 3.00 | 56.92 |
| 2001 | 25.20 | 5.10 | 77.86 | 26.65 | 4.96 | 83.42 | 18.94 | 3.96 | 58.16 |
| 2002 | 24.75 | 4.99 | 76.49 | 25.17 | 4.53 | 79.23 | 22.93 | 4.66 | 70.79 |
| 2003 | 24.13 | 5.21 | 73.65 | 25.29 | 5.22 | 77.87 | 19.96 | 3.68 | 62.57 |
| 2004 | 24.71 | 5.40 | 75.29 | 25.64 | 5.10 | 79.42 | 20.10 | 3.68 | 63.09 |
| 2005 | 24.50 | 5.41 | 74.47 | 25.28 | 5.25 | 77.75 | 22.70 | 4.00 | 71.67 |
| 2006 | 24.08 | 5.01 | 74.03 | 25.30 | 4.91 | 78.70 | 18.90 | 3.57 | 59.05 |
| 1975-2006 | 26.12 | 5.04 | 81.30 | 27.48 | 5.06 | 86.21 | 18.41 | 3.10 | 58.49 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 7.5
Cancer of the Corpus and Uterus, NOS (Invasive)
Age-adjusted U.S. Death^a Rates by Year, Race and Age

| Year of Death: | All Races, Females | | | White Females | | | Black Females | | |
|----------------|--------------------|------|-------|---------------|------|-------|---------------|------|-------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| 1975 | 5.28 | 0.52 | 17.74 | 5.05 | 0.47 | 17.06 | 7.80 | 0.93 | 25.81 |
| 1976 | 5.33 | 0.53 | 17.90 | 5.11 | 0.49 | 17.19 | 8.07 | 0.86 | 26.96 |
| 1977 | 5.22 | 0.52 | 17.54 | 4.96 | 0.48 | 16.67 | 8.35 | 0.83 | 28.04 |
| 1978 | 5.10 | 0.52 | 17.08 | 4.84 | 0.48 | 16.23 | 8.16 | 0.85 | 27.30 |
| 1979 | 4.99 | 0.40 | 17.01 | 4.74 | 0.36 | 16.19 | 7.90 | 0.73 | 26.66 |
| 1980 | 5.12 | 0.44 | 17.38 | 4.84 | 0.39 | 16.50 | 8.39 | 0.85 | 28.12 |
| 1981 | 4.88 | 0.41 | 16.60 | 4.68 | 0.38 | 15.91 | 7.34 | 0.56 | 25.11 |
| 1982 | 4.85 | 0.41 | 16.49 | 4.61 | 0.40 | 15.65 | 7.61 | 0.43 | 26.40 |
| 1983 | 4.80 | 0.36 | 16.43 | 4.59 | 0.32 | 15.78 | 7.29 | 0.60 | 24.82 |
| 1984 | 4.68 | 0.36 | 15.99 | 4.42 | 0.31 | 15.17 | 7.70 | 0.69 | 26.07 |
| 1985 | 4.61 | 0.36 | 15.74 | 4.39 | 0.34 | 14.99 | 7.37 | 0.51 | 25.33 |
| 1986 | 4.44 | 0.35 | 15.17 | 4.20 | 0.33 | 14.36 | 7.27 | 0.56 | 24.85 |
| 1987 | 4.42 | 0.34 | 15.10 | 4.18 | 0.32 | 14.28 | 7.32 | 0.52 | 25.12 |
| 1988 | 4.35 | 0.31 | 14.90 | 4.09 | 0.30 | 14.01 | 7.40 | 0.46 | 25.57 |
| 1989 | 4.26 | 0.35 | 14.52 | 4.01 | 0.33 | 13.63 | 7.25 | 0.48 | 24.99 |
| 1990 | 4.34 | 0.33 | 14.85 | 4.08 | 0.32 | 13.94 | 7.36 | 0.42 | 25.56 |
| 1991 | 4.18 | 0.33 | 14.24 | 3.90 | 0.31 | 13.30 | 7.29 | 0.49 | 25.08 |
| 1992 | 4.21 | 0.32 | 14.40 | 3.97 | 0.29 | 13.59 | 6.98 | 0.44 | 24.11 |
| 1993 | 4.17 | 0.32 | 14.26 | 3.91 | 0.30 | 13.35 | 7.28 | 0.44 | 25.17 |
| 1994 | 4.15 | 0.28 | 14.29 | 3.91 | 0.28 | 13.41 | 7.06 | 0.34 | 24.67 |
| 1995 | 4.15 | 0.34 | 14.12 | 3.97 | 0.34 | 13.47 | 6.46 | 0.35 | 22.47 |
| 1996 | 4.13 | 0.31 | 14.14 | 3.86 | 0.30 | 13.18 | 7.18 | 0.37 | 25.03 |
| 1997 | 4.00 | 0.31 | 13.67 | 3.78 | 0.31 | 12.85 | 6.74 | 0.30 | 23.62 |
| 1998 | 4.10 | 0.32 | 14.00 | 3.86 | 0.29 | 13.19 | 6.85 | 0.45 | 23.60 |
| 1999 | 4.08 | 0.35 | 13.87 | 3.86 | 0.34 | 13.10 | 6.84 | 0.39 | 23.71 |
| 2000 | 4.11 | 0.35 | 13.94 | 3.84 | 0.34 | 12.99 | 7.23 | 0.40 | 25.11 |
| 2001 | 4.17 | 0.36 | 14.16 | 3.93 | 0.34 | 13.35 | 6.97 | 0.48 | 23.96 |
| 2002 | 4.16 | 0.34 | 14.15 | 3.87 | 0.33 | 13.13 | 7.32 | 0.40 | 25.43 |
| 2003 | 4.13 | 0.34 | 14.05 | 3.87 | 0.30 | 13.21 | 7.05 | 0.49 | 24.22 |
| 2004 | 4.12 | 0.38 | 13.93 | 3.87 | 0.37 | 13.03 | 6.97 | 0.45 | 24.06 |
| 2005 | 4.10 | 0.36 | 13.91 | 3.83 | 0.34 | 12.97 | 7.15 | 0.49 | 24.60 |
| 2006 | 4.21 | 0.37 | 14.27 | 3.92 | 0.36 | 13.27 | 7.37 | 0.46 | 25.48 |
| 1975-2006 | 4.42 | 0.36 | 15.06 | 4.18 | 0.34 | 14.24 | 7.29 | 0.50 | 25.09 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 8.4
Cancer of the Esophagus (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 4.14 | 6.77 | 2.12 | 3.55 | 5.64 | 1.99 | 10.88 | 19.90 | 3.70 |
| 1976 | 4.30 | 6.88 | 2.34 | 3.72 | 5.94 | 2.09 | 10.74 | 16.89 | 5.61 |
| 1977 | 4.06 | 6.57 | 2.19 | 3.33 | 5.26 | 1.92 | 12.02 | 20.89 | 5.44 |
| 1978 | 4.12 | 6.95 | 1.91 | 3.41 | 5.66 | 1.69 | 13.05 | 24.41 | 4.33 |
| 1979 | 4.42 | 7.43 | 2.21 | 3.73 | 6.23 | 1.98 | 12.94 | 23.04 | 5.24 |
| 1980 | 4.27 | 6.87 | 2.28 | 3.54 | 5.75 | 1.86 | 12.32 | 18.79 | 7.41 |
| 1981 | 4.14 | 6.47 | 2.36 | 3.31 | 5.18 | 1.94 | 13.53 | 21.28 | 7.46 |
| 1982 | 4.26 | 6.81 | 2.28 | 3.46 | 5.46 | 1.96 | 13.61 | 22.72 | 6.41 |
| 1983 | 4.29 | 7.42 | 1.95 | 3.57 | 6.11 | 1.77 | 12.23 | 22.22 | 4.26 |
| 1984 | 4.19 | 6.66 | 2.25 | 3.53 | 5.60 | 1.93 | 12.00 | 19.85 | 5.98 |
| 1985 | 4.45 | 7.42 | 2.22 | 3.74 | 6.26 | 1.89 | 12.68 | 22.07 | 5.81 |
| 1986 | 4.64 | 7.50 | 2.46 | 3.90 | 6.25 | 2.19 | 14.06 | 24.02 | 6.30 |
| 1987 | 4.55 | 7.54 | 2.25 | 3.91 | 6.41 | 2.02 | 11.44 | 20.13 | 4.81 |
| 1988 | 4.44 | 7.35 | 2.21 | 3.88 | 6.32 | 2.02 | 11.18 | 19.49 | 4.99 |
| 1989 | 4.40 | 7.30 | 2.21 | 3.78 | 6.20 | 1.97 | 10.49 | 17.29 | 5.39 |
| 1990 | 4.82 | 8.25 | 2.19 | 4.19 | 7.24 | 1.90 | 12.53 | 21.45 | 5.83 |
| 1991 | 4.55 | 7.44 | 2.35 | 4.15 | 6.74 | 2.19 | 10.21 | 17.78 | 4.88 |
| 1992 | 4.59 | 7.99 | 2.00 | 4.21 | 7.39 | 1.80 | 10.62 | 18.05 | 5.32 |
| 1993 | 4.61 | 7.66 | 2.25 | 4.20 | 6.91 | 2.10 | 10.03 | 17.50 | 4.64 |
| 1994 | 4.50 | 7.70 | 2.06 | 4.16 | 7.16 | 1.87 | 9.20 | 14.68 | 5.13 |
| 1995 | 4.35 | 7.06 | 2.20 | 4.08 | 6.62 | 2.07 | 8.40 | 13.69 | 4.37 |
| 1996 | 4.80 | 8.12 | 2.20 | 4.41 | 7.49 | 1.99 | 9.69 | 16.36 | 4.81 |
| 1997 | 4.68 | 7.77 | 2.22 | 4.58 | 7.72 | 2.02 | 7.10 | 10.85 | 4.46 |
| 1998 | 4.72 | 8.16 | 2.04 | 4.51 | 7.91 | 1.87 | 7.80 | 12.52 | 4.32 |
| 1999 | 4.91 | 8.38 | 2.14 | 4.88 | 8.38 | 2.04 | 7.17 | 11.52 | 4.08 |
| 2000 | 4.76 | 7.98 | 2.16 | 4.72 | 8.01 | 2.02 | 7.04 | 10.58 | 4.43 |
| 2001 | 4.81 | 8.44 | 1.92 | 4.83 | 8.42 | 1.93 | 6.33 | 11.43 | 2.60 |
| 2002 | 4.60 | 7.88 | 1.99 | 4.59 | 8.02 | 1.82 | 6.36 | 10.03 | 3.64 |
| 2003 | 4.62 | 7.99 | 1.95 | 4.73 | 8.19 | 1.93 | 6.27 | 10.02 | 3.67 |
| 2004 | 5.23 | 8.84 | 2.35 | 5.36 | 9.12 | 2.32 | 6.06 | 8.85 | 4.01 |
| 2005 | 4.53 | 7.94 | 1.80 | 4.70 | 8.34 | 1.74 | 4.94 | 8.01 | 2.83 |
| 2006 | 4.56 | 7.60 | 2.06 | 4.67 | 7.80 | 2.05 | 5.28 | 8.69 | 2.81 |
| 1975-2006 | 4.54 | 7.62 | 2.16 | 4.17 | 7.03 | 1.97 | 9.30 | 15.56 | 4.64 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 8.5
Cancer of the Esophagus (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 3.69 | 6.36 | 1.71 | 3.11 | 5.37 | 1.47 | 9.70 | 17.01 | 4.06 |
| 1976 | 3.74 | 6.39 | 1.76 | 3.11 | 5.32 | 1.50 | 10.18 | 17.73 | 4.35 |
| 1977 | 3.70 | 6.26 | 1.78 | 3.07 | 5.20 | 1.51 | 10.08 | 17.57 | 4.35 |
| 1978 | 3.79 | 6.41 | 1.82 | 3.17 | 5.35 | 1.55 | 10.03 | 17.59 | 4.30 |
| 1979 | 3.76 | 6.40 | 1.79 | 3.12 | 5.28 | 1.53 | 10.32 | 18.55 | 4.17 |
| 1980 | 3.85 | 6.60 | 1.82 | 3.17 | 5.49 | 1.50 | 10.60 | 18.30 | 4.88 |
| 1981 | 3.82 | 6.58 | 1.78 | 3.16 | 5.44 | 1.51 | 10.45 | 18.72 | 4.35 |
| 1982 | 3.84 | 6.52 | 1.84 | 3.17 | 5.37 | 1.56 | 10.49 | 18.60 | 4.51 |
| 1983 | 3.85 | 6.64 | 1.79 | 3.22 | 5.59 | 1.50 | 10.21 | 17.95 | 4.52 |
| 1984 | 3.93 | 6.74 | 1.86 | 3.28 | 5.63 | 1.56 | 10.60 | 18.80 | 4.65 |
| 1985 | 3.86 | 6.72 | 1.72 | 3.26 | 5.71 | 1.46 | 9.93 | 17.72 | 4.26 |
| 1986 | 3.94 | 6.76 | 1.84 | 3.34 | 5.76 | 1.57 | 10.19 | 17.96 | 4.55 |
| 1987 | 3.97 | 6.88 | 1.80 | 3.41 | 5.96 | 1.53 | 9.67 | 17.13 | 4.37 |
| 1988 | 4.05 | 7.04 | 1.80 | 3.47 | 6.01 | 1.58 | 9.95 | 18.40 | 3.89 |
| 1989 | 4.11 | 7.15 | 1.82 | 3.56 | 6.23 | 1.57 | 9.75 | 17.32 | 4.29 |
| 1990 | 4.10 | 7.16 | 1.81 | 3.58 | 6.32 | 1.55 | 9.35 | 16.33 | 4.37 |
| 1991 | 4.13 | 7.24 | 1.81 | 3.64 | 6.43 | 1.57 | 9.23 | 16.54 | 4.14 |
| 1992 | 4.17 | 7.30 | 1.81 | 3.74 | 6.61 | 1.57 | 8.65 | 15.10 | 4.11 |
| 1993 | 4.19 | 7.33 | 1.82 | 3.75 | 6.60 | 1.59 | 8.99 | 16.00 | 4.07 |
| 1994 | 4.27 | 7.59 | 1.77 | 3.87 | 6.94 | 1.56 | 8.57 | 15.28 | 3.88 |
| 1995 | 4.26 | 7.56 | 1.76 | 3.92 | 7.00 | 1.58 | 8.04 | 14.45 | 3.51 |
| 1996 | 4.31 | 7.52 | 1.86 | 4.01 | 7.06 | 1.67 | 7.77 | 13.57 | 3.72 |
| 1997 | 4.25 | 7.50 | 1.77 | 4.00 | 7.11 | 1.60 | 7.26 | 12.84 | 3.39 |
| 1998 | 4.36 | 7.65 | 1.84 | 4.13 | 7.31 | 1.66 | 7.11 | 12.21 | 3.56 |
| 1999 | 4.35 | 7.69 | 1.77 | 4.19 | 7.44 | 1.65 | 6.54 | 11.56 | 3.06 |
| 2000 | 4.39 | 7.70 | 1.83 | 4.27 | 7.56 | 1.69 | 6.44 | 10.94 | 3.26 |
| 2001 | 4.43 | 7.84 | 1.79 | 4.30 | 7.65 | 1.67 | 6.43 | 11.33 | 3.02 |
| 2002 | 4.41 | 7.81 | 1.75 | 4.35 | 7.76 | 1.64 | 5.92 | 10.21 | 2.93 |
| 2003 | 4.38 | 7.74 | 1.74 | 4.33 | 7.70 | 1.65 | 5.72 | 9.72 | 2.87 |
| 2004 | 4.36 | 7.73 | 1.72 | 4.34 | 7.76 | 1.62 | 5.40 | 9.06 | 2.86 |
| 2005 | 4.44 | 7.95 | 1.66 | 4.48 | 8.07 | 1.60 | 5.04 | 8.90 | 2.34 |
| 2006 | 4.41 | 7.85 | 1.68 | 4.47 | 7.99 | 1.63 | 4.83 | 8.39 | 2.35 |
| 1975-2006 | 4.14 | 7.24 | 1.79 | 3.76 | 6.63 | 1.59 | 8.16 | 14.36 | 3.71 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 9.4
Hodgkin Lymphoma

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 3.08 | 3.86 | 2.39 | 3.23 | 3.91 | 2.63 | 2.58 | 4.41 | - |
| 1976 | 2.76 | 3.37 | 2.24 | 2.87 | 3.43 | 2.41 | 2.57 | 3.91 | - |
| 1977 | 2.98 | 3.60 | 2.46 | 3.26 | 3.92 | 2.71 | 1.68 | 1.91 | - |
| 1978 | 2.79 | 3.40 | 2.32 | 3.06 | 3.69 | 2.58 | 1.47 | 2.01 | - |
| 1979 | 2.93 | 3.91 | 2.08 | 3.08 | 4.05 | 2.26 | 2.86 | 4.68 | - |
| 1980 | 2.77 | 3.31 | 2.27 | 2.99 | 3.54 | 2.49 | 1.79 | 2.01 | 1.54 |
| 1981 | 2.91 | 3.58 | 2.33 | 3.22 | 3.91 | 2.61 | 1.39 | 1.86 | - |
| 1982 | 2.93 | 3.50 | 2.46 | 3.13 | 3.68 | 2.70 | 2.58 | 3.15 | 2.04 |
| 1983 | 3.02 | 3.72 | 2.39 | 3.33 | 3.99 | 2.70 | 2.11 | 3.85 | - |
| 1984 | 3.06 | 3.71 | 2.47 | 3.38 | 4.01 | 2.79 | 2.07 | 3.31 | - |
| 1985 | 2.97 | 3.49 | 2.51 | 3.31 | 3.83 | 2.83 | 1.74 | 2.26 | - |
| 1986 | 2.73 | 3.34 | 2.22 | 2.92 | 3.51 | 2.41 | 2.27 | 2.80 | 1.90 |
| 1987 | 3.05 | 3.72 | 2.50 | 3.35 | 4.00 | 2.80 | 2.05 | 2.78 | 1.41 |
| 1988 | 3.09 | 3.59 | 2.66 | 3.43 | 4.00 | 2.94 | 1.90 | 1.93 | 1.86 |
| 1989 | 3.05 | 3.67 | 2.54 | 3.35 | 3.97 | 2.84 | 2.16 | 3.15 | 1.37 |
| 1990 | 3.07 | 3.50 | 2.64 | 3.31 | 3.81 | 2.82 | 2.99 | 3.23 | 2.75 |
| 1991 | 3.01 | 3.48 | 2.59 | 3.34 | 3.83 | 2.90 | 2.38 | 2.65 | 2.08 |
| 1992 | 2.86 | 3.10 | 2.68 | 3.04 | 3.22 | 2.91 | 2.64 | 2.66 | 2.61 |
| 1993 | 2.85 | 3.26 | 2.49 | 3.08 | 3.48 | 2.74 | 2.92 | 3.70 | 2.35 |
| 1994 | 2.84 | 3.14 | 2.58 | 3.14 | 3.49 | 2.81 | 2.06 | 2.30 | 1.91 |
| 1995 | 2.76 | 3.22 | 2.35 | 3.05 | 3.52 | 2.64 | 2.29 | 2.84 | 1.78 |
| 1996 | 2.85 | 3.31 | 2.42 | 3.13 | 3.61 | 2.68 | 2.46 | 3.11 | 1.90 |
| 1997 | 2.81 | 2.94 | 2.69 | 3.07 | 3.23 | 2.91 | 2.54 | 2.38 | 2.67 |
| 1998 | 2.79 | 3.10 | 2.54 | 3.13 | 3.46 | 2.86 | 2.11 | 2.27 | 1.98 |
| 1999 | 2.83 | 3.06 | 2.66 | 3.08 | 3.22 | 2.99 | 2.46 | 3.01 | 2.01 |
| 2000 | 2.79 | 3.28 | 2.34 | 3.04 | 3.51 | 2.61 | 2.42 | 3.03 | 1.87 |
| 2001 | 2.55 | 2.90 | 2.22 | 2.71 | 3.08 | 2.35 | 2.41 | 3.11 | 1.92 |
| 2002 | 2.93 | 3.23 | 2.68 | 3.11 | 3.38 | 2.89 | 2.85 | 3.20 | 2.62 |
| 2003 | 2.69 | 3.08 | 2.37 | 2.97 | 3.37 | 2.60 | 2.37 | 2.72 | 2.22 |
| 2004 | 2.90 | 2.98 | 2.86 | 3.07 | 3.12 | 3.07 | 2.93 | 3.12 | 2.80 |
| 2005 | 2.98 | 3.42 | 2.61 | 3.19 | 3.64 | 2.82 | 3.06 | 3.29 | 2.86 |
| 2006 | 2.93 | 3.22 | 2.64 | 3.12 | 3.38 | 2.84 | 2.92 | 3.61 | 2.33 |
| 1975-2006 | 2.89 | 3.35 | 2.49 | 3.14 | 3.60 | 2.74 | 2.39 | 2.95 | 1.94 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 9.5
Hodgkin Lymphoma

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 1.31 | 1.69 | 1.00 | 1.35 | 1.73 | 1.04 | 1.07 | 1.50 | 0.72 |
| 1976 | 1.17 | 1.50 | 0.90 | 1.20 | 1.52 | 0.94 | 0.97 | 1.44 | 0.60 |
| 1977 | 1.14 | 1.45 | 0.87 | 1.18 | 1.49 | 0.90 | 0.92 | 1.20 | 0.69 |
| 1978 | 1.06 | 1.35 | 0.82 | 1.10 | 1.38 | 0.87 | 0.76 | 1.17 | 0.44 |
| 1979 | 0.97 | 1.27 | 0.71 | 1.00 | 1.31 | 0.74 | 0.76 | 1.06 | 0.51 |
| 1980 | 0.98 | 1.23 | 0.76 | 1.01 | 1.27 | 0.80 | 0.75 | 1.09 | 0.48 |
| 1981 | 0.93 | 1.15 | 0.74 | 0.96 | 1.19 | 0.76 | 0.74 | 0.88 | 0.62 |
| 1982 | 0.88 | 1.09 | 0.70 | 0.92 | 1.15 | 0.72 | 0.70 | 0.78 | 0.62 |
| 1983 | 0.85 | 1.09 | 0.65 | 0.88 | 1.13 | 0.68 | 0.64 | 0.90 | 0.45 |
| 1984 | 0.86 | 1.18 | 0.60 | 0.88 | 1.19 | 0.63 | 0.79 | 1.18 | 0.47 |
| 1985 | 0.77 | 0.98 | 0.58 | 0.78 | 1.00 | 0.60 | 0.71 | 0.96 | 0.51 |
| 1986 | 0.78 | 1.00 | 0.58 | 0.80 | 1.04 | 0.60 | 0.65 | 0.85 | 0.49 |
| 1987 | 0.74 | 0.96 | 0.56 | 0.77 | 0.98 | 0.58 | 0.67 | 0.94 | 0.46 |
| 1988 | 0.67 | 0.88 | 0.50 | 0.69 | 0.91 | 0.51 | 0.59 | 0.75 | 0.48 |
| 1989 | 0.71 | 0.92 | 0.54 | 0.73 | 0.94 | 0.56 | 0.66 | 0.92 | 0.47 |
| 1990 | 0.66 | 0.85 | 0.50 | 0.69 | 0.88 | 0.52 | 0.56 | 0.81 | 0.36 |
| 1991 | 0.65 | 0.83 | 0.51 | 0.66 | 0.83 | 0.51 | 0.67 | 0.89 | 0.49 |
| 1992 | 0.65 | 0.83 | 0.50 | 0.67 | 0.86 | 0.52 | 0.58 | 0.77 | 0.45 |
| 1993 | 0.61 | 0.79 | 0.47 | 0.63 | 0.80 | 0.50 | 0.57 | 0.83 | 0.39 |
| 1994 | 0.56 | 0.66 | 0.47 | 0.56 | 0.66 | 0.48 | 0.55 | 0.73 | 0.41 |
| 1995 | 0.55 | 0.67 | 0.44 | 0.56 | 0.68 | 0.46 | 0.53 | 0.73 | 0.36 |
| 1996 | 0.53 | 0.64 | 0.44 | 0.55 | 0.65 | 0.46 | 0.47 | 0.63 | 0.36 |
| 1997 | 0.53 | 0.62 | 0.46 | 0.54 | 0.63 | 0.47 | 0.54 | 0.67 | 0.43 |
| 1998 | 0.48 | 0.61 | 0.36 | 0.49 | 0.62 | 0.37 | 0.48 | 0.66 | 0.33 |
| 1999 | 0.51 | 0.63 | 0.41 | 0.53 | 0.66 | 0.43 | 0.41 | 0.55 | 0.30 |
| 2000 | 0.46 | 0.55 | 0.39 | 0.48 | 0.57 | 0.41 | 0.35 | 0.43 | 0.30 |
| 2001 | 0.46 | 0.56 | 0.38 | 0.48 | 0.59 | 0.39 | 0.42 | 0.52 | 0.35 |
| 2002 | 0.47 | 0.58 | 0.38 | 0.49 | 0.59 | 0.40 | 0.40 | 0.57 | 0.27 |
| 2003 | 0.46 | 0.55 | 0.39 | 0.48 | 0.57 | 0.41 | 0.36 | 0.48 | 0.28 |
| 2004 | 0.43 | 0.54 | 0.34 | 0.44 | 0.56 | 0.35 | 0.40 | 0.47 | 0.34 |
| 2005 | 0.42 | 0.50 | 0.36 | 0.43 | 0.50 | 0.37 | 0.42 | 0.56 | 0.32 |
| 2006 | 0.44 | 0.56 | 0.33 | 0.45 | 0.57 | 0.35 | 0.41 | 0.56 | 0.30 |
| 1975-2006 | 0.68 | 0.85 | 0.53 | 0.71 | 0.88 | 0.56 | 0.58 | 0.78 | 0.42 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 10.4
Kaposi Sarcoma

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 0.23 | 0.56 | - | 0.22 | 0.56 | - | - | - | - |
| 1976 | 0.25 | 0.47 | - | 0.24 | 0.48 | - | - | - | - |
| 1977 | 0.29 | 0.54 | - | 0.30 | 0.55 | - | - | - | - |
| 1978 | 0.24 | 0.35 | 0.17 | 0.23 | 0.34 | - | - | - | - |
| 1979 | 0.18 | 0.30 | - | 0.18 | 0.30 | - | - | - | - |
| 1980 | 0.28 | 0.56 | - | 0.30 | 0.60 | - | - | - | - |
| 1981 | 0.31 | 0.54 | - | 0.34 | 0.61 | - | - | - | - |
| 1982 | 0.65 | 1.20 | 0.18 | 0.71 | 1.32 | 0.18 | - | - | - |
| 1983 | 1.07 | 2.19 | - | 1.19 | 2.40 | - | - | - | - |
| 1984 | 1.81 | 3.58 | - | 1.98 | 3.90 | - | 1.27 | 2.89 | - |
| 1985 | 2.46 | 4.86 | 0.16 | 2.79 | 5.47 | 0.18 | 1.09 | 2.24 | - |
| 1986 | 3.27 | 6.54 | - | 3.71 | 7.34 | - | 1.65 | 3.59 | - |
| 1987 | 3.80 | 7.61 | - | 4.25 | 8.44 | - | 2.09 | 4.29 | - |
| 1988 | 4.42 | 8.86 | 0.15 | 4.82 | 9.56 | 0.15 | 3.40 | 7.28 | - |
| 1989 | 4.76 | 9.52 | 0.14 | 5.13 | 10.17 | - | 4.57 | 9.68 | - |
| 1990 | 4.61 | 9.21 | 0.18 | 5.03 | 9.98 | 0.15 | 3.65 | 7.54 | - |
| 1991 | 4.70 | 9.40 | 0.19 | 5.04 | 10.00 | 0.18 | 4.59 | 9.54 | - |
| 1992 | 4.37 | 8.68 | 0.18 | 4.59 | 9.01 | 0.18 | 4.97 | 10.48 | - |
| 1993 | 3.86 | 7.64 | 0.17 | 4.05 | 7.96 | 0.15 | 4.28 | 8.59 | - |
| 1994 | 3.79 | 7.47 | 0.17 | 3.88 | 7.59 | 0.13 | 4.15 | 8.42 | - |
| 1995 | 3.29 | 6.45 | 0.20 | 3.28 | 6.37 | 0.17 | 4.90 | 10.19 | - |
| 1996 | 2.23 | 4.40 | 0.16 | 2.01 | 3.98 | - | 4.61 | 9.09 | - |
| 1997 | 1.07 | 2.08 | 0.13 | 0.91 | 1.77 | - | 2.67 | 5.31 | - |
| 1998 | 0.81 | 1.57 | - | 0.69 | 1.31 | - | 1.79 | 3.78 | - |
| 1999 | 0.78 | 1.47 | 0.14 | 0.66 | 1.24 | - | 1.94 | 3.72 | - |
| 2000 | 0.80 | 1.54 | - | 0.66 | 1.29 | - | 1.98 | 4.00 | - |
| 2001 | 0.82 | 1.48 | 0.18 | 0.66 | 1.18 | 0.15 | 1.94 | 3.80 | - |
| 2002 | 0.67 | 1.25 | 0.11 | 0.54 | 1.01 | - | 1.49 | 3.01 | - |
| 2003 | 0.65 | 1.24 | - | 0.59 | 1.09 | - | 1.47 | 3.08 | - |
| 2004 | 0.67 | 1.29 | - | 0.57 | 1.09 | - | 1.63 | 3.26 | - |
| 2005 | 0.73 | 1.38 | 0.11 | 0.56 | 1.06 | - | 1.97 | 3.90 | - |
| 2006 | 0.55 | 1.07 | - | 0.45 | 0.85 | - | 1.26 | 2.62 | - |
| 1975-2006 | 1.88 | 3.71 | 0.13 | 1.95 | 3.82 | 0.12 | 2.25 | 4.60 | 0.24 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 11.4
Cancer of the Kidney and Renal Pelvis (Invasive)
Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 7.08 | 10.34 | 4.52 | 7.33 | 10.82 | 4.63 | 6.17 | 8.24 | 4.31 |
| 1976 | 7.96 | 11.20 | 5.47 | 8.13 | 11.48 | 5.59 | 7.75 | 11.68 | 4.69 |
| 1977 | 8.06 | 11.42 | 5.49 | 8.13 | 11.68 | 5.48 | 8.51 | 11.10 | 6.39 |
| 1978 | 7.84 | 11.81 | 4.83 | 7.96 | 12.08 | 4.88 | 8.01 | 11.01 | 5.65 |
| 1979 | 7.63 | 11.11 | 5.05 | 7.84 | 11.58 | 5.10 | 7.26 | 8.22 | 6.30 |
| 1980 | 8.06 | 11.69 | 5.38 | 8.36 | 12.21 | 5.55 | 5.97 | 7.95 | 4.36 |
| 1981 | 8.49 | 12.84 | 5.29 | 8.57 | 13.04 | 5.31 | 10.00 | 15.77 | 5.82 |
| 1982 | 8.35 | 11.82 | 5.72 | 8.53 | 12.03 | 5.87 | 7.59 | 11.20 | 4.90 |
| 1983 | 8.94 | 13.24 | 5.67 | 9.24 | 13.77 | 5.81 | 8.81 | 13.53 | 5.52 |
| 1984 | 9.19 | 13.12 | 6.20 | 9.47 | 13.61 | 6.33 | 9.02 | 12.01 | 6.90 |
| 1985 | 8.95 | 13.13 | 5.77 | 9.23 | 13.61 | 5.93 | 8.60 | 11.63 | 6.22 |
| 1986 | 9.66 | 13.84 | 6.56 | 9.83 | 14.09 | 6.67 | 10.12 | 14.97 | 6.62 |
| 1987 | 9.90 | 14.07 | 6.73 | 10.09 | 14.37 | 6.88 | 11.06 | 15.60 | 7.51 |
| 1988 | 9.93 | 13.99 | 6.95 | 10.10 | 14.31 | 7.03 | 11.59 | 15.36 | 8.77 |
| 1989 | 10.29 | 14.47 | 7.14 | 10.58 | 15.02 | 7.21 | 10.78 | 15.06 | 7.82 |
| 1990 | 10.43 | 14.71 | 7.14 | 10.70 | 14.97 | 7.45 | 10.51 | 15.94 | 6.61 |
| 1991 | 10.64 | 15.06 | 7.18 | 10.82 | 15.24 | 7.35 | 12.29 | 18.30 | 7.79 |
| 1992 | 10.78 | 15.29 | 7.41 | 11.08 | 15.60 | 7.73 | 10.98 | 16.39 | 7.18 |
| 1993 | 10.74 | 14.67 | 7.63 | 10.84 | 14.73 | 7.73 | 12.77 | 17.29 | 9.44 |
| 1994 | 11.29 | 15.65 | 7.86 | 11.54 | 15.98 | 8.04 | 12.71 | 18.01 | 8.66 |
| 1995 | 11.11 | 15.49 | 7.65 | 11.12 | 15.40 | 7.70 | 14.26 | 21.62 | 9.19 |
| 1996 | 11.35 | 15.84 | 7.91 | 11.46 | 16.23 | 7.77 | 13.78 | 17.38 | 11.13 |
| 1997 | 10.92 | 15.08 | 7.60 | 10.99 | 15.09 | 7.69 | 13.56 | 19.47 | 9.29 |
| 1998 | 11.79 | 16.40 | 8.16 | 12.07 | 16.70 | 8.43 | 12.62 | 17.22 | 9.05 |
| 1999 | 11.42 | 15.94 | 7.81 | 11.63 | 16.24 | 7.88 | 13.71 | 18.82 | 10.02 |
| 2000 | 12.45 | 17.37 | 8.54 | 12.71 | 17.90 | 8.46 | 14.78 | 19.24 | 11.84 |
| 2001 | 12.52 | 17.42 | 8.57 | 12.72 | 17.62 | 8.74 | 15.53 | 22.91 | 10.25 |
| 2002 | 12.76 | 17.60 | 8.82 | 13.00 | 17.82 | 9.03 | 15.14 | 22.05 | 10.29 |
| 2003 | 13.35 | 18.00 | 9.58 | 13.68 | 18.20 | 9.97 | 14.37 | 21.41 | 9.16 |
| 2004 | 13.34 | 18.07 | 9.47 | 13.66 | 18.46 | 9.68 | 15.52 | 21.74 | 10.82 |
| 2005 | 13.60 | 18.62 | 9.51 | 13.78 | 18.82 | 9.62 | 16.01 | 23.19 | 10.91 |
| 2006 | 13.90 | 18.11 | 10.38 | 14.14 | 18.40 | 10.51 | 15.08 | 19.65 | 11.60 |
| 1975-2006 | 10.68 | 14.98 | 7.33 | 10.82 | 15.21 | 7.41 | 12.05 | 16.97 | 8.43 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 11.5
Cancer of the Kidney and Renal Pelvis (Invasive)
Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 3.61 | 5.18 | 2.43 | 3.70 | 5.30 | 2.49 | 2.81 | 4.15 | 1.82 |
| 1976 | 3.61 | 5.23 | 2.41 | 3.69 | 5.35 | 2.45 | 2.92 | 4.24 | 1.92 |
| 1977 | 3.68 | 5.35 | 2.45 | 3.78 | 5.51 | 2.50 | 2.88 | 4.03 | 1.99 |
| 1978 | 3.69 | 5.43 | 2.41 | 3.76 | 5.52 | 2.46 | 3.17 | 4.83 | 1.93 |
| 1979 | 3.63 | 5.32 | 2.40 | 3.71 | 5.46 | 2.43 | 2.95 | 4.17 | 2.04 |
| 1980 | 3.68 | 5.46 | 2.38 | 3.77 | 5.61 | 2.44 | 2.96 | 4.19 | 2.01 |
| 1981 | 3.73 | 5.40 | 2.51 | 3.80 | 5.50 | 2.56 | 3.18 | 4.60 | 2.14 |
| 1982 | 3.85 | 5.68 | 2.54 | 3.92 | 5.80 | 2.56 | 3.40 | 4.83 | 2.37 |
| 1983 | 3.85 | 5.57 | 2.60 | 3.93 | 5.69 | 2.67 | 3.26 | 4.85 | 2.15 |
| 1984 | 3.90 | 5.75 | 2.57 | 3.96 | 5.88 | 2.60 | 3.45 | 4.86 | 2.45 |
| 1985 | 3.96 | 5.76 | 2.66 | 4.00 | 5.79 | 2.71 | 3.83 | 6.00 | 2.35 |
| 1986 | 4.05 | 5.83 | 2.78 | 4.11 | 5.94 | 2.82 | 3.68 | 5.34 | 2.52 |
| 1987 | 4.14 | 6.04 | 2.77 | 4.23 | 6.15 | 2.83 | 3.66 | 5.49 | 2.42 |
| 1988 | 4.03 | 5.86 | 2.71 | 4.09 | 5.93 | 2.76 | 3.78 | 5.82 | 2.41 |
| 1989 | 4.16 | 6.08 | 2.79 | 4.22 | 6.15 | 2.83 | 3.99 | 6.05 | 2.64 |
| 1990 | 4.19 | 6.16 | 2.77 | 4.23 | 6.22 | 2.79 | 4.20 | 6.35 | 2.76 |
| 1991 | 4.30 | 6.18 | 2.95 | 4.34 | 6.24 | 2.98 | 4.34 | 6.35 | 2.97 |
| 1992 | 4.29 | 6.24 | 2.88 | 4.37 | 6.33 | 2.94 | 3.97 | 5.94 | 2.61 |
| 1993 | 4.16 | 6.09 | 2.75 | 4.22 | 6.18 | 2.77 | 4.07 | 6.09 | 2.75 |
| 1994 | 4.27 | 6.20 | 2.89 | 4.32 | 6.25 | 2.92 | 4.24 | 6.40 | 2.81 |
| 1995 | 4.34 | 6.19 | 2.99 | 4.39 | 6.28 | 3.01 | 4.38 | 6.26 | 3.14 |
| 1996 | 4.27 | 6.19 | 2.84 | 4.33 | 6.29 | 2.88 | 4.15 | 5.93 | 2.93 |
| 1997 | 4.27 | 6.19 | 2.86 | 4.32 | 6.23 | 2.91 | 4.32 | 6.55 | 2.84 |
| 1998 | 4.26 | 6.19 | 2.81 | 4.34 | 6.27 | 2.88 | 4.01 | 6.28 | 2.46 |
| 1999 | 4.06 | 5.90 | 2.68 | 4.10 | 5.96 | 2.71 | 4.15 | 6.10 | 2.82 |
| 2000 | 4.21 | 6.14 | 2.79 | 4.29 | 6.23 | 2.84 | 4.13 | 6.28 | 2.74 |
| 2001 | 4.27 | 6.22 | 2.80 | 4.33 | 6.31 | 2.83 | 4.26 | 6.43 | 2.82 |
| 2002 | 4.22 | 6.14 | 2.76 | 4.32 | 6.28 | 2.80 | 4.06 | 5.84 | 2.84 |
| 2003 | 4.18 | 6.07 | 2.73 | 4.27 | 6.18 | 2.78 | 4.10 | 6.09 | 2.73 |
| 2004 | 4.12 | 5.92 | 2.72 | 4.22 | 6.06 | 2.78 | 3.98 | 5.86 | 2.70 |
| 2005 | 4.11 | 5.93 | 2.72 | 4.20 | 6.01 | 2.80 | 4.04 | 6.11 | 2.59 |
| 2006 | 3.99 | 5.74 | 2.63 | 4.07 | 5.83 | 2.70 | 3.87 | 5.87 | 2.49 |
| 1975-2006 | 4.07 | 5.91 | 2.71 | 4.14 | 6.01 | 2.75 | 3.83 | 5.66 | 2.57 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 12.4
Cancer of the Larynx (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 5.02 | 9.47 | 1.48 | 4.96 | 9.43 | 1.46 | 7.08 | 13.36 | - |
| 1976 | 5.22 | 10.01 | 1.39 | 5.19 | 9.95 | 1.44 | 7.31 | 14.61 | - |
| 1977 | 5.03 | 9.72 | 1.35 | 4.96 | 9.68 | 1.31 | 7.46 | 13.89 | 2.45 |
| 1978 | 5.32 | 9.80 | 1.84 | 5.32 | 9.80 | 1.85 | 6.95 | 13.20 | 2.22 |
| 1979 | 5.38 | 10.08 | 1.68 | 5.39 | 10.20 | 1.64 | 7.50 | 13.22 | 3.07 |
| 1980 | 5.25 | 10.01 | 1.54 | 5.16 | 9.88 | 1.52 | 7.34 | 14.47 | - |
| 1981 | 5.39 | 9.82 | 1.92 | 5.31 | 9.56 | 2.03 | 8.09 | 16.33 | - |
| 1982 | 5.30 | 9.76 | 1.80 | 5.27 | 9.75 | 1.79 | 7.91 | 14.49 | 2.88 |
| 1983 | 5.34 | 9.93 | 1.73 | 5.43 | 10.18 | 1.69 | 6.82 | 12.20 | 2.63 |
| 1984 | 5.10 | 9.62 | 1.58 | 4.99 | 9.58 | 1.46 | 8.46 | 14.85 | 3.44 |
| 1985 | 5.41 | 9.93 | 1.92 | 5.41 | 10.01 | 1.88 | 8.41 | 15.40 | 3.28 |
| 1986 | 5.00 | 9.39 | 1.61 | 4.93 | 9.19 | 1.67 | 7.74 | 15.36 | 2.08 |
| 1987 | 5.19 | 9.41 | 1.92 | 5.13 | 9.33 | 1.89 | 8.25 | 15.20 | 3.12 |
| 1988 | 5.20 | 9.58 | 1.87 | 5.19 | 9.57 | 1.85 | 8.17 | 14.93 | 3.21 |
| 1989 | 5.02 | 8.90 | 2.00 | 4.98 | 8.88 | 1.92 | 7.89 | 14.25 | 3.28 |
| 1990 | 5.10 | 9.17 | 1.92 | 4.99 | 8.96 | 1.87 | 9.46 | 17.72 | 3.59 |
| 1991 | 4.73 | 8.46 | 1.80 | 4.79 | 8.53 | 1.87 | 6.59 | 12.27 | 2.28 |
| 1992 | 4.97 | 9.06 | 1.77 | 4.87 | 8.88 | 1.72 | 8.64 | 15.66 | 3.40 |
| 1993 | 4.38 | 8.13 | 1.48 | 4.38 | 8.04 | 1.55 | 6.94 | 14.10 | 1.71 |
| 1994 | 4.66 | 8.33 | 1.76 | 4.53 | 8.13 | 1.67 | 8.91 | 15.96 | 3.71 |
| 1995 | 4.41 | 7.89 | 1.62 | 4.42 | 7.86 | 1.63 | 6.82 | 13.15 | 2.25 |
| 1996 | 4.27 | 7.58 | 1.64 | 4.27 | 7.58 | 1.61 | 6.48 | 11.51 | 2.78 |
| 1997 | 4.22 | 7.53 | 1.62 | 4.14 | 7.31 | 1.62 | 7.21 | 13.46 | 2.74 |
| 1998 | 3.86 | 6.78 | 1.59 | 3.72 | 6.51 | 1.55 | 6.60 | 12.03 | 2.73 |
| 1999 | 4.19 | 7.07 | 1.85 | 4.16 | 7.02 | 1.78 | 7.10 | 12.17 | 3.55 |
| 2000 | 4.05 | 7.27 | 1.44 | 4.05 | 7.21 | 1.47 | 6.05 | 11.39 | 2.18 |
| 2001 | 3.64 | 6.54 | 1.26 | 3.58 | 6.32 | 1.29 | 6.18 | 12.17 | 1.82 |
| 2002 | 3.54 | 6.29 | 1.32 | 3.45 | 6.11 | 1.29 | 6.30 | 11.23 | 2.67 |
| 2003 | 3.33 | 5.94 | 1.23 | 3.33 | 5.84 | 1.28 | 5.02 | 9.50 | 1.81 |
| 2004 | 3.45 | 6.07 | 1.36 | 3.42 | 5.93 | 1.38 | 5.82 | 10.99 | 2.27 |
| 2005 | 3.44 | 6.14 | 1.28 | 3.47 | 6.10 | 1.34 | 5.53 | 10.35 | 1.95 |
| 2006 | 3.24 | 5.71 | 1.22 | 3.28 | 5.65 | 1.31 | 5.17 | 10.22 | 1.50 |
| 1975-2006 | 4.52 | 8.21 | 1.60 | 4.51 | 8.17 | 1.60 | 6.99 | 13.09 | 2.49 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 12.5
Cancer of the Larynx (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 1.68 | 3.35 | 0.46 | 1.60 | 3.22 | 0.43 | 2.61 | 4.97 | 0.78 |
| 1976 | 1.72 | 3.41 | 0.49 | 1.63 | 3.26 | 0.47 | 2.63 | 5.13 | 0.71 |
| 1977 | 1.70 | 3.41 | 0.47 | 1.61 | 3.25 | 0.44 | 2.74 | 5.47 | 0.72 |
| 1978 | 1.70 | 3.40 | 0.49 | 1.59 | 3.18 | 0.46 | 2.90 | 5.88 | 0.71 |
| 1979 | 1.67 | 3.34 | 0.46 | 1.59 | 3.19 | 0.44 | 2.59 | 5.21 | 0.63 |
| 1980 | 1.61 | 3.17 | 0.48 | 1.49 | 2.95 | 0.44 | 2.91 | 5.73 | 0.87 |
| 1981 | 1.66 | 3.28 | 0.52 | 1.54 | 3.09 | 0.46 | 2.95 | 5.55 | 1.06 |
| 1982 | 1.67 | 3.23 | 0.54 | 1.57 | 3.05 | 0.51 | 2.79 | 5.51 | 0.83 |
| 1983 | 1.64 | 3.18 | 0.54 | 1.54 | 3.01 | 0.50 | 2.73 | 5.28 | 0.89 |
| 1984 | 1.63 | 3.16 | 0.53 | 1.50 | 2.90 | 0.51 | 3.05 | 6.21 | 0.75 |
| 1985 | 1.56 | 3.01 | 0.52 | 1.42 | 2.77 | 0.47 | 2.97 | 5.79 | 0.94 |
| 1986 | 1.59 | 3.06 | 0.55 | 1.45 | 2.81 | 0.50 | 3.07 | 6.11 | 0.96 |
| 1987 | 1.60 | 3.10 | 0.53 | 1.48 | 2.87 | 0.50 | 2.93 | 5.88 | 0.86 |
| 1988 | 1.58 | 3.04 | 0.53 | 1.43 | 2.73 | 0.51 | 3.21 | 6.73 | 0.76 |
| 1989 | 1.59 | 3.01 | 0.58 | 1.44 | 2.71 | 0.55 | 3.20 | 6.59 | 0.83 |
| 1990 | 1.55 | 2.97 | 0.54 | 1.42 | 2.73 | 0.49 | 3.00 | 5.79 | 1.08 |
| 1991 | 1.65 | 3.07 | 0.63 | 1.50 | 2.80 | 0.58 | 3.33 | 6.43 | 1.20 |
| 1992 | 1.61 | 3.00 | 0.61 | 1.46 | 2.73 | 0.57 | 3.26 | 6.39 | 1.07 |
| 1993 | 1.59 | 3.01 | 0.57 | 1.45 | 2.74 | 0.53 | 3.24 | 6.38 | 1.06 |
| 1994 | 1.55 | 2.92 | 0.57 | 1.41 | 2.63 | 0.53 | 3.20 | 6.41 | 1.04 |
| 1995 | 1.50 | 2.82 | 0.54 | 1.37 | 2.56 | 0.50 | 3.03 | 5.97 | 1.03 |
| 1996 | 1.50 | 2.78 | 0.56 | 1.36 | 2.49 | 0.53 | 3.16 | 6.35 | 0.97 |
| 1997 | 1.44 | 2.67 | 0.53 | 1.30 | 2.40 | 0.50 | 2.98 | 6.08 | 0.89 |
| 1998 | 1.43 | 2.58 | 0.57 | 1.31 | 2.34 | 0.54 | 2.83 | 5.55 | 0.90 |
| 1999 | 1.39 | 2.54 | 0.53 | 1.29 | 2.34 | 0.50 | 2.59 | 5.13 | 0.85 |
| 2000 | 1.39 | 2.55 | 0.52 | 1.27 | 2.32 | 0.49 | 2.79 | 5.46 | 0.95 |
| 2001 | 1.34 | 2.48 | 0.49 | 1.24 | 2.27 | 0.46 | 2.56 | 5.01 | 0.88 |
| 2002 | 1.29 | 2.33 | 0.50 | 1.20 | 2.15 | 0.47 | 2.48 | 4.85 | 0.85 |
| 2003 | 1.29 | 2.36 | 0.47 | 1.20 | 2.16 | 0.45 | 2.51 | 5.13 | 0.75 |
| 2004 | 1.22 | 2.25 | 0.44 | 1.15 | 2.09 | 0.43 | 2.25 | 4.60 | 0.66 |
| 2005 | 1.25 | 2.24 | 0.48 | 1.16 | 2.07 | 0.47 | 2.32 | 4.62 | 0.77 |
| 2006 | 1.23 | 2.19 | 0.48 | 1.16 | 2.04 | 0.48 | 2.10 | 4.17 | 0.68 |
| 1975-2006 | 1.51 | 2.84 | 0.52 | 1.40 | 2.63 | 0.49 | 2.79 | 5.54 | 0.86 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 13.4
LeukemiaAge-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 12.78 | 16.91 | 9.95 | 13.09 | 17.48 | 10.10 | 11.52 | 16.21 | 8.47 |
| 1976 | 13.55 | 18.48 | 10.21 | 13.84 | 19.13 | 10.31 | 11.30 | 14.19 | 9.39 |
| 1977 | 12.74 | 16.34 | 10.22 | 13.20 | 17.04 | 10.52 | 10.15 | 13.61 | 7.72 |
| 1978 | 12.87 | 17.25 | 9.78 | 13.54 | 18.30 | 10.24 | 9.16 | 11.42 | 7.47 |
| 1979 | 12.56 | 16.80 | 9.69 | 13.02 | 17.55 | 10.01 | 10.63 | 13.59 | 8.64 |
| 1980 | 12.91 | 17.35 | 9.84 | 13.26 | 17.90 | 10.10 | 12.18 | 16.04 | 9.54 |
| 1981 | 12.38 | 16.49 | 9.64 | 12.65 | 17.05 | 9.81 | 12.24 | 14.47 | 10.34 |
| 1982 | 13.17 | 18.25 | 9.72 | 13.52 | 18.80 | 9.96 | 10.75 | 15.16 | 7.91 |
| 1983 | 13.15 | 17.82 | 9.91 | 13.49 | 18.44 | 10.07 | 12.48 | 15.95 | 10.06 |
| 1984 | 12.99 | 17.08 | 10.10 | 13.44 | 17.94 | 10.34 | 10.68 | 11.87 | 9.61 |
| 1985 | 13.35 | 17.71 | 10.38 | 13.72 | 18.22 | 10.65 | 12.23 | 16.82 | 9.30 |
| 1986 | 12.83 | 17.53 | 9.50 | 13.39 | 18.37 | 9.92 | 10.23 | 13.20 | 8.09 |
| 1987 | 13.34 | 17.72 | 10.31 | 13.77 | 18.44 | 10.58 | 12.57 | 17.60 | 9.26 |
| 1988 | 12.94 | 17.47 | 9.68 | 13.38 | 18.25 | 9.96 | 11.55 | 14.03 | 9.42 |
| 1989 | 13.41 | 17.63 | 10.42 | 13.92 | 18.26 | 10.93 | 11.26 | 15.91 | 8.02 |
| 1990 | 12.85 | 17.25 | 9.74 | 13.47 | 18.20 | 10.18 | 11.31 | 14.78 | 8.82 |
| 1991 | 13.03 | 16.76 | 10.20 | 13.64 | 17.70 | 10.54 | 10.81 | 11.97 | 10.06 |
| 1992 | 12.99 | 17.48 | 9.75 | 13.56 | 18.61 | 9.95 | 10.68 | 13.91 | 8.42 |
| 1993 | 12.99 | 16.97 | 9.99 | 13.54 | 17.89 | 10.25 | 11.57 | 15.12 | 9.09 |
| 1994 | 12.96 | 16.85 | 10.09 | 13.61 | 17.75 | 10.57 | 9.82 | 12.61 | 7.78 |
| 1995 | 13.45 | 17.62 | 10.35 | 14.31 | 18.71 | 11.04 | 9.82 | 13.35 | 7.42 |
| 1996 | 13.02 | 16.85 | 10.28 | 13.58 | 17.60 | 10.72 | 10.75 | 13.91 | 8.79 |
| 1997 | 13.06 | 17.47 | 9.85 | 13.87 | 18.56 | 10.45 | 10.44 | 13.88 | 8.19 |
| 1998 | 13.01 | 16.97 | 10.09 | 13.79 | 17.85 | 10.78 | 9.73 | 14.20 | 6.96 |
| 1999 | 12.61 | 16.88 | 9.45 | 13.22 | 17.64 | 9.94 | 10.19 | 13.69 | 7.93 |
| 2000 | 13.28 | 17.19 | 10.36 | 13.93 | 18.07 | 10.84 | 11.30 | 13.07 | 9.97 |
| 2001 | 13.86 | 18.04 | 10.70 | 14.71 | 19.03 | 11.44 | 11.42 | 13.97 | 9.64 |
| 2002 | 13.00 | 17.11 | 9.88 | 13.99 | 18.43 | 10.60 | 8.39 | 11.10 | 6.54 |
| 2003 | 12.87 | 16.82 | 9.85 | 13.55 | 17.85 | 10.28 | 10.57 | 13.43 | 8.46 |
| 2004 | 12.92 | 16.72 | 10.04 | 13.51 | 17.42 | 10.47 | 11.25 | 15.38 | 8.70 |
| 2005 | 12.46 | 16.19 | 9.57 | 13.17 | 17.36 | 9.91 | 10.15 | 11.37 | 9.25 |
| 2006 | 11.89 | 14.83 | 9.70 | 12.59 | 15.56 | 10.38 | 9.18 | 12.90 | 6.67 |
| 1975-2006 | 12.93 | 17.04 | 9.95 | 13.51 | 17.89 | 10.35 | 10.71 | 13.85 | 8.55 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 13.5
Acute Lymphocytic Leukemia

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 0.93 | 1.12 | 0.76 | 1.00 | 1.23 | 0.79 | - | - | - |
| 1976 | 1.08 | 1.29 | 0.93 | 1.09 | 1.34 | 0.91 | 0.79 | - | - |
| 1977 | 1.10 | 1.30 | 0.91 | 1.17 | 1.38 | 0.96 | - | - | - |
| 1978 | 1.03 | 1.24 | 0.83 | 1.11 | 1.33 | 0.90 | - | - | - |
| 1979 | 1.02 | 1.18 | 0.88 | 1.08 | 1.20 | 0.97 | - | - | - |
| 1980 | 1.12 | 1.41 | 0.88 | 1.16 | 1.47 | 0.90 | 0.73 | - | - |
| 1981 | 1.24 | 1.31 | 1.18 | 1.30 | 1.39 | 1.23 | 0.95 | - | - |
| 1982 | 1.23 | 1.50 | 0.99 | 1.30 | 1.56 | 1.05 | - | - | - |
| 1983 | 1.29 | 1.53 | 1.08 | 1.39 | 1.65 | 1.17 | 0.80 | - | - |
| 1984 | 1.32 | 1.49 | 1.20 | 1.35 | 1.50 | 1.24 | 0.92 | - | - |
| 1985 | 1.30 | 1.55 | 1.09 | 1.35 | 1.62 | 1.12 | - | - | - |
| 1986 | 1.30 | 1.67 | 0.96 | 1.38 | 1.82 | 0.97 | 0.77 | - | - |
| 1987 | 1.32 | 1.48 | 1.19 | 1.35 | 1.55 | 1.19 | 0.89 | - | 1.15 |
| 1988 | 1.40 | 1.78 | 1.07 | 1.44 | 1.83 | 1.12 | 0.98 | - | - |
| 1989 | 1.62 | 1.81 | 1.44 | 1.69 | 1.89 | 1.50 | 1.03 | - | - |
| 1990 | 1.41 | 1.75 | 1.16 | 1.52 | 1.85 | 1.28 | 0.57 | - | - |
| 1991 | 1.38 | 1.73 | 1.05 | 1.52 | 1.95 | 1.09 | 0.86 | - | - |
| 1992 | 1.35 | 1.59 | 1.14 | 1.39 | 1.72 | 1.09 | 0.91 | - | - |
| 1993 | 1.26 | 1.41 | 1.09 | 1.37 | 1.57 | 1.13 | 0.82 | - | - |
| 1994 | 1.19 | 1.48 | 0.94 | 1.29 | 1.59 | 1.02 | 0.68 | - | - |
| 1995 | 1.49 | 1.80 | 1.20 | 1.60 | 1.87 | 1.34 | 0.96 | 1.27 | - |
| 1996 | 1.38 | 1.56 | 1.20 | 1.46 | 1.65 | 1.25 | 0.85 | - | - |
| 1997 | 1.34 | 1.52 | 1.16 | 1.45 | 1.63 | 1.28 | 0.75 | 0.89 | - |
| 1998 | 1.47 | 1.63 | 1.30 | 1.58 | 1.74 | 1.41 | 0.93 | 1.16 | - |
| 1999 | 1.31 | 1.49 | 1.13 | 1.40 | 1.58 | 1.23 | 0.68 | - | - |
| 2000 | 1.39 | 1.49 | 1.30 | 1.52 | 1.59 | 1.46 | 0.85 | - | - |
| 2001 | 1.45 | 1.64 | 1.26 | 1.58 | 1.80 | 1.37 | 0.65 | - | - |
| 2002 | 1.49 | 1.78 | 1.20 | 1.64 | 1.99 | 1.28 | 0.61 | - | - |
| 2003 | 1.41 | 1.63 | 1.21 | 1.47 | 1.68 | 1.27 | 0.72 | - | - |
| 2004 | 1.53 | 1.67 | 1.37 | 1.63 | 1.90 | 1.36 | 1.06 | - | 1.34 |
| 2005 | 1.55 | 1.85 | 1.25 | 1.65 | 1.95 | 1.34 | 0.89 | 1.00 | - |
| 2006 | 1.42 | 1.58 | 1.26 | 1.46 | 1.59 | 1.34 | 0.95 | 1.35 | - |
| 1975-2006 | 1.32 | 1.54 | 1.12 | 1.40 | 1.63 | 1.18 | 0.79 | 0.86 | 0.72 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 13.6
Chronic Lymphocytic Leukemia

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 4.33 | 5.90 | 3.26 | 4.47 | 6.11 | 3.37 | 4.76 | 6.94 | 3.23 |
| 1976 | 4.57 | 6.77 | 3.10 | 4.77 | 7.18 | 3.17 | 4.26 | 6.58 | 2.95 |
| 1977 | 4.18 | 5.72 | 3.13 | 4.37 | 5.96 | 3.31 | 3.81 | 5.78 | 2.46 |
| 1978 | 4.46 | 5.94 | 3.43 | 4.75 | 6.43 | 3.61 | 3.33 | 3.64 | 3.07 |
| 1979 | 4.05 | 5.69 | 2.93 | 4.31 | 6.09 | 3.12 | 3.06 | 4.16 | - |
| 1980 | 4.51 | 6.30 | 3.32 | 4.68 | 6.54 | 3.45 | 4.51 | 7.06 | 2.89 |
| 1981 | 4.26 | 6.30 | 2.88 | 4.46 | 6.63 | 3.03 | 3.79 | 5.41 | 2.44 |
| 1982 | 4.31 | 6.16 | 3.02 | 4.52 | 6.42 | 3.19 | 3.77 | 6.65 | - |
| 1983 | 4.46 | 6.40 | 3.14 | 4.68 | 6.77 | 3.26 | 4.21 | 6.11 | 3.03 |
| 1984 | 4.41 | 6.27 | 3.11 | 4.70 | 6.79 | 3.28 | 3.45 | 4.46 | 2.65 |
| 1985 | 4.45 | 6.14 | 3.30 | 4.75 | 6.53 | 3.52 | 4.11 | 6.15 | 2.87 |
| 1986 | 4.49 | 6.31 | 3.23 | 4.85 | 6.82 | 3.53 | 2.79 | 4.18 | - |
| 1987 | 4.63 | 6.77 | 3.05 | 4.92 | 7.21 | 3.26 | 4.03 | 6.14 | 2.45 |
| 1988 | 4.51 | 6.42 | 3.11 | 4.72 | 6.83 | 3.22 | 4.43 | 5.32 | 3.53 |
| 1989 | 4.64 | 6.39 | 3.40 | 4.93 | 6.84 | 3.59 | 3.91 | 5.26 | 2.95 |
| 1990 | 4.55 | 6.25 | 3.36 | 4.85 | 6.72 | 3.54 | 4.51 | 6.13 | 3.43 |
| 1991 | 4.55 | 6.13 | 3.39 | 4.94 | 6.74 | 3.59 | 3.40 | 3.63 | 3.24 |
| 1992 | 4.55 | 6.65 | 3.11 | 4.97 | 7.32 | 3.37 | 3.21 | 4.96 | 2.08 |
| 1993 | 4.37 | 6.03 | 3.12 | 4.63 | 6.42 | 3.29 | 4.44 | 6.61 | 2.89 |
| 1994 | 4.48 | 5.99 | 3.33 | 4.87 | 6.52 | 3.65 | 3.19 | 4.13 | 2.32 |
| 1995 | 4.42 | 6.30 | 3.06 | 4.80 | 6.84 | 3.34 | 3.34 | 5.22 | 2.06 |
| 1996 | 4.56 | 6.17 | 3.41 | 4.96 | 6.64 | 3.76 | 3.55 | 5.36 | 2.31 |
| 1997 | 4.36 | 5.95 | 3.15 | 4.74 | 6.46 | 3.45 | 3.39 | 4.96 | 2.37 |
| 1998 | 4.07 | 5.76 | 2.84 | 4.45 | 6.28 | 3.13 | 2.72 | 3.56 | 2.21 |
| 1999 | 3.96 | 5.80 | 2.60 | 4.27 | 6.23 | 2.81 | 3.56 | 5.61 | 2.35 |
| 2000 | 4.23 | 5.95 | 2.93 | 4.67 | 6.61 | 3.18 | 2.95 | 3.61 | 2.54 |
| 2001 | 4.82 | 6.76 | 3.37 | 5.27 | 7.38 | 3.70 | 3.56 | 4.60 | 2.82 |
| 2002 | 4.56 | 6.42 | 3.15 | 5.09 | 7.17 | 3.51 | 2.63 | 3.77 | 1.82 |
| 2003 | 4.75 | 6.57 | 3.36 | 5.19 | 7.23 | 3.64 | 3.69 | 5.15 | 2.64 |
| 2004 | 4.66 | 6.45 | 3.31 | 5.04 | 6.82 | 3.68 | 3.52 | 6.22 | 1.76 |
| 2005 | 4.32 | 6.20 | 2.88 | 4.74 | 6.88 | 3.06 | 3.16 | 3.71 | 2.78 |
| 2006 | 4.29 | 5.79 | 3.19 | 4.65 | 6.23 | 3.49 | 3.12 | 5.03 | 1.91 |
| 1975-2006 | 4.43 | 6.19 | 3.15 | 4.76 | 6.68 | 3.38 | 3.54 | 5.05 | 2.51 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 13.7
Acute Myeloid Leukemia

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 3.43 | 4.17 | 2.98 | 3.48 | 4.28 | 2.99 | 2.49 | 3.37 | 2.02 |
| 1976 | 3.50 | 4.43 | 2.86 | 3.53 | 4.46 | 2.91 | 2.56 | - | 2.64 |
| 1977 | 3.51 | 4.19 | 3.05 | 3.64 | 4.40 | 3.14 | 1.87 | - | - |
| 1978 | 3.42 | 4.43 | 2.69 | 3.53 | 4.59 | 2.77 | 2.70 | 3.47 | 2.13 |
| 1979 | 3.32 | 4.05 | 2.85 | 3.36 | 4.17 | 2.85 | 3.43 | 3.47 | 3.37 |
| 1980 | 3.14 | 4.10 | 2.43 | 3.16 | 4.11 | 2.48 | 2.88 | 3.28 | 2.53 |
| 1981 | 3.16 | 3.99 | 2.64 | 3.22 | 4.11 | 2.69 | 3.07 | 3.22 | 2.89 |
| 1982 | 3.42 | 4.80 | 2.59 | 3.47 | 4.91 | 2.62 | 2.50 | 3.22 | 2.09 |
| 1983 | 3.21 | 4.22 | 2.53 | 3.18 | 4.29 | 2.41 | 2.93 | 3.02 | 2.92 |
| 1984 | 3.30 | 4.06 | 2.73 | 3.35 | 4.16 | 2.77 | 2.35 | 2.10 | 2.44 |
| 1985 | 3.33 | 4.15 | 2.87 | 3.25 | 4.02 | 2.84 | 3.71 | 4.91 | 2.99 |
| 1986 | 3.00 | 3.91 | 2.40 | 2.95 | 3.85 | 2.38 | 3.21 | 4.04 | 2.68 |
| 1987 | 3.02 | 3.98 | 2.45 | 3.03 | 4.01 | 2.46 | 3.01 | 4.73 | 2.04 |
| 1988 | 3.03 | 3.79 | 2.51 | 3.07 | 3.87 | 2.56 | 2.79 | 3.14 | 2.49 |
| 1989 | 3.19 | 4.00 | 2.61 | 3.29 | 4.02 | 2.78 | 2.22 | 3.79 | - |
| 1990 | 3.04 | 3.89 | 2.43 | 3.11 | 4.03 | 2.47 | 2.51 | 3.02 | 2.12 |
| 1991 | 3.32 | 3.91 | 2.88 | 3.37 | 3.89 | 2.98 | 3.28 | 4.38 | 2.59 |
| 1992 | 3.22 | 3.86 | 2.76 | 3.28 | 4.05 | 2.74 | 3.05 | 2.99 | 3.05 |
| 1993 | 3.45 | 4.31 | 2.86 | 3.52 | 4.45 | 2.86 | 2.48 | 2.62 | 2.55 |
| 1994 | 3.45 | 4.14 | 3.00 | 3.48 | 4.25 | 2.96 | 2.88 | 3.24 | 2.67 |
| 1995 | 3.66 | 4.16 | 3.31 | 3.81 | 4.31 | 3.45 | 2.60 | 2.56 | 2.73 |
| 1996 | 3.47 | 4.33 | 2.90 | 3.46 | 4.27 | 2.94 | 3.24 | 4.10 | 2.73 |
| 1997 | 3.52 | 4.59 | 2.78 | 3.66 | 4.81 | 2.85 | 3.02 | 3.78 | 2.60 |
| 1998 | 3.90 | 4.92 | 3.18 | 4.04 | 5.04 | 3.31 | 3.05 | 4.64 | 2.11 |
| 1999 | 3.72 | 4.51 | 3.18 | 3.84 | 4.62 | 3.29 | 3.03 | 3.45 | 2.76 |
| 2000 | 4.04 | 4.93 | 3.40 | 4.04 | 4.95 | 3.37 | 4.40 | 4.87 | 4.00 |
| 2001 | 4.03 | 4.91 | 3.38 | 4.17 | 4.96 | 3.59 | 3.43 | 4.07 | 2.98 |
| 2002 | 3.61 | 4.54 | 2.96 | 3.75 | 4.74 | 3.08 | 2.88 | 3.24 | 2.65 |
| 2003 | 3.56 | 4.47 | 2.91 | 3.64 | 4.59 | 2.98 | 3.26 | 3.44 | 2.93 |
| 2004 | 3.36 | 4.05 | 2.90 | 3.45 | 4.13 | 2.97 | 3.25 | 4.04 | 2.90 |
| 2005 | 3.51 | 4.12 | 3.13 | 3.56 | 4.25 | 3.14 | 3.41 | 3.34 | 3.51 |
| 2006 | 3.38 | 3.85 | 3.06 | 3.47 | 3.95 | 3.13 | 2.72 | 3.18 | 2.46 |
| 1975-2006 | 3.43 | 4.25 | 2.86 | 3.48 | 4.33 | 2.90 | 2.99 | 3.53 | 2.64 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 13.8
Chronic Myeloid Leukemia

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 1.98 | 2.93 | 1.33 | 2.00 | 3.01 | 1.31 | 1.94 | - | - |
| 1976 | 1.93 | 2.48 | 1.55 | 1.90 | 2.45 | 1.52 | 1.86 | 2.58 | - |
| 1977 | 1.68 | 2.08 | 1.39 | 1.69 | 2.14 | 1.37 | 1.84 | - | - |
| 1978 | 1.72 | 2.26 | 1.34 | 1.74 | 2.28 | 1.39 | 1.43 | - | - |
| 1979 | 1.87 | 2.48 | 1.43 | 1.89 | 2.49 | 1.45 | 1.94 | 2.80 | - |
| 1980 | 1.70 | 2.05 | 1.46 | 1.73 | 2.09 | 1.51 | 1.48 | - | - |
| 1981 | 1.64 | 1.97 | 1.41 | 1.59 | 1.90 | 1.39 | 2.38 | 3.00 | 1.81 |
| 1982 | 1.68 | 2.16 | 1.36 | 1.66 | 2.19 | 1.32 | 1.75 | - | 1.79 |
| 1983 | 1.69 | 2.23 | 1.33 | 1.64 | 2.15 | 1.32 | 2.51 | 3.44 | 1.85 |
| 1984 | 1.67 | 2.40 | 1.18 | 1.62 | 2.40 | 1.10 | 2.58 | 2.61 | 2.59 |
| 1985 | 1.98 | 2.76 | 1.44 | 2.01 | 2.80 | 1.48 | 2.06 | 2.89 | - |
| 1986 | 1.68 | 2.20 | 1.29 | 1.71 | 2.26 | 1.31 | 1.54 | - | - |
| 1987 | 1.88 | 2.32 | 1.58 | 1.90 | 2.38 | 1.58 | 2.14 | 2.88 | - |
| 1988 | 1.91 | 2.54 | 1.50 | 1.92 | 2.59 | 1.49 | 2.08 | 2.88 | - |
| 1989 | 1.69 | 2.22 | 1.34 | 1.67 | 2.18 | 1.34 | 2.21 | 3.12 | - |
| 1990 | 1.75 | 2.46 | 1.24 | 1.75 | 2.50 | 1.20 | 2.57 | 3.53 | 1.95 |
| 1991 | 1.88 | 2.44 | 1.47 | 1.91 | 2.55 | 1.46 | 1.56 | - | 1.75 |
| 1992 | 1.76 | 2.16 | 1.47 | 1.80 | 2.22 | 1.49 | 1.49 | 1.73 | - |
| 1993 | 1.92 | 2.53 | 1.48 | 1.95 | 2.52 | 1.56 | 2.26 | 3.56 | - |
| 1994 | 1.86 | 2.46 | 1.42 | 1.90 | 2.48 | 1.50 | 1.65 | 2.47 | - |
| 1995 | 1.89 | 2.55 | 1.37 | 1.96 | 2.69 | 1.39 | 1.54 | 1.98 | - |
| 1996 | 1.81 | 2.45 | 1.35 | 1.81 | 2.53 | 1.28 | 1.99 | 2.39 | 1.83 |
| 1997 | 1.86 | 2.67 | 1.27 | 1.92 | 2.73 | 1.34 | 1.67 | 2.53 | - |
| 1998 | 1.63 | 1.92 | 1.41 | 1.71 | 1.94 | 1.54 | 1.48 | 2.31 | - |
| 1999 | 1.78 | 2.45 | 1.28 | 1.76 | 2.41 | 1.27 | 1.68 | 2.20 | 1.26 |
| 2000 | 1.77 | 2.31 | 1.35 | 1.79 | 2.32 | 1.38 | 1.55 | 1.72 | 1.33 |
| 2001 | 1.75 | 2.28 | 1.36 | 1.79 | 2.30 | 1.41 | 1.87 | 2.58 | 1.35 |
| 2002 | 1.57 | 2.01 | 1.24 | 1.61 | 2.07 | 1.28 | 1.00 | - | - |
| 2003 | 1.54 | 2.08 | 1.12 | 1.63 | 2.21 | 1.16 | 1.36 | 1.92 | - |
| 2004 | 1.73 | 2.42 | 1.20 | 1.79 | 2.52 | 1.23 | 1.39 | 1.84 | 1.08 |
| 2005 | 1.52 | 2.04 | 1.11 | 1.60 | 2.11 | 1.18 | 1.47 | 2.30 | - |
| 2006 | 1.55 | 2.03 | 1.17 | 1.64 | 2.13 | 1.26 | 1.18 | 1.63 | - |
| 1975-2006 | 1.75 | 2.31 | 1.34 | 1.77 | 2.35 | 1.36 | 1.75 | 2.30 | 1.35 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 13.9
LeukemiaAge-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 8.09 | 10.98 | 6.16 | 8.27 | 11.28 | 6.29 | 6.58 | 8.57 | 5.15 |
| 1976 | 8.10 | 10.88 | 6.19 | 8.27 | 11.19 | 6.28 | 6.76 | 8.49 | 5.48 |
| 1977 | 8.11 | 10.84 | 6.26 | 8.27 | 11.11 | 6.34 | 6.82 | 8.76 | 5.51 |
| 1978 | 8.02 | 10.83 | 6.15 | 8.19 | 11.06 | 6.28 | 6.70 | 9.25 | 4.94 |
| 1979 | 8.26 | 11.06 | 6.41 | 8.43 | 11.36 | 6.52 | 6.99 | 8.90 | 5.68 |
| 1980 | 8.36 | 11.21 | 6.46 | 8.54 | 11.51 | 6.56 | 7.05 | 9.09 | 5.67 |
| 1981 | 8.10 | 10.89 | 6.24 | 8.24 | 11.16 | 6.31 | 7.11 | 8.92 | 5.86 |
| 1982 | 8.19 | 11.04 | 6.31 | 8.33 | 11.29 | 6.39 | 7.14 | 9.36 | 5.67 |
| 1983 | 8.11 | 10.93 | 6.26 | 8.24 | 11.19 | 6.32 | 7.22 | 9.21 | 5.90 |
| 1984 | 8.08 | 10.68 | 6.37 | 8.23 | 10.91 | 6.47 | 7.08 | 9.16 | 5.69 |
| 1985 | 8.07 | 10.77 | 6.31 | 8.23 | 10.97 | 6.44 | 7.08 | 9.62 | 5.42 |
| 1986 | 7.96 | 10.71 | 6.16 | 8.10 | 10.97 | 6.23 | 7.25 | 9.26 | 5.91 |
| 1987 | 7.80 | 10.31 | 6.14 | 7.91 | 10.50 | 6.20 | 7.24 | 9.41 | 5.85 |
| 1988 | 7.83 | 10.60 | 6.00 | 7.95 | 10.79 | 6.08 | 7.24 | 9.83 | 5.58 |
| 1989 | 7.95 | 10.62 | 6.15 | 8.07 | 10.75 | 6.26 | 7.36 | 10.34 | 5.45 |
| 1990 | 7.98 | 10.71 | 6.18 | 8.10 | 10.90 | 6.26 | 7.44 | 9.96 | 5.83 |
| 1991 | 7.99 | 10.50 | 6.32 | 8.12 | 10.64 | 6.44 | 7.45 | 10.05 | 5.80 |
| 1992 | 7.96 | 10.71 | 6.13 | 8.11 | 10.94 | 6.24 | 7.16 | 9.41 | 5.68 |
| 1993 | 7.93 | 10.66 | 6.09 | 8.04 | 10.84 | 6.16 | 7.36 | 9.76 | 5.77 |
| 1994 | 7.85 | 10.57 | 6.03 | 8.00 | 10.78 | 6.13 | 7.12 | 9.59 | 5.54 |
| 1995 | 7.90 | 10.80 | 5.98 | 8.04 | 11.03 | 6.05 | 7.17 | 9.75 | 5.53 |
| 1996 | 7.84 | 10.53 | 6.05 | 7.97 | 10.71 | 6.13 | 7.46 | 10.09 | 5.90 |
| 1997 | 7.71 | 10.42 | 5.89 | 7.89 | 10.68 | 6.01 | 6.83 | 9.23 | 5.29 |
| 1998 | 7.58 | 10.11 | 5.82 | 7.72 | 10.31 | 5.91 | 7.04 | 9.37 | 5.55 |
| 1999 | 7.68 | 10.22 | 5.96 | 7.87 | 10.47 | 6.09 | 6.74 | 8.78 | 5.43 |
| 2000 | 7.68 | 10.27 | 5.90 | 7.88 | 10.53 | 6.04 | 6.71 | 9.06 | 5.27 |
| 2001 | 7.61 | 10.14 | 5.87 | 7.81 | 10.42 | 6.00 | 6.72 | 8.76 | 5.40 |
| 2002 | 7.51 | 10.10 | 5.70 | 7.71 | 10.39 | 5.82 | 6.69 | 8.81 | 5.35 |
| 2003 | 7.40 | 9.93 | 5.59 | 7.62 | 10.28 | 5.71 | 6.50 | 8.37 | 5.26 |
| 2004 | 7.24 | 9.71 | 5.47 | 7.41 | 9.95 | 5.59 | 6.73 | 8.84 | 5.35 |
| 2005 | 7.19 | 9.72 | 5.40 | 7.43 | 10.04 | 5.57 | 6.03 | 8.03 | 4.74 |
| 2006 | 7.16 | 9.61 | 5.40 | 7.41 | 9.89 | 5.59 | 6.24 | 8.71 | 4.67 |
| 1975-2006 | 7.82 | 10.49 | 6.02 | 7.99 | 10.74 | 6.13 | 6.94 | 9.17 | 5.49 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 14.4
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)
Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 2.64 | 3.93 | 1.64 | 2.19 | 3.23 | 1.43 | 4.74 | 6.46 | 3.19 |
| 1976 | 2.66 | 3.83 | 1.76 | 2.29 | 3.16 | 1.62 | 4.03 | 6.23 | - |
| 1977 | 2.68 | 4.25 | 1.56 | 2.24 | 3.57 | 1.33 | 3.47 | 5.02 | - |
| 1978 | 2.67 | 4.15 | 1.56 | 2.17 | 3.27 | 1.39 | 4.23 | 6.82 | 2.13 |
| 1979 | 2.68 | 4.06 | 1.67 | 2.23 | 3.30 | 1.50 | 4.07 | 7.10 | - |
| 1980 | 2.62 | 3.94 | 1.66 | 2.25 | 3.29 | 1.54 | 3.62 | 5.01 | 2.47 |
| 1981 | 2.93 | 4.65 | 1.66 | 2.49 | 3.88 | 1.51 | 4.59 | 7.09 | 2.53 |
| 1982 | 2.94 | 4.37 | 1.88 | 2.41 | 3.48 | 1.64 | 3.85 | 5.40 | 2.63 |
| 1983 | 2.91 | 4.68 | 1.58 | 2.38 | 3.74 | 1.41 | 4.11 | 6.30 | 2.33 |
| 1984 | 2.86 | 4.27 | 1.87 | 2.32 | 3.41 | 1.60 | 3.34 | 5.26 | 1.91 |
| 1985 | 3.16 | 5.10 | 1.73 | 2.61 | 4.15 | 1.51 | 3.96 | 7.64 | - |
| 1986 | 3.19 | 4.78 | 2.01 | 2.60 | 3.89 | 1.68 | 5.03 | 7.56 | 3.20 |
| 1987 | 3.39 | 4.84 | 2.21 | 2.66 | 3.64 | 1.88 | 5.35 | 7.69 | 3.57 |
| 1988 | 3.38 | 5.44 | 1.81 | 2.70 | 4.24 | 1.59 | 4.84 | 8.86 | - |
| 1989 | 3.60 | 5.58 | 2.17 | 2.86 | 4.28 | 1.89 | 5.28 | 9.07 | 2.60 |
| 1990 | 3.93 | 6.03 | 2.30 | 3.05 | 4.65 | 1.86 | 6.50 | 10.61 | 3.47 |
| 1991 | 4.32 | 6.68 | 2.57 | 3.53 | 5.55 | 2.12 | 5.95 | 9.24 | 3.48 |
| 1992 | 3.99 | 6.06 | 2.39 | 3.18 | 4.81 | 1.94 | 5.63 | 8.42 | 3.43 |
| 1993 | 4.50 | 7.13 | 2.48 | 3.68 | 5.87 | 2.07 | 6.19 | 10.12 | 3.09 |
| 1994 | 4.45 | 6.98 | 2.48 | 3.59 | 5.59 | 2.08 | 5.05 | 9.52 | 1.86 |
| 1995 | 4.55 | 6.81 | 2.80 | 3.58 | 5.37 | 2.25 | 6.05 | 8.58 | 3.94 |
| 1996 | 5.41 | 8.27 | 3.14 | 4.41 | 6.75 | 2.63 | 6.66 | 10.82 | 3.55 |
| 1997 | 5.37 | 8.05 | 3.17 | 4.38 | 6.54 | 2.60 | 7.48 | 11.51 | 4.43 |
| 1998 | 5.47 | 8.31 | 3.15 | 4.55 | 6.95 | 2.58 | 6.51 | 9.64 | 4.06 |
| 1999 | 5.82 | 8.86 | 3.33 | 4.74 | 7.21 | 2.74 | 7.73 | 12.21 | 4.39 |
| 2000 | 5.61 | 8.50 | 3.22 | 4.68 | 7.02 | 2.74 | 6.92 | 11.63 | 3.44 |
| 2001 | 5.59 | 8.59 | 3.16 | 4.32 | 6.51 | 2.56 | 7.92 | 13.72 | 3.48 |
| 2002 | 5.72 | 8.69 | 3.24 | 4.59 | 6.94 | 2.57 | 7.31 | 11.97 | 3.75 |
| 2003 | 6.10 | 9.38 | 3.29 | 4.95 | 7.66 | 2.64 | 7.60 | 12.38 | 3.84 |
| 2004 | 6.30 | 9.82 | 3.29 | 5.03 | 7.79 | 2.65 | 9.26 | 15.50 | 4.47 |
| 2005 | 6.61 | 10.03 | 3.65 | 5.29 | 8.25 | 2.76 | 9.19 | 14.34 | 4.98 |
| 2006 | 6.69 | 10.19 | 3.62 | 5.42 | 8.40 | 2.80 | 8.96 | 14.15 | 4.80 |
| 1975-2006 | 4.44 | 6.84 | 2.53 | 3.57 | 5.48 | 2.09 | 6.27 | 10.05 | 3.34 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 14.5
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 2.81 | 3.85 | 2.05 | 2.62 | 3.52 | 1.99 | 4.30 | 6.34 | 2.69 |
| 1976 | 2.75 | 3.82 | 1.96 | 2.55 | 3.48 | 1.88 | 4.46 | 6.81 | 2.67 |
| 1977 | 2.78 | 3.84 | 1.99 | 2.56 | 3.44 | 1.92 | 4.52 | 7.19 | 2.49 |
| 1978 | 2.74 | 3.86 | 1.92 | 2.50 | 3.48 | 1.81 | 4.60 | 6.99 | 2.84 |
| 1979 | 2.82 | 3.88 | 2.04 | 2.62 | 3.54 | 1.95 | 4.36 | 6.64 | 2.70 |
| 1980 | 2.78 | 3.88 | 2.00 | 2.56 | 3.51 | 1.91 | 4.49 | 6.78 | 2.78 |
| 1981 | 2.87 | 4.11 | 1.99 | 2.67 | 3.76 | 1.91 | 4.40 | 6.67 | 2.73 |
| 1982 | 2.94 | 4.20 | 2.05 | 2.69 | 3.79 | 1.93 | 4.69 | 7.16 | 2.93 |
| 1983 | 2.92 | 4.20 | 2.00 | 2.63 | 3.72 | 1.87 | 4.80 | 7.27 | 3.00 |
| 1984 | 3.02 | 4.35 | 2.07 | 2.74 | 3.85 | 1.98 | 4.82 | 7.93 | 2.65 |
| 1985 | 3.11 | 4.40 | 2.19 | 2.85 | 3.94 | 2.08 | 4.57 | 7.18 | 2.74 |
| 1986 | 3.15 | 4.53 | 2.15 | 2.88 | 4.08 | 2.03 | 4.72 | 7.26 | 2.90 |
| 1987 | 3.20 | 4.56 | 2.20 | 2.92 | 4.14 | 2.04 | 4.76 | 6.90 | 3.23 |
| 1988 | 3.30 | 4.82 | 2.21 | 3.02 | 4.38 | 2.08 | 4.76 | 7.24 | 3.03 |
| 1989 | 3.44 | 5.02 | 2.29 | 3.13 | 4.51 | 2.15 | 5.10 | 8.00 | 3.09 |
| 1990 | 3.61 | 5.27 | 2.43 | 3.27 | 4.74 | 2.25 | 5.29 | 7.81 | 3.59 |
| 1991 | 3.69 | 5.32 | 2.51 | 3.35 | 4.80 | 2.32 | 5.29 | 7.97 | 3.48 |
| 1992 | 3.91 | 5.71 | 2.59 | 3.56 | 5.17 | 2.40 | 5.45 | 8.50 | 3.32 |
| 1993 | 4.04 | 5.82 | 2.74 | 3.71 | 5.32 | 2.56 | 5.49 | 8.10 | 3.59 |
| 1994 | 4.13 | 6.02 | 2.73 | 3.77 | 5.42 | 2.55 | 5.72 | 8.90 | 3.53 |
| 1995 | 4.36 | 6.31 | 2.91 | 3.97 | 5.71 | 2.69 | 6.21 | 9.30 | 4.11 |
| 1996 | 4.44 | 6.46 | 2.91 | 4.08 | 5.87 | 2.73 | 5.90 | 9.07 | 3.68 |
| 1997 | 4.50 | 6.53 | 2.97 | 4.11 | 5.91 | 2.75 | 6.02 | 8.89 | 3.92 |
| 1998 | 4.59 | 6.74 | 2.92 | 4.19 | 6.08 | 2.73 | 6.38 | 10.13 | 3.70 |
| 1999 | 4.52 | 6.50 | 2.95 | 4.15 | 5.95 | 2.73 | 5.95 | 8.76 | 3.82 |
| 2000 | 4.64 | 6.80 | 2.94 | 4.22 | 6.15 | 2.71 | 6.08 | 9.54 | 3.58 |
| 2001 | 4.71 | 6.92 | 2.97 | 4.32 | 6.35 | 2.73 | 6.25 | 9.14 | 4.08 |
| 2002 | 4.87 | 7.16 | 3.02 | 4.48 | 6.51 | 2.83 | 6.33 | 9.99 | 3.64 |
| 2003 | 5.01 | 7.38 | 3.11 | 4.57 | 6.72 | 2.84 | 6.66 | 10.32 | 4.01 |
| 2004 | 5.11 | 7.42 | 3.21 | 4.67 | 6.72 | 2.96 | 6.95 | 10.76 | 4.10 |
| 2005 | 5.26 | 7.77 | 3.19 | 4.82 | 7.08 | 2.97 | 7.03 | 11.37 | 3.78 |
| 2006 | 5.29 | 7.73 | 3.27 | 4.87 | 7.04 | 3.07 | 7.03 | 11.16 | 3.90 |
| 1975-2006 | 3.94 | 5.73 | 2.58 | 3.59 | 5.18 | 2.41 | 5.63 | 8.65 | 3.43 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 15.5
Cancer of the Lung and Bronchus (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|--------|---------|--------|--------|---------|--------|--------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 52.26 | 89.54 | 24.53 | 51.96 | 89.16 | 24.86 | 64.56 | 114.92 | 24.66 |
| 1976 | 55.42 | 93.81 | 27.31 | 54.64 | 92.77 | 27.28 | 72.34 | 131.00 | 28.63 |
| 1977 | 56.70 | 95.52 | 28.32 | 55.88 | 94.77 | 28.05 | 73.57 | 124.84 | 33.88 |
| 1978 | 57.84 | 96.67 | 29.65 | 57.27 | 95.86 | 29.90 | 74.47 | 130.51 | 31.58 |
| 1979 | 58.61 | 95.81 | 31.56 | 58.02 | 95.18 | 31.63 | 74.52 | 127.59 | 34.11 |
| 1980 | 60.65 | 99.96 | 32.21 | 59.42 | 97.71 | 32.36 | 86.57 | 151.30 | 38.25 |
| 1981 | 62.04 | 99.42 | 35.12 | 61.55 | 98.47 | 35.52 | 82.31 | 142.96 | 37.45 |
| 1982 | 63.30 | 100.71 | 36.73 | 63.06 | 99.70 | 37.68 | 80.37 | 141.06 | 35.68 |
| 1983 | 63.45 | 99.45 | 37.89 | 63.05 | 97.66 | 39.10 | 86.62 | 151.34 | 39.49 |
| 1984 | 65.50 | 102.08 | 39.45 | 64.35 | 99.94 | 39.63 | 93.06 | 159.21 | 44.82 |
| 1985 | 64.63 | 98.65 | 40.23 | 64.01 | 96.91 | 40.88 | 89.56 | 149.68 | 46.04 |
| 1986 | 65.78 | 99.08 | 42.28 | 65.18 | 97.41 | 42.94 | 92.99 | 156.37 | 48.67 |
| 1987 | 67.90 | 101.52 | 44.15 | 68.07 | 100.64 | 45.50 | 84.78 | 143.70 | 43.22 |
| 1988 | 68.01 | 98.70 | 46.26 | 68.24 | 97.94 | 47.68 | 88.12 | 144.93 | 48.06 |
| 1989 | 67.54 | 97.95 | 46.09 | 67.56 | 97.04 | 47.23 | 89.04 | 142.77 | 51.37 |
| 1990 | 68.06 | 96.95 | 47.74 | 68.21 | 96.16 | 48.91 | 86.74 | 137.17 | 52.06 |
| 1991 | 69.21 | 97.27 | 49.56 | 68.98 | 95.93 | 50.62 | 93.10 | 144.90 | 56.87 |
| 1992 | 69.43 | 97.15 | 49.84 | 68.89 | 94.82 | 51.06 | 93.92 | 148.84 | 55.86 |
| 1993 | 67.75 | 93.92 | 49.18 | 67.82 | 92.49 | 50.68 | 86.30 | 135.35 | 52.43 |
| 1994 | 67.14 | 90.92 | 50.49 | 67.23 | 90.09 | 51.57 | 87.15 | 131.58 | 56.36 |
| 1995 | 66.80 | 89.74 | 50.42 | 67.08 | 87.89 | 52.44 | 85.63 | 138.54 | 50.00 |
| 1996 | 66.38 | 87.95 | 51.12 | 66.75 | 86.68 | 52.90 | 84.20 | 128.30 | 55.15 |
| 1997 | 66.56 | 86.22 | 52.56 | 67.19 | 84.80 | 54.93 | 82.03 | 129.27 | 50.30 |
| 1998 | 67.48 | 87.95 | 52.93 | 68.02 | 86.58 | 55.07 | 86.95 | 130.36 | 57.67 |
| 1999 | 65.77 | 84.50 | 52.44 | 66.26 | 83.25 | 54.38 | 84.27 | 121.64 | 58.92 |
| 2000 | 64.01 | 81.91 | 51.14 | 64.58 | 80.63 | 53.18 | 79.37 | 113.33 | 56.59 |
| 2001 | 63.91 | 80.94 | 51.52 | 64.82 | 80.62 | 53.40 | 79.75 | 113.40 | 56.99 |
| 2002 | 63.67 | 79.87 | 52.17 | 64.42 | 79.34 | 53.91 | 80.50 | 111.25 | 60.11 |
| 2003 | 64.22 | 80.28 | 52.65 | 65.03 | 78.96 | 55.22 | 79.77 | 117.20 | 54.57 |
| 2004 | 61.44 | 75.17 | 51.45 | 62.13 | 74.41 | 53.25 | 74.01 | 101.90 | 55.53 |
| 2005 | 61.75 | 74.14 | 52.82 | 62.77 | 73.99 | 54.77 | 72.37 | 94.23 | 57.84 |
| 2006 | 59.97 | 71.28 | 51.63 | 60.45 | 70.11 | 53.46 | 73.03 | 95.71 | 57.23 |
| 1975-2006 | 64.13 | 90.32 | 45.12 | 64.16 | 89.40 | 46.09 | 82.27 | 128.76 | 49.60 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 15.6
 Small Cell Cancer of the Lung and Bronchus (Invasive)
 Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 6.65 | 10.34 | 3.79 | 6.89 | 10.70 | 3.97 | 5.87 | 8.94 | 3.48 |
| 1976 | 7.06 | 11.36 | 3.84 | 7.12 | 11.52 | 3.89 | 8.23 | 12.82 | 4.76 |
| 1977 | 7.70 | 12.38 | 4.22 | 7.84 | 12.79 | 4.21 | 8.13 | 11.19 | 5.78 |
| 1978 | 8.64 | 13.39 | 5.03 | 8.87 | 13.69 | 5.28 | 7.81 | 13.89 | 3.13 |
| 1979 | 8.68 | 13.09 | 5.36 | 9.03 | 13.63 | 5.61 | 7.63 | 11.83 | 4.47 |
| 1980 | 9.10 | 14.20 | 5.23 | 9.33 | 14.40 | 5.54 | 9.99 | 17.53 | 4.38 |
| 1981 | 9.71 | 14.55 | 6.12 | 10.01 | 14.84 | 6.48 | 9.21 | 15.87 | 4.21 |
| 1982 | 10.37 | 14.92 | 7.12 | 10.73 | 15.28 | 7.55 | 9.63 | 15.94 | 4.92 |
| 1983 | 10.36 | 14.86 | 7.08 | 10.74 | 15.04 | 7.64 | 9.12 | 15.08 | 4.85 |
| 1984 | 10.69 | 15.18 | 7.41 | 11.05 | 15.64 | 7.77 | 10.41 | 14.99 | 6.86 |
| 1985 | 10.44 | 14.96 | 7.08 | 10.78 | 15.25 | 7.49 | 11.07 | 17.77 | 6.27 |
| 1986 | 11.20 | 15.52 | 7.98 | 11.54 | 15.99 | 8.27 | 12.26 | 17.70 | 8.35 |
| 1987 | 10.65 | 14.13 | 8.19 | 11.17 | 14.56 | 8.84 | 9.53 | 14.49 | 6.03 |
| 1988 | 11.37 | 15.28 | 8.50 | 11.89 | 15.62 | 9.19 | 10.49 | 17.86 | 5.30 |
| 1989 | 11.24 | 14.84 | 8.68 | 11.68 | 15.21 | 9.21 | 12.07 | 17.67 | 8.35 |
| 1990 | 10.65 | 13.84 | 8.38 | 11.26 | 14.46 | 9.05 | 9.35 | 13.70 | 6.21 |
| 1991 | 11.28 | 14.00 | 9.35 | 11.68 | 14.34 | 9.84 | 12.34 | 15.92 | 10.00 |
| 1992 | 10.49 | 13.39 | 8.44 | 10.96 | 13.84 | 8.96 | 10.16 | 13.32 | 7.95 |
| 1993 | 10.02 | 12.33 | 8.34 | 10.48 | 12.64 | 8.94 | 9.38 | 13.25 | 6.74 |
| 1994 | 9.77 | 11.69 | 8.42 | 10.24 | 11.98 | 9.05 | 9.95 | 13.62 | 7.43 |
| 1995 | 9.85 | 12.22 | 8.14 | 10.35 | 12.58 | 8.76 | 9.43 | 13.72 | 6.61 |
| 1996 | 9.70 | 12.06 | 7.96 | 10.26 | 12.45 | 8.63 | 8.80 | 12.02 | 6.84 |
| 1997 | 9.53 | 10.85 | 8.64 | 10.11 | 11.11 | 9.49 | 7.86 | 10.82 | 6.05 |
| 1998 | 9.28 | 10.83 | 8.14 | 9.85 | 11.26 | 8.83 | 9.37 | 12.10 | 7.45 |
| 1999 | 8.80 | 10.03 | 7.96 | 9.33 | 10.25 | 8.78 | 8.14 | 10.95 | 6.05 |
| 2000 | 8.76 | 10.38 | 7.56 | 9.25 | 10.79 | 8.12 | 9.41 | 11.54 | 7.86 |
| 2001 | 8.33 | 9.77 | 7.32 | 8.86 | 10.24 | 7.91 | 8.50 | 10.78 | 6.87 |
| 2002 | 8.24 | 9.35 | 7.48 | 8.92 | 9.81 | 8.32 | 7.21 | 9.74 | 5.71 |
| 2003 | 8.17 | 9.06 | 7.54 | 8.67 | 9.32 | 8.28 | 7.85 | 9.46 | 6.72 |
| 2004 | 8.01 | 8.87 | 7.40 | 8.41 | 9.12 | 7.90 | 8.52 | 9.69 | 7.77 |
| 2005 | 7.47 | 8.52 | 6.72 | 8.08 | 9.07 | 7.39 | 6.44 | 7.51 | 5.75 |
| 2006 | 7.51 | 8.28 | 6.97 | 8.05 | 8.56 | 7.72 | 6.72 | 8.09 | 5.81 |
| 1975-2006 | 9.34 | 12.09 | 7.28 | 9.79 | 12.50 | 7.79 | 8.99 | 12.73 | 6.35 |

Note: Small Cell Cancer of the Lung and Bronchus includes histologies 8041-8045.
^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
 Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 15.7
 Non-Small Cell Cancer of the Lung and Bronchus (Invasive)
 Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|--------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 45.61 | 79.20 | 20.74 | 45.07 | 78.46 | 20.89 | 58.68 | 105.98 | 21.18 |
| 1976 | 48.36 | 82.45 | 23.47 | 47.52 | 81.25 | 23.39 | 64.11 | 118.18 | 23.87 |
| 1977 | 49.00 | 83.14 | 24.10 | 48.04 | 81.98 | 23.84 | 65.44 | 113.65 | 28.10 |
| 1978 | 49.20 | 83.28 | 24.62 | 48.40 | 82.17 | 24.62 | 66.65 | 116.63 | 28.45 |
| 1979 | 49.92 | 82.72 | 26.21 | 48.99 | 81.55 | 26.02 | 66.89 | 115.76 | 29.64 |
| 1980 | 51.56 | 85.76 | 26.98 | 50.09 | 83.31 | 26.82 | 76.58 | 133.78 | 33.87 |
| 1981 | 52.33 | 84.88 | 29.00 | 51.54 | 83.63 | 29.05 | 73.10 | 127.09 | 33.24 |
| 1982 | 52.93 | 85.79 | 29.62 | 52.33 | 84.42 | 30.13 | 70.74 | 125.12 | 30.76 |
| 1983 | 53.09 | 84.59 | 30.81 | 52.30 | 82.62 | 31.46 | 77.50 | 136.26 | 34.64 |
| 1984 | 54.81 | 86.90 | 32.05 | 53.31 | 84.29 | 31.85 | 82.64 | 144.23 | 37.96 |
| 1985 | 54.18 | 83.69 | 33.15 | 53.22 | 81.66 | 33.39 | 78.49 | 131.91 | 39.78 |
| 1986 | 54.58 | 83.56 | 34.30 | 53.64 | 81.42 | 34.67 | 80.73 | 138.68 | 40.31 |
| 1987 | 57.25 | 87.39 | 35.96 | 56.91 | 86.08 | 36.65 | 75.25 | 129.21 | 37.19 |
| 1988 | 56.65 | 83.42 | 37.76 | 56.35 | 82.32 | 38.49 | 77.63 | 127.07 | 42.76 |
| 1989 | 56.30 | 83.11 | 37.41 | 55.88 | 81.83 | 38.02 | 76.97 | 125.10 | 43.02 |
| 1990 | 57.40 | 83.11 | 39.35 | 56.95 | 81.70 | 39.85 | 77.38 | 123.46 | 45.84 |
| 1991 | 57.93 | 83.27 | 40.22 | 57.30 | 81.59 | 40.79 | 80.76 | 128.98 | 46.87 |
| 1992 | 58.93 | 83.75 | 41.40 | 57.92 | 80.98 | 42.09 | 83.76 | 135.52 | 47.91 |
| 1993 | 57.73 | 81.59 | 40.84 | 57.34 | 79.85 | 41.75 | 76.92 | 122.10 | 45.69 |
| 1994 | 57.37 | 79.23 | 42.07 | 56.99 | 78.12 | 42.51 | 77.20 | 117.96 | 48.93 |
| 1995 | 56.95 | 77.51 | 42.28 | 56.72 | 75.31 | 43.68 | 76.20 | 124.83 | 43.39 |
| 1996 | 56.68 | 75.89 | 43.16 | 56.49 | 74.23 | 44.26 | 75.40 | 116.28 | 48.31 |
| 1997 | 57.04 | 75.37 | 43.92 | 57.09 | 73.69 | 45.44 | 74.17 | 118.45 | 44.25 |
| 1998 | 58.21 | 77.12 | 44.79 | 58.17 | 75.32 | 46.23 | 77.58 | 118.26 | 50.22 |
| 1999 | 56.97 | 74.47 | 44.48 | 56.93 | 73.00 | 45.60 | 76.13 | 110.70 | 52.87 |
| 2000 | 55.26 | 71.52 | 43.57 | 55.33 | 69.83 | 45.06 | 69.96 | 101.79 | 48.72 |
| 2001 | 55.58 | 71.17 | 44.21 | 55.97 | 70.38 | 45.50 | 71.24 | 102.62 | 50.12 |
| 2002 | 55.43 | 70.52 | 44.69 | 55.50 | 69.54 | 45.59 | 73.29 | 101.50 | 54.41 |
| 2003 | 56.05 | 71.22 | 45.11 | 56.35 | 69.63 | 46.94 | 71.92 | 107.74 | 47.84 |
| 2004 | 53.43 | 66.30 | 44.06 | 53.72 | 65.29 | 45.36 | 65.50 | 92.21 | 47.76 |
| 2005 | 54.28 | 65.62 | 46.09 | 54.69 | 64.92 | 47.37 | 65.93 | 86.71 | 52.09 |
| 2006 | 52.46 | 63.00 | 44.66 | 52.40 | 61.55 | 45.74 | 66.31 | 87.62 | 51.42 |
| 1975-2006 | 54.79 | 78.23 | 37.83 | 54.37 | 76.90 | 38.31 | 73.28 | 116.04 | 43.24 |

Note: Non-Small Cell Cancer of the Lung and Bronchus includes histologies 8000-8040, 8046-9049, 9056-9139, 9141-9589.
^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
 Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 15.8
Cancer of the Lung and Bronchus (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|--------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 42.56 | 76.39 | 17.56 | 42.08 | 75.47 | 17.67 | 49.33 | 91.01 | 17.34 |
| 1976 | 44.20 | 78.52 | 18.97 | 43.72 | 77.55 | 19.10 | 51.07 | 93.96 | 18.51 |
| 1977 | 45.49 | 80.45 | 20.01 | 44.82 | 79.09 | 20.09 | 54.56 | 100.80 | 19.91 |
| 1978 | 46.88 | 82.06 | 21.40 | 46.29 | 80.76 | 21.58 | 55.26 | 101.65 | 20.66 |
| 1979 | 47.69 | 82.94 | 22.32 | 47.08 | 81.64 | 22.42 | 56.43 | 102.89 | 22.24 |
| 1980 | 49.41 | 84.71 | 24.13 | 48.75 | 83.27 | 24.22 | 59.16 | 106.67 | 24.57 |
| 1981 | 49.99 | 85.16 | 24.95 | 49.28 | 83.52 | 25.10 | 60.57 | 109.71 | 25.10 |
| 1982 | 51.43 | 86.79 | 26.47 | 50.78 | 85.18 | 26.72 | 62.27 | 112.55 | 26.30 |
| 1983 | 52.40 | 87.04 | 28.10 | 51.70 | 85.40 | 28.28 | 63.68 | 112.76 | 28.73 |
| 1984 | 53.36 | 88.22 | 28.94 | 52.64 | 86.24 | 29.27 | 65.07 | 117.98 | 27.98 |
| 1985 | 54.32 | 88.54 | 30.41 | 53.68 | 86.69 | 30.81 | 65.76 | 117.50 | 29.64 |
| 1986 | 55.04 | 88.95 | 31.41 | 54.36 | 87.03 | 31.76 | 66.95 | 119.18 | 31.00 |
| 1987 | 56.24 | 90.05 | 32.72 | 55.56 | 88.09 | 33.04 | 68.65 | 121.77 | 32.48 |
| 1988 | 56.97 | 89.87 | 34.14 | 56.35 | 87.86 | 34.57 | 69.18 | 122.15 | 33.54 |
| 1989 | 57.90 | 89.77 | 35.77 | 57.21 | 87.57 | 36.27 | 71.04 | 124.72 | 34.87 |
| 1990 | 58.85 | 90.56 | 36.85 | 58.18 | 88.40 | 37.31 | 72.06 | 125.17 | 36.48 |
| 1991 | 58.99 | 89.88 | 37.61 | 58.38 | 87.76 | 38.12 | 71.64 | 123.58 | 37.16 |
| 1992 | 58.90 | 88.01 | 38.70 | 58.34 | 85.98 | 39.21 | 71.35 | 121.06 | 38.55 |
| 1993 | 59.13 | 87.60 | 39.28 | 58.66 | 85.52 | 39.97 | 71.12 | 121.98 | 37.72 |
| 1994 | 58.54 | 85.69 | 39.62 | 58.15 | 83.78 | 40.30 | 70.15 | 118.63 | 38.61 |
| 1995 | 58.38 | 84.37 | 40.26 | 58.08 | 82.62 | 40.97 | 69.28 | 116.08 | 38.84 |
| 1996 | 57.91 | 82.82 | 40.41 | 57.69 | 81.15 | 41.22 | 68.30 | 113.18 | 39.08 |
| 1997 | 57.51 | 81.33 | 40.82 | 57.31 | 79.83 | 41.52 | 67.89 | 110.21 | 40.53 |
| 1998 | 57.08 | 79.89 | 40.98 | 56.93 | 78.47 | 41.69 | 66.86 | 107.40 | 40.67 |
| 1999 | 55.42 | 76.97 | 40.15 | 55.31 | 75.67 | 40.86 | 65.00 | 103.61 | 40.00 |
| 2000 | 55.84 | 76.45 | 41.12 | 55.99 | 75.39 | 42.08 | 63.81 | 100.91 | 39.66 |
| 2001 | 55.30 | 75.22 | 41.00 | 55.53 | 74.25 | 42.03 | 62.65 | 99.35 | 38.84 |
| 2002 | 54.97 | 73.55 | 41.56 | 55.28 | 72.78 | 42.59 | 62.24 | 95.70 | 40.26 |
| 2003 | 54.16 | 71.92 | 41.24 | 54.48 | 71.23 | 42.23 | 61.16 | 93.08 | 40.31 |
| 2004 | 53.34 | 70.37 | 40.89 | 53.67 | 69.69 | 41.90 | 60.09 | 90.67 | 40.01 |
| 2005 | 52.83 | 69.47 | 40.62 | 53.28 | 69.09 | 41.59 | 58.88 | 87.32 | 40.21 |
| 2006 | 51.74 | 67.45 | 40.17 | 52.26 | 67.13 | 41.25 | 57.21 | 84.57 | 39.18 |
| 1975-2006 | 54.30 | 81.74 | 34.65 | 54.01 | 80.41 | 35.13 | 63.80 | 107.17 | 34.22 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 16.4
Melanoma of the Skin (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 7.89 | 8.54 | 7.44 | 8.70 | 9.41 | 8.21 | - | - | - |
| 1976 | 8.16 | 8.68 | 7.69 | 9.00 | 9.55 | 8.51 | 1.75 | - | - |
| 1977 | 8.86 | 9.77 | 8.15 | 9.87 | 10.83 | 9.17 | - | - | - |
| 1978 | 8.95 | 9.49 | 8.72 | 10.10 | 10.72 | 9.87 | - | - | - |
| 1979 | 9.54 | 10.56 | 8.90 | 10.76 | 11.88 | 10.09 | - | - | - |
| 1980 | 10.50 | 11.77 | 9.63 | 12.01 | 13.37 | 11.12 | - | - | - |
| 1981 | 11.09 | 12.10 | 10.44 | 12.57 | 13.56 | 11.99 | 1.35 | - | - |
| 1982 | 11.17 | 12.49 | 10.27 | 12.77 | 14.15 | 11.85 | - | - | - |
| 1983 | 11.08 | 12.65 | 10.01 | 12.60 | 14.38 | 11.42 | - | - | - |
| 1984 | 11.39 | 12.91 | 10.42 | 13.05 | 14.79 | 11.95 | - | - | - |
| 1985 | 12.75 | 15.12 | 11.14 | 14.44 | 17.12 | 12.67 | - | - | - |
| 1986 | 13.29 | 15.67 | 11.59 | 15.15 | 17.82 | 13.28 | - | - | - |
| 1987 | 13.66 | 15.92 | 12.16 | 15.60 | 18.09 | 13.96 | 1.30 | - | - |
| 1988 | 12.86 | 14.97 | 11.43 | 14.90 | 17.39 | 13.23 | 1.11 | - | - |
| 1989 | 13.73 | 16.41 | 11.89 | 15.90 | 18.93 | 13.83 | 1.19 | - | - |
| 1990 | 13.83 | 16.63 | 11.84 | 16.13 | 19.25 | 13.92 | 1.16 | - | - |
| 1991 | 14.60 | 17.58 | 12.51 | 17.05 | 20.42 | 14.72 | 1.17 | - | - |
| 1992 | 14.78 | 18.30 | 12.24 | 17.19 | 21.17 | 14.37 | - | - | - |
| 1993 | 14.58 | 18.08 | 12.08 | 16.99 | 20.91 | 14.19 | - | - | - |
| 1994 | 15.61 | 19.76 | 12.57 | 18.31 | 23.09 | 14.85 | 1.11 | - | - |
| 1995 | 16.40 | 20.23 | 13.75 | 19.23 | 23.41 | 16.39 | 1.65 | 2.97 | - |
| 1996 | 17.28 | 21.64 | 14.13 | 20.40 | 25.26 | 16.92 | 1.06 | - | - |
| 1997 | 17.72 | 21.69 | 14.85 | 20.87 | 25.17 | 17.83 | 1.10 | - | - |
| 1998 | 17.87 | 22.11 | 14.92 | 21.34 | 26.11 | 18.05 | 1.19 | - | - |
| 1999 | 18.22 | 22.83 | 15.00 | 21.77 | 26.86 | 18.25 | 1.14 | - | - |
| 2000 | 18.81 | 23.73 | 15.38 | 22.66 | 28.14 | 18.88 | 0.80 | - | - |
| 2001 | 19.55 | 24.33 | 16.21 | 23.72 | 29.04 | 20.05 | 0.81 | - | - |
| 2002 | 19.06 | 23.67 | 15.90 | 23.03 | 28.06 | 19.63 | 1.04 | - | - |
| 2003 | 19.28 | 23.77 | 16.13 | 23.34 | 28.22 | 19.97 | 0.83 | - | - |
| 2004 | 20.21 | 24.89 | 16.98 | 24.60 | 29.95 | 20.92 | 0.77 | - | - |
| 2005 | 21.74 | 27.10 | 18.00 | 26.64 | 32.64 | 22.50 | 0.99 | - | 1.00 |
| 2006 | 21.14 | 26.14 | 17.63 | 25.63 | 31.16 | 21.82 | 0.89 | - | - |
| 1975-2006 | 15.17 | 18.47 | 12.93 | 17.70 | 21.38 | 15.24 | 0.99 | 1.18 | 0.87 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 16.5
Melanoma of the Skin (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 2.07 | 2.64 | 1.60 | 2.25 | 2.88 | 1.74 | 0.50 | 0.56 | 0.44 |
| 1976 | 2.24 | 2.86 | 1.76 | 2.45 | 3.10 | 1.94 | 0.55 | 0.74 | 0.39 |
| 1977 | 2.27 | 2.91 | 1.76 | 2.48 | 3.17 | 1.93 | 0.53 | 0.69 | 0.41 |
| 1978 | 2.31 | 3.04 | 1.74 | 2.53 | 3.32 | 1.90 | 0.52 | 0.61 | 0.46 |
| 1979 | 2.42 | 3.16 | 1.84 | 2.66 | 3.46 | 2.02 | 0.53 | 0.64 | 0.47 |
| 1980 | 2.34 | 3.10 | 1.74 | 2.57 | 3.41 | 1.90 | 0.54 | 0.55 | 0.56 |
| 1981 | 2.43 | 3.13 | 1.88 | 2.68 | 3.45 | 2.07 | 0.58 | 0.56 | 0.57 |
| 1982 | 2.46 | 3.31 | 1.81 | 2.71 | 3.64 | 1.99 | 0.52 | 0.56 | 0.49 |
| 1983 | 2.48 | 3.29 | 1.85 | 2.74 | 3.63 | 2.04 | 0.50 | 0.63 | 0.40 |
| 1984 | 2.53 | 3.42 | 1.85 | 2.80 | 3.79 | 2.03 | 0.51 | 0.45 | 0.54 |
| 1985 | 2.56 | 3.46 | 1.86 | 2.83 | 3.82 | 2.06 | 0.53 | 0.58 | 0.49 |
| 1986 | 2.59 | 3.55 | 1.88 | 2.88 | 3.92 | 2.09 | 0.47 | 0.70 | 0.31 |
| 1987 | 2.65 | 3.73 | 1.83 | 2.93 | 4.13 | 2.02 | 0.55 | 0.57 | 0.54 |
| 1988 | 2.65 | 3.67 | 1.90 | 2.95 | 4.07 | 2.13 | 0.45 | 0.53 | 0.40 |
| 1989 | 2.69 | 3.74 | 1.89 | 2.99 | 4.15 | 2.10 | 0.55 | 0.57 | 0.53 |
| 1990 | 2.75 | 3.80 | 1.97 | 3.06 | 4.21 | 2.20 | 0.52 | 0.65 | 0.45 |
| 1991 | 2.71 | 3.89 | 1.82 | 3.01 | 4.31 | 2.02 | 0.59 | 0.65 | 0.54 |
| 1992 | 2.71 | 3.85 | 1.85 | 3.01 | 4.27 | 2.06 | 0.51 | 0.57 | 0.44 |
| 1993 | 2.71 | 3.86 | 1.86 | 3.03 | 4.31 | 2.06 | 0.53 | 0.42 | 0.59 |
| 1994 | 2.66 | 3.82 | 1.81 | 2.98 | 4.26 | 2.03 | 0.45 | 0.44 | 0.43 |
| 1995 | 2.70 | 3.89 | 1.81 | 3.03 | 4.35 | 2.02 | 0.46 | 0.51 | 0.42 |
| 1996 | 2.80 | 4.04 | 1.90 | 3.14 | 4.50 | 2.14 | 0.44 | 0.51 | 0.38 |
| 1997 | 2.73 | 3.94 | 1.84 | 3.06 | 4.41 | 2.05 | 0.56 | 0.49 | 0.58 |
| 1998 | 2.75 | 4.09 | 1.77 | 3.10 | 4.59 | 1.98 | 0.51 | 0.45 | 0.54 |
| 1999 | 2.63 | 3.81 | 1.74 | 2.97 | 4.27 | 1.98 | 0.40 | 0.52 | 0.33 |
| 2000 | 2.66 | 3.82 | 1.81 | 3.00 | 4.28 | 2.03 | 0.54 | 0.59 | 0.50 |
| 2001 | 2.66 | 3.88 | 1.75 | 3.02 | 4.37 | 1.99 | 0.42 | 0.35 | 0.45 |
| 2002 | 2.60 | 3.77 | 1.72 | 2.95 | 4.24 | 1.96 | 0.39 | 0.49 | 0.32 |
| 2003 | 2.66 | 3.88 | 1.74 | 3.02 | 4.38 | 1.97 | 0.44 | 0.43 | 0.44 |
| 2004 | 2.66 | 3.94 | 1.70 | 3.03 | 4.46 | 1.94 | 0.44 | 0.55 | 0.38 |
| 2005 | 2.74 | 4.00 | 1.81 | 3.12 | 4.52 | 2.06 | 0.44 | 0.48 | 0.40 |
| 2006 | 2.72 | 4.07 | 1.72 | 3.10 | 4.60 | 1.97 | 0.44 | 0.60 | 0.35 |
| 1975-2006 | 2.60 | 3.68 | 1.80 | 2.91 | 4.09 | 2.02 | 0.49 | 0.54 | 0.45 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 17.2
MesotheliomaAge-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 0.63 | 1.00 | 0.31 | 0.63 | 1.00 | 0.33 | - | - | - |
| 1976 | 0.67 | 1.13 | 0.35 | 0.71 | 1.21 | 0.36 | - | - | - |
| 1977 | 0.62 | 1.05 | 0.31 | 0.65 | 1.06 | 0.34 | - | - | - |
| 1978 | 0.73 | 1.37 | 0.25 | 0.79 | 1.50 | 0.25 | - | - | - |
| 1979 | 0.78 | 1.51 | 0.27 | 0.80 | 1.60 | 0.25 | - | - | - |
| 1980 | 0.98 | 1.83 | 0.35 | 1.05 | 1.97 | 0.39 | - | - | - |
| 1981 | 0.75 | 1.42 | 0.27 | 0.80 | 1.50 | 0.29 | - | - | - |
| 1982 | 0.83 | 1.42 | 0.38 | 0.90 | 1.53 | 0.42 | - | - | - |
| 1983 | 1.05 | 1.86 | 0.49 | 1.09 | 1.95 | 0.51 | - | - | - |
| 1984 | 1.06 | 2.04 | 0.36 | 1.15 | 2.24 | 0.37 | - | - | - |
| 1985 | 0.92 | 1.67 | 0.37 | 0.97 | 1.73 | 0.39 | - | - | - |
| 1986 | 0.94 | 1.67 | 0.43 | 1.01 | 1.79 | 0.49 | - | - | - |
| 1987 | 0.97 | 1.73 | 0.43 | 1.03 | 1.82 | 0.47 | - | - | - |
| 1988 | 0.98 | 1.85 | 0.39 | 1.05 | 1.98 | 0.41 | - | - | - |
| 1989 | 1.13 | 2.20 | 0.37 | 1.20 | 2.35 | 0.39 | - | - | - |
| 1990 | 1.15 | 2.15 | 0.46 | 1.25 | 2.32 | 0.50 | - | - | - |
| 1991 | 1.11 | 2.10 | 0.42 | 1.18 | 2.21 | 0.46 | - | - | - |
| 1992 | 1.23 | 2.48 | 0.39 | 1.33 | 2.71 | 0.42 | - | - | - |
| 1993 | 1.06 | 2.04 | 0.38 | 1.14 | 2.24 | 0.37 | - | - | - |
| 1994 | 1.16 | 2.17 | 0.43 | 1.24 | 2.32 | 0.46 | 1.06 | - | - |
| 1995 | 1.16 | 2.17 | 0.46 | 1.23 | 2.30 | 0.48 | - | - | - |
| 1996 | 1.15 | 2.14 | 0.45 | 1.27 | 2.38 | 0.50 | - | - | - |
| 1997 | 1.01 | 1.92 | 0.37 | 1.06 | 2.06 | 0.35 | 0.92 | - | - |
| 1998 | 1.08 | 2.06 | 0.40 | 1.18 | 2.23 | 0.47 | - | - | - |
| 1999 | 1.12 | 2.17 | 0.37 | 1.20 | 2.30 | 0.40 | - | - | - |
| 2000 | 1.08 | 2.11 | 0.34 | 1.22 | 2.37 | 0.38 | - | - | - |
| 2001 | 1.06 | 1.97 | 0.41 | 1.17 | 2.17 | 0.44 | - | - | - |
| 2002 | 1.02 | 1.93 | 0.38 | 1.15 | 2.16 | 0.43 | - | - | - |
| 2003 | 0.95 | 1.71 | 0.39 | 1.03 | 1.86 | 0.43 | - | - | - |
| 2004 | 1.10 | 2.03 | 0.44 | 1.22 | 2.26 | 0.47 | - | - | - |
| 2005 | 1.02 | 1.87 | 0.42 | 1.17 | 2.13 | 0.49 | - | - | - |
| 2006 | 0.93 | 1.57 | 0.44 | 1.02 | 1.73 | 0.47 | - | - | - |
| 1975-2006 | 1.00 | 1.87 | 0.39 | 1.08 | 2.02 | 0.42 | 0.57 | 1.03 | 0.26 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
 Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 18.4
MyelomaAge-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 4.91 | 6.43 | 3.86 | 4.61 | 6.17 | 3.57 | 9.72 | 10.86 | 8.61 |
| 1976 | 5.02 | 6.08 | 4.37 | 4.76 | 5.77 | 4.14 | 9.20 | 11.22 | 7.90 |
| 1977 | 5.06 | 6.38 | 4.16 | 4.85 | 6.19 | 3.95 | 9.09 | 11.00 | 7.75 |
| 1978 | 4.74 | 5.93 | 3.94 | 4.43 | 5.60 | 3.65 | 9.61 | 10.97 | 8.66 |
| 1979 | 4.88 | 6.00 | 4.09 | 4.50 | 5.41 | 3.83 | 10.99 | 15.26 | 8.23 |
| 1980 | 4.79 | 5.77 | 4.12 | 4.43 | 5.44 | 3.75 | 9.89 | 10.83 | 9.05 |
| 1981 | 5.02 | 6.00 | 4.33 | 4.62 | 5.46 | 4.00 | 11.01 | 15.23 | 8.19 |
| 1982 | 5.29 | 6.28 | 4.67 | 4.89 | 5.87 | 4.32 | 11.57 | 13.93 | 9.72 |
| 1983 | 5.31 | 6.93 | 4.22 | 4.74 | 6.30 | 3.73 | 13.72 | 16.22 | 11.92 |
| 1984 | 5.41 | 6.98 | 4.33 | 5.05 | 6.44 | 4.11 | 10.84 | 15.35 | 7.71 |
| 1985 | 5.20 | 6.32 | 4.35 | 4.94 | 5.95 | 4.19 | 9.38 | 12.58 | 7.13 |
| 1986 | 5.33 | 6.31 | 4.65 | 4.89 | 6.02 | 4.13 | 11.01 | 11.57 | 10.42 |
| 1987 | 5.93 | 7.37 | 5.02 | 5.66 | 7.18 | 4.71 | 11.13 | 13.48 | 9.69 |
| 1988 | 5.33 | 6.57 | 4.51 | 5.03 | 6.37 | 4.18 | 10.81 | 12.17 | 9.72 |
| 1989 | 5.44 | 6.73 | 4.49 | 5.10 | 6.35 | 4.17 | 10.74 | 13.46 | 8.96 |
| 1990 | 5.65 | 7.25 | 4.56 | 5.25 | 6.93 | 4.10 | 11.96 | 14.74 | 10.18 |
| 1991 | 6.03 | 7.67 | 4.85 | 5.57 | 7.18 | 4.41 | 12.78 | 15.92 | 10.64 |
| 1992 | 5.96 | 7.28 | 5.07 | 5.63 | 6.99 | 4.76 | 11.51 | 13.04 | 10.40 |
| 1993 | 5.60 | 7.33 | 4.34 | 5.15 | 6.82 | 3.97 | 12.74 | 15.72 | 10.50 |
| 1994 | 5.72 | 6.80 | 5.03 | 5.20 | 6.34 | 4.44 | 12.76 | 15.16 | 11.82 |
| 1995 | 5.73 | 7.23 | 4.69 | 5.32 | 6.79 | 4.32 | 12.18 | 15.13 | 10.30 |
| 1996 | 5.81 | 7.18 | 4.77 | 5.42 | 6.87 | 4.29 | 12.54 | 13.39 | 11.87 |
| 1997 | 6.11 | 7.36 | 5.24 | 5.68 | 7.02 | 4.74 | 12.71 | 14.42 | 11.94 |
| 1998 | 5.90 | 7.29 | 4.87 | 5.56 | 7.08 | 4.41 | 11.73 | 13.20 | 10.82 |
| 1999 | 5.58 | 7.06 | 4.49 | 5.12 | 6.56 | 4.06 | 12.10 | 15.35 | 10.06 |
| 2000 | 6.02 | 7.49 | 4.92 | 5.72 | 7.49 | 4.40 | 10.81 | 10.47 | 10.81 |
| 2001 | 5.85 | 7.22 | 4.82 | 5.44 | 6.81 | 4.38 | 12.05 | 14.58 | 10.53 |
| 2002 | 5.91 | 7.67 | 4.59 | 5.58 | 7.42 | 4.20 | 10.78 | 13.87 | 8.77 |
| 2003 | 5.70 | 7.18 | 4.58 | 5.44 | 6.95 | 4.27 | 10.49 | 13.68 | 8.36 |
| 2004 | 5.66 | 7.10 | 4.59 | 5.15 | 6.61 | 4.08 | 12.17 | 14.96 | 10.20 |
| 2005 | 5.79 | 7.32 | 4.65 | 5.36 | 6.96 | 4.14 | 11.90 | 13.89 | 10.60 |
| 2006 | 5.37 | 6.48 | 4.54 | 5.04 | 6.25 | 4.09 | 11.42 | 12.98 | 10.43 |
| 1975-2006 | 5.55 | 6.90 | 4.59 | 5.17 | 6.54 | 4.19 | 11.40 | 13.68 | 9.88 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 18.5

Myeloma

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 2.94 | 3.70 | 2.42 | 2.73 | 3.47 | 2.23 | 5.26 | 6.44 | 4.41 |
| 1976 | 3.10 | 3.89 | 2.57 | 2.89 | 3.64 | 2.38 | 5.56 | 6.92 | 4.58 |
| 1977 | 3.13 | 3.92 | 2.60 | 2.91 | 3.64 | 2.43 | 5.55 | 7.24 | 4.42 |
| 1978 | 3.17 | 3.94 | 2.66 | 2.95 | 3.69 | 2.45 | 5.74 | 6.97 | 4.93 |
| 1979 | 3.26 | 4.12 | 2.68 | 3.03 | 3.85 | 2.48 | 5.91 | 7.42 | 4.89 |
| 1980 | 3.27 | 4.12 | 2.72 | 3.02 | 3.83 | 2.49 | 6.10 | 7.37 | 5.23 |
| 1981 | 3.29 | 4.05 | 2.79 | 3.01 | 3.70 | 2.54 | 6.57 | 8.16 | 5.50 |
| 1982 | 3.38 | 4.17 | 2.85 | 3.10 | 3.81 | 2.62 | 6.68 | 8.56 | 5.45 |
| 1983 | 3.44 | 4.34 | 2.87 | 3.15 | 3.97 | 2.63 | 6.82 | 8.73 | 5.58 |
| 1984 | 3.47 | 4.26 | 2.95 | 3.20 | 3.92 | 2.73 | 6.64 | 8.38 | 5.47 |
| 1985 | 3.58 | 4.46 | 3.02 | 3.29 | 4.11 | 2.78 | 6.90 | 8.57 | 5.82 |
| 1986 | 3.61 | 4.51 | 3.03 | 3.29 | 4.12 | 2.76 | 7.16 | 9.10 | 5.95 |
| 1987 | 3.61 | 4.57 | 2.97 | 3.32 | 4.21 | 2.72 | 7.06 | 9.18 | 5.71 |
| 1988 | 3.63 | 4.61 | 3.02 | 3.36 | 4.35 | 2.73 | 6.89 | 8.14 | 6.12 |
| 1989 | 3.70 | 4.64 | 3.10 | 3.42 | 4.30 | 2.85 | 7.08 | 8.91 | 5.92 |
| 1990 | 3.80 | 4.83 | 3.13 | 3.49 | 4.45 | 2.87 | 7.32 | 9.64 | 5.92 |
| 1991 | 3.87 | 4.83 | 3.26 | 3.54 | 4.46 | 2.95 | 7.77 | 9.59 | 6.64 |
| 1992 | 3.80 | 4.72 | 3.19 | 3.48 | 4.36 | 2.90 | 7.45 | 9.33 | 6.32 |
| 1993 | 3.96 | 4.89 | 3.35 | 3.64 | 4.56 | 3.04 | 7.82 | 9.32 | 6.88 |
| 1994 | 3.97 | 5.03 | 3.26 | 3.67 | 4.69 | 2.99 | 7.68 | 9.68 | 6.46 |
| 1995 | 3.98 | 4.94 | 3.36 | 3.66 | 4.61 | 3.02 | 7.79 | 9.41 | 6.90 |
| 1996 | 3.91 | 4.94 | 3.25 | 3.59 | 4.60 | 2.94 | 7.70 | 9.63 | 6.54 |
| 1997 | 3.90 | 4.93 | 3.22 | 3.60 | 4.61 | 2.93 | 7.49 | 9.24 | 6.45 |
| 1998 | 3.83 | 4.76 | 3.21 | 3.50 | 4.41 | 2.88 | 7.75 | 9.49 | 6.77 |
| 1999 | 3.84 | 4.72 | 3.24 | 3.52 | 4.41 | 2.91 | 7.45 | 8.84 | 6.64 |
| 2000 | 3.83 | 4.66 | 3.27 | 3.53 | 4.38 | 2.96 | 7.51 | 8.75 | 6.73 |
| 2001 | 3.80 | 4.68 | 3.18 | 3.52 | 4.40 | 2.89 | 7.22 | 8.71 | 6.28 |
| 2002 | 3.80 | 4.76 | 3.14 | 3.56 | 4.54 | 2.86 | 6.88 | 8.27 | 6.02 |
| 2003 | 3.70 | 4.58 | 3.08 | 3.43 | 4.33 | 2.78 | 7.11 | 8.36 | 6.29 |
| 2004 | 3.56 | 4.40 | 2.97 | 3.30 | 4.14 | 2.69 | 6.85 | 8.22 | 6.02 |
| 2005 | 3.56 | 4.49 | 2.89 | 3.32 | 4.24 | 2.65 | 6.47 | 8.01 | 5.48 |
| 2006 | 3.48 | 4.36 | 2.85 | 3.24 | 4.09 | 2.61 | 6.53 | 8.23 | 5.44 |
| 1975-2006 | 3.64 | 4.54 | 3.03 | 3.36 | 4.23 | 2.77 | 6.97 | 8.57 | 5.96 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 19.4
Non-Hodgkin Lymphoma

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 11.06 | 12.89 | 9.58 | 11.61 | 13.49 | 10.11 | 6.36 | 7.31 | 5.31 |
| 1976 | 11.20 | 12.82 | 10.04 | 11.65 | 13.27 | 10.50 | 5.95 | 7.45 | 4.87 |
| 1977 | 11.18 | 12.92 | 9.82 | 11.64 | 13.50 | 10.20 | 6.64 | 7.23 | 6.10 |
| 1978 | 11.93 | 14.30 | 10.12 | 12.49 | 14.79 | 10.74 | 7.15 | 9.46 | 5.17 |
| 1979 | 12.51 | 14.19 | 11.20 | 12.98 | 14.58 | 11.75 | 8.39 | 11.81 | 5.91 |
| 1980 | 12.61 | 15.01 | 10.78 | 12.95 | 15.26 | 11.19 | 8.77 | 10.89 | 7.20 |
| 1981 | 13.58 | 15.90 | 11.81 | 14.05 | 16.33 | 12.35 | 7.39 | 10.52 | 4.98 |
| 1982 | 13.39 | 15.55 | 11.61 | 14.01 | 16.39 | 12.05 | 10.27 | 10.22 | 10.01 |
| 1983 | 13.99 | 16.61 | 11.96 | 14.64 | 17.52 | 12.42 | 10.28 | 10.88 | 9.62 |
| 1984 | 15.19 | 18.32 | 12.77 | 15.95 | 19.11 | 13.51 | 9.46 | 12.52 | 6.92 |
| 1985 | 15.49 | 18.36 | 13.17 | 16.26 | 19.22 | 13.84 | 9.89 | 11.96 | 8.32 |
| 1986 | 15.91 | 19.42 | 13.02 | 16.82 | 20.53 | 13.77 | 10.63 | 13.13 | 8.46 |
| 1987 | 16.74 | 20.79 | 13.43 | 17.71 | 22.14 | 14.02 | 10.43 | 10.43 | 10.09 |
| 1988 | 17.24 | 21.32 | 14.08 | 18.20 | 22.40 | 14.90 | 11.57 | 15.79 | 8.47 |
| 1989 | 17.33 | 21.62 | 13.82 | 18.33 | 22.99 | 14.48 | 11.67 | 14.17 | 9.82 |
| 1990 | 18.52 | 22.99 | 14.81 | 19.36 | 23.96 | 15.54 | 13.78 | 17.51 | 10.86 |
| 1991 | 18.79 | 23.74 | 14.57 | 19.67 | 24.81 | 15.26 | 14.03 | 18.72 | 10.37 |
| 1992 | 18.62 | 23.03 | 14.95 | 19.57 | 23.95 | 15.86 | 13.61 | 18.73 | 9.63 |
| 1993 | 18.85 | 23.45 | 15.00 | 19.64 | 24.30 | 15.70 | 14.04 | 19.37 | 9.67 |
| 1994 | 19.92 | 24.96 | 15.74 | 20.64 | 25.56 | 16.50 | 14.35 | 21.50 | 8.65 |
| 1995 | 19.90 | 25.33 | 15.35 | 20.61 | 26.09 | 15.95 | 16.49 | 23.26 | 11.25 |
| 1996 | 19.36 | 24.20 | 15.40 | 20.23 | 25.20 | 16.13 | 15.16 | 18.92 | 11.97 |
| 1997 | 19.91 | 24.40 | 16.41 | 20.80 | 25.11 | 17.41 | 16.23 | 23.93 | 10.84 |
| 1998 | 19.51 | 22.86 | 16.72 | 20.50 | 23.94 | 17.62 | 14.05 | 16.38 | 12.19 |
| 1999 | 19.82 | 24.47 | 16.16 | 20.78 | 25.38 | 17.13 | 14.10 | 18.21 | 10.97 |
| 2000 | 19.61 | 24.06 | 16.00 | 20.73 | 25.35 | 16.96 | 13.91 | 18.36 | 10.63 |
| 2001 | 19.87 | 24.25 | 16.31 | 20.65 | 25.26 | 16.88 | 15.65 | 18.06 | 13.55 |
| 2002 | 19.93 | 23.87 | 16.81 | 20.99 | 25.11 | 17.67 | 14.24 | 17.09 | 12.07 |
| 2003 | 20.43 | 24.49 | 17.32 | 21.51 | 25.73 | 18.22 | 15.82 | 19.22 | 13.36 |
| 2004 | 20.90 | 25.29 | 17.50 | 21.80 | 26.21 | 18.35 | 17.78 | 23.12 | 13.86 |
| 2005 | 20.20 | 24.79 | 16.54 | 21.32 | 25.92 | 17.59 | 15.94 | 18.86 | 13.51 |
| 2006 | 19.52 | 23.33 | 16.56 | 20.57 | 24.38 | 17.58 | 15.54 | 19.83 | 12.38 |
| 1975-2006 | 17.47 | 21.34 | 14.40 | 18.20 | 22.16 | 15.04 | 12.99 | 16.51 | 10.22 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 19.5
Non-Hodgkin Lymphoma

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 5.63 | 6.92 | 4.65 | 5.83 | 7.15 | 4.82 | 3.74 | 4.83 | 2.92 |
| 1976 | 5.71 | 6.87 | 4.83 | 5.94 | 7.14 | 5.03 | 3.54 | 4.54 | 2.76 |
| 1977 | 5.77 | 6.88 | 4.92 | 5.96 | 7.10 | 5.10 | 3.89 | 4.89 | 3.12 |
| 1978 | 5.93 | 7.30 | 4.90 | 6.15 | 7.53 | 5.12 | 3.90 | 5.40 | 2.81 |
| 1979 | 5.95 | 7.40 | 4.88 | 6.18 | 7.70 | 5.08 | 3.65 | 4.50 | 2.98 |
| 1980 | 6.23 | 7.47 | 5.29 | 6.47 | 7.74 | 5.52 | 3.99 | 5.15 | 3.12 |
| 1981 | 6.16 | 7.38 | 5.27 | 6.40 | 7.66 | 5.47 | 3.98 | 5.00 | 3.25 |
| 1982 | 6.55 | 7.98 | 5.49 | 6.83 | 8.29 | 5.74 | 4.12 | 5.23 | 3.29 |
| 1983 | 6.66 | 8.11 | 5.58 | 6.92 | 8.39 | 5.85 | 4.30 | 5.79 | 3.23 |
| 1984 | 6.76 | 8.15 | 5.75 | 7.05 | 8.50 | 6.01 | 4.19 | 5.28 | 3.39 |
| 1985 | 7.06 | 8.80 | 5.80 | 7.34 | 9.17 | 6.02 | 4.62 | 5.57 | 3.87 |
| 1986 | 7.31 | 8.98 | 6.05 | 7.59 | 9.25 | 6.33 | 4.95 | 6.64 | 3.70 |
| 1987 | 7.26 | 8.81 | 6.10 | 7.56 | 9.12 | 6.38 | 4.73 | 6.30 | 3.62 |
| 1988 | 7.52 | 9.28 | 6.24 | 7.82 | 9.59 | 6.53 | 5.01 | 6.82 | 3.72 |
| 1989 | 7.83 | 9.73 | 6.43 | 8.12 | 10.04 | 6.71 | 5.33 | 7.10 | 4.07 |
| 1990 | 7.87 | 9.97 | 6.34 | 8.16 | 10.32 | 6.58 | 5.39 | 7.13 | 4.17 |
| 1991 | 8.19 | 10.15 | 6.74 | 8.50 | 10.52 | 7.00 | 5.70 | 7.17 | 4.61 |
| 1992 | 8.22 | 10.30 | 6.69 | 8.53 | 10.66 | 6.96 | 5.64 | 7.23 | 4.45 |
| 1993 | 8.25 | 10.14 | 6.86 | 8.57 | 10.45 | 7.17 | 5.69 | 7.53 | 4.40 |
| 1994 | 8.63 | 10.75 | 7.10 | 9.00 | 11.20 | 7.41 | 5.54 | 6.82 | 4.56 |
| 1995 | 8.72 | 10.81 | 7.16 | 9.09 | 11.21 | 7.49 | 5.79 | 7.62 | 4.45 |
| 1996 | 8.75 | 10.94 | 7.14 | 9.09 | 11.29 | 7.46 | 6.21 | 8.25 | 4.74 |
| 1997 | 8.88 | 11.03 | 7.25 | 9.21 | 11.35 | 7.57 | 6.34 | 8.45 | 4.87 |
| 1998 | 8.69 | 10.85 | 7.10 | 9.03 | 11.27 | 7.38 | 5.99 | 7.50 | 4.89 |
| 1999 | 8.32 | 10.36 | 6.85 | 8.67 | 10.71 | 7.18 | 5.65 | 7.44 | 4.41 |
| 2000 | 8.16 | 10.19 | 6.69 | 8.53 | 10.62 | 7.01 | 5.20 | 6.61 | 4.17 |
| 2001 | 7.89 | 9.94 | 6.40 | 8.19 | 10.27 | 6.66 | 5.61 | 7.26 | 4.46 |
| 2002 | 7.62 | 9.60 | 6.17 | 7.93 | 9.99 | 6.41 | 5.41 | 6.86 | 4.41 |
| 2003 | 7.33 | 9.32 | 5.88 | 7.64 | 9.74 | 6.09 | 4.93 | 5.82 | 4.26 |
| 2004 | 7.04 | 8.86 | 5.68 | 7.32 | 9.20 | 5.91 | 5.00 | 6.21 | 4.11 |
| 2005 | 6.89 | 8.78 | 5.50 | 7.18 | 9.14 | 5.73 | 4.76 | 6.10 | 3.79 |
| 2006 | 6.68 | 8.41 | 5.38 | 6.95 | 8.72 | 5.60 | 4.92 | 6.28 | 3.91 |
| 1975-2006 | 7.46 | 9.28 | 6.13 | 7.76 | 9.62 | 6.39 | 5.04 | 6.47 | 3.99 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 20.4
Cancer of the Oral Cavity and Pharynx (Invasive)
Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 13.21 | 21.19 | 7.10 | 13.30 | 21.72 | 6.93 | 13.38 | 20.16 | 8.25 |
| 1976 | 13.28 | 21.01 | 7.38 | 13.23 | 21.03 | 7.37 | 15.18 | 23.76 | 8.37 |
| 1977 | 12.73 | 20.11 | 6.95 | 12.59 | 19.85 | 7.01 | 14.45 | 23.86 | 6.65 |
| 1978 | 13.38 | 20.86 | 7.71 | 13.21 | 20.76 | 7.58 | 15.83 | 25.97 | 8.05 |
| 1979 | 14.04 | 21.94 | 8.01 | 13.72 | 21.63 | 7.74 | 18.39 | 28.16 | 10.64 |
| 1980 | 13.34 | 20.27 | 7.91 | 12.98 | 19.98 | 7.60 | 16.82 | 25.39 | 9.59 |
| 1981 | 13.59 | 20.85 | 7.92 | 13.40 | 20.51 | 7.95 | 16.65 | 28.16 | 7.46 |
| 1982 | 13.34 | 20.33 | 7.93 | 13.09 | 19.95 | 7.85 | 17.12 | 27.39 | 8.86 |
| 1983 | 13.29 | 21.21 | 7.24 | 13.25 | 21.21 | 7.29 | 16.12 | 26.08 | 8.32 |
| 1984 | 13.46 | 20.75 | 7.86 | 12.99 | 20.11 | 7.64 | 18.91 | 29.99 | 10.07 |
| 1985 | 13.33 | 19.84 | 8.24 | 13.27 | 19.60 | 8.35 | 16.07 | 25.87 | 8.53 |
| 1986 | 12.47 | 19.13 | 7.21 | 12.36 | 18.87 | 7.30 | 16.18 | 28.09 | 6.97 |
| 1987 | 13.21 | 20.55 | 7.51 | 13.11 | 20.19 | 7.66 | 16.81 | 30.00 | 6.99 |
| 1988 | 12.13 | 18.15 | 7.30 | 11.78 | 17.70 | 7.09 | 16.43 | 25.56 | 9.08 |
| 1989 | 12.12 | 18.30 | 7.26 | 11.84 | 17.71 | 7.30 | 15.75 | 27.16 | 7.05 |
| 1990 | 12.64 | 19.37 | 7.34 | 12.37 | 18.78 | 7.39 | 15.99 | 27.77 | 6.92 |
| 1991 | 12.16 | 18.35 | 7.25 | 12.08 | 18.11 | 7.30 | 14.41 | 23.71 | 7.32 |
| 1992 | 11.96 | 18.32 | 6.92 | 11.81 | 18.02 | 6.93 | 14.79 | 25.08 | 6.94 |
| 1993 | 12.43 | 18.71 | 7.29 | 12.23 | 18.41 | 7.21 | 15.54 | 24.85 | 8.10 |
| 1994 | 11.71 | 17.67 | 6.83 | 11.39 | 16.84 | 6.92 | 15.62 | 26.64 | 6.99 |
| 1995 | 11.54 | 16.93 | 7.20 | 11.53 | 16.73 | 7.34 | 13.47 | 22.46 | 6.66 |
| 1996 | 11.76 | 17.73 | 6.99 | 11.45 | 17.23 | 6.87 | 14.97 | 24.34 | 7.63 |
| 1997 | 11.51 | 17.33 | 6.76 | 11.52 | 17.19 | 6.87 | 11.45 | 18.24 | 6.19 |
| 1998 | 11.23 | 16.66 | 6.76 | 11.23 | 16.41 | 6.91 | 13.00 | 22.44 | 6.22 |
| 1999 | 10.57 | 15.55 | 6.45 | 10.46 | 15.42 | 6.33 | 11.74 | 19.78 | 5.82 |
| 2000 | 10.73 | 16.02 | 6.28 | 10.65 | 15.61 | 6.41 | 11.68 | 20.70 | 4.87 |
| 2001 | 10.77 | 15.40 | 6.89 | 10.83 | 15.46 | 6.90 | 11.88 | 18.61 | 6.91 |
| 2002 | 11.02 | 16.18 | 6.66 | 11.21 | 16.42 | 6.75 | 10.97 | 16.79 | 6.41 |
| 2003 | 10.40 | 15.43 | 6.12 | 10.33 | 15.41 | 5.95 | 11.57 | 16.69 | 7.51 |
| 2004 | 10.63 | 15.74 | 6.20 | 10.81 | 15.99 | 6.22 | 9.87 | 15.65 | 5.56 |
| 2005 | 10.41 | 15.15 | 6.37 | 10.52 | 15.48 | 6.20 | 10.78 | 16.10 | 6.87 |
| 2006 | 10.29 | 14.97 | 6.24 | 10.42 | 14.97 | 6.39 | 9.61 | 15.85 | 4.96 |
| 1975-2006 | 12.01 | 18.11 | 7.08 | 11.95 | 17.99 | 7.09 | 13.76 | 22.39 | 7.10 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 20.5
Cancer of the Oral Cavity and Pharynx (Invasive)
Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 4.30 | 6.93 | 2.33 | 4.12 | 6.64 | 2.29 | 5.79 | 9.93 | 2.52 |
| 1976 | 4.22 | 6.89 | 2.22 | 4.03 | 6.55 | 2.17 | 6.08 | 10.55 | 2.62 |
| 1977 | 4.32 | 7.03 | 2.30 | 4.11 | 6.67 | 2.25 | 6.22 | 10.87 | 2.65 |
| 1978 | 4.19 | 6.78 | 2.26 | 3.99 | 6.41 | 2.21 | 6.15 | 10.69 | 2.68 |
| 1979 | 4.24 | 6.71 | 2.39 | 3.99 | 6.25 | 2.32 | 6.65 | 11.47 | 2.97 |
| 1980 | 4.16 | 6.69 | 2.27 | 3.87 | 6.12 | 2.22 | 6.97 | 12.48 | 2.76 |
| 1981 | 4.05 | 6.46 | 2.24 | 3.78 | 5.96 | 2.17 | 6.52 | 11.36 | 2.84 |
| 1982 | 4.02 | 6.46 | 2.22 | 3.74 | 5.96 | 2.13 | 6.62 | 11.46 | 2.94 |
| 1983 | 3.94 | 6.27 | 2.20 | 3.64 | 5.73 | 2.10 | 6.65 | 11.83 | 2.76 |
| 1984 | 3.88 | 6.18 | 2.17 | 3.61 | 5.64 | 2.12 | 6.37 | 11.70 | 2.38 |
| 1985 | 3.75 | 5.93 | 2.11 | 3.49 | 5.44 | 2.05 | 6.23 | 11.03 | 2.62 |
| 1986 | 3.72 | 5.80 | 2.16 | 3.42 | 5.27 | 2.07 | 6.51 | 11.35 | 2.93 |
| 1987 | 3.53 | 5.46 | 2.05 | 3.29 | 5.01 | 2.00 | 5.70 | 9.95 | 2.54 |
| 1988 | 3.59 | 5.63 | 2.05 | 3.33 | 5.14 | 1.98 | 5.99 | 10.67 | 2.55 |
| 1989 | 3.50 | 5.34 | 2.07 | 3.23 | 4.84 | 2.00 | 5.92 | 10.44 | 2.59 |
| 1990 | 3.58 | 5.61 | 2.03 | 3.29 | 5.07 | 1.96 | 6.12 | 11.05 | 2.52 |
| 1991 | 3.47 | 5.34 | 2.03 | 3.21 | 4.87 | 1.95 | 5.84 | 10.06 | 2.70 |
| 1992 | 3.34 | 5.15 | 1.94 | 3.09 | 4.66 | 1.89 | 5.56 | 9.97 | 2.31 |
| 1993 | 3.34 | 5.19 | 1.90 | 3.07 | 4.68 | 1.83 | 5.68 | 10.09 | 2.44 |
| 1994 | 3.15 | 4.84 | 1.83 | 2.92 | 4.41 | 1.77 | 5.33 | 9.26 | 2.39 |
| 1995 | 3.16 | 4.87 | 1.84 | 2.91 | 4.42 | 1.75 | 5.37 | 9.28 | 2.54 |
| 1996 | 3.03 | 4.64 | 1.75 | 2.82 | 4.24 | 1.71 | 4.86 | 8.69 | 2.07 |
| 1997 | 2.98 | 4.53 | 1.76 | 2.79 | 4.17 | 1.70 | 4.68 | 8.10 | 2.19 |
| 1998 | 2.96 | 4.53 | 1.72 | 2.80 | 4.20 | 1.68 | 4.53 | 7.91 | 2.09 |
| 1999 | 2.73 | 4.15 | 1.60 | 2.55 | 3.80 | 1.54 | 4.37 | 7.49 | 2.12 |
| 2000 | 2.69 | 4.06 | 1.58 | 2.52 | 3.72 | 1.55 | 4.12 | 7.31 | 1.76 |
| 2001 | 2.72 | 4.14 | 1.56 | 2.58 | 3.87 | 1.52 | 3.88 | 6.68 | 1.80 |
| 2002 | 2.67 | 4.07 | 1.54 | 2.56 | 3.86 | 1.50 | 3.67 | 6.30 | 1.73 |
| 2003 | 2.63 | 4.06 | 1.47 | 2.50 | 3.78 | 1.44 | 3.85 | 6.92 | 1.64 |
| 2004 | 2.60 | 3.99 | 1.47 | 2.48 | 3.71 | 1.45 | 3.84 | 6.91 | 1.63 |
| 2005 | 2.53 | 3.84 | 1.43 | 2.42 | 3.59 | 1.43 | 3.53 | 6.52 | 1.37 |
| 2006 | 2.47 | 3.78 | 1.39 | 2.39 | 3.60 | 1.37 | 3.27 | 5.70 | 1.50 |
| 1975-2006 | 3.34 | 5.18 | 1.90 | 3.14 | 4.80 | 1.85 | 5.15 | 9.05 | 2.26 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 21.4
Cancer of the Ovary (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

| | All Races, Females | | | White Females | | | Black Females | | |
|--------------------|--------------------|-------|-------|---------------|-------|-------|---------------|------|-------|
| | All | <65 | 65+ | All | <65 | 65+ | All | <65 | 65+ |
| Year of Diagnosis: | | | | | | | | | |
| 1975 | 16.32 | 11.61 | 48.88 | 16.77 | 12.00 | 49.79 | 11.72 | 7.40 | 41.58 |
| 1976 | 15.85 | 11.22 | 47.86 | 16.63 | 11.87 | 49.55 | 11.41 | 7.37 | 39.32 |
| 1977 | 15.62 | 11.20 | 46.15 | 16.19 | 11.58 | 48.07 | 11.34 | 8.69 | 29.65 |
| 1978 | 15.38 | 10.78 | 47.18 | 16.16 | 11.36 | 49.31 | 9.96 | 6.07 | 36.83 |
| 1979 | 15.38 | 10.69 | 47.79 | 15.83 | 10.90 | 49.91 | 11.94 | 9.07 | 31.79 |
| 1980 | 15.44 | 10.64 | 48.61 | 16.20 | 11.18 | 50.88 | 11.74 | 7.42 | 41.60 |
| 1981 | 15.42 | 10.48 | 49.53 | 16.00 | 10.87 | 51.47 | 11.68 | 7.58 | 40.08 |
| 1982 | 15.56 | 10.08 | 53.43 | 16.13 | 10.48 | 55.14 | 12.33 | 7.43 | 46.23 |
| 1983 | 15.97 | 10.44 | 54.18 | 16.41 | 10.71 | 55.78 | 13.06 | 9.18 | 39.89 |
| 1984 | 16.25 | 10.96 | 52.79 | 17.15 | 11.71 | 54.71 | 10.98 | 5.87 | 46.31 |
| 1985 | 16.53 | 10.76 | 56.41 | 17.43 | 11.40 | 59.13 | 11.30 | 7.25 | 39.27 |
| 1986 | 15.01 | 9.87 | 50.55 | 15.71 | 10.47 | 51.91 | 10.69 | 6.24 | 41.47 |
| 1987 | 16.15 | 10.15 | 57.58 | 17.04 | 10.85 | 59.80 | 11.82 | 6.26 | 50.27 |
| 1988 | 15.26 | 9.56 | 54.69 | 16.01 | 10.08 | 56.95 | 11.56 | 6.63 | 45.65 |
| 1989 | 15.51 | 9.53 | 56.80 | 16.49 | 10.27 | 59.53 | 10.99 | 5.42 | 49.45 |
| 1990 | 15.39 | 9.44 | 56.48 | 16.26 | 10.05 | 59.15 | 10.23 | 5.68 | 41.65 |
| 1991 | 15.75 | 9.69 | 57.66 | 16.54 | 10.29 | 59.74 | 11.03 | 5.48 | 49.47 |
| 1992 | 14.90 | 9.33 | 53.40 | 15.65 | 9.73 | 56.55 | 11.23 | 7.77 | 35.16 |
| 1993 | 15.22 | 9.13 | 57.35 | 16.04 | 9.62 | 60.42 | 11.62 | 6.59 | 46.34 |
| 1994 | 14.47 | 8.87 | 53.14 | 15.00 | 9.16 | 55.32 | 12.44 | 6.49 | 53.54 |
| 1995 | 14.56 | 8.74 | 54.80 | 15.33 | 9.30 | 57.03 | 10.30 | 4.89 | 47.70 |
| 1996 | 14.10 | 8.40 | 53.51 | 15.26 | 9.22 | 57.05 | 9.03 | 4.53 | 40.11 |
| 1997 | 14.71 | 9.20 | 52.79 | 15.42 | 9.68 | 55.09 | 10.61 | 5.65 | 44.91 |
| 1998 | 14.31 | 8.80 | 52.43 | 15.08 | 9.24 | 55.50 | 10.76 | 5.93 | 44.19 |
| 1999 | 14.65 | 8.68 | 55.89 | 15.39 | 8.92 | 60.13 | 10.69 | 6.25 | 41.36 |
| 2000 | 14.26 | 9.01 | 50.53 | 15.17 | 9.62 | 53.53 | 10.78 | 5.49 | 47.33 |
| 2001 | 14.48 | 9.12 | 51.53 | 15.51 | 9.71 | 55.62 | 9.36 | 6.19 | 31.29 |
| 2002 | 13.73 | 7.94 | 53.78 | 14.35 | 8.28 | 56.30 | 10.35 | 5.54 | 43.60 |
| 2003 | 13.48 | 8.32 | 49.21 | 14.02 | 8.78 | 50.23 | 11.92 | 5.95 | 53.21 |
| 2004 | 12.99 | 7.86 | 48.42 | 13.63 | 8.06 | 52.13 | 9.87 | 6.15 | 35.57 |
| 2005 | 12.67 | 7.91 | 45.52 | 13.23 | 8.02 | 49.27 | 10.29 | 6.91 | 33.59 |
| 2006 | 12.48 | 7.41 | 47.52 | 13.10 | 7.73 | 50.21 | 8.44 | 5.16 | 31.09 |
| 1975-2006 | 14.82 | 9.39 | 52.38 | 15.59 | 9.90 | 54.93 | 10.90 | 6.38 | 42.12 |

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 21.5
Cancer of the Ovary (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Age

| | All Races, Females | | | White Females | | | Black Females | | |
|----------------|--------------------|------|-------|---------------|------|-------|---------------|------|-------|
| | All | <65 | 65+ | All | <65 | 65+ | All | <65 | 65+ |
| Year of Death: | | | | | | | | | |
| 1975 | 9.84 | 5.83 | 37.55 | 10.16 | 6.01 | 38.81 | 7.16 | 4.64 | 24.58 |
| 1976 | 10.02 | 5.73 | 39.66 | 10.28 | 5.90 | 40.58 | 8.01 | 4.61 | 31.52 |
| 1977 | 9.58 | 5.47 | 38.05 | 9.87 | 5.65 | 39.08 | 7.41 | 4.25 | 29.21 |
| 1978 | 9.55 | 5.44 | 37.97 | 9.83 | 5.62 | 38.88 | 7.35 | 4.17 | 29.33 |
| 1979 | 9.33 | 5.10 | 38.58 | 9.56 | 5.20 | 39.70 | 7.53 | 4.43 | 28.94 |
| 1980 | 9.29 | 4.97 | 39.15 | 9.52 | 5.11 | 39.96 | 7.85 | 4.18 | 33.21 |
| 1981 | 9.23 | 4.94 | 38.89 | 9.48 | 5.10 | 39.75 | 7.36 | 3.74 | 32.36 |
| 1982 | 9.22 | 4.80 | 39.81 | 9.49 | 4.97 | 40.72 | 7.41 | 3.64 | 33.45 |
| 1983 | 9.20 | 4.70 | 40.34 | 9.43 | 4.80 | 41.38 | 7.67 | 4.11 | 32.25 |
| 1984 | 9.11 | 4.63 | 40.08 | 9.36 | 4.79 | 40.93 | 7.55 | 3.76 | 33.80 |
| 1985 | 9.08 | 4.57 | 40.23 | 9.34 | 4.72 | 41.29 | 7.30 | 3.74 | 31.87 |
| 1986 | 9.24 | 4.57 | 41.55 | 9.52 | 4.73 | 42.63 | 7.42 | 3.66 | 33.43 |
| 1987 | 9.18 | 4.46 | 41.86 | 9.46 | 4.64 | 42.82 | 7.55 | 3.52 | 35.45 |
| 1988 | 9.31 | 4.41 | 43.16 | 9.55 | 4.56 | 44.08 | 7.87 | 3.67 | 36.88 |
| 1989 | 9.21 | 4.30 | 43.20 | 9.52 | 4.49 | 44.27 | 7.41 | 3.26 | 36.16 |
| 1990 | 9.33 | 4.26 | 44.35 | 9.61 | 4.46 | 45.23 | 7.86 | 3.32 | 39.24 |
| 1991 | 9.51 | 4.32 | 45.37 | 9.74 | 4.48 | 46.13 | 8.36 | 3.43 | 42.46 |
| 1992 | 9.46 | 4.24 | 45.51 | 9.70 | 4.38 | 46.49 | 8.19 | 3.59 | 39.99 |
| 1993 | 9.08 | 4.02 | 44.06 | 9.34 | 4.19 | 44.93 | 7.74 | 3.02 | 40.42 |
| 1994 | 9.38 | 4.14 | 45.64 | 9.67 | 4.32 | 46.70 | 7.82 | 3.20 | 39.72 |
| 1995 | 9.12 | 3.96 | 44.77 | 9.47 | 4.17 | 46.07 | 7.37 | 2.93 | 38.04 |
| 1996 | 8.86 | 3.82 | 43.70 | 9.13 | 3.97 | 44.80 | 7.32 | 2.89 | 37.92 |
| 1997 | 8.94 | 3.77 | 44.70 | 9.21 | 3.94 | 45.59 | 7.93 | 3.11 | 41.19 |
| 1998 | 8.73 | 3.68 | 43.64 | 9.02 | 3.81 | 45.03 | 7.37 | 3.22 | 36.06 |
| 1999 | 8.77 | 3.73 | 43.57 | 9.07 | 3.90 | 44.83 | 7.38 | 3.05 | 37.31 |
| 2000 | 8.89 | 3.67 | 44.98 | 9.25 | 3.85 | 46.56 | 7.15 | 2.91 | 36.48 |
| 2001 | 8.99 | 3.73 | 45.37 | 9.31 | 3.87 | 46.86 | 7.52 | 3.20 | 37.36 |
| 2002 | 9.02 | 3.79 | 45.18 | 9.37 | 3.97 | 46.73 | 7.43 | 3.11 | 37.29 |
| 2003 | 8.85 | 3.68 | 44.60 | 9.17 | 3.83 | 46.06 | 7.45 | 3.12 | 37.39 |
| 2004 | 8.76 | 3.54 | 44.88 | 9.06 | 3.65 | 46.44 | 7.65 | 3.13 | 38.90 |
| 2005 | 8.63 | 3.57 | 43.64 | 8.96 | 3.73 | 45.09 | 7.28 | 2.90 | 37.57 |
| 2006 | 8.54 | 3.47 | 43.59 | 8.91 | 3.62 | 45.45 | 6.85 | 2.92 | 34.03 |
| 1975-2006 | 9.15 | 4.26 | 42.97 | 9.44 | 4.43 | 44.11 | 7.56 | 3.43 | 36.13 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 22.4
Cancer of the Pancreas (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 11.84 | 15.61 | 9.12 | 11.69 | 15.65 | 8.89 | 15.90 | 18.08 | 13.91 |
| 1976 | 11.99 | 14.82 | 9.91 | 11.76 | 14.50 | 9.75 | 17.61 | 22.68 | 13.97 |
| 1977 | 11.90 | 14.82 | 9.83 | 11.62 | 14.59 | 9.54 | 17.12 | 19.44 | 15.25 |
| 1978 | 11.07 | 14.06 | 8.85 | 10.83 | 13.90 | 8.60 | 15.14 | 20.02 | 11.53 |
| 1979 | 11.52 | 14.15 | 9.65 | 11.21 | 13.84 | 9.39 | 16.08 | 19.14 | 13.68 |
| 1980 | 11.52 | 14.20 | 9.61 | 11.07 | 13.70 | 9.19 | 17.94 | 21.76 | 15.53 |
| 1981 | 11.69 | 13.55 | 10.17 | 11.39 | 13.26 | 9.91 | 17.18 | 20.86 | 14.10 |
| 1982 | 11.71 | 13.81 | 10.17 | 11.41 | 13.47 | 9.94 | 16.86 | 20.99 | 13.96 |
| 1983 | 12.19 | 14.66 | 10.44 | 11.75 | 14.07 | 10.12 | 18.13 | 23.05 | 14.90 |
| 1984 | 12.22 | 13.94 | 10.93 | 12.03 | 13.87 | 10.64 | 17.19 | 18.36 | 16.34 |
| 1985 | 12.10 | 14.34 | 10.50 | 11.70 | 13.60 | 10.34 | 18.70 | 24.98 | 14.57 |
| 1986 | 11.68 | 13.78 | 10.21 | 11.37 | 13.54 | 9.82 | 17.32 | 20.16 | 15.59 |
| 1987 | 11.61 | 13.81 | 10.10 | 11.09 | 13.38 | 9.54 | 18.43 | 18.98 | 17.93 |
| 1988 | 11.65 | 13.70 | 10.20 | 11.11 | 13.35 | 9.57 | 18.48 | 19.55 | 17.32 |
| 1989 | 11.34 | 12.88 | 10.08 | 10.98 | 12.66 | 9.64 | 14.99 | 16.05 | 13.97 |
| 1990 | 11.28 | 12.81 | 10.06 | 11.02 | 12.46 | 9.84 | 15.20 | 18.82 | 12.63 |
| 1991 | 11.40 | 12.90 | 10.23 | 11.04 | 12.61 | 9.79 | 17.06 | 18.20 | 16.24 |
| 1992 | 11.62 | 13.22 | 10.34 | 11.33 | 12.90 | 10.02 | 17.34 | 19.77 | 15.75 |
| 1993 | 10.96 | 12.53 | 9.69 | 10.51 | 11.98 | 9.30 | 16.82 | 18.71 | 15.18 |
| 1994 | 11.37 | 13.25 | 9.90 | 10.89 | 12.34 | 9.73 | 17.45 | 21.53 | 14.49 |
| 1995 | 11.13 | 12.72 | 9.96 | 10.71 | 12.31 | 9.58 | 18.11 | 20.31 | 16.31 |
| 1996 | 11.29 | 12.92 | 9.92 | 10.90 | 12.48 | 9.56 | 17.06 | 19.68 | 14.88 |
| 1997 | 11.49 | 13.06 | 10.19 | 11.04 | 12.57 | 9.74 | 17.83 | 19.35 | 16.48 |
| 1998 | 11.47 | 13.11 | 10.22 | 11.25 | 13.00 | 9.85 | 14.75 | 16.22 | 13.89 |
| 1999 | 11.09 | 12.59 | 9.81 | 10.69 | 12.22 | 9.33 | 16.67 | 19.46 | 14.69 |
| 2000 | 11.41 | 13.31 | 9.83 | 11.07 | 12.92 | 9.58 | 15.67 | 18.16 | 13.24 |
| 2001 | 11.29 | 13.08 | 9.97 | 11.03 | 13.00 | 9.53 | 14.19 | 15.37 | 13.27 |
| 2002 | 11.68 | 12.97 | 10.50 | 11.53 | 13.22 | 10.01 | 15.97 | 14.89 | 16.07 |
| 2003 | 11.52 | 12.75 | 10.49 | 11.32 | 12.55 | 10.25 | 15.90 | 16.86 | 15.16 |
| 2004 | 11.82 | 13.73 | 10.22 | 11.61 | 13.43 | 10.08 | 15.65 | 18.92 | 13.09 |
| 2005 | 11.76 | 13.42 | 10.40 | 11.47 | 13.13 | 10.08 | 16.54 | 17.77 | 15.85 |
| 2006 | 12.01 | 13.56 | 10.71 | 11.86 | 13.65 | 10.36 | 15.96 | 17.26 | 14.76 |
| 1975-2006 | 11.57 | 13.47 | 10.10 | 11.25 | 13.17 | 9.76 | 16.62 | 18.92 | 14.87 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 22.5
Cancer of the Pancreas (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 10.66 | 13.80 | 8.42 | 10.48 | 13.61 | 8.27 | 12.81 | 16.32 | 10.15 |
| 1976 | 10.68 | 13.62 | 8.57 | 10.50 | 13.43 | 8.41 | 12.86 | 16.05 | 10.47 |
| 1977 | 10.87 | 13.86 | 8.76 | 10.68 | 13.61 | 8.62 | 13.20 | 17.14 | 10.35 |
| 1978 | 10.79 | 13.63 | 8.76 | 10.62 | 13.47 | 8.59 | 12.90 | 15.93 | 10.69 |
| 1979 | 10.73 | 13.39 | 8.81 | 10.53 | 13.25 | 8.57 | 13.39 | 15.85 | 11.55 |
| 1980 | 10.62 | 13.10 | 8.83 | 10.36 | 12.85 | 8.58 | 13.78 | 16.47 | 11.80 |
| 1981 | 10.70 | 13.16 | 8.92 | 10.50 | 12.91 | 8.77 | 13.40 | 16.96 | 10.83 |
| 1982 | 10.59 | 13.06 | 8.85 | 10.32 | 12.76 | 8.60 | 14.09 | 17.07 | 11.91 |
| 1983 | 10.69 | 12.82 | 9.13 | 10.36 | 12.40 | 8.87 | 14.78 | 18.10 | 12.45 |
| 1984 | 10.80 | 12.96 | 9.22 | 10.45 | 12.61 | 8.89 | 14.96 | 17.52 | 13.08 |
| 1985 | 10.63 | 12.86 | 9.03 | 10.33 | 12.53 | 8.76 | 14.43 | 17.29 | 12.40 |
| 1986 | 10.60 | 12.46 | 9.23 | 10.30 | 12.16 | 8.93 | 14.60 | 16.72 | 13.04 |
| 1987 | 10.59 | 12.52 | 9.18 | 10.28 | 12.16 | 8.90 | 14.58 | 17.36 | 12.61 |
| 1988 | 10.49 | 12.51 | 8.98 | 10.15 | 12.16 | 8.65 | 14.52 | 17.05 | 12.66 |
| 1989 | 10.62 | 12.61 | 9.17 | 10.25 | 12.22 | 8.82 | 15.02 | 17.71 | 13.15 |
| 1990 | 10.69 | 12.59 | 9.25 | 10.34 | 12.19 | 8.92 | 15.18 | 18.08 | 13.21 |
| 1991 | 10.69 | 12.58 | 9.28 | 10.40 | 12.27 | 8.99 | 14.55 | 16.93 | 12.79 |
| 1992 | 10.70 | 12.61 | 9.27 | 10.35 | 12.23 | 8.92 | 15.40 | 18.22 | 13.44 |
| 1993 | 10.66 | 12.41 | 9.37 | 10.33 | 12.06 | 9.05 | 15.05 | 17.59 | 13.31 |
| 1994 | 10.64 | 12.37 | 9.31 | 10.28 | 11.98 | 8.95 | 15.16 | 17.69 | 13.38 |
| 1995 | 10.45 | 12.13 | 9.18 | 10.21 | 11.91 | 8.92 | 13.90 | 15.78 | 12.54 |
| 1996 | 10.47 | 12.08 | 9.23 | 10.17 | 11.79 | 8.91 | 14.43 | 16.45 | 12.97 |
| 1997 | 10.45 | 12.22 | 9.07 | 10.16 | 11.97 | 8.76 | 14.57 | 16.39 | 13.14 |
| 1998 | 10.52 | 12.29 | 9.15 | 10.23 | 12.02 | 8.83 | 14.52 | 16.49 | 13.07 |
| 1999 | 10.62 | 12.32 | 9.27 | 10.38 | 12.07 | 9.03 | 14.18 | 16.58 | 12.40 |
| 2000 | 10.54 | 12.14 | 9.26 | 10.32 | 11.97 | 8.99 | 14.04 | 15.79 | 12.70 |
| 2001 | 10.55 | 12.12 | 9.28 | 10.37 | 12.02 | 9.05 | 13.67 | 15.05 | 12.53 |
| 2002 | 10.53 | 12.18 | 9.20 | 10.32 | 12.04 | 8.93 | 13.68 | 15.44 | 12.34 |
| 2003 | 10.50 | 12.05 | 9.23 | 10.30 | 11.89 | 8.99 | 13.79 | 15.49 | 12.45 |
| 2004 | 10.65 | 12.36 | 9.25 | 10.46 | 12.19 | 9.02 | 13.67 | 15.58 | 12.13 |
| 2005 | 10.77 | 12.37 | 9.46 | 10.61 | 12.25 | 9.25 | 13.73 | 15.47 | 12.39 |
| 2006 | 10.80 | 12.40 | 9.47 | 10.66 | 12.36 | 9.24 | 13.67 | 14.96 | 12.54 |
| 1975-2006 | 10.65 | 12.61 | 9.16 | 10.39 | 12.36 | 8.89 | 14.21 | 16.54 | 12.48 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 23.4
Cancer of the Prostate (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

| | All Races, Males | | | White Males | | | Black Males | | |
|--------------------|------------------|-------|----------|-------------|-------|----------|-------------|--------|----------|
| | All | <65 | 65+ | All | <65 | 65+ | All | <65 | 65+ |
| Year of Diagnosis: | | | | | | | | | |
| 1975 | 94.02 | 13.66 | 649.53 | 92.21 | 13.02 | 639.63 | 140.97 | 26.76 | 930.45 |
| 1976 | 97.92 | 14.57 | 674.08 | 97.31 | 13.90 | 673.91 | 140.79 | 29.38 | 910.86 |
| 1977 | 100.45 | 15.03 | 690.95 | 98.54 | 14.34 | 680.54 | 159.16 | 29.24 | 1,057.18 |
| 1978 | 99.41 | 15.23 | 681.22 | 97.72 | 14.80 | 670.92 | 148.11 | 26.83 | 986.42 |
| 1979 | 103.45 | 14.93 | 715.26 | 102.32 | 14.64 | 708.40 | 162.31 | 26.35 | 1,102.12 |
| 1980 | 105.99 | 15.47 | 731.75 | 104.95 | 14.64 | 729.19 | 160.99 | 33.62 | 1,041.42 |
| 1981 | 108.90 | 16.81 | 745.49 | 107.91 | 15.95 | 743.54 | 161.86 | 34.55 | 1,041.86 |
| 1982 | 108.18 | 16.30 | 743.32 | 107.30 | 15.75 | 740.10 | 167.63 | 31.32 | 1,109.83 |
| 1983 | 111.59 | 17.12 | 764.61 | 110.80 | 16.56 | 762.23 | 170.84 | 34.05 | 1,116.37 |
| 1984 | 111.69 | 17.26 | 764.41 | 110.10 | 16.22 | 759.05 | 178.77 | 37.08 | 1,158.14 |
| 1985 | 115.49 | 17.84 | 790.49 | 114.63 | 17.56 | 785.59 | 170.18 | 31.41 | 1,129.40 |
| 1986 | 119.02 | 18.51 | 813.79 | 118.98 | 18.27 | 815.14 | 168.59 | 34.08 | 1,098.34 |
| 1987 | 133.66 | 21.50 | 908.96 | 134.50 | 21.09 | 918.49 | 189.18 | 36.06 | 1,247.58 |
| 1988 | 137.59 | 22.18 | 935.31 | 138.69 | 22.30 | 943.22 | 190.19 | 34.71 | 1,264.94 |
| 1989 | 145.26 | 24.09 | 982.81 | 146.00 | 23.93 | 989.80 | 191.98 | 37.28 | 1,261.29 |
| 1990 | 170.89 | 28.77 | 1,153.26 | 172.66 | 28.86 | 1,166.62 | 221.75 | 44.28 | 1,448.44 |
| 1991 | 214.71 | 38.82 | 1,430.48 | 216.27 | 39.20 | 1,440.23 | 288.39 | 57.32 | 1,885.65 |
| 1992 | 237.21 | 49.36 | 1,535.71 | 238.18 | 49.44 | 1,542.78 | 328.10 | 77.26 | 2,061.99 |
| 1993 | 209.25 | 50.45 | 1,306.93 | 204.25 | 49.00 | 1,277.39 | 344.14 | 94.13 | 2,072.27 |
| 1994 | 180.04 | 48.60 | 1,088.63 | 174.05 | 46.82 | 1,053.50 | 312.82 | 95.00 | 1,818.52 |
| 1995 | 168.68 | 49.86 | 990.03 | 163.83 | 47.99 | 964.52 | 280.53 | 98.00 | 1,542.27 |
| 1996 | 168.90 | 53.33 | 967.76 | 164.43 | 51.82 | 942.80 | 282.46 | 100.46 | 1,540.47 |
| 1997 | 173.07 | 55.10 | 988.46 | 169.43 | 53.93 | 967.78 | 281.66 | 96.59 | 1,560.89 |
| 1998 | 170.10 | 54.72 | 967.67 | 166.43 | 52.87 | 951.39 | 284.38 | 101.94 | 1,545.49 |
| 1999 | 182.79 | 60.67 | 1,026.94 | 178.19 | 58.75 | 1,003.82 | 288.02 | 111.18 | 1,510.38 |
| 2000 | 181.84 | 61.55 | 1,013.37 | 178.16 | 59.62 | 997.53 | 288.86 | 113.14 | 1,503.51 |
| 2001 | 183.45 | 64.08 | 1,008.54 | 181.24 | 62.02 | 1,005.35 | 270.58 | 114.05 | 1,352.54 |
| 2002 | 180.45 | 64.81 | 979.78 | 177.70 | 63.02 | 970.41 | 284.75 | 116.53 | 1,447.49 |
| 2003 | 167.45 | 61.07 | 902.82 | 163.79 | 58.63 | 890.70 | 254.58 | 114.69 | 1,221.51 |
| 2004 | 162.62 | 60.63 | 867.63 | 158.87 | 58.64 | 851.65 | 247.19 | 103.69 | 1,239.11 |
| 2005 | 151.96 | 55.96 | 815.51 | 146.65 | 52.97 | 794.21 | 235.94 | 101.29 | 1,166.71 |
| 2006 | 163.06 | 61.18 | 867.29 | 157.10 | 58.07 | 841.60 | 232.77 | 100.69 | 1,145.73 |
| 1975-2006 | 151.48 | 39.28 | 927.05 | 148.48 | 37.52 | 915.49 | 233.86 | 73.37 | 1,343.23 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 23.5
Cancer of the Prostate (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Age

| Year of Death: | All Races, Males | | | White Males | | | Black Males | | |
|----------------|------------------|------|--------|-------------|------|--------|-------------|------|--------|
| | All | <65 | 65+ | All | <65 | 65+ | All | <65 | 65+ |
| 1975 | 30.97 | 2.54 | 227.48 | 29.06 | 2.18 | 214.88 | 55.52 | 6.56 | 393.98 |
| 1976 | 31.78 | 2.69 | 232.91 | 29.93 | 2.31 | 220.84 | 55.60 | 6.90 | 392.19 |
| 1977 | 31.83 | 2.69 | 233.28 | 29.90 | 2.31 | 220.57 | 56.44 | 6.93 | 398.67 |
| 1978 | 32.66 | 2.61 | 240.33 | 30.62 | 2.24 | 226.79 | 58.49 | 6.76 | 416.09 |
| 1979 | 32.84 | 2.56 | 242.13 | 30.77 | 2.21 | 228.17 | 59.24 | 6.63 | 422.90 |
| 1980 | 33.05 | 2.67 | 243.08 | 30.77 | 2.33 | 227.39 | 61.99 | 6.57 | 445.03 |
| 1981 | 33.17 | 2.68 | 243.95 | 30.76 | 2.30 | 227.52 | 63.61 | 7.06 | 454.45 |
| 1982 | 33.36 | 2.68 | 245.47 | 31.09 | 2.34 | 229.85 | 62.36 | 6.62 | 447.67 |
| 1983 | 33.92 | 2.76 | 249.30 | 31.57 | 2.40 | 233.18 | 64.55 | 6.99 | 462.46 |
| 1984 | 34.06 | 2.71 | 250.77 | 31.58 | 2.37 | 233.49 | 65.93 | 6.74 | 475.13 |
| 1985 | 33.91 | 2.73 | 249.43 | 31.28 | 2.34 | 231.34 | 67.26 | 7.16 | 482.68 |
| 1986 | 34.93 | 2.80 | 256.99 | 32.46 | 2.47 | 239.72 | 67.39 | 6.63 | 487.36 |
| 1987 | 35.11 | 2.71 | 259.06 | 32.57 | 2.38 | 241.27 | 68.88 | 6.66 | 499.03 |
| 1988 | 35.88 | 2.83 | 264.40 | 33.38 | 2.47 | 247.08 | 69.57 | 6.98 | 502.28 |
| 1989 | 37.10 | 2.97 | 273.01 | 34.54 | 2.62 | 255.20 | 71.71 | 7.06 | 518.54 |
| 1990 | 38.56 | 3.02 | 284.23 | 35.67 | 2.65 | 263.98 | 77.98 | 7.43 | 565.65 |
| 1991 | 39.31 | 2.89 | 291.08 | 36.50 | 2.52 | 271.39 | 78.24 | 7.06 | 570.21 |
| 1992 | 39.22 | 2.86 | 290.58 | 36.29 | 2.44 | 270.28 | 79.84 | 7.66 | 578.73 |
| 1993 | 39.34 | 2.72 | 292.48 | 36.26 | 2.33 | 270.82 | 81.86 | 7.04 | 599.08 |
| 1994 | 38.54 | 2.67 | 286.48 | 35.64 | 2.31 | 266.04 | 79.96 | 6.76 | 585.93 |
| 1995 | 37.29 | 2.58 | 277.23 | 34.41 | 2.22 | 256.93 | 78.20 | 6.70 | 572.47 |
| 1996 | 36.00 | 2.50 | 267.61 | 33.00 | 2.13 | 246.38 | 78.84 | 6.45 | 579.20 |
| 1997 | 34.15 | 2.31 | 254.26 | 31.35 | 1.94 | 234.58 | 74.30 | 6.32 | 544.22 |
| 1998 | 32.63 | 2.27 | 242.49 | 29.89 | 1.90 | 223.36 | 72.77 | 6.28 | 532.41 |
| 1999 | 31.56 | 2.06 | 235.48 | 28.91 | 1.72 | 216.88 | 70.13 | 5.80 | 514.78 |
| 2000 | 30.31 | 1.98 | 226.13 | 27.70 | 1.66 | 207.70 | 68.65 | 5.41 | 505.78 |
| 2001 | 29.17 | 1.94 | 217.38 | 26.67 | 1.64 | 199.73 | 66.51 | 5.40 | 488.94 |
| 2002 | 28.19 | 1.93 | 209.67 | 25.91 | 1.61 | 193.89 | 62.98 | 5.36 | 461.24 |
| 2003 | 26.57 | 1.91 | 197.00 | 24.49 | 1.60 | 182.73 | 58.01 | 5.44 | 421.37 |
| 2004 | 25.51 | 1.83 | 189.20 | 23.48 | 1.56 | 175.01 | 56.20 | 4.96 | 410.42 |
| 2005 | 24.70 | 1.81 | 182.92 | 22.75 | 1.52 | 169.50 | 54.17 | 5.04 | 393.79 |
| 2006 | 23.56 | 1.86 | 173.56 | 21.76 | 1.59 | 161.16 | 51.07 | 4.79 | 370.97 |
| 1975-2006 | 32.78 | 2.45 | 242.40 | 30.34 | 2.11 | 225.44 | 67.12 | 6.33 | 487.32 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 24.4
Cancer of the Stomach (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 11.68 | 17.14 | 7.85 | 10.66 | 15.98 | 6.99 | 17.05 | 24.76 | 11.46 |
| 1976 | 12.21 | 17.52 | 8.33 | 11.18 | 15.99 | 7.68 | 15.82 | 23.55 | 10.12 |
| 1977 | 11.54 | 16.58 | 7.87 | 10.34 | 15.02 | 6.98 | 17.34 | 24.68 | 11.63 |
| 1978 | 11.49 | 16.49 | 7.95 | 10.29 | 15.00 | 6.98 | 17.59 | 22.94 | 13.46 |
| 1979 | 12.00 | 17.40 | 8.25 | 10.76 | 15.58 | 7.49 | 18.51 | 28.20 | 11.73 |
| 1980 | 11.29 | 17.13 | 7.18 | 10.33 | 15.68 | 6.57 | 15.12 | 26.32 | 7.32 |
| 1981 | 11.14 | 16.60 | 7.39 | 10.06 | 15.07 | 6.70 | 17.61 | 27.84 | 10.42 |
| 1982 | 10.92 | 16.29 | 7.12 | 9.71 | 14.58 | 6.33 | 19.03 | 31.17 | 10.33 |
| 1983 | 10.95 | 15.86 | 7.46 | 9.54 | 13.69 | 6.60 | 18.01 | 27.78 | 11.16 |
| 1984 | 10.57 | 15.32 | 7.31 | 9.43 | 13.90 | 6.42 | 16.47 | 21.84 | 12.35 |
| 1985 | 10.22 | 15.09 | 6.88 | 8.95 | 13.35 | 5.98 | 16.44 | 23.50 | 11.34 |
| 1986 | 10.21 | 15.00 | 6.81 | 9.06 | 13.52 | 5.88 | 16.47 | 23.17 | 11.88 |
| 1987 | 10.20 | 15.06 | 6.81 | 8.99 | 13.27 | 5.96 | 17.06 | 26.68 | 10.69 |
| 1988 | 10.19 | 15.49 | 6.47 | 8.80 | 13.58 | 5.47 | 16.22 | 24.75 | 10.09 |
| 1989 | 10.00 | 15.15 | 6.33 | 8.73 | 13.47 | 5.35 | 15.78 | 23.16 | 10.66 |
| 1990 | 9.27 | 13.73 | 6.19 | 7.95 | 11.92 | 5.20 | 14.44 | 21.78 | 9.43 |
| 1991 | 9.70 | 14.19 | 6.58 | 8.30 | 12.32 | 5.51 | 16.49 | 25.62 | 10.52 |
| 1992 | 9.19 | 13.37 | 6.14 | 7.96 | 11.66 | 5.23 | 14.02 | 19.90 | 9.84 |
| 1993 | 9.01 | 13.61 | 5.78 | 7.66 | 11.61 | 4.87 | 14.37 | 23.70 | 8.38 |
| 1994 | 9.01 | 13.60 | 5.70 | 7.70 | 11.76 | 4.76 | 15.03 | 24.74 | 8.76 |
| 1995 | 8.33 | 12.57 | 5.23 | 7.25 | 11.24 | 4.35 | 13.33 | 18.88 | 9.50 |
| 1996 | 8.49 | 12.60 | 5.53 | 7.32 | 11.02 | 4.69 | 14.48 | 22.59 | 8.91 |
| 1997 | 8.61 | 12.71 | 5.62 | 7.16 | 10.80 | 4.50 | 15.68 | 22.35 | 10.92 |
| 1998 | 8.57 | 12.22 | 5.91 | 7.16 | 10.54 | 4.71 | 14.98 | 20.16 | 11.45 |
| 1999 | 8.60 | 12.07 | 6.03 | 7.40 | 10.52 | 5.05 | 13.00 | 18.17 | 9.84 |
| 2000 | 8.10 | 11.68 | 5.42 | 6.79 | 9.87 | 4.47 | 12.83 | 19.79 | 8.16 |
| 2001 | 7.73 | 11.20 | 5.11 | 6.44 | 9.59 | 4.02 | 12.91 | 18.30 | 9.01 |
| 2002 | 7.91 | 11.21 | 5.46 | 6.68 | 9.83 | 4.32 | 12.84 | 15.44 | 10.81 |
| 2003 | 7.69 | 10.73 | 5.37 | 6.50 | 9.24 | 4.35 | 13.03 | 17.90 | 9.72 |
| 2004 | 7.79 | 11.13 | 5.26 | 6.90 | 9.87 | 4.58 | 10.78 | 16.14 | 7.26 |
| 2005 | 7.31 | 10.45 | 4.89 | 6.33 | 9.07 | 4.14 | 10.94 | 17.14 | 7.01 |
| 2006 | 7.34 | 10.30 | 5.05 | 6.31 | 8.99 | 4.18 | 11.88 | 16.37 | 8.75 |
| 1975-2006 | 9.34 | 13.61 | 6.24 | 8.17 | 12.04 | 5.35 | 14.74 | 21.75 | 9.90 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 24.5
Cancer of the Stomach (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 8.51 | 12.27 | 5.94 | 7.93 | 11.37 | 5.62 | 13.87 | 21.05 | 8.61 |
| 1976 | 8.30 | 12.02 | 5.73 | 7.74 | 11.17 | 5.41 | 13.52 | 20.22 | 8.59 |
| 1977 | 7.91 | 11.48 | 5.45 | 7.33 | 10.68 | 5.05 | 13.36 | 19.46 | 9.00 |
| 1978 | 7.73 | 11.02 | 5.46 | 7.19 | 10.24 | 5.12 | 12.61 | 18.67 | 8.31 |
| 1979 | 7.56 | 10.98 | 5.23 | 6.99 | 10.17 | 4.85 | 12.86 | 18.73 | 8.71 |
| 1980 | 7.36 | 10.57 | 5.15 | 6.82 | 9.76 | 4.80 | 12.16 | 18.27 | 7.96 |
| 1981 | 7.25 | 10.58 | 4.98 | 6.68 | 9.80 | 4.56 | 12.48 | 18.06 | 8.60 |
| 1982 | 7.02 | 10.26 | 4.86 | 6.46 | 9.37 | 4.53 | 12.17 | 19.24 | 7.44 |
| 1983 | 6.83 | 9.90 | 4.76 | 6.24 | 9.02 | 4.38 | 12.27 | 18.39 | 8.08 |
| 1984 | 6.78 | 9.89 | 4.68 | 6.21 | 9.07 | 4.28 | 11.92 | 17.95 | 7.93 |
| 1985 | 6.51 | 9.54 | 4.44 | 5.96 | 8.75 | 4.07 | 11.54 | 17.03 | 7.82 |
| 1986 | 6.36 | 9.44 | 4.24 | 5.77 | 8.58 | 3.83 | 11.39 | 17.39 | 7.42 |
| 1987 | 6.18 | 9.11 | 4.18 | 5.62 | 8.32 | 3.77 | 11.29 | 16.97 | 7.55 |
| 1988 | 6.08 | 8.81 | 4.17 | 5.51 | 7.92 | 3.82 | 11.05 | 17.02 | 7.07 |
| 1989 | 6.21 | 9.18 | 4.16 | 5.60 | 8.28 | 3.77 | 11.49 | 17.73 | 7.33 |
| 1990 | 6.07 | 8.86 | 4.16 | 5.45 | 7.95 | 3.73 | 11.46 | 17.24 | 7.71 |
| 1991 | 6.00 | 8.86 | 4.01 | 5.39 | 7.97 | 3.59 | 11.55 | 17.64 | 7.55 |
| 1992 | 5.63 | 8.14 | 3.90 | 5.01 | 7.30 | 3.43 | 10.71 | 15.55 | 7.53 |
| 1993 | 5.62 | 8.19 | 3.83 | 5.03 | 7.34 | 3.41 | 10.70 | 16.26 | 7.11 |
| 1994 | 5.41 | 7.87 | 3.70 | 4.81 | 7.04 | 3.26 | 10.26 | 15.37 | 6.97 |
| 1995 | 5.35 | 7.70 | 3.71 | 4.72 | 6.89 | 3.21 | 10.36 | 14.89 | 7.37 |
| 1996 | 5.13 | 7.43 | 3.54 | 4.53 | 6.60 | 3.10 | 9.99 | 15.09 | 6.80 |
| 1997 | 4.94 | 7.16 | 3.37 | 4.34 | 6.33 | 2.92 | 9.66 | 14.64 | 6.50 |
| 1998 | 4.81 | 6.83 | 3.39 | 4.25 | 6.06 | 2.97 | 9.29 | 13.72 | 6.42 |
| 1999 | 4.64 | 6.59 | 3.25 | 4.05 | 5.78 | 2.79 | 9.25 | 13.73 | 6.47 |
| 2000 | 4.54 | 6.38 | 3.23 | 3.97 | 5.62 | 2.78 | 8.94 | 12.75 | 6.49 |
| 2001 | 4.36 | 6.08 | 3.09 | 3.83 | 5.37 | 2.70 | 8.29 | 12.00 | 5.83 |
| 2002 | 4.24 | 5.94 | 3.00 | 3.69 | 5.26 | 2.55 | 8.42 | 12.04 | 6.10 |
| 2003 | 4.13 | 5.71 | 2.96 | 3.62 | 5.01 | 2.58 | 7.82 | 11.43 | 5.41 |
| 2004 | 3.98 | 5.54 | 2.84 | 3.49 | 4.86 | 2.48 | 7.56 | 11.19 | 5.21 |
| 2005 | 3.79 | 5.25 | 2.71 | 3.30 | 4.56 | 2.36 | 7.22 | 10.71 | 4.96 |
| 2006 | 3.67 | 5.04 | 2.65 | 3.20 | 4.44 | 2.27 | 6.84 | 9.86 | 4.89 |
| 1975-2006 | 5.67 | 8.13 | 3.94 | 5.12 | 7.35 | 3.54 | 10.37 | 15.46 | 7.00 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 25.4
Cancer of the Testis (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

| Year of Diagnosis: | All Races, Males | | | White Males | | | Black Males | | |
|--------------------|------------------|------|------|-------------|------|------|-------------|------|------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| 1975 | 3.73 | 4.23 | 2.41 | 4.16 | 4.74 | 2.65 | - | - | - |
| 1976 | 3.44 | 4.18 | 1.51 | 3.76 | 4.54 | 1.72 | - | - | - |
| 1977 | 4.30 | 5.09 | 2.24 | 4.88 | 5.82 | 2.41 | - | - | - |
| 1978 | 3.58 | 4.25 | 1.84 | 3.96 | 4.66 | 2.11 | - | - | - |
| 1979 | 3.88 | 4.53 | 2.15 | 4.33 | 5.10 | 2.31 | - | - | - |
| 1980 | 4.35 | 5.28 | 1.89 | 4.92 | 5.99 | 2.11 | - | - | - |
| 1981 | 4.22 | 5.12 | 1.85 | 4.77 | 5.79 | 2.07 | - | - | - |
| 1982 | 4.40 | 5.22 | 2.25 | 5.00 | 5.98 | 2.44 | - | - | - |
| 1983 | 4.59 | 5.50 | 2.21 | 5.24 | 6.32 | 2.43 | - | - | - |
| 1984 | 4.39 | 5.36 | 1.86 | 5.03 | 6.15 | 2.08 | - | - | - |
| 1985 | 4.47 | 5.47 | 1.83 | 5.14 | 6.32 | 2.05 | - | - | - |
| 1986 | 4.82 | 5.83 | 2.17 | 5.59 | 6.82 | 2.39 | - | - | - |
| 1987 | 5.04 | 6.26 | 1.86 | 5.68 | 7.11 | 1.94 | - | - | - |
| 1988 | 4.63 | 5.83 | 1.46 | 5.45 | 6.88 | 1.72 | - | - | - |
| 1989 | 5.48 | 6.71 | 2.24 | 6.32 | 7.79 | 2.46 | - | - | - |
| 1990 | 5.10 | 6.26 | 2.06 | 6.00 | 7.39 | 2.35 | - | - | - |
| 1991 | 5.10 | 6.24 | 2.12 | 5.74 | 7.05 | 2.31 | - | - | - |
| 1992 | 5.15 | 6.42 | 1.81 | 6.07 | 7.62 | 2.03 | - | - | - |
| 1993 | 5.07 | 6.38 | 1.65 | 5.98 | 7.53 | 1.93 | - | - | - |
| 1994 | 5.47 | 6.75 | 2.12 | 6.27 | 7.78 | 2.30 | 1.23 | 1.54 | - |
| 1995 | 4.55 | 5.58 | 1.84 | 5.24 | 6.45 | 2.07 | 1.46 | 1.87 | - |
| 1996 | 5.25 | 6.60 | 1.73 | 6.19 | 7.83 | 1.89 | - | - | - |
| 1997 | 5.39 | 6.52 | 2.43 | 6.35 | 7.74 | 2.68 | 1.27 | - | - |
| 1998 | 5.59 | 7.11 | 1.62 | 6.60 | 8.44 | 1.79 | 1.52 | 1.97 | - |
| 1999 | 5.43 | 6.81 | 1.81 | 6.31 | 7.95 | 2.02 | 1.63 | 2.02 | - |
| 2000 | 5.73 | 7.16 | 1.99 | 6.74 | 8.42 | 2.33 | 1.52 | 1.86 | - |
| 2001 | 5.49 | 6.81 | 2.03 | 6.48 | 8.10 | 2.23 | 1.67 | 1.81 | - |
| 2002 | 5.72 | 7.16 | 1.94 | 6.86 | 8.60 | 2.29 | 1.37 | 1.78 | - |
| 2003 | 5.27 | 6.60 | 1.79 | 6.37 | 8.06 | 1.93 | 1.12 | - | - |
| 2004 | 5.85 | 7.23 | 2.25 | 7.18 | 8.94 | 2.59 | 1.03 | - | - |
| 2005 | 5.92 | 7.38 | 2.09 | 7.08 | 8.93 | 2.25 | 1.05 | 1.35 | - |
| 2006 | 5.52 | 6.85 | 2.05 | 6.56 | 8.17 | 2.33 | 1.57 | 1.65 | - |
| 1975-2006 | 5.01 | 6.16 | 1.98 | 5.79 | 7.16 | 2.20 | 1.10 | 1.29 | 0.62 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 25.5
Cancer of the Testis (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Age

| Year of Death: | All Races, Males | | | White Males | | | Black Males | | |
|----------------|------------------|------|------|-------------|------|------|-------------|------|------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| 1975 | 0.74 | 0.75 | 0.72 | 0.81 | 0.83 | 0.76 | 0.20 | - | - |
| 1976 | 0.71 | 0.72 | 0.69 | 0.77 | 0.80 | 0.69 | 0.36 | 0.21 | - |
| 1977 | 0.65 | 0.66 | 0.60 | 0.71 | 0.73 | 0.65 | 0.20 | - | - |
| 1978 | 0.56 | 0.57 | 0.53 | 0.60 | 0.61 | 0.57 | 0.22 | 0.22 | - |
| 1979 | 0.52 | 0.49 | 0.59 | 0.54 | 0.52 | 0.60 | 0.38 | 0.29 | - |
| 1980 | 0.45 | 0.44 | 0.47 | 0.49 | 0.49 | 0.50 | 0.18 | - | - |
| 1981 | 0.37 | 0.38 | 0.36 | 0.40 | 0.41 | 0.36 | 0.23 | - | - |
| 1982 | 0.40 | 0.39 | 0.42 | 0.43 | 0.43 | 0.44 | 0.19 | - | - |
| 1983 | 0.39 | 0.37 | 0.44 | 0.41 | 0.40 | 0.44 | 0.23 | 0.18 | - |
| 1984 | 0.35 | 0.33 | 0.39 | 0.37 | 0.36 | 0.40 | 0.19 | - | - |
| 1985 | 0.36 | 0.34 | 0.40 | 0.39 | 0.38 | 0.40 | 0.20 | - | - |
| 1986 | 0.32 | 0.32 | 0.34 | 0.35 | 0.35 | 0.35 | 0.17 | - | - |
| 1987 | 0.34 | 0.35 | 0.32 | 0.37 | 0.38 | 0.33 | 0.19 | 0.15 | - |
| 1988 | 0.32 | 0.29 | 0.40 | 0.35 | 0.32 | 0.42 | 0.16 | - | - |
| 1989 | 0.32 | 0.31 | 0.33 | 0.34 | 0.34 | 0.33 | 0.23 | 0.19 | - |
| 1990 | 0.27 | 0.30 | 0.21 | 0.30 | 0.33 | 0.22 | - | - | - |
| 1991 | 0.29 | 0.29 | 0.28 | 0.31 | 0.32 | 0.29 | 0.15 | - | - |
| 1992 | 0.28 | 0.28 | 0.30 | 0.31 | 0.31 | 0.32 | 0.14 | 0.15 | - |
| 1993 | 0.30 | 0.29 | 0.30 | 0.33 | 0.33 | 0.32 | 0.17 | 0.14 | - |
| 1994 | 0.27 | 0.28 | 0.23 | 0.29 | 0.31 | 0.24 | 0.14 | 0.14 | - |
| 1995 | 0.24 | 0.24 | 0.23 | 0.26 | 0.27 | 0.25 | - | - | - |
| 1996 | 0.26 | 0.26 | 0.27 | 0.29 | 0.29 | 0.29 | 0.12 | - | - |
| 1997 | 0.24 | 0.24 | 0.25 | 0.26 | 0.26 | 0.27 | 0.16 | 0.18 | - |
| 1998 | 0.28 | 0.27 | 0.29 | 0.30 | 0.30 | 0.30 | 0.20 | 0.13 | - |
| 1999 | 0.28 | 0.28 | 0.26 | 0.31 | 0.32 | 0.28 | 0.14 | 0.14 | - |
| 2000 | 0.25 | 0.25 | 0.23 | 0.27 | 0.28 | 0.24 | 0.16 | 0.16 | - |
| 2001 | 0.24 | 0.24 | 0.25 | 0.26 | 0.27 | 0.25 | 0.16 | - | - |
| 2002 | 0.28 | 0.27 | 0.33 | 0.31 | 0.30 | 0.34 | 0.17 | 0.16 | - |
| 2003 | 0.25 | 0.24 | 0.27 | 0.27 | 0.26 | 0.29 | 0.18 | 0.12 | - |
| 2004 | 0.25 | 0.25 | 0.25 | 0.27 | 0.27 | 0.27 | 0.15 | 0.15 | - |
| 2005 | 0.25 | 0.24 | 0.28 | 0.26 | 0.25 | 0.29 | 0.19 | 0.18 | - |
| 2006 | 0.24 | 0.23 | 0.27 | 0.27 | 0.26 | 0.30 | 0.13 | 0.15 | - |
| 1975-2006 | 0.34 | 0.34 | 0.34 | 0.37 | 0.38 | 0.36 | 0.18 | 0.15 | 0.26 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 26.4
Cancer of the Thyroid (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 4.85 | 3.14 | 6.45 | 4.66 | 2.80 | 6.37 | 2.97 | - | 4.21 |
| 1976 | 4.80 | 2.95 | 6.57 | 4.67 | 2.88 | 6.39 | 3.41 | - | 5.03 |
| 1977 | 5.44 | 3.49 | 7.32 | 5.35 | 3.43 | 7.22 | 3.74 | - | 5.40 |
| 1978 | 5.09 | 3.14 | 6.94 | 4.95 | 2.97 | 6.83 | 2.92 | - | 4.00 |
| 1979 | 4.48 | 2.67 | 6.17 | 4.43 | 2.57 | 6.19 | 3.24 | - | 4.30 |
| 1980 | 4.33 | 2.39 | 6.14 | 4.31 | 2.28 | 6.22 | 2.37 | - | 3.43 |
| 1981 | 4.41 | 2.51 | 6.23 | 4.38 | 2.47 | 6.27 | 2.38 | - | 3.41 |
| 1982 | 4.63 | 3.00 | 6.15 | 4.55 | 2.87 | 6.13 | 3.29 | - | 4.34 |
| 1983 | 4.70 | 2.80 | 6.48 | 4.73 | 2.61 | 6.71 | 3.02 | 2.15 | 3.77 |
| 1984 | 4.84 | 2.65 | 6.91 | 4.72 | 2.57 | 6.79 | 3.02 | - | 4.08 |
| 1985 | 5.12 | 3.08 | 7.09 | 5.09 | 3.10 | 7.05 | 2.85 | - | 3.64 |
| 1986 | 5.32 | 3.07 | 7.48 | 5.22 | 3.10 | 7.32 | 3.35 | - | 5.09 |
| 1987 | 5.04 | 2.80 | 7.13 | 5.04 | 2.87 | 7.07 | 2.82 | - | 4.30 |
| 1988 | 4.94 | 2.94 | 6.90 | 4.90 | 3.00 | 6.81 | 2.53 | - | 3.21 |
| 1989 | 5.35 | 2.97 | 7.64 | 5.47 | 3.21 | 7.71 | 2.51 | - | 4.01 |
| 1990 | 5.49 | 2.91 | 7.96 | 5.55 | 2.96 | 8.10 | 3.76 | 2.28 | 4.91 |
| 1991 | 5.46 | 3.18 | 7.65 | 5.66 | 3.30 | 7.97 | 3.19 | - | 4.20 |
| 1992 | 5.86 | 3.52 | 8.09 | 5.88 | 3.53 | 8.17 | 4.20 | 3.09 | 5.13 |
| 1993 | 5.64 | 3.58 | 7.64 | 5.75 | 3.68 | 7.79 | 3.38 | 2.07 | 4.56 |
| 1994 | 6.07 | 3.38 | 8.72 | 6.21 | 3.43 | 8.98 | 3.75 | 1.78 | 5.35 |
| 1995 | 6.20 | 3.37 | 8.95 | 6.38 | 3.59 | 9.16 | 3.06 | - | 4.50 |
| 1996 | 6.49 | 3.47 | 9.45 | 6.70 | 3.61 | 9.80 | 3.83 | - | 5.75 |
| 1997 | 6.76 | 3.65 | 9.76 | 7.02 | 3.76 | 10.28 | 3.98 | 2.44 | 5.24 |
| 1998 | 6.94 | 3.71 | 10.04 | 7.16 | 3.95 | 10.33 | 3.85 | 1.50 | 5.79 |
| 1999 | 7.32 | 3.86 | 10.63 | 7.54 | 4.08 | 10.97 | 4.57 | 1.54 | 7.03 |
| 2000 | 7.58 | 4.01 | 11.06 | 8.01 | 4.34 | 11.72 | 4.60 | 2.18 | 6.60 |
| 2001 | 8.21 | 4.30 | 12.03 | 8.57 | 4.55 | 12.63 | 4.82 | 2.67 | 6.57 |
| 2002 | 9.13 | 4.86 | 13.30 | 9.62 | 5.18 | 14.12 | 5.42 | 2.77 | 7.56 |
| 2003 | 9.51 | 4.69 | 14.21 | 10.16 | 5.02 | 15.33 | 5.21 | 2.74 | 7.37 |
| 2004 | 9.95 | 5.13 | 14.68 | 10.59 | 5.44 | 15.82 | 6.16 | 3.24 | 8.61 |
| 2005 | 10.68 | 5.63 | 15.61 | 11.58 | 6.24 | 16.96 | 5.75 | 2.76 | 8.35 |
| 2006 | 11.03 | 5.66 | 16.29 | 11.79 | 6.18 | 17.49 | 6.22 | 2.62 | 9.10 |
| 1975-2006 | 6.56 | 3.63 | 9.37 | 6.68 | 3.73 | 9.58 | 3.97 | 2.01 | 5.60 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 26.5
Cancer of the Thyroid (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 0.55 | 0.39 | 0.65 | 0.55 | 0.40 | 0.65 | 0.47 | 0.35 | 0.56 |
| 1976 | 0.56 | 0.49 | 0.61 | 0.56 | 0.49 | 0.60 | 0.52 | 0.42 | 0.60 |
| 1977 | 0.57 | 0.46 | 0.63 | 0.56 | 0.46 | 0.62 | 0.67 | 0.53 | 0.74 |
| 1978 | 0.55 | 0.44 | 0.61 | 0.55 | 0.45 | 0.61 | 0.48 | 0.33 | 0.60 |
| 1979 | 0.53 | 0.43 | 0.58 | 0.53 | 0.45 | 0.58 | 0.47 | 0.28 | 0.60 |
| 1980 | 0.48 | 0.38 | 0.53 | 0.49 | 0.40 | 0.54 | 0.42 | 0.29 | 0.51 |
| 1981 | 0.50 | 0.42 | 0.55 | 0.50 | 0.43 | 0.54 | 0.48 | 0.26 | 0.62 |
| 1982 | 0.48 | 0.37 | 0.53 | 0.47 | 0.37 | 0.53 | 0.48 | 0.38 | 0.54 |
| 1983 | 0.44 | 0.34 | 0.50 | 0.45 | 0.35 | 0.50 | 0.41 | 0.28 | 0.49 |
| 1984 | 0.47 | 0.38 | 0.52 | 0.47 | 0.39 | 0.51 | 0.47 | - | 0.64 |
| 1985 | 0.45 | 0.39 | 0.48 | 0.44 | 0.39 | 0.46 | 0.49 | 0.34 | 0.58 |
| 1986 | 0.46 | 0.39 | 0.49 | 0.45 | 0.39 | 0.49 | 0.48 | 0.34 | 0.57 |
| 1987 | 0.45 | 0.40 | 0.48 | 0.45 | 0.41 | 0.47 | 0.48 | 0.33 | 0.58 |
| 1988 | 0.44 | 0.41 | 0.45 | 0.44 | 0.42 | 0.44 | 0.44 | 0.33 | 0.50 |
| 1989 | 0.44 | 0.41 | 0.44 | 0.43 | 0.41 | 0.43 | 0.51 | 0.40 | 0.56 |
| 1990 | 0.44 | 0.37 | 0.48 | 0.44 | 0.38 | 0.47 | 0.45 | 0.30 | 0.54 |
| 1991 | 0.43 | 0.38 | 0.46 | 0.42 | 0.39 | 0.43 | 0.50 | 0.37 | 0.58 |
| 1992 | 0.46 | 0.38 | 0.50 | 0.46 | 0.39 | 0.48 | 0.41 | 0.30 | 0.48 |
| 1993 | 0.46 | 0.39 | 0.50 | 0.46 | 0.40 | 0.48 | 0.46 | 0.26 | 0.58 |
| 1994 | 0.42 | 0.38 | 0.44 | 0.42 | 0.38 | 0.44 | 0.37 | 0.30 | 0.40 |
| 1995 | 0.44 | 0.41 | 0.44 | 0.44 | 0.43 | 0.44 | 0.40 | 0.27 | 0.46 |
| 1996 | 0.45 | 0.42 | 0.47 | 0.45 | 0.43 | 0.46 | 0.44 | 0.33 | 0.49 |
| 1997 | 0.46 | 0.45 | 0.46 | 0.45 | 0.45 | 0.44 | 0.49 | 0.37 | 0.55 |
| 1998 | 0.44 | 0.38 | 0.47 | 0.43 | 0.39 | 0.45 | 0.43 | 0.27 | 0.53 |
| 1999 | 0.45 | 0.44 | 0.45 | 0.45 | 0.45 | 0.44 | 0.44 | 0.39 | 0.47 |
| 2000 | 0.48 | 0.47 | 0.48 | 0.47 | 0.48 | 0.45 | 0.51 | 0.34 | 0.59 |
| 2001 | 0.48 | 0.47 | 0.48 | 0.47 | 0.48 | 0.46 | 0.51 | 0.41 | 0.58 |
| 2002 | 0.47 | 0.45 | 0.48 | 0.48 | 0.46 | 0.48 | 0.47 | 0.39 | 0.52 |
| 2003 | 0.45 | 0.42 | 0.46 | 0.46 | 0.45 | 0.46 | 0.37 | 0.28 | 0.44 |
| 2004 | 0.47 | 0.47 | 0.47 | 0.47 | 0.48 | 0.46 | 0.44 | 0.38 | 0.47 |
| 2005 | 0.48 | 0.47 | 0.49 | 0.47 | 0.49 | 0.46 | 0.51 | 0.30 | 0.63 |
| 2006 | 0.49 | 0.46 | 0.51 | 0.49 | 0.48 | 0.49 | 0.47 | 0.27 | 0.59 |
| 1975-2006 | 0.47 | 0.42 | 0.50 | 0.47 | 0.43 | 0.48 | 0.47 | 0.33 | 0.55 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 27.4
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 19.31 | 34.35 | 8.88 | 20.21 | 36.37 | 9.16 | 10.56 | 16.34 | 6.26 |
| 1976 | 19.70 | 34.39 | 9.44 | 20.73 | 36.71 | 9.73 | 12.48 | 18.16 | 8.59 |
| 1977 | 18.99 | 33.39 | 8.88 | 20.00 | 35.58 | 9.18 | 13.45 | 22.67 | 7.06 |
| 1978 | 19.99 | 35.37 | 9.44 | 21.08 | 37.86 | 9.73 | 11.73 | 19.36 | 6.39 |
| 1979 | 19.98 | 35.35 | 9.38 | 21.30 | 38.18 | 9.84 | 10.35 | 15.61 | 6.54 |
| 1980 | 20.42 | 37.14 | 8.91 | 21.54 | 39.85 | 9.06 | 13.33 | 18.23 | 9.98 |
| 1981 | 20.69 | 36.34 | 9.75 | 21.95 | 38.76 | 10.31 | 11.60 | 20.72 | 5.48 |
| 1982 | 20.08 | 35.81 | 9.52 | 21.38 | 38.32 | 10.15 | 12.19 | 21.77 | 5.60 |
| 1983 | 20.05 | 36.02 | 9.01 | 21.33 | 38.70 | 9.43 | 12.17 | 18.70 | 7.57 |
| 1984 | 20.85 | 37.71 | 9.17 | 22.35 | 40.89 | 9.63 | 11.42 | 18.77 | 6.24 |
| 1985 | 20.68 | 36.38 | 9.87 | 22.09 | 39.29 | 10.34 | 12.99 | 20.33 | 8.02 |
| 1986 | 21.04 | 37.94 | 9.43 | 22.44 | 40.82 | 9.95 | 12.94 | 22.17 | 6.63 |
| 1987 | 21.66 | 39.52 | 9.24 | 23.18 | 42.83 | 9.58 | 13.28 | 21.78 | 7.50 |
| 1988 | 20.85 | 38.08 | 8.99 | 22.72 | 41.85 | 9.68 | 11.29 | 17.99 | 6.45 |
| 1989 | 21.03 | 37.42 | 9.64 | 22.74 | 40.94 | 10.20 | 11.32 | 17.49 | 7.16 |
| 1990 | 21.07 | 37.48 | 9.63 | 22.60 | 40.75 | 10.01 | 12.69 | 18.59 | 8.75 |
| 1991 | 20.91 | 37.62 | 9.32 | 22.69 | 41.25 | 9.88 | 12.92 | 19.69 | 8.50 |
| 1992 | 21.25 | 37.50 | 9.82 | 22.95 | 40.72 | 10.51 | 12.48 | 20.82 | 6.89 |
| 1993 | 21.23 | 37.63 | 9.70 | 22.85 | 40.57 | 10.38 | 14.11 | 24.40 | 7.76 |
| 1994 | 20.79 | 36.85 | 9.45 | 22.60 | 40.23 | 10.18 | 12.28 | 20.01 | 6.91 |
| 1995 | 20.64 | 36.24 | 9.55 | 22.44 | 39.50 | 10.28 | 11.87 | 18.30 | 7.39 |
| 1996 | 20.77 | 36.58 | 9.55 | 22.62 | 39.96 | 10.28 | 12.21 | 19.78 | 7.28 |
| 1997 | 21.03 | 37.06 | 9.55 | 22.91 | 40.59 | 10.19 | 12.54 | 19.97 | 7.98 |
| 1998 | 21.56 | 38.44 | 9.45 | 23.58 | 42.11 | 10.18 | 11.80 | 20.18 | 6.38 |
| 1999 | 21.73 | 38.15 | 9.97 | 23.60 | 41.55 | 10.60 | 14.58 | 22.63 | 9.44 |
| 2000 | 21.76 | 38.66 | 9.49 | 23.80 | 42.28 | 10.25 | 13.08 | 20.13 | 8.53 |
| 2001 | 21.58 | 38.13 | 9.61 | 23.85 | 42.03 | 10.58 | 11.50 | 18.44 | 7.01 |
| 2002 | 21.11 | 36.76 | 9.68 | 23.18 | 40.28 | 10.54 | 12.42 | 17.74 | 8.76 |
| 2003 | 21.30 | 37.19 | 9.65 | 23.49 | 40.93 | 10.53 | 12.52 | 20.51 | 7.17 |
| 2004 | 21.15 | 37.22 | 9.40 | 23.30 | 40.94 | 10.22 | 13.01 | 21.16 | 7.95 |
| 2005 | 21.38 | 37.57 | 9.47 | 23.39 | 41.29 | 10.06 | 13.13 | 21.02 | 7.80 |
| 2006 | 20.46 | 36.05 | 8.94 | 22.38 | 39.45 | 9.61 | 11.14 | 16.85 | 7.29 |
| 1975-2006 | 20.85 | 37.03 | 9.42 | 22.50 | 40.25 | 10.00 | 12.43 | 19.78 | 7.51 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 27.5
Cancer of the Urinary Bladder (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Sex

| Year of Death: | All Races | | | Whites | | | Blacks | | |
|----------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975 | 5.50 | 9.77 | 2.76 | 5.56 | 10.11 | 2.68 | 4.96 | 6.85 | 3.63 |
| 1976 | 5.59 | 9.79 | 2.88 | 5.64 | 10.05 | 2.83 | 5.29 | 7.80 | 3.54 |
| 1977 | 5.52 | 9.67 | 2.89 | 5.58 | 9.94 | 2.83 | 5.19 | 7.57 | 3.57 |
| 1978 | 5.44 | 9.54 | 2.86 | 5.51 | 9.83 | 2.82 | 4.94 | 7.22 | 3.40 |
| 1979 | 5.25 | 9.34 | 2.71 | 5.30 | 9.59 | 2.66 | 5.00 | 7.61 | 3.28 |
| 1980 | 5.17 | 9.16 | 2.73 | 5.24 | 9.51 | 2.66 | 4.72 | 6.21 | 3.71 |
| 1981 | 5.07 | 9.02 | 2.66 | 5.13 | 9.28 | 2.61 | 4.89 | 7.20 | 3.40 |
| 1982 | 4.99 | 8.95 | 2.58 | 5.04 | 9.23 | 2.50 | 4.86 | 6.83 | 3.56 |
| 1983 | 4.88 | 8.79 | 2.54 | 4.93 | 9.04 | 2.50 | 4.72 | 7.17 | 3.19 |
| 1984 | 4.74 | 8.49 | 2.49 | 4.78 | 8.73 | 2.43 | 4.74 | 6.89 | 3.39 |
| 1985 | 4.67 | 8.43 | 2.44 | 4.73 | 8.67 | 2.40 | 4.48 | 6.86 | 3.02 |
| 1986 | 4.53 | 8.10 | 2.41 | 4.60 | 8.41 | 2.34 | 4.25 | 5.83 | 3.29 |
| 1987 | 4.41 | 7.87 | 2.34 | 4.46 | 8.09 | 2.28 | 4.37 | 6.50 | 3.11 |
| 1988 | 4.41 | 7.76 | 2.41 | 4.47 | 8.02 | 2.34 | 4.30 | 5.89 | 3.33 |
| 1989 | 4.48 | 7.99 | 2.37 | 4.52 | 8.18 | 2.31 | 4.61 | 7.02 | 3.20 |
| 1990 | 4.49 | 7.97 | 2.42 | 4.55 | 8.22 | 2.37 | 4.44 | 6.44 | 3.24 |
| 1991 | 4.42 | 7.89 | 2.34 | 4.48 | 8.14 | 2.28 | 4.40 | 6.58 | 3.12 |
| 1992 | 4.46 | 7.89 | 2.43 | 4.52 | 8.11 | 2.38 | 4.35 | 6.66 | 3.08 |
| 1993 | 4.47 | 8.05 | 2.31 | 4.55 | 8.32 | 2.27 | 4.17 | 6.36 | 2.93 |
| 1994 | 4.47 | 7.86 | 2.42 | 4.54 | 8.13 | 2.37 | 4.34 | 6.36 | 3.19 |
| 1995 | 4.35 | 7.78 | 2.28 | 4.45 | 8.08 | 2.23 | 4.05 | 5.77 | 3.09 |
| 1996 | 4.41 | 7.79 | 2.37 | 4.51 | 8.08 | 2.33 | 4.16 | 5.89 | 3.14 |
| 1997 | 4.39 | 7.59 | 2.42 | 4.50 | 7.88 | 2.40 | 4.00 | 5.88 | 2.98 |
| 1998 | 4.38 | 7.71 | 2.32 | 4.49 | 8.02 | 2.29 | 3.95 | 5.58 | 3.03 |
| 1999 | 4.36 | 7.66 | 2.28 | 4.47 | 7.95 | 2.24 | 4.01 | 5.84 | 2.96 |
| 2000 | 4.32 | 7.59 | 2.26 | 4.47 | 7.93 | 2.27 | 3.73 | 5.66 | 2.69 |
| 2001 | 4.33 | 7.56 | 2.25 | 4.46 | 7.91 | 2.23 | 3.71 | 5.10 | 2.91 |
| 2002 | 4.39 | 7.51 | 2.37 | 4.53 | 7.85 | 2.35 | 3.93 | 5.53 | 3.04 |
| 2003 | 4.26 | 7.38 | 2.20 | 4.40 | 7.71 | 2.20 | 3.70 | 5.44 | 2.68 |
| 2004 | 4.37 | 7.59 | 2.25 | 4.55 | 8.00 | 2.25 | 3.60 | 5.02 | 2.76 |
| 2005 | 4.35 | 7.58 | 2.22 | 4.52 | 7.94 | 2.22 | 3.70 | 5.74 | 2.55 |
| 2006 | 4.34 | 7.57 | 2.18 | 4.51 | 7.95 | 2.17 | 3.79 | 5.55 | 2.76 |
| 1975-2006 | 4.61 | 8.12 | 2.42 | 4.70 | 8.42 | 2.38 | 4.30 | 6.27 | 3.12 |

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.