

Table 1.9
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 1997-2006

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.4*	-2.4*	-0.4*	-2.0*	-2.6*	-1.4*
Oral Cavity & Pharynx:	-2.7*	-3.6*	-1.1	-3.5*	-3.0*	-4.7*
Lip	-	-	-	-	-	-
Tongue	-2.2	-3.0*	-0.7	-4.5*	-4.3*	-4.4*
Salivary gland	1.3	-1.0	3.1	-2.2	-2.5	-2.8
Floor of mouth	-9.1*	-10.0*	-	-11.1*	-9.3*	-
Gum & other oral cavity	-2.3	-4.9*	1.8	-6.2*	-6.1*	-6.8*
Nasopharynx	-0.6	-	-	-2.3	-0.9	-4.6
Tonsil	-3.4	-3.5	-	-3.1*	-3.0*	-3.3
Oropharynx	0.9	1.9	-	-1.2	-0.7	-1.8
Hypopharynx	-2.5	-3.2	-	-5.2*	-4.0	-
Other oral cavity & pharynx	-	-	-	-2.0	-1.6	-3.2
Digestive System:	-1.1*	-1.4*	-0.9*	-1.7*	-1.7*	-1.8*
Esophagus	-4.3*	-4.1*	-4.7*	-4.4*	-4.6*	-4.1*
Stomach	-3.3*	-3.2*	-3.1*	-3.7*	-3.9*	-3.5*
Small intestine	-0.6	-2.1	0.2	-1.2	-1.2	-1.1
Colon & Rectum:	-1.5*	-1.8*	-1.3*	-2.0*	-1.7*	-2.4*
Colon	-1.6*	-1.9*	-1.3*	-	-	-
Rectum	-1.4*	-1.6	-1.3	-	-	-
Anus, anal canal & anorectum	3.0	1.5	4.1*	1.0	-0.3	2.4
Liver & intrahep. bile duct:	3.3*	3.4*	2.4	2.0*	2.6*	0.4
Liver	3.9*	4.3*	2.3	1.6*	2.5*	-0.6
Intrahepatic bile duct	-3.8	-	-	4.2*	4.0*	4.2*
Gallbladder	1.6	-	0.6	-0.3	0.6	-0.5
Other biliary	4.2*	3.9	4.0*	-1.9	-2.6	-1.6
Pancreas	-0.2	-0.8	0.2	-0.7*	-1.0*	-0.6*
Retroperitoneum	-	-	-	-4.0	-	-
Peritoneum, omentum & mesentery	-	-	-	2.3	-	2.4
Other digestive system	6.1	-	-	-0.5	0.5	-1.4
Respiratory System:	-1.9*	-3.3*	0.0	-1.9*	-2.9*	-0.3
Nose, nasal cavity & middle ear	-3.7	-	-	-5.7*	-4.7	-6.1
Larynx	-3.8*	-3.2*	-5.8*	-3.3*	-3.2*	-3.2*
Lung & bronchus	-1.8*	-3.3*	0.3	-1.8*	-2.9*	-0.2
Pleura	-	-	-	-7.7*	-	-
Trachea & other respiratory organs	-	-	-	-8.1*	-	-
Bones & joints	0.5	-2.3	4.1	0.8	-0.8	2.5
Soft tissue (including heart)	-0.9	-2.8	0.6	-3.7*	-3.6*	-3.6*
Skin (excl. basal & squamous):	-3.0	-7.3*	0.8	-2.9*	-1.9	-4.1*
Melanoma of the skin	-4.4*	-8.8*	-	-2.1	1.2	-4.3*
Other non-epithelial skin	-2.0	-5.4	1.5	-3.7*	-3.5*	-3.5
Breast	-0.7*	-	-0.6*	-1.6*	-3.5*	-1.5*
Breast (<i>in situ</i>)	1.5*	-	1.4*	-	-	-

^a The APC is the Annual Percent Change over the time interval.
SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.9 - continued
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 1997-2006

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	-1.1	-	-1.0	-1.3*	-	-1.2*
Cervix uteri	-5.3*	-	-5.2*	-4.4*	-	-4.3*
Corpus uteri	0.9	-	1.0	-0.2	-	0.0
Uterus, NOS	-	-	-	1.1*	-	1.3*
Ovary ^c	-1.2	-	-1.0	-0.7	-	-0.7
Vagina	-1.1	-	-0.7	-1.5	-	-1.6
Vulva	-0.2	-	0.0	-2.4	-	-2.1
Other female genital system	-1.6	-	-1.7	-2.8	-	-2.6
Male Genital System:	-2.7*	-3.0*	-	-3.8*	-4.2*	-
Prostate	-2.7*	-3.0*	-	-3.9*	-4.2*	-
Testis	-1.8	-2.3	-	-0.2	-0.6	-
Penis	-	-	-	-1.0	-1.5	-
Other male genital system	-	-	-	-	-	-
Urinary System:	0.6	0.2	0.9	-0.8*	-0.7*	-1.0*
Urinary bladder	-0.1	-0.6	0.3	-0.9*	-0.6	-1.3*
Kidney & renal pelvis	1.5	1.4	1.6	-0.8*	-0.9*	-0.7
Ureter	-	-	-	-	-	-
Other urinary system	-	-	-	0.5	-	0.8
Eye & Orbit	-	-	-	-	-	-
Brain & Nervous System: ^d	-0.4	-1.0	0.1	-1.7*	-0.9	-2.4*
Brain	-0.4	-1.3	0.5	-	-	-
Cranial nerves & other nervous system	-	-	-	-	-	-
Endocrine System:	6.1*	5.4*	6.4*	-0.4	-1.3	0.0
Thyroid	6.4*	5.3*	6.8*	0.0	-1.7	0.7
Other endocrine & thymus	4.3*	-	-	-0.9	-0.9	-1.1
Lymphoma:	0.5	0.1	0.9	-2.8*	-3.2*	-2.4*
Hodgkin lymphoma	2.0	1.9	2.3	-2.2	-2.1	-2.3
Non-Hodgkin lymphoma	0.3	-0.2	0.7	-2.8*	-3.3*	-2.4*
Myeloma	-0.2	0.0	-0.4	-1.9*	-1.7*	-2.3*
Leukemia:	-0.7	-1.1	-0.3	-1.2*	-1.1*	-1.4*
Lymphocytic:	-0.1	-0.3	0.2	-1.4*	-0.8	-2.3*
Acute lymphocytic	1.0	-0.7	2.7	-0.8	-1.5	-0.5
Chronic lymphocytic	-0.5	-0.6	-0.1	-1.3*	-0.4	-2.8*
Other lymphocytic	-	-	-	-2.6	-2.6	-
Myeloid & Monocytic:	-1.5	-2.6*	-0.6	-1.0*	-1.1	-0.7
Acute myeloid	-0.7	-1.9	0.0	1.2	1.2	1.4*
Chronic myeloid	-2.8*	-4.4*	-0.7	-9.0*	-8.6*	-9.3*
Acute monocytic	-	-	-	-	-	-
Other myeloid & monocytic	-	-	-	4.2	1.3	-
Other leukemia:	1.2	-	-	-1.3*	-1.2	-1.7
Other acute leukemia	-	-	-	-4.5*	-3.4*	-6.0*
Aleukemic, subleukemic & NOS	6.7	-	-	0.5	0.1	0.9
Kaposi Sarcoma ^e	-5.7*	-5.5*	-	-	-	-
Mesothelioma ^e	-5.9*	-	-	-	-	-
Ill-defined & unspecified	-4.6*	-5.7*	-4.0*	-1.8*	-1.6	-2.0*

^a The APC is the Annual Percent Change over the time interval.
SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.