


Table 2.18
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	173.79	0.10		
High Five States				
Kentucky	207.09 ^b	0.95	(01)	19.16 ^c
Mississippi	200.31 ^b	1.15	(02)	15.26 ^c
West Virginia	200.29 ^b	1.32	(03)	15.25 ^c
Louisiana	198.64 ^b	0.94	(04)	14.30 ^c
Arkansas	196.92 ^b	1.10	(05)	13.31 ^c
Low Five States				
New Mexico	153.00 ^b	1.21	(47)	-11.96 ^c
Arizona	152.89 ^b	0.68	(48)	-12.02 ^c
Colorado	149.94 ^b	0.82	(49)	-13.72 ^c
Hawaii	141.63 ^b	1.36	(50)	-18.50 ^c
Utah	128.51 ^b	1.13	(51)	-26.05 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	193.97 ^b	0.87	(07)	11.61 ^c	Montana	163.76 ^b	1.69	(42)	-5.77 ^c
Alaska	181.58	2.98	(18)	4.49	Nebraska	168.23 ^b	1.30	(35)	-3.19
Arizona	152.89 ^b	0.68	(48)	-12.02 ^c	Nevada	176.03	1.21	(25)	1.29
Arkansas	196.92 ^b	1.10	(05)	13.31 ^c	New Hampshire	176.21	1.57	(24)	1.39
California	157.70 ^b	0.30	(46)	-9.26 ^c	New Jersey	171.48 ^b	0.60	(34)	-1.32
Colorado	149.94 ^b	0.82	(49)	-13.72 ^c	New Mexico	153.00 ^b	1.21	(47)	-11.96 ^c
Connecticut	162.88 ^b	0.90	(43)	-6.27 ^c	New York	164.30 ^b	0.39	(40)	-5.46 ^c
Delaware	185.45 ^b	1.93	(14)	6.71 ^c	North Carolina	179.48 ^b	0.61	(20)	3.27
D.C.	192.87 ^b	2.63	(08)	10.98 ^c	North Dakota	161.01 ^b	2.04	(45)	-7.35 ^c
Florida	165.51 ^b	0.37	(39)	-4.76	Ohio	189.73 ^b	0.54	(11)	9.18 ^c
Georgia	176.39 ^b	0.65	(22)	1.50	Oklahoma	192.45 ^b	0.98	(09)	10.74 ^c
Hawaii	141.63 ^b	1.36	(50)	-18.50 ^c	Oregon	174.51	0.91	(29)	0.42
Idaho	161.61 ^b	1.46	(44)	-7.00 ^c	Pennsylvania	181.97 ^b	0.48	(17)	4.71
Illinois	181.11 ^b	0.53	(19)	4.21	Rhode Island	174.68	1.69	(27)	0.51
Indiana	190.08 ^b	0.75	(10)	9.37 ^c	South Carolina	185.11 ^b	0.88	(15)	6.52 ^c
Iowa	173.06	0.98	(32)	-0.42	South Dakota	167.62	1.90	(36)	-3.55
Kansas	172.67	1.07	(33)	-0.64	Tennessee	195.93 ^b	0.77	(06)	12.74 ^c
Kentucky	207.09 ^b	0.95	(01)	19.16 ^c	Texas	166.91 ^b	0.40	(37)	-3.96
Louisiana	198.64 ^b	0.94	(04)	14.30 ^c	Utah	128.51 ^b	1.13	(51)	-26.05 ^c
Maine	185.55 ^b	1.50	(13)	6.77 ^c	Vermont	178.31	2.23	(21)	2.60
Maryland	174.69	0.78	(26)	0.52	Virginia	176.32 ^b	0.67	(23)	1.46
Massachusetts	173.44	0.69	(30)	-0.20	Washington	173.10	0.73	(31)	-0.39
Michigan	182.41 ^b	0.58	(16)	4.97	West Virginia	200.29 ^b	1.32	(03)	15.25 ^c
Minnesota	166.72 ^b	0.78	(38)	-4.07	Wisconsin	174.62	0.75	(28)	0.48
Mississippi	200.31 ^b	1.15	(02)	15.26 ^c	Wyoming	164.12 ^b	2.43	(41)	-5.56 ^c
Missouri	185.91 ^b	0.75	(12)	6.98 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.19
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	211.61	0.18		
High Five States				
Mississippi	264.72 ^b	2.10	(01)	25.10 ^c
Kentucky	257.53 ^b	1.66	(02)	21.70 ^c
Alabama	251.64 ^b	1.55	(03)	18.92 ^c
Tennessee	251.15 ^b	1.36	(04)	18.68 ^c
Louisiana	250.75 ^b	1.65	(05)	18.50 ^c
Low Five States				
New Mexico	183.76 ^b	2.00	(47)	-13.16 ^c
Arizona	182.62 ^b	1.11	(48)	-13.70 ^c
Colorado	177.70 ^b	1.39	(49)	-16.02 ^c
Hawaii	175.77 ^b	2.27	(50)	-16.94 ^c
Utah	154.64 ^b	1.88	(51)	-26.92 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	251.64 ^b	1.55	(03)	18.92 ^c	Montana	192.45 ^b	2.74	(43)	-9.05 ^c
Alaska	215.43	4.99	(23)	1.81	Nebraska	204.18	2.19	(35)	-3.51
Arizona	182.62 ^b	1.11	(48)	-13.70 ^c	Nevada	206.94	1.98	(31)	-2.20
Arkansas	248.67 ^b	1.91	(06)	17.52 ^c	New Hampshire	211.07	2.64	(29)	-0.26
California	186.79 ^b	0.50	(46)	-11.73 ^c	New Jersey	203.67 ^b	1.02	(36)	-3.75
Colorado	177.70 ^b	1.39	(49)	-16.02 ^c	New Mexico	183.76 ^b	2.00	(47)	-13.16 ^c
Connecticut	195.69 ^b	1.51	(42)	-7.52 ^c	New York	196.58 ^b	0.67	(41)	-7.10 ^c
Delaware	222.77	3.25	(17)	5.28 ^c	North Carolina	227.72 ^b	1.08	(14)	7.61 ^c
D.C.	234.84 ^b	4.60	(11)	10.98 ^c	North Dakota	200.04	3.43	(40)	-5.47 ^c
Florida	200.30 ^b	0.61	(39)	-5.35 ^c	Ohio	232.75 ^b	0.93	(12)	9.99 ^c
Georgia	222.94 ^b	1.18	(16)	5.36 ^c	Oklahoma	238.19 ^b	1.68	(08)	12.56 ^c
Hawaii	175.77 ^b	2.27	(50)	-16.94 ^c	Oregon	206.57	1.50	(32)	-2.38
Idaho	192.21 ^b	2.40	(45)	-9.17 ^c	Pennsylvania	222.21 ^b	0.82	(18)	5.01 ^c
Illinois	218.90 ^b	0.90	(20)	3.44	Rhode Island	217.48	2.93	(21)	2.77
Indiana	235.76 ^b	1.30	(10)	11.41 ^c	South Carolina	236.13 ^b	1.55	(09)	11.59 ^c
Iowa	211.07	1.65	(28)	-0.26	South Dakota	201.40	3.15	(37)	-4.82
Kansas	211.27	1.80	(27)	-0.16	Tennessee	251.15 ^b	1.36	(04)	18.68 ^c
Kentucky	257.53 ^b	1.66	(02)	21.70 ^c	Texas	205.16 ^b	0.69	(34)	-3.05
Louisiana	250.75 ^b	1.65	(05)	18.50 ^c	Utah	154.64 ^b	1.88	(51)	-26.92 ^c
Maine	227.73 ^b	2.54	(13)	7.62 ^c	Vermont	213.57	3.74	(24)	0.93
Maryland	211.78	1.35	(26)	0.08	Virginia	216.71 ^b	1.17	(22)	2.41
Massachusetts	210.79	1.18	(30)	-0.39	Washington	205.71 ^b	1.20	(33)	-2.79
Michigan	219.43 ^b	0.98	(19)	3.69	West Virginia	246.11 ^b	2.25	(07)	16.30 ^c
Minnesota	201.15 ^b	1.30	(38)	-4.94	Wisconsin	212.58	1.26	(25)	0.46
Mississippi	264.72 ^b	2.10	(01)	25.10 ^c	Wyoming	192.34 ^b	3.97	(44)	-9.10 ^c
Missouri	225.94 ^b	1.27	(15)	6.77 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.20
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	147.45	0.13		
High Five States				
Kentucky	172.23 ^b	1.16	(01)	16.81 ^c
West Virginia	167.80 ^b	1.63	(02)	13.81 ^c
D.C.	166.83 ^b	3.22	(03)	13.14 ^c
Louisiana	162.16 ^b	1.13	(04)	9.98 ^c
Ohio	160.38 ^b	0.66	(05)	8.77 ^c
Low Five States				
Colorado	130.53 ^b	1.03	(47)	-11.47 ^c
Arizona	129.96 ^b	0.85	(48)	-11.86 ^c
New Mexico	129.84 ^b	1.50	(49)	-11.94 ^c
Hawaii	115.58 ^b	1.67	(50)	-21.61 ^c
Utah	109.06 ^b	1.41	(51)	-26.03 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	154.28 ^b	1.03	(18)	4.63	Montana	142.17	2.16	(40)	-3.58
Alaska	155.03	3.71	(16)	5.14 ^c	Nebraska	142.53	1.62	(39)	-3.34
Arizona	129.96 ^b	0.85	(48)	-11.86 ^c	Nevada	151.73	1.52	(21)	2.90
Arkansas	159.81 ^b	1.34	(06)	8.39 ^c	New Hampshire	152.07	1.96	(20)	3.14
California	137.30 ^b	0.38	(45)	-6.88 ^c	New Jersey	151.01 ^b	0.74	(23)	2.42
Colorado	130.53 ^b	1.03	(47)	-11.47 ^c	New Mexico	129.84 ^b	1.50	(49)	-11.94 ^c
Connecticut	141.26 ^b	1.11	(41)	-4.20	New York	143.44 ^b	0.49	(36)	-2.72
Delaware	158.91 ^b	2.40	(09)	7.78 ^c	North Carolina	147.18	0.73	(31)	-0.18
D.C.	166.83 ^b	3.22	(03)	13.14 ^c	North Dakota	133.07 ^b	2.53	(46)	-9.75 ^c
Florida	138.76 ^b	0.46	(43)	-5.89 ^c	Ohio	160.38 ^b	0.66	(05)	8.77 ^c
Georgia	146.26	0.78	(33)	-0.81	Oklahoma	159.70 ^b	1.21	(07)	8.31 ^c
Hawaii	115.58 ^b	1.67	(50)	-21.61 ^c	Oregon	151.44	1.15	(22)	2.71
Idaho	138.58 ^b	1.85	(44)	-6.01 ^c	Pennsylvania	154.91 ^b	0.60	(17)	5.06 ^c
Illinois	156.44 ^b	0.65	(15)	6.10 ^c	Rhode Island	147.45	2.07	(30)	0.01
Indiana	159.05 ^b	0.92	(08)	7.87 ^c	South Carolina	149.97	1.05	(25)	1.71
Iowa	147.08	1.23	(32)	-0.25	South Dakota	143.90	2.40	(35)	-2.41
Kansas	145.00	1.32	(34)	-1.66	Tennessee	158.15 ^b	0.92	(10)	7.26 ^c
Kentucky	172.23 ^b	1.16	(01)	16.81 ^c	Texas	139.66 ^b	0.49	(42)	-5.28 ^c
Louisiana	162.16 ^b	1.13	(04)	9.98 ^c	Utah	109.06 ^b	1.41	(51)	-26.03 ^c
Maine	156.47 ^b	1.85	(14)	6.12 ^c	Vermont	153.15	2.78	(19)	3.87
Maryland	150.15	0.96	(24)	1.84	Virginia	149.10	0.82	(28)	1.12
Massachusetts	149.16	0.85	(27)	1.16	Washington	149.70	0.91	(26)	1.53
Michigan	157.09 ^b	0.72	(12)	6.54 ^c	West Virginia	167.80 ^b	1.63	(02)	13.81 ^c
Minnesota	142.98 ^b	0.97	(38)	-3.03	Wisconsin	148.45	0.93	(29)	0.68
Mississippi	156.96 ^b	1.35	(13)	6.45 ^c	Wyoming	143.42	3.11	(37)	-2.73
Missouri	157.80 ^b	0.93	(11)	7.02 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.18
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.27	0.02		
High Five States				
South Dakota	5.55 ^b	0.36	(01)	30.03 ^c
Nebraska	5.26 ^b	0.24	(02)	23.24 ^c
Iowa	5.21 ^b	0.18	(03)	22.14 ^c
Washington	5.11 ^b	0.12	(04)	19.64 ^c
Wisconsin	5.01 ^b	0.13	(05)	17.46 ^c
Low Five States				
New Jersey	3.73 ^b	0.09	(47)	-12.64
New York	3.71 ^b	0.06	(48)	-13.04
New Mexico	3.69	0.19	(49)	-13.54
D.C.	3.43	0.35	(50)	-19.63 ^c
Hawaii	2.31 ^b	0.18	(51)	-45.85 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.93 ^b	0.14	(09)	15.55 ^c	Montana	4.66	0.29	(20)	9.15
Alaska	4.67	0.44	(19)	9.48	Nebraska	5.26 ^b	0.24	(02)	23.24 ^c
Arizona	4.03	0.11	(38)	-5.59	Nevada	4.01	0.18	(39)	-6.02
Arkansas	4.98 ^b	0.18	(08)	16.74 ^c	New Hampshire	4.75	0.26	(15)	11.40
California	4.26	0.05	(28)	-0.21	New Jersey	3.73 ^b	0.09	(47)	-12.64
Colorado	4.43	0.14	(25)	3.70	New Mexico	3.69	0.19	(49)	-13.54
Connecticut	4.24	0.15	(30)	-0.56	New York	3.71 ^b	0.06	(48)	-13.04
Delaware	4.10	0.29	(36)	-3.89	North Carolina	4.20	0.09	(34)	-1.59
D.C.	3.43	0.35	(50)	-19.63 ^c	North Dakota	4.00	0.34	(40)	-6.30
Florida	3.97 ^b	0.06	(43)	-7.09	Ohio	4.41	0.08	(26)	3.40
Georgia	4.06	0.10	(37)	-4.92	Oklahoma	4.79	0.16	(14)	12.20
Hawaii	2.31 ^b	0.18	(51)	-45.85 ^c	Oregon	4.99 ^b	0.16	(07)	17.00 ^c
Idaho	4.82	0.25	(12)	13.05	Pennsylvania	4.14	0.08	(35)	-2.92
Illinois	3.99 ^b	0.08	(42)	-6.55	Rhode Island	3.76	0.25	(46)	-11.93
Indiana	4.53	0.12	(22)	6.19	South Carolina	4.35	0.13	(27)	2.05
Iowa	5.21 ^b	0.18	(03)	22.14 ^c	South Dakota	5.55 ^b	0.36	(01)	30.03 ^c
Kansas	4.92 ^b	0.18	(10)	15.35 ^c	Tennessee	4.73 ^b	0.12	(16)	10.73
Kentucky	4.72	0.14	(17)	10.63	Texas	4.00 ^b	0.06	(41)	-6.37
Louisiana	4.21	0.14	(33)	-1.44	Utah	4.25	0.20	(29)	-0.49
Maine	4.99	0.25	(06)	17.01 ^c	Vermont	4.88	0.37	(11)	14.33
Maryland	3.92	0.12	(45)	-8.22	Virginia	3.94	0.10	(44)	-7.73
Massachusetts	4.24	0.11	(31)	-0.62	Washington	5.11 ^b	0.12	(04)	19.64 ^c
Michigan	4.82 ^b	0.10	(13)	13.05	West Virginia	4.22	0.20	(32)	-1.02
Minnesota	4.56	0.13	(21)	6.89	Wisconsin	5.01 ^b	0.13	(05)	17.46 ^c
Mississippi	4.50	0.17	(24)	5.33	Wyoming	4.71	0.41	(18)	10.28
Missouri	4.53	0.12	(23)	6.07					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.19
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.21	0.03		
High Five States				
Iowa	6.50 ^b	0.29	(01)	24.74 ^c
Oregon	6.28 ^b	0.26	(02)	20.67 ^c
Washington	6.26 ^b	0.20	(03)	20.25 ^c
Wisconsin	6.26 ^b	0.21	(04)	20.16 ^c
Kansas	6.25	0.31	(05)	19.94 ^c
Low Five States				
New York	4.60 ^b	0.10	(47)	-11.59
New Mexico	4.52	0.30	(48)	-13.16
North Dakota	4.46	0.52	(49)	-14.42
D.C.	4.28	0.61	(50)	-17.86 ^c
Hawaii	2.95 ^b	0.29	(51)	-43.39 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.96	0.23	(13)	14.44	Montana	5.41	0.45	(24)	3.90
Alaska	5.66	0.73	(15)	8.78	Nebraska	6.19	0.38	(08)	18.84 ^c
Arizona	4.77	0.18	(41)	-8.38	Nevada	5.30	0.30	(28)	1.84
Arkansas	6.23	0.30	(06)	19.65 ^c	New Hampshire	6.13	0.43	(10)	17.66 ^c
California	5.18	0.08	(33)	-0.62	New Jersey	4.68 ^b	0.15	(46)	-10.19
Colorado	5.48	0.23	(20)	5.15	New Mexico	4.52	0.30	(48)	-13.16
Connecticut	5.32	0.25	(27)	2.14	New York	4.60 ^b	0.10	(47)	-11.59
Delaware	4.95	0.48	(37)	-4.98	North Carolina	5.33	0.16	(26)	2.29
D.C.	4.28	0.61	(50)	-17.86 ^c	North Dakota	4.46	0.52	(49)	-14.42
Florida	4.93	0.10	(38)	-5.41	Ohio	5.25	0.14	(30)	0.80
Georgia	4.95	0.16	(36)	-4.86	Oklahoma	5.59	0.25	(18)	7.25
Hawaii	2.95 ^b	0.29	(51)	-43.39 ^c	Oregon	6.28 ^b	0.26	(02)	20.67 ^c
Idaho	6.05	0.41	(12)	16.21 ^c	Pennsylvania	5.07	0.12	(35)	-2.61
Illinois	4.83	0.13	(39)	-7.29	Rhode Island	4.73	0.43	(44)	-9.24
Indiana	5.46	0.19	(21)	4.78	South Carolina	5.59	0.23	(17)	7.34
Iowa	6.50 ^b	0.29	(01)	24.74 ^c	South Dakota	6.19	0.55	(07)	18.86 ^c
Kansas	6.25	0.31	(05)	19.94 ^c	Tennessee	5.65	0.20	(16)	8.58
Kentucky	5.55	0.23	(19)	6.51	Texas	4.72 ^b	0.10	(45)	-9.45
Louisiana	5.08	0.23	(34)	-2.39	Utah	5.23	0.32	(31)	0.43
Maine	6.13	0.41	(11)	17.66 ^c	Vermont	6.17	0.61	(09)	18.47 ^c
Maryland	4.74	0.19	(42)	-9.03	Virginia	4.80	0.16	(40)	-7.91
Massachusetts	5.22	0.18	(32)	0.16	Washington	6.26 ^b	0.20	(03)	20.25 ^c
Michigan	5.89 ^b	0.16	(14)	13.06	West Virginia	4.73	0.31	(43)	-9.13
Minnesota	5.39	0.21	(25)	3.58	Wisconsin	6.26 ^b	0.21	(04)	20.16 ^c
Mississippi	5.30	0.29	(29)	1.78	Wyoming	5.45	0.63	(23)	4.70
Missouri	5.45	0.19	(22)	4.70					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.20
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.47	0.02		
High Five States				
South Dakota	5.04	0.48	(01)	45.13 ^c
Nebraska	4.38	0.29	(02)	26.17 ^c
Iowa	4.13	0.22	(03)	18.82 ^c
Oklahoma	4.06	0.20	(04)	16.97 ^c
Washington	4.06 ^b	0.15	(05)	16.89 ^c
Low Five States				
New Mexico	2.95	0.23	(47)	-15.05 ^c
New Jersey	2.95 ^b	0.11	(48)	-15.16 ^c
Nevada	2.84	0.21	(49)	-18.14 ^c
D.C.	2.82	0.42	(50)	-18.71 ^c
Hawaii	1.75 ^b	0.21	(51)	-49.49 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.06	0.17	(06)	16.86 ^c	Montana	3.99	0.38	(09)	15.00 ^c
Alaska	3.81	0.55	(18)	9.72	Nebraska	4.38	0.29	(02)	26.17 ^c
Arizona	3.36	0.14	(34)	-3.14	Nevada	2.84	0.21	(49)	-18.14 ^c
Arkansas	3.93	0.21	(14)	13.11	New Hampshire	3.59	0.31	(26)	3.34
California	3.47	0.06	(29)	-0.03	New Jersey	2.95 ^b	0.11	(48)	-15.16 ^c
Colorado	3.55	0.17	(27)	2.15	New Mexico	2.95	0.23	(47)	-15.05 ^c
Connecticut	3.33	0.18	(39)	-4.06	New York	3.00 ^b	0.07	(45)	-13.62
Delaware	3.41	0.36	(32)	-1.75	North Carolina	3.30	0.11	(41)	-4.97
D.C.	2.82	0.42	(50)	-18.71 ^c	North Dakota	3.52	0.43	(28)	1.42
Florida	3.13 ^b	0.07	(44)	-9.82	Ohio	3.68	0.10	(24)	5.95
Georgia	3.35	0.12	(36)	-3.51	Oklahoma	4.06	0.20	(04)	16.97 ^c
Hawaii	1.75 ^b	0.21	(51)	-49.49 ^c	Oregon	3.86	0.19	(16)	11.22
Idaho	3.69	0.31	(23)	6.18	Pennsylvania	3.36	0.09	(35)	-3.16
Illinois	3.30	0.10	(40)	-4.89	Rhode Island	2.99	0.30	(46)	-14.00
Indiana	3.76	0.14	(21)	8.26	South Carolina	3.35	0.16	(37)	-3.53
Iowa	4.13	0.22	(03)	18.82 ^c	South Dakota	5.04	0.48	(01)	45.13 ^c
Kansas	3.78	0.22	(20)	8.70	Tennessee	3.96	0.15	(10)	14.04
Kentucky	4.01	0.18	(07)	15.55 ^c	Texas	3.39	0.08	(33)	-2.40
Louisiana	3.45	0.17	(30)	-0.66	Utah	3.35	0.24	(38)	-3.68
Maine	4.00	0.31	(08)	15.10 ^c	Vermont	3.66	0.44	(25)	5.40
Maryland	3.24	0.14	(42)	-6.68	Virginia	3.20	0.12	(43)	-7.83
Massachusetts	3.45	0.13	(31)	-0.75	Washington	4.06 ^b	0.15	(05)	16.89 ^c
Michigan	3.94 ^b	0.12	(13)	13.49	West Virginia	3.81	0.26	(19)	9.65
Minnesota	3.84	0.16	(17)	10.51	Wisconsin	3.96	0.16	(11)	13.97
Mississippi	3.89	0.22	(15)	12.00	Wyoming	3.95	0.52	(12)	13.81
Missouri	3.73	0.15	(22)	7.41					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 4.24
Cancer of the Female Breast (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	22.23	0.05		
High Five States				
D.C.	29.40 ^b	1.36	(01)	32.22 ^c
Louisiana	25.03 ^b	0.45	(02)	12.59 ^c
New Jersey	24.61 ^b	0.30	(03)	10.68 ^c
Ohio	24.22 ^b	0.26	(04)	8.92
Mississippi	24.18 ^b	0.54	(05)	8.77
Low Five States				
Maine	19.99	0.67	(47)	-10.10 ^c
Nebraska	19.78 ^b	0.61	(48)	-11.02 ^c
Vermont	19.69	1.00	(49)	-11.42 ^c
Colorado	19.35 ^b	0.39	(50)	-12.97 ^c
Hawaii	15.16 ^b	0.62	(51)	-31.83 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	22.88	0.40	(18)	2.92	Montana	20.09	0.82	(45)	-9.65
Alaska	22.70	1.38	(23)	2.08	Nebraska	19.78 ^b	0.61	(48)	-11.02 ^c
Arizona	20.07 ^b	0.34	(46)	-9.72	Nevada	23.51	0.59	(09)	5.74
Arkansas	22.83	0.51	(19)	2.70	New Hampshire	21.16	0.73	(30)	-4.83
California	21.46 ^b	0.15	(27)	-3.47	New Jersey	24.61 ^b	0.30	(03)	10.68 ^c
Colorado	19.35 ^b	0.39	(50)	-12.97 ^c	New Mexico	20.84	0.60	(38)	-6.24
Connecticut	20.77	0.43	(40)	-6.56	New York	21.96	0.19	(25)	-1.21
Delaware	22.91	0.92	(17)	3.05	North Carolina	22.74	0.29	(22)	2.27
D.C.	29.40 ^b	1.36	(01)	32.22 ^c	North Dakota	21.12	1.02	(32)	-5.02
Florida	21.04 ^b	0.18	(37)	-5.38	Ohio	24.22 ^b	0.26	(04)	8.92
Georgia	23.15	0.31	(16)	4.10	Oklahoma	23.53	0.47	(08)	5.85
Hawaii	15.16 ^b	0.62	(51)	-31.83 ^c	Oregon	21.05	0.43	(35)	-5.30
Idaho	21.90	0.74	(26)	-1.50	Pennsylvania	23.47 ^b	0.24	(13)	5.55
Illinois	23.38 ^b	0.25	(14)	5.18	Rhode Island	20.56	0.78	(43)	-7.53
Indiana	23.20	0.35	(15)	4.37	South Carolina	23.49	0.42	(12)	5.64
Iowa	20.72	0.47	(41)	-6.82	South Dakota	21.13	0.93	(31)	-4.96
Kansas	21.41	0.51	(28)	-3.71	Tennessee	22.74	0.35	(21)	2.29
Kentucky	22.83	0.42	(20)	2.70	Texas	21.35 ^b	0.19	(29)	-3.99
Louisiana	25.03 ^b	0.45	(02)	12.59 ^c	Utah	20.83	0.62	(39)	-6.31
Maine	19.99	0.67	(47)	-10.10 ^c	Vermont	19.69	1.00	(49)	-11.42 ^c
Maryland	23.90 ^b	0.38	(06)	7.48	Virginia	23.51 ^b	0.32	(10)	5.74
Massachusetts	20.44 ^b	0.32	(44)	-8.06	Washington	21.10	0.34	(34)	-5.08
Michigan	23.49 ^b	0.28	(11)	5.66	West Virginia	22.52	0.61	(24)	1.30
Minnesota	20.66 ^b	0.37	(42)	-7.07	Wisconsin	21.05	0.35	(36)	-5.32
Mississippi	24.18 ^b	0.54	(05)	8.77	Wyoming	21.11	1.18	(33)	-5.04
Missouri	23.77 ^b	0.36	(07)	6.89					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 5.14
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.33	0.02		
High Five States				
Mississippi	3.46 ^b	0.21	(01)	48.24 ^c
Arkansas	3.39 ^b	0.21	(02)	45.57 ^c
West Virginia	3.27 ^b	0.26	(03)	40.28 ^c
Louisiana	3.13 ^b	0.16	(04)	34.36 ^c
Kentucky	3.06 ^b	0.16	(05)	31.28 ^c
Low Five States				
Rhode Island	1.39 ^b	0.21	(47)	-40.25 ^c
Massachusetts	1.38 ^b	0.08	(48)	-40.94 ^c
North Dakota	1.35 ^b	0.27	(49)	-42.28 ^c
Vermont	1.27 ^b	0.27	(50)	-45.54 ^c
Utah	1.21 ^b	0.15	(51)	-47.95 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.05 ^b	0.15	(06)	30.71 ^c	Montana	1.49 ^b	0.24	(45)	-35.94 ^c
Alaska	2.35	0.40	(20)	0.97	Nebraska	1.87	0.20	(37)	-19.78 ^c
Arizona	1.97	0.11	(35)	-15.63 ^c	Nevada	2.14	0.18	(26)	-8.16
Arkansas	3.39 ^b	0.21	(02)	45.57 ^c	New Hampshire	1.77	0.22	(40)	-23.89 ^c
California	2.29	0.05	(23)	-1.73	New Jersey	2.33	0.10	(21)	-0.17
Colorado	1.62 ^b	0.11	(41)	-30.60 ^c	New Mexico	2.10	0.20	(28)	-9.83
Connecticut	1.56 ^b	0.12	(43)	-33.30 ^c	New York	2.31	0.06	(22)	-1.02
Delaware	2.53	0.32	(18)	8.54	North Carolina	2.10	0.09	(29)	-9.89
D.C.	2.55	0.41	(16)	9.58	North Dakota	1.35 ^b	0.27	(49)	-42.28 ^c
Florida	2.63 ^b	0.07	(15)	12.68	Ohio	2.65	0.09	(13)	13.58
Georgia	2.71 ^b	0.11	(12)	16.28 ^c	Oklahoma	2.82	0.17	(08)	21.13 ^c
Hawaii	1.83	0.22	(39)	-21.67 ^c	Oregon	1.98	0.14	(34)	-15.04 ^c
Idaho	2.13	0.24	(27)	-8.54	Pennsylvania	2.14	0.08	(25)	-8.15
Illinois	2.63 ^b	0.09	(14)	12.98	Rhode Island	1.39 ^b	0.21	(47)	-40.25 ^c
Indiana	2.39	0.12	(19)	2.72	South Carolina	2.85	0.15	(07)	22.07 ^c
Iowa	2.05	0.16	(33)	-11.95	South Dakota	2.07	0.31	(31)	-11.20
Kansas	1.88	0.16	(36)	-19.26 ^c	Tennessee	2.82 ^b	0.13	(09)	20.87 ^c
Kentucky	3.06 ^b	0.16	(05)	31.28 ^c	Texas	2.80 ^b	0.07	(10)	20.28 ^c
Louisiana	3.13 ^b	0.16	(04)	34.36 ^c	Utah	1.21 ^b	0.15	(51)	-47.95 ^c
Maine	1.60 ^b	0.19	(42)	-31.36 ^c	Vermont	1.27 ^b	0.27	(50)	-45.54 ^c
Maryland	2.23	0.12	(24)	-4.40	Virginia	2.07	0.10	(32)	-11.21
Massachusetts	1.38 ^b	0.08	(48)	-40.94 ^c	Washington	1.86 ^b	0.10	(38)	-20.03 ^c
Michigan	2.08	0.09	(30)	-10.84	West Virginia	3.27 ^b	0.26	(03)	40.28 ^c
Minnesota	1.41 ^b	0.10	(46)	-39.43 ^c	Wisconsin	1.55 ^b	0.10	(44)	-33.63 ^c
Mississippi	3.46 ^b	0.21	(01)	48.24 ^c	Wyoming	2.79	0.46	(11)	19.55 ^c
Missouri	2.53	0.13	(17)	8.64					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.25
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	15.94	0.03		
High Five States				
Mississippi	19.96 ^b	0.36	(01)	25.23 ^c
D.C.	19.46 ^b	0.83	(02)	22.10 ^c
West Virginia	19.07 ^b	0.41	(03)	19.66 ^c
Kentucky	18.92 ^b	0.29	(04)	18.71 ^c
Louisiana	18.87 ^b	0.29	(05)	18.40 ^c
Low Five States				
Arizona	13.90 ^b	0.20	(47)	-12.78 ^c
Hawaii	13.78 ^b	0.42	(48)	-13.53 ^c
Colorado	13.70 ^b	0.25	(49)	-14.05 ^c
Connecticut	13.11 ^b	0.25	(50)	-17.76 ^c
Utah	12.06 ^b	0.35	(51)	-24.34 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	17.42 ^b	0.26	(11)	9.30	Montana	14.18 ^b	0.49	(45)	-11.01 ^c
Alaska	16.63	0.91	(19)	4.32	Nebraska	17.48 ^b	0.42	(10)	9.64
Arizona	13.90 ^b	0.20	(47)	-12.78 ^c	Nevada	17.37 ^b	0.38	(12)	8.96
Arkansas	18.80 ^b	0.34	(06)	17.95 ^c	New Hampshire	14.75	0.45	(41)	-7.49
California	14.53 ^b	0.09	(42)	-8.87	New Jersey	16.81 ^b	0.19	(18)	5.44
Colorado	13.70 ^b	0.25	(49)	-14.05 ^c	New Mexico	15.25	0.38	(37)	-4.30
Connecticut	13.11 ^b	0.25	(50)	-17.76 ^c	New York	15.54	0.12	(33)	-2.52
Delaware	15.98	0.57	(27)	0.28	North Carolina	15.29 ^b	0.18	(36)	-4.09
D.C.	19.46 ^b	0.83	(02)	22.10 ^c	North Dakota	16.56	0.65	(20)	3.89
Florida	14.81 ^b	0.11	(40)	-7.09	Ohio	17.53 ^b	0.16	(09)	9.97
Georgia	16.12	0.20	(25)	1.10	Oklahoma	17.70 ^b	0.30	(08)	11.06 ^c
Hawaii	13.78 ^b	0.42	(48)	-13.53 ^c	Oregon	15.54	0.27	(32)	-2.49
Idaho	13.99 ^b	0.43	(46)	-12.22 ^c	Pennsylvania	17.18 ^b	0.15	(14)	7.80
Illinois	17.32 ^b	0.16	(13)	8.68	Rhode Island	15.41	0.50	(35)	-3.35
Indiana	17.06 ^b	0.22	(17)	7.00	South Carolina	16.47	0.26	(21)	3.30
Iowa	17.07 ^b	0.31	(15)	7.11	South Dakota	16.12	0.58	(24)	1.12
Kansas	16.13	0.33	(22)	1.21	Tennessee	18.02 ^b	0.23	(07)	13.07 ^c
Kentucky	18.92 ^b	0.29	(04)	18.71 ^c	Texas	15.75	0.12	(29)	-1.20
Louisiana	18.87 ^b	0.29	(05)	18.40 ^c	Utah	12.06 ^b	0.35	(51)	-24.34 ^c
Maine	16.12	0.44	(23)	1.13	Vermont	15.51	0.66	(34)	-2.72
Maryland	15.92	0.24	(28)	-0.12	Virginia	15.60	0.20	(31)	-2.15
Massachusetts	14.82 ^b	0.20	(39)	-7.04	Washington	14.48 ^b	0.21	(43)	-9.16
Michigan	16.05	0.17	(26)	0.72	West Virginia	19.07 ^b	0.41	(03)	19.66 ^c
Minnesota	14.42 ^b	0.23	(44)	-9.53	Wisconsin	14.82 ^b	0.22	(38)	-7.03
Mississippi	19.96 ^b	0.36	(01)	25.23 ^c	Wyoming	15.63	0.75	(30)	-1.96
Missouri	17.06 ^b	0.23	(16)	7.01					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.26
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	19.12	0.05		
High Five States				
Mississippi	24.56 ^b	0.64	(01)	28.43 ^c
West Virginia	23.66 ^b	0.70	(02)	23.72 ^c
Louisiana	23.33 ^b	0.50	(03)	22.00 ^c
Arkansas	22.88 ^b	0.58	(04)	19.66 ^c
Kentucky	22.60 ^b	0.49	(05)	18.17 ^c
Low Five States				
Idaho	16.01 ^b	0.69	(47)	-16.25 ^c
Colorado	15.84 ^b	0.41	(48)	-17.18 ^c
Montana	15.78 ^b	0.78	(49)	-17.50 ^c
Connecticut	15.28 ^b	0.42	(50)	-20.10 ^c
Utah	14.05 ^b	0.56	(51)	-26.54 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	21.45 ^b	0.45	(08)	12.19 ^c	Montana	15.78 ^b	0.78	(49)	-17.50 ^c
Alaska	19.57	1.48	(24)	2.33	Nebraska	20.42	0.69	(17)	6.81
Arizona	16.50 ^b	0.33	(46)	-13.73 ^c	Nevada	20.96	0.63	(10)	9.59
Arkansas	22.88 ^b	0.58	(04)	19.66 ^c	New Hampshire	16.62	0.74	(45)	-13.11 ^c
California	17.19 ^b	0.15	(42)	-10.13 ^c	New Jersey	20.25 ^b	0.32	(18)	5.88
Colorado	15.84 ^b	0.41	(48)	-17.18 ^c	New Mexico	18.66	0.64	(32)	-2.40
Connecticut	15.28 ^b	0.42	(50)	-20.10 ^c	New York	18.54	0.21	(33)	-3.07
Delaware	19.02	0.96	(29)	-0.51	North Carolina	18.77	0.31	(31)	-1.86
D.C.	20.58	1.35	(16)	7.64	North Dakota	20.81	1.10	(13)	8.84
Florida	17.65 ^b	0.18	(39)	-7.70	Ohio	21.40 ^b	0.28	(09)	11.92 ^c
Georgia	19.83	0.35	(21)	3.71	Oklahoma	22.14 ^b	0.51	(06)	15.79 ^c
Hawaii	17.59	0.71	(40)	-8.00	Oregon	18.22	0.45	(35)	-4.72
Idaho	16.01 ^b	0.69	(47)	-16.25 ^c	Pennsylvania	20.70 ^b	0.25	(15)	8.24
Illinois	20.71 ^b	0.28	(14)	8.31	Rhode Island	18.10	0.85	(36)	-5.36
Indiana	20.82 ^b	0.39	(12)	8.86	South Carolina	19.91	0.45	(20)	4.11
Iowa	19.96	0.51	(19)	4.39	South Dakota	18.86	0.96	(30)	-1.35
Kansas	19.80	0.55	(22)	3.53	Tennessee	21.72 ^b	0.40	(07)	13.59 ^c
Kentucky	22.60 ^b	0.49	(05)	18.17 ^c	Texas	19.43	0.21	(25)	1.60
Louisiana	23.33 ^b	0.50	(03)	22.00 ^c	Utah	14.05 ^b	0.56	(51)	-26.54 ^c
Maine	19.40	0.75	(26)	1.48	Vermont	16.75	1.05	(43)	-12.40 ^c
Maryland	19.61	0.41	(23)	2.57	Virginia	18.31	0.34	(34)	-4.27
Massachusetts	17.74 ^b	0.34	(38)	-7.22	Washington	16.75 ^b	0.34	(44)	-12.42 ^c
Michigan	19.11	0.29	(27)	-0.04	West Virginia	23.66 ^b	0.70	(02)	23.72 ^c
Minnesota	17.25 ^b	0.38	(41)	-9.80	Wisconsin	17.93	0.37	(37)	-6.24
Mississippi	24.56 ^b	0.64	(01)	28.43 ^c	Wyoming	19.06	1.23	(28)	-0.34
Missouri	20.88 ^b	0.39	(11)	9.22					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.27
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	13.46	0.04		
High Five States				
D.C.	18.77 ^b	1.07	(01)	39.47 ^c
Mississippi	16.50 ^b	0.44	(02)	22.59 ^c
Kentucky	16.05 ^b	0.35	(03)	19.23 ^c
West Virginia	15.62 ^b	0.49	(04)	16.01 ^c
Arkansas	15.61 ^b	0.42	(05)	15.95 ^c
Low Five States				
Colorado	12.03 ^b	0.31	(47)	-10.63 ^c
Arizona	11.68 ^b	0.25	(48)	-13.20 ^c
Connecticut	11.42 ^b	0.31	(49)	-15.18 ^c
Hawaii	10.66 ^b	0.50	(50)	-20.84 ^c
Utah	10.33 ^b	0.44	(51)	-23.25 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	14.42	0.31	(14)	7.14	Montana	12.72	0.64	(37)	-5.53
Alaska	13.88	1.13	(20)	3.10	Nebraska	14.97	0.51	(08)	11.22 ^c
Arizona	11.68 ^b	0.25	(48)	-13.20 ^c	Nevada	14.29	0.47	(15)	6.18
Arkansas	15.61 ^b	0.42	(05)	15.95 ^c	New Hampshire	13.25	0.57	(31)	-1.54
California	12.43 ^b	0.11	(43)	-7.64	New Jersey	14.29 ^b	0.23	(16)	6.14
Colorado	12.03 ^b	0.31	(47)	-10.63 ^c	New Mexico	12.47	0.46	(41)	-7.34
Connecticut	11.42 ^b	0.31	(49)	-15.18 ^c	New York	13.33	0.15	(29)	-0.97
Delaware	13.47	0.69	(27)	0.07	North Carolina	12.69 ^b	0.21	(39)	-5.70
D.C.	18.77 ^b	1.07	(01)	39.47 ^c	North Dakota	13.14	0.77	(33)	-2.39
Florida	12.45 ^b	0.14	(42)	-7.54	Ohio	14.61 ^b	0.20	(11)	8.51
Georgia	13.49	0.24	(26)	0.22	Oklahoma	14.28	0.36	(17)	6.06
Hawaii	10.66 ^b	0.50	(50)	-20.84 ^c	Oregon	13.30	0.34	(30)	-1.17
Idaho	12.30	0.55	(45)	-8.60	Pennsylvania	14.55 ^b	0.18	(12)	8.10
Illinois	14.83 ^b	0.20	(09)	10.20 ^c	Rhode Island	13.41	0.61	(28)	-0.40
Indiana	14.25	0.27	(18)	5.82	South Carolina	13.68	0.31	(23)	1.61
Iowa	14.83 ^b	0.38	(10)	10.17 ^c	South Dakota	13.61	0.71	(24)	1.07
Kansas	13.08	0.39	(34)	-2.80	Tennessee	15.25 ^b	0.28	(07)	13.29 ^c
Kentucky	16.05 ^b	0.35	(03)	19.23 ^c	Texas	12.83 ^b	0.15	(36)	-4.69
Louisiana	15.37 ^b	0.35	(06)	14.21 ^c	Utah	10.33 ^b	0.44	(51)	-23.25 ^c
Maine	13.73	0.54	(21)	1.99	Vermont	14.48	0.85	(13)	7.58
Maryland	13.19	0.28	(32)	-2.03	Virginia	13.52	0.25	(25)	0.44
Massachusetts	12.70	0.24	(38)	-5.68	Washington	12.57	0.26	(40)	-6.62
Michigan	13.73	0.21	(22)	1.98	West Virginia	15.62 ^b	0.49	(04)	16.01 ^c
Minnesota	12.12 ^b	0.28	(46)	-9.94	Wisconsin	12.42 ^b	0.26	(44)	-7.73
Mississippi	16.50 ^b	0.44	(02)	22.59 ^c	Wyoming	12.84	0.93	(35)	-4.60
Missouri	14.09	0.27	(19)	4.64					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 7.14
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.31	0.02		
High Five States				
D.C.	7.15 ^b	0.66	(01)	66.06 ^c
New Jersey	5.33 ^b	0.14	(02)	23.78 ^c
New York	5.27 ^b	0.09	(03)	22.38 ^c
Vermont	5.22	0.51	(04)	21.14 ^c
Michigan	5.07 ^b	0.13	(05)	17.68 ^c
Low Five States				
Arizona	3.33 ^b	0.13	(47)	-22.71 ^c
Nevada	3.29 ^b	0.22	(48)	-23.69 ^c
Arkansas	3.28 ^b	0.19	(49)	-23.78 ^c
New Mexico	3.06 ^b	0.23	(50)	-28.85 ^c
Alaska	2.83	0.50	(51)	-34.20 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.55 ^b	0.15	(44)	-17.61 ^c	Montana	3.86	0.35	(34)	-10.33
Alaska	2.83	0.50	(51)	-34.20 ^c	Nebraska	4.45	0.29	(16)	3.25
Arizona	3.33 ^b	0.13	(47)	-22.71 ^c	Nevada	3.29 ^b	0.22	(48)	-23.69 ^c
Arkansas	3.28 ^b	0.19	(49)	-23.78 ^c	New Hampshire	5.05	0.35	(06)	17.32 ^c
California	4.27	0.07	(25)	-0.83	New Jersey	5.33 ^b	0.14	(02)	23.78 ^c
Colorado	3.61 ^b	0.17	(40)	-16.27 ^c	New Mexico	3.06 ^b	0.23	(50)	-28.85 ^c
Connecticut	4.32	0.20	(23)	0.23	New York	5.27 ^b	0.09	(03)	22.38 ^c
Delaware	4.73	0.41	(12)	9.77	North Carolina	4.04	0.12	(29)	-6.30
D.C.	7.15 ^b	0.66	(01)	66.06 ^c	North Dakota	3.53	0.41	(45)	-17.96 ^c
Florida	3.90 ^b	0.08	(33)	-9.36	Ohio	4.86 ^b	0.11	(10)	12.85
Georgia	3.92	0.13	(32)	-8.94	Oklahoma	3.80	0.19	(37)	-11.84
Hawaii	4.39	0.33	(19)	1.91	Oregon	4.00	0.19	(30)	-7.11
Idaho	4.36	0.33	(21)	1.26	Pennsylvania	4.98 ^b	0.11	(08)	15.59 ^c
Illinois	4.83 ^b	0.11	(11)	12.21	Rhode Island	3.84	0.33	(35)	-10.84
Indiana	4.39	0.15	(18)	2.03	South Carolina	4.43	0.18	(17)	2.78
Iowa	4.98	0.23	(09)	15.50 ^c	South Dakota	4.70	0.44	(13)	9.16
Kansas	3.99	0.22	(31)	-7.30	Tennessee	3.65 ^b	0.14	(38)	-15.30 ^c
Kentucky	3.53 ^b	0.16	(46)	-18.09 ^c	Texas	3.59 ^b	0.08	(41)	-16.61 ^c
Louisiana	3.56 ^b	0.17	(42)	-17.27 ^c	Utah	3.64	0.26	(39)	-15.61 ^c
Maine	4.51	0.31	(15)	4.64	Vermont	5.22	0.51	(04)	21.14 ^c
Maryland	5.03 ^b	0.17	(07)	16.87 ^c	Virginia	4.36	0.14	(22)	1.22
Massachusetts	4.29	0.14	(24)	-0.31	Washington	4.09	0.15	(27)	-5.15
Michigan	5.07 ^b	0.13	(05)	17.68 ^c	West Virginia	4.37	0.26	(20)	1.53
Minnesota	4.06	0.16	(28)	-5.81	Wisconsin	4.68	0.16	(14)	8.76
Mississippi	3.84	0.21	(36)	-10.90	Wyoming	3.56	0.49	(43)	-17.31 ^c
Missouri	4.25	0.15	(26)	-1.43					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.18
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.22	0.02		
High Five States				
New Hampshire	5.73 ^b	0.28	(01)	35.83 ^c
Maine	5.69 ^b	0.26	(02)	34.94 ^c
Rhode Island	5.12	0.29	(03)	21.56 ^c
Ohio	5.02 ^b	0.09	(04)	19.15 ^c
Indiana	4.99 ^b	0.12	(05)	18.28 ^c
Low Five States				
Texas	3.49 ^b	0.06	(47)	-17.21 ^c
California	3.46 ^b	0.04	(48)	-17.97 ^c
Hawaii	3.23 ^b	0.20	(49)	-23.38 ^c
New Mexico	3.19 ^b	0.17	(50)	-24.22 ^c
Utah	2.87 ^b	0.17	(51)	-31.86 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.03	0.12	(40)	-4.38	Montana	4.47	0.28	(24)	5.96
Alaska	4.53	0.46	(19)	7.52	Nebraska	4.49	0.21	(23)	6.42
Arizona	3.96	0.11	(42)	-6.08	Nevada	4.80	0.20	(15)	13.76
Arkansas	3.72	0.15	(46)	-11.85	New Hampshire	5.73 ^b	0.28	(01)	35.83 ^c
California	3.46 ^b	0.04	(48)	-17.97 ^c	New Jersey	4.01	0.09	(41)	-4.97
Colorado	3.93	0.13	(43)	-6.79	New Mexico	3.19 ^b	0.17	(50)	-24.22 ^c
Connecticut	4.24	0.15	(30)	0.51	New York	4.14	0.06	(38)	-1.82
Delaware	4.29	0.29	(28)	1.71	North Carolina	3.81 ^b	0.09	(44)	-9.55
D.C.	4.85	0.42	(11)	15.07 ^c	North Dakota	4.24	0.34	(29)	0.55
Florida	4.10	0.06	(39)	-2.84	Ohio	5.02 ^b	0.09	(04)	19.15 ^c
Georgia	3.80 ^b	0.09	(45)	-9.81	Oklahoma	4.57	0.15	(18)	8.41
Hawaii	3.23 ^b	0.20	(49)	-23.38 ^c	Oregon	4.42	0.14	(25)	4.93
Idaho	4.21	0.23	(32)	-0.10	Pennsylvania	4.89 ^b	0.08	(09)	15.98 ^c
Illinois	4.52 ^b	0.08	(21)	7.25	Rhode Island	5.12	0.29	(03)	21.56 ^c
Indiana	4.99 ^b	0.12	(05)	18.28 ^c	South Carolina	4.23	0.13	(31)	0.36
Iowa	4.71	0.16	(17)	11.67	South Dakota	4.96	0.33	(07)	17.72 ^c
Kansas	4.15	0.16	(36)	-1.50	Tennessee	4.17	0.11	(34)	-1.10
Kentucky	4.53	0.14	(20)	7.40	Texas	3.49 ^b	0.06	(47)	-17.21 ^c
Louisiana	4.32	0.14	(26)	2.50	Utah	2.87 ^b	0.17	(51)	-31.86 ^c
Maine	5.69 ^b	0.26	(02)	34.94 ^c	Vermont	4.80	0.36	(14)	13.84
Maryland	4.17	0.12	(35)	-1.11	Virginia	4.20	0.10	(33)	-4.38
Massachusetts	4.98 ^b	0.12	(06)	18.05 ^c	Washington	4.82 ^b	0.12	(13)	14.36
Michigan	4.83 ^b	0.09	(12)	14.67	West Virginia	4.95 ^b	0.20	(08)	17.53 ^c
Minnesota	4.50	0.13	(22)	6.65	Wisconsin	4.78 ^b	0.12	(16)	13.38
Mississippi	4.15	0.16	(37)	-1.56	Wyoming	4.87	0.42	(10)	15.52 ^c
Missouri	4.31	0.11	(27)	2.19					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.19
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.53	0.03		
High Five States				
Maine	10.52 ^b	0.53	(01)	39.82 ^c
New Hampshire	10.41 ^b	0.57	(02)	38.29 ^c
Rhode Island	9.25	0.60	(03)	22.97 ^c
Indiana	9.19 ^b	0.25	(04)	22.13 ^c
Ohio	9.12 ^b	0.18	(05)	21.14 ^c
Low Five States				
Texas	6.28 ^b	0.12	(47)	-16.55 ^c
California	6.01 ^b	0.09	(48)	-20.11 ^c
Hawaii	5.88 ^b	0.41	(49)	-21.87 ^c
New Mexico	5.55 ^b	0.34	(50)	-26.30 ^c
Utah	5.02 ^b	0.33	(51)	-33.34 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.38	0.26	(38)	-1.96	Montana	7.72	0.54	(25)	2.55
Alaska	7.63	0.89	(28)	1.42	Nebraska	8.11	0.43	(21)	7.71
Arizona	6.96	0.21	(44)	-7.50	Nevada	8.19	0.38	(18)	8.81
Arkansas	6.46 ^b	0.30	(46)	-14.21	New Hampshire	10.41 ^b	0.57	(02)	38.29 ^c
California	6.01 ^b	0.09	(48)	-20.11 ^c	New Jersey	7.01	0.18	(42)	-6.83
Colorado	6.98	0.27	(43)	-7.29	New Mexico	5.55 ^b	0.34	(50)	-26.30 ^c
Connecticut	7.50	0.29	(34)	-0.29	New York	7.35	0.13	(39)	-2.39
Delaware	7.66	0.59	(26)	1.84	North Carolina	7.03	0.18	(41)	-6.59
D.C.	8.30	0.85	(15)	10.32	North Dakota	7.87	0.68	(22)	4.55
Florida	7.28	0.11	(40)	-3.23	Ohio	9.12 ^b	0.18	(05)	21.14 ^c
Georgia	6.75 ^b	0.19	(45)	-10.30	Oklahoma	8.19	0.30	(19)	8.81
Hawaii	5.88 ^b	0.41	(49)	-21.87 ^c	Oregon	7.80	0.29	(23)	3.59
Idaho	7.45	0.46	(36)	-1.06	Pennsylvania	8.91 ^b	0.16	(09)	18.34 ^c
Illinois	8.17 ^b	0.17	(20)	8.62	Rhode Island	9.25	0.60	(03)	22.97 ^c
Indiana	9.19 ^b	0.25	(04)	22.13 ^c	South Carolina	7.53	0.27	(33)	0.06
Iowa	8.77 ^b	0.33	(10)	16.59 ^c	South Dakota	8.93	0.66	(07)	18.68 ^c
Kansas	7.75	0.34	(24)	2.94	Tennessee	7.50	0.23	(35)	-0.36
Kentucky	8.19	0.28	(17)	8.84	Texas	6.28 ^b	0.12	(47)	-16.55 ^c
Louisiana	7.56	0.28	(31)	0.42	Utah	5.02 ^b	0.33	(51)	-33.34 ^c
Maine	10.52 ^b	0.53	(01)	39.82 ^c	Vermont	8.63	0.74	(12)	14.74
Maryland	7.41	0.24	(37)	-1.50	Virginia	7.64	0.21	(27)	1.48
Massachusetts	8.92 ^b	0.24	(08)	18.56 ^c	Washington	8.33	0.24	(14)	10.74
Michigan	8.66 ^b	0.19	(11)	15.08 ^c	West Virginia	9.06 ^b	0.42	(06)	20.34 ^c
Minnesota	7.58	0.25	(30)	0.66	Wisconsin	8.47 ^b	0.25	(13)	12.55
Mississippi	7.61	0.34	(29)	1.09	Wyoming	8.20	0.81	(16)	8.96
Missouri	7.54	0.23	(32)	0.19					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 8.20
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.57	0.01		
High Five States				
D.C.	2.33	0.38	(01)	48.72 ^c
Minnesota	1.99 ^b	0.11	(02)	26.79 ^c
Rhode Island	1.99	0.24	(03)	26.58 ^c
Massachusetts	1.95 ^b	0.10	(04)	24.09 ^c
Wyoming	1.94	0.36	(05)	23.92 ^c
Low Five States				
Kansas	1.22	0.12	(47)	-22.20 ^c
New Mexico	1.20	0.15	(48)	-23.47 ^c
North Dakota	1.18	0.24	(49)	-24.89 ^c
Utah	1.02 ^b	0.14	(50)	-34.69 ^c
Hawaii	1.01 ^b	0.16	(51)	-35.65 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.37	0.10	(43)	-12.81	Montana	1.59	0.22	(27)	1.65
Alaska	1.59	0.38	(30)	1.39	Nebraska	1.57	0.17	(31)	-0.12
Arizona	1.40	0.09	(41)	-10.48	Nevada	1.75	0.16	(10)	11.41
Arkansas	1.45	0.13	(37)	-7.53	New Hampshire	1.89	0.22	(07)	20.72 ^c
California	1.41 ^b	0.04	(40)	-9.84	New Jersey	1.68	0.08	(21)	7.40
Colorado	1.42	0.11	(39)	-9.36	New Mexico	1.20	0.15	(48)	-23.47 ^c
Connecticut	1.73	0.12	(14)	10.34	New York	1.70	0.05	(19)	8.23
Delaware	1.62	0.24	(26)	3.19	North Carolina	1.33	0.07	(45)	-14.94
D.C.	2.33	0.38	(01)	48.72 ^c	North Dakota	1.18	0.24	(49)	-24.89 ^c
Florida	1.46	0.05	(36)	-6.99	Ohio	1.83 ^b	0.07	(08)	16.43 ^c
Georgia	1.49	0.08	(34)	-5.20	Oklahoma	1.59	0.12	(28)	1.58
Hawaii	1.01 ^b	0.16	(51)	-35.65 ^c	Oregon	1.65	0.12	(24)	5.45
Idaho	1.35	0.18	(44)	-13.81	Pennsylvania	1.73	0.06	(12)	10.58
Illinois	1.69	0.07	(20)	7.90	Rhode Island	1.99	0.24	(03)	26.58 ^c
Indiana	1.66	0.09	(22)	5.62	South Carolina	1.64	0.11	(25)	4.29
Iowa	1.38	0.12	(42)	-12.17	South Dakota	1.72	0.26	(18)	9.62
Kansas	1.22	0.12	(47)	-22.20 ^c	Tennessee	1.49	0.09	(33)	-5.10
Kentucky	1.52	0.11	(32)	-2.95	Texas	1.24 ^b	0.05	(46)	-21.07 ^c
Louisiana	1.72	0.12	(16)	9.84	Utah	1.02 ^b	0.14	(50)	-34.69 ^c
Maine	1.73	0.19	(13)	10.55	Vermont	1.76	0.30	(09)	12.17
Maryland	1.65	0.10	(23)	5.50	Virginia	1.46	0.08	(35)	-6.94
Massachusetts	1.95 ^b	0.10	(04)	24.09 ^c	Washington	1.91	0.10	(06)	21.52 ^c
Michigan	1.74	0.08	(11)	11.17	West Virginia	1.59	0.16	(29)	1.53
Minnesota	1.99 ^b	0.11	(02)	26.79 ^c	Wisconsin	1.72	0.10	(17)	9.69
Mississippi	1.45	0.13	(38)	-7.85	Wyoming	1.94	0.36	(05)	23.92 ^c
Missouri	1.73	0.10	(15)	10.06					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.18
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.38	0.00		
High Five States				
Vermont	0.68	0.15	(01)	76.83 ^c
Louisiana	0.49	0.05	(02)	28.45 ^c
Maine	0.49	0.08	(03)	26.71 ^c
Nebraska	0.47	0.07	(04)	22.31 ^c
Oregon	0.45	0.05	(05)	17.94 ^c
Low Five States				
Arizona	0.32	0.03	(42)	-15.99 ^c
Maryland	0.32	0.03	(43)	-17.02 ^c
Montana	0.29	0.08	(44)	-23.34 ^c
Utah	0.26	0.05	(45)	-32.42 ^c
Hawaii	0.21	0.05	(46)	-45.74 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.40	0.04	(18)	4.93	Montana	0.29	0.08	(44)	-23.34 ^c
Alaska	-	-	(-)	-	Nebraska	0.47	0.07	(04)	22.31 ^c
Arizona	0.32	0.03	(42)	-15.99 ^c	Nevada	0.42	0.06	(12)	8.80
Arkansas	0.42	0.05	(13)	8.68	New Hampshire	0.37	0.08	(32)	-2.71
California	0.40	0.02	(20)	3.10	New Jersey	0.41	0.03	(17)	5.87
Colorado	0.35	0.04	(36)	-9.87	New Mexico	0.34	0.06	(37)	-10.61
Connecticut	0.40	0.05	(19)	3.84	New York	0.42	0.02	(10)	10.05
Delaware	0.34	0.09	(38)	-10.67	North Carolina	0.38	0.03	(30)	-1.56
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.38	0.02	(25)	-0.42	Ohio	0.43	0.03	(07)	12.99
Georgia	0.35	0.03	(35)	-9.15	Oklahoma	0.42	0.05	(11)	9.13
Hawaii	0.21	0.05	(46)	-45.74 ^c	Oregon	0.45	0.05	(05)	17.94 ^c
Idaho	0.43	0.08	(09)	10.95	Pennsylvania	0.38	0.02	(24)	-0.09
Illinois	0.34	0.02	(39)	-11.51	Rhode Island	0.42	0.08	(14)	8.51
Indiana	0.34	0.03	(40)	-11.53	South Carolina	0.39	0.04	(21)	2.49
Iowa	0.38	0.05	(29)	-1.21	South Dakota	-	-	(-)	-
Kansas	0.38	0.05	(26)	-0.66	Tennessee	0.42	0.04	(15)	8.43
Kentucky	0.35	0.04	(33)	-7.60	Texas	0.38	0.02	(27)	-0.88
Louisiana	0.49	0.05	(02)	28.45 ^c	Utah	0.26	0.05	(45)	-32.42 ^c
Maine	0.49	0.08	(03)	26.71 ^c	Vermont	0.68	0.15	(01)	76.83 ^c
Maryland	0.32	0.03	(43)	-17.02 ^c	Virginia	0.39	0.03	(23)	0.33
Massachusetts	0.37	0.03	(31)	-2.48	Washington	0.33	0.03	(41)	-14.40
Michigan	0.41	0.03	(16)	7.47	West Virginia	0.45	0.07	(06)	17.93 ^c
Minnesota	0.39	0.04	(22)	1.19	Wisconsin	0.43	0.04	(08)	11.02
Mississippi	0.35	0.05	(34)	-8.93	Wyoming	-	-	(-)	-
Missouri	0.38	0.03	(28)	-0.90					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.19
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.48	0.01		
High Five States				
Rhode Island	0.76	0.17	(01)	58.13 ^c
Nebraska	0.76	0.13	(02)	57.79 ^c
Maine	0.62	0.14	(03)	30.13 ^c
Louisiana	0.60	0.08	(04)	25.66 ^c
Oregon	0.59	0.08	(05)	23.37 ^c
Low Five States				
Illinois	0.40	0.04	(38)	-16.57 ^c
Kansas	0.40	0.08	(39)	-17.11 ^c
Maryland	0.39	0.06	(40)	-17.87 ^c
Arizona	0.39	0.05	(41)	-19.33 ^c
Utah	0.33	0.08	(42)	-30.10 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.49	0.07	(21)	1.45	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.76	0.13	(02)	57.79 ^c
Arizona	0.39	0.05	(41)	-19.33 ^c	Nevada	0.58	0.10	(07)	22.15 ^c
Arkansas	0.45	0.08	(30)	-6.80	New Hampshire	0.59	0.14	(06)	22.29 ^c
California	0.49	0.03	(19)	2.14	New Jersey	0.51	0.05	(10)	7.17
Colorado	0.41	0.06	(37)	-15.10 ^c	New Mexico	0.49	0.10	(20)	1.93
Connecticut	0.44	0.07	(32)	-8.93	New York	0.53	0.03	(09)	10.01
Delaware	-	-	(-)	-	North Carolina	0.48	0.05	(23)	-0.75
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.47	0.03	(25)	-1.57	Ohio	0.55	0.05	(08)	14.69
Georgia	0.50	0.05	(17)	3.92	Oklahoma	0.48	0.08	(22)	0.14
Hawaii	-	-	(-)	-	Oregon	0.59	0.08	(05)	23.37 ^c
Idaho	0.43	0.11	(34)	-10.85	Pennsylvania	0.46	0.04	(26)	-4.57
Illinois	0.40	0.04	(38)	-16.57 ^c	Rhode Island	0.76	0.17	(01)	58.13 ^c
Indiana	0.45	0.06	(28)	-6.17	South Carolina	0.47	0.07	(24)	-1.48
Iowa	0.42	0.07	(35)	-12.35	South Dakota	-	-	(-)	-
Kansas	0.40	0.08	(39)	-17.11 ^c	Tennessee	0.51	0.06	(13)	6.14
Kentucky	0.44	0.07	(31)	-7.55	Texas	0.51	0.03	(15)	5.49
Louisiana	0.60	0.08	(04)	25.66 ^c	Utah	0.33	0.08	(42)	-30.10 ^c
Maine	0.62	0.14	(03)	30.13 ^c	Vermont	-	-	(-)	-
Maryland	0.39	0.06	(40)	-17.87 ^c	Virginia	0.50	0.05	(16)	4.68
Massachusetts	0.50	0.06	(18)	3.60	Washington	0.41	0.05	(36)	-13.96
Michigan	0.51	0.05	(12)	6.18	West Virginia	0.46	0.10	(27)	-4.73
Minnesota	0.51	0.06	(14)	6.11	Wisconsin	0.51	0.06	(11)	6.39
Mississippi	0.43	0.09	(33)	-9.29	Wyoming	-	-	(-)	-
Missouri	0.45	0.06	(29)	-6.49					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 9.20
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.31	0.01		
High Five States				
West Virginia	0.44	0.09	(01)	44.48 ^c
Idaho	0.41	0.10	(02)	34.82 ^c
Arkansas	0.40	0.07	(03)	32.32 ^c
Louisiana	0.40	0.06	(04)	32.28 ^c
Oklahoma	0.39	0.06	(05)	27.13 ^c
Low Five States				
Arizona	0.27	0.04	(33)	-13.04
Indiana	0.26	0.04	(34)	-13.26
Washington	0.26	0.04	(35)	-14.03
Maryland	0.25	0.04	(36)	-17.02 ^c
Georgia	0.25	0.03	(37)	-19.68 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.33	0.05	(16)	8.36	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.27	0.04	(33)	-13.04	Nevada	0.27	0.07	(31)	-9.88
Arkansas	0.40	0.07	(03)	32.32 ^c	New Hampshire	-	-	(-)	-
California	0.32	0.02	(20)	4.70	New Jersey	0.32	0.04	(21)	3.58
Colorado	0.28	0.05	(28)	-6.76	New Mexico	-	-	(-)	-
Connecticut	0.36	0.06	(08)	18.08 ^c	New York	0.34	0.02	(11)	11.17
Delaware	-	-	(-)	-	North Carolina	0.30	0.03	(24)	-1.81
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.30	0.02	(23)	-1.12	Ohio	0.33	0.03	(12)	9.77
Georgia	0.25	0.03	(37)	-19.68 ^c	Oklahoma	0.39	0.06	(05)	27.13 ^c
Hawaii	-	-	(-)	-	Oregon	0.33	0.05	(13)	9.72
Idaho	0.41	0.10	(02)	34.82 ^c	Pennsylvania	0.33	0.03	(19)	6.62
Illinois	0.29	0.03	(25)	-3.86	Rhode Island	-	-	(-)	-
Indiana	0.26	0.04	(34)	-13.26	South Carolina	0.33	0.05	(17)	7.92
Iowa	0.35	0.06	(10)	14.09	South Dakota	-	-	(-)	-
Kansas	0.37	0.07	(06)	22.17 ^c	Tennessee	0.33	0.04	(15)	8.42
Kentucky	0.27	0.05	(32)	-11.49	Texas	0.28	0.02	(30)	-9.03
Louisiana	0.40	0.06	(04)	32.28 ^c	Utah	-	-	(-)	-
Maine	0.37	0.10	(07)	19.77 ^c	Vermont	-	-	(-)	-
Maryland	0.25	0.04	(36)	-17.02 ^c	Virginia	0.29	0.04	(26)	-4.49
Massachusetts	0.28	0.04	(29)	-8.25	Washington	0.26	0.04	(35)	-14.03
Michigan	0.33	0.03	(14)	8.70	West Virginia	0.44	0.09	(01)	44.48 ^c
Minnesota	0.29	0.04	(27)	-5.49	Wisconsin	0.35	0.05	(09)	16.21 ^c
Mississippi	0.30	0.06	(22)	-0.89	Wyoming	-	-	(-)	-
Missouri	0.33	0.04	(18)	7.81					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 11.18
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.96	0.02		
High Five States				
Oklahoma	5.18 ^b	0.16	(01)	30.91 ^c
Louisiana	5.14 ^b	0.15	(02)	29.91 ^c
Alaska	5.07	0.50	(03)	27.94 ^c
North Dakota	5.03	0.36	(04)	27.14 ^c
West Virginia	4.83 ^b	0.21	(05)	22.08 ^c
Low Five States				
Connecticut	3.41 ^b	0.13	(47)	-13.86
Montana	3.39	0.25	(48)	-14.39
New York	3.34 ^b	0.06	(49)	-15.78 ^c
Hawaii	2.71 ^b	0.19	(50)	-31.57 ^c
D.C.	2.71 ^b	0.31	(51)	-31.69 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.87	0.12	(32)	-2.20	Montana	3.39	0.25	(48)	-14.39
Alaska	5.07	0.50	(03)	27.94 ^c	Nebraska	4.41	0.21	(14)	11.40
Arizona	3.80	0.11	(35)	-3.95	Nevada	3.93	0.18	(27)	-0.66
Arkansas	4.43	0.16	(13)	11.76	New Hampshire	3.59	0.22	(39)	-9.42
California	3.53 ^b	0.05	(41)	-10.96	New Jersey	3.42 ^b	0.08	(45)	-13.57
Colorado	3.66	0.13	(36)	-7.58	New Mexico	4.57	0.21	(09)	15.31 ^c
Connecticut	3.41 ^b	0.13	(47)	-13.86	New York	3.34 ^b	0.06	(49)	-15.78 ^c
Delaware	3.81	0.28	(34)	-3.73	North Carolina	4.02	0.09	(26)	1.40
D.C.	2.71 ^b	0.31	(51)	-31.69 ^c	North Dakota	5.03	0.36	(04)	27.14 ^c
Florida	3.52 ^b	0.05	(43)	-11.24	Ohio	4.41 ^b	0.08	(15)	11.28
Georgia	3.62 ^b	0.09	(38)	-8.58	Oklahoma	5.18 ^b	0.16	(01)	30.91 ^c
Hawaii	2.71 ^b	0.19	(50)	-31.57 ^c	Oregon	3.87	0.13	(33)	-2.25
Idaho	4.06	0.23	(25)	2.53	Pennsylvania	3.93	0.07	(29)	-0.80
Illinois	4.13	0.08	(23)	4.38	Rhode Island	3.55	0.24	(40)	-10.37
Indiana	4.55 ^b	0.12	(10)	14.79	South Carolina	4.15	0.13	(22)	4.85
Iowa	4.75 ^b	0.16	(06)	19.84 ^c	South Dakota	4.37	0.31	(17)	10.45
Kansas	4.57 ^b	0.17	(08)	15.51 ^c	Tennessee	4.27	0.11	(20)	7.74
Kentucky	4.72 ^b	0.14	(07)	19.09 ^c	Texas	4.52 ^b	0.07	(12)	14.05
Louisiana	5.14 ^b	0.15	(02)	29.91 ^c	Utah	3.50	0.19	(44)	-11.75
Maine	4.10	0.22	(24)	3.47	Vermont	3.64	0.32	(37)	-7.99
Maryland	3.89	0.12	(31)	-1.76	Virginia	3.89	0.10	(30)	-1.72
Massachusetts	3.41 ^b	0.10	(46)	-13.78	Washington	3.93	0.11	(28)	-0.77
Michigan	4.17	0.09	(21)	5.22	West Virginia	4.83 ^b	0.21	(05)	22.08 ^c
Minnesota	4.37	0.13	(18)	10.40	Wisconsin	4.35	0.12	(19)	9.92
Mississippi	4.53	0.17	(11)	14.45	Wyoming	3.52	0.35	(42)	-11.08
Missouri	4.40 ^b	0.11	(16)	11.03					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.19
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.76	0.03		
High Five States				
Oklahoma	7.60 ^b	0.30	(01)	31.94 ^c
North Dakota	7.29	0.66	(02)	26.63 ^c
Iowa	7.03 ^b	0.30	(03)	22.13 ^c
Louisiana	7.03 ^b	0.27	(04)	22.11 ^c
Alaska	6.98	0.89	(05)	21.15 ^c
Low Five States				
Massachusetts	4.92 ^b	0.18	(47)	-14.65
Utah	4.79	0.33	(48)	-16.80 ^c
Montana	4.78	0.43	(49)	-16.92 ^c
D.C.	4.55	0.63	(50)	-20.99 ^c
Hawaii	4.24 ^b	0.35	(51)	-26.38 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.46	0.23	(37)	-5.21	Montana	4.78	0.43	(49)	-16.92 ^c
Alaska	6.98	0.89	(05)	21.15 ^c	Nebraska	6.32	0.38	(17)	9.73
Arizona	5.48	0.19	(34)	-4.77	Nevada	5.55	0.32	(32)	-3.56
Arkansas	6.33	0.30	(16)	9.84	New Hampshire	5.47	0.41	(35)	-4.98
California	5.13 ^b	0.08	(42)	-10.98	New Jersey	5.24	0.16	(40)	-8.97
Colorado	5.36	0.23	(38)	-6.88	New Mexico	6.20	0.36	(20)	7.66
Connecticut	5.00	0.24	(46)	-13.21	New York	5.04 ^b	0.11	(45)	-12.50
Delaware	5.21	0.50	(41)	-9.55	North Carolina	6.06	0.18	(21)	5.31
D.C.	4.55	0.63	(50)	-20.99 ^c	North Dakota	7.29	0.66	(02)	26.63 ^c
Florida	5.11 ^b	0.10	(43)	-11.33	Ohio	6.22	0.15	(19)	8.02
Georgia	5.47	0.18	(36)	-5.06	Oklahoma	7.60 ^b	0.30	(01)	31.94 ^c
Hawaii	4.24 ^b	0.35	(51)	-26.38 ^c	Oregon	5.60	0.24	(31)	-2.79
Idaho	5.66	0.40	(29)	-1.65	Pennsylvania	5.60	0.13	(30)	-2.71
Illinois	5.93	0.15	(24)	2.95	Rhode Island	5.53	0.47	(33)	-3.92
Indiana	6.42	0.21	(13)	11.46	South Carolina	6.38	0.25	(15)	10.82
Iowa	7.03 ^b	0.30	(03)	22.13 ^c	South Dakota	5.90	0.54	(25)	2.53
Kansas	6.72	0.32	(07)	16.61 ^c	Tennessee	6.41	0.21	(14)	11.38
Kentucky	6.61	0.26	(09)	14.81	Texas	6.49 ^b	0.12	(11)	12.79
Louisiana	7.03 ^b	0.27	(04)	22.11 ^c	Utah	4.79	0.33	(48)	-16.80 ^c
Maine	6.01	0.41	(22)	4.36	Vermont	5.10	0.57	(44)	-11.39
Maryland	5.93	0.22	(23)	3.00	Virginia	5.87	0.19	(26)	1.93
Massachusetts	4.92 ^b	0.18	(47)	-14.65	Washington	5.69	0.20	(28)	-1.22
Michigan	5.81	0.16	(27)	0.95	West Virginia	6.75	0.37	(06)	17.24 ^c
Minnesota	6.47	0.23	(12)	12.28	Wisconsin	6.32	0.21	(18)	9.67
Mississippi	6.68	0.33	(08)	15.92 ^c	Wyoming	5.31	0.64	(39)	-7.81
Missouri	6.50	0.21	(10)	12.85					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 11.20
Cancer of the Kidney And Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.56	0.02		
High Five States				
Louisiana	3.63 ^b	0.17	(01)	42.12 ^c
Alaska	3.43	0.56	(02)	34.21 ^c
West Virginia	3.38 ^b	0.23	(03)	32.21 ^c
Oklahoma	3.27 ^b	0.17	(04)	27.94 ^c
Kentucky	3.25 ^b	0.16	(05)	27.25 ^c
Low Five States				
New Jersey	2.08 ^b	0.09	(47)	-18.79 ^c
New Hampshire	2.02	0.22	(48)	-21.18 ^c
Wyoming	1.99	0.36	(49)	-22.07 ^c
Hawaii	1.44 ^b	0.18	(50)	-43.88 ^c
D.C.	1.34 ^b	0.29	(51)	-47.41 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.64	0.13	(25)	3.21	Montana	2.16	0.26	(44)	-15.34 ^c
Alaska	3.43	0.56	(02)	34.21 ^c	Nebraska	2.80	0.22	(18)	9.40
Arizona	2.39	0.11	(36)	-6.66	Nevada	2.48	0.20	(30)	-3.01
Arkansas	2.92	0.18	(13)	14.19	New Hampshire	2.02	0.22	(48)	-21.18 ^c
California	2.25 ^b	0.05	(41)	-12.20	New Jersey	2.08 ^b	0.09	(47)	-18.79 ^c
Colorado	2.28	0.14	(39)	-10.74	New Mexico	3.21	0.23	(06)	23.23 ^c
Connecticut	2.20	0.14	(43)	-13.96	New York	2.09 ^b	0.06	(46)	-18.34 ^c
Delaware	2.76	0.32	(20)	7.88	North Carolina	2.54	0.10	(28)	-0.80
D.C.	1.34 ^b	0.29	(51)	-47.41 ^c	North Dakota	3.15	0.38	(07)	23.23 ^c
Florida	2.23 ^b	0.06	(42)	-12.70	Ohio	3.02 ^b	0.09	(10)	17.99 ^c
Georgia	2.26	0.10	(40)	-11.78	Oklahoma	3.27 ^b	0.17	(04)	27.94 ^c
Hawaii	1.44 ^b	0.18	(50)	-43.88 ^c	Oregon	2.48	0.15	(29)	-2.90
Idaho	2.66	0.26	(24)	4.12	Pennsylvania	2.63	0.08	(26)	2.97
Illinois	2.77	0.09	(19)	8.22	Rhode Island	2.16	0.25	(45)	-15.56 ^c
Indiana	3.06 ^b	0.13	(09)	19.61 ^c	South Carolina	2.46	0.13	(33)	-3.72
Iowa	3.00	0.18	(11)	17.20 ^c	South Dakota	3.10	0.35	(08)	21.31 ^c
Kansas	2.88	0.19	(15)	12.78	Tennessee	2.63	0.12	(27)	2.94
Kentucky	3.25 ^b	0.16	(05)	27.25 ^c	Texas	2.95 ^b	0.07	(12)	15.19 ^c
Louisiana	3.63 ^b	0.17	(01)	42.12 ^c	Utah	2.41	0.21	(34)	-5.82
Maine	2.73	0.24	(22)	6.60	Vermont	2.47	0.36	(32)	-3.53
Maryland	2.38	0.12	(37)	-6.95	Virginia	2.39	0.10	(35)	-6.52
Massachusetts	2.31	0.10	(38)	-9.51	Washington	2.47	0.12	(31)	-3.23
Michigan	2.89	0.10	(14)	13.06	West Virginia	3.38 ^b	0.23	(03)	32.21 ^c
Minnesota	2.71	0.13	(23)	5.82	Wisconsin	2.75	0.13	(21)	7.71
Mississippi	2.87	0.18	(16)	12.20	Wyoming	1.99	0.36	(49)	-22.07 ^c
Missouri	2.80	0.12	(17)	9.47					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.18
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.10	0.01		
High Five States				
D.C.	1.79	0.25	(01)	62.63 ^c
West Virginia	1.49	0.11	(02)	34.70 ^c
Kentucky	1.40 ^b	0.08	(03)	27.23 ^c
Louisiana	1.39 ^b	0.08	(04)	26.44 ^c
Alabama	1.39 ^b	0.07	(05)	26.03 ^c
Low Five States				
Oregon	0.78 ^b	0.06	(47)	-29.63 ^c
Idaho	0.75 ^b	0.10	(48)	-31.74 ^c
Wyoming	0.67	0.15	(49)	-39.26 ^c
Colorado	0.67 ^b	0.05	(50)	-39.27 ^c
Utah	0.37 ^b	0.06	(51)	-66.86 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.39 ^b	0.07	(05)	26.03 ^c	Montana	1.14	0.14	(22)	3.36
Alaska	1.01	0.22	(34)	-8.26	Nebraska	0.86	0.09	(42)	-21.61 ^c
Arizona	0.86 ^b	0.05	(43)	-21.87 ^c	Nevada	1.11	0.09	(26)	0.24
Arkansas	1.03	0.08	(33)	-6.31	New Hampshire	1.04	0.12	(31)	-5.62
California	0.83 ^b	0.02	(44)	-24.63 ^c	New Jersey	1.20	0.05	(19)	8.42
Colorado	0.67 ^b	0.05	(50)	-39.27 ^c	New Mexico	1.00	0.10	(35)	-9.69
Connecticut	0.90	0.07	(39)	-18.07 ^c	New York	1.16	0.03	(21)	4.96
Delaware	1.29	0.16	(12)	16.52 ^c	North Carolina	1.22	0.05	(16)	10.28
D.C.	1.79	0.25	(01)	62.63 ^c	North Dakota	1.03	0.16	(32)	-6.30
Florida	1.22 ^b	0.03	(15)	10.79	Ohio	1.36 ^b	0.05	(07)	23.52 ^c
Georgia	1.08	0.05	(28)	-2.50	Oklahoma	1.09	0.07	(27)	-1.07
Hawaii	0.82	0.10	(45)	-25.74 ^c	Oregon	0.78 ^b	0.06	(47)	-29.63 ^c
Idaho	0.75 ^b	0.10	(48)	-31.74 ^c	Pennsylvania	1.21	0.04	(17)	9.84
Illinois	1.19	0.04	(20)	7.45	Rhode Island	1.25	0.14	(13)	13.38
Indiana	1.20	0.06	(18)	9.24	South Carolina	1.35	0.07	(08)	22.73 ^c
Iowa	0.96	0.07	(36)	-12.81	South Dakota	0.94	0.14	(37)	-15.16 ^c
Kansas	0.78 ^b	0.07	(46)	-28.90 ^c	Tennessee	1.36 ^b	0.06	(06)	23.72 ^c
Kentucky	1.40 ^b	0.08	(03)	27.23 ^c	Texas	1.11	0.03	(24)	0.65
Louisiana	1.39 ^b	0.08	(04)	26.44 ^c	Utah	0.37 ^b	0.06	(51)	-66.86 ^c
Maine	1.35	0.13	(09)	22.06 ^c	Vermont	1.07	0.17	(29)	-3.14
Maryland	1.30	0.07	(11)	17.57 ^c	Virginia	1.05	0.05	(30)	-5.08
Massachusetts	1.11	0.06	(25)	0.45	Washington	0.90 ^b	0.05	(40)	-18.52 ^c
Michigan	1.24	0.05	(14)	12.68	West Virginia	1.49	0.11	(02)	34.70 ^c
Minnesota	0.87 ^b	0.06	(41)	-21.22 ^c	Wisconsin	0.93	0.05	(38)	-15.56 ^c
Mississippi	1.32	0.09	(10)	19.99 ^c	Wyoming	0.67	0.15	(49)	-39.26 ^c
Missouri	1.12	0.06	(23)	1.72					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.19
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	1.97	0.02		
High Five States				
D.C.	3.40	0.54	(01)	72.30 ^c
Alabama	2.55 ^b	0.15	(02)	29.48 ^c
West Virginia	2.55	0.22	(03)	29.04 ^c
Tennessee	2.48 ^b	0.13	(04)	25.81 ^c
Mississippi	2.48	0.20	(05)	25.77 ^c
Low Five States				
Oregon	1.31 ^b	0.12	(47)	-33.70 ^c
Idaho	1.30 ^b	0.19	(48)	-34.30 ^c
Wyoming	1.12	0.29	(49)	-42.98 ^c
Colorado	1.11 ^b	0.11	(50)	-43.94 ^c
Utah	0.60 ^b	0.11	(51)	-69.38 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.55 ^b	0.15	(02)	29.48 ^c	Montana	2.05	0.28	(21)	4.10
Alaska	1.85	0.46	(31)	-6.11	Nebraska	1.61	0.19	(40)	-18.56 ^c
Arizona	1.49 ^b	0.10	(45)	-24.65 ^c	Nevada	1.76	0.18	(35)	-10.78
Arkansas	1.83	0.16	(32)	-7.16	New Hampshire	1.91	0.25	(30)	-3.06
California	1.52 ^b	0.04	(44)	-23.05 ^c	New Jersey	2.06	0.10	(20)	4.58
Colorado	1.11 ^b	0.11	(50)	-43.94 ^c	New Mexico	1.71	0.19	(36)	-13.22
Connecticut	1.71	0.14	(37)	-13.43	New York	2.08	0.07	(19)	5.52
Delaware	2.04	0.30	(22)	3.52	North Carolina	2.24	0.10	(13)	13.62
D.C.	3.40	0.54	(01)	72.30 ^c	North Dakota	1.94	0.33	(28)	-1.86
Florida	2.17	0.06	(16)	9.96	Ohio	2.42 ^b	0.09	(08)	22.85 ^c
Georgia	2.02	0.11	(24)	2.23	Oklahoma	1.93	0.15	(29)	-2.17
Hawaii	1.53	0.21	(42)	-22.20 ^c	Oregon	1.31 ^b	0.12	(47)	-33.70 ^c
Idaho	1.30 ^b	0.19	(48)	-34.30 ^c	Pennsylvania	2.20	0.08	(14)	11.31
Illinois	2.16	0.09	(17)	9.47	Rhode Island	2.24	0.30	(12)	13.66
Indiana	1.99	0.12	(25)	0.97	South Carolina	2.47	0.15	(06)	25.22 ^c
Iowa	1.64	0.14	(39)	-16.69 ^c	South Dakota	1.77	0.30	(34)	-10.34
Kansas	1.38 ^b	0.14	(46)	-30.00 ^c	Tennessee	2.48 ^b	0.13	(04)	25.81 ^c
Kentucky	2.40	0.15	(10)	21.48 ^c	Texas	2.09	0.07	(18)	5.83
Louisiana	2.45	0.16	(07)	24.26 ^c	Utah	0.60 ^b	0.11	(51)	-69.38 ^c
Maine	2.41	0.25	(09)	22.24 ^c	Vermont	1.78	0.33	(33)	-9.97
Maryland	2.30	0.14	(11)	16.70 ^c	Virginia	1.95	0.11	(27)	-0.94
Massachusetts	1.98	0.11	(26)	0.44	Washington	1.53 ^b	0.10	(43)	-22.44 ^c
Michigan	2.19	0.10	(15)	10.90	West Virginia	2.55	0.22	(03)	29.04 ^c
Minnesota	1.57 ^b	0.11	(41)	-20.38 ^c	Wisconsin	1.70	0.11	(38)	-13.86
Mississippi	2.48	0.20	(05)	25.77 ^c	Wyoming	1.12	0.29	(49)	-42.98 ^c
Missouri	2.04	0.12	(23)	3.20					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.20
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.42	0.01		
High Five States				
Delaware	0.66	0.15	(01)	56.70 ^c
Kentucky	0.64	0.07	(02)	51.95 ^c
West Virginia	0.60	0.10	(03)	43.21 ^c
Louisiana	0.58	0.07	(04)	37.30 ^c
Indiana	0.56	0.05	(05)	34.24 ^c
Low Five States				
Wisconsin	0.33	0.04	(35)	-22.62 ^c
Minnesota	0.32	0.05	(36)	-24.31 ^c
Connecticut	0.30	0.05	(37)	-27.40 ^c
California	0.30 ^b	0.02	(38)	-29.09 ^c
Kansas	0.30	0.06	(39)	-29.73 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.47	0.06	(15)	12.28	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.34	0.04	(32)	-17.99 ^c	Nevada	0.54	0.09	(08)	27.95 ^c
Arkansas	0.39	0.07	(28)	-6.32	New Hampshire	-	-	(-)	-
California	0.30 ^b	0.02	(38)	-29.09 ^c	New Jersey	0.54	0.04	(07)	29.74 ^c
Colorado	0.33	0.05	(34)	-20.88 ^c	New Mexico	0.40	0.08	(27)	-4.83
Connecticut	0.30	0.05	(37)	-27.40 ^c	New York	0.47	0.03	(16)	11.60
Delaware	0.66	0.15	(01)	56.70 ^c	North Carolina	0.44	0.04	(22)	4.12
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.43	0.03	(23)	2.60	Ohio	0.53	0.04	(09)	26.57 ^c
Georgia	0.37	0.04	(30)	-12.67	Oklahoma	0.43	0.06	(24)	2.35
Hawaii	-	-	(-)	-	Oregon	0.36	0.06	(31)	-15.00 ^c
Idaho	-	-	(-)	-	Pennsylvania	0.45	0.03	(17)	8.28
Illinois	0.44	0.03	(20)	5.88	Rhode Island	0.55	0.13	(06)	31.56 ^c
Indiana	0.56	0.05	(05)	34.24 ^c	South Carolina	0.50	0.06	(13)	18.72 ^c
Iowa	0.45	0.07	(19)	6.01	South Dakota	-	-	(-)	-
Kansas	0.30	0.06	(39)	-29.73 ^c	Tennessee	0.50	0.05	(12)	19.73 ^c
Kentucky	0.64	0.07	(02)	51.95 ^c	Texas	0.33 ^b	0.02	(33)	-20.76 ^c
Louisiana	0.58	0.07	(04)	37.30 ^c	Utah	-	-	(-)	-
Maine	0.50	0.11	(14)	18.54 ^c	Vermont	-	-	(-)	-
Maryland	0.51	0.06	(10)	22.60 ^c	Virginia	0.37	0.04	(29)	-11.90
Massachusetts	0.45	0.05	(18)	7.69	Washington	0.40	0.05	(26)	-4.72
Michigan	0.50	0.04	(11)	20.12 ^c	West Virginia	0.60	0.10	(03)	43.21 ^c
Minnesota	0.32	0.05	(36)	-24.31 ^c	Wisconsin	0.33	0.04	(35)	-22.62 ^c
Mississippi	0.44	0.07	(21)	4.27	Wyoming	-	-	(-)	-
Missouri	0.42	0.05	(25)	0.78					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.26
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.01	0.02		
High Five States				
Wyoming	8.39	0.55	(01)	19.63 ^c
Wisconsin	7.97 ^b	0.16	(02)	13.61
Minnesota	7.96 ^b	0.17	(03)	13.43
West Virginia	7.94	0.27	(04)	13.24
Kansas	7.92 ^b	0.23	(05)	12.91
Low Five States				
Nevada	6.47	0.23	(47)	-7.71
Arizona	6.42 ^b	0.14	(48)	-8.50
New Mexico	6.17	0.24	(49)	-11.99
D.C.	5.67	0.45	(50)	-19.14 ^c
Hawaii	5.49 ^b	0.27	(51)	-21.70 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.53	0.17	(12)	7.40	Montana	7.22	0.36	(22)	2.88
Alaska	7.01	0.60	(27)	-0.04	Nebraska	6.82	0.26	(32)	-2.76
Arizona	6.42 ^b	0.14	(48)	-8.50	Nevada	6.47	0.23	(47)	-7.71
Arkansas	7.63	0.22	(08)	8.75	New Hampshire	6.91	0.31	(29)	-1.50
California	6.63 ^b	0.06	(42)	-5.54	New Jersey	6.66	0.12	(37)	-5.05
Colorado	6.64	0.18	(39)	-5.36	New Mexico	6.17	0.24	(49)	-11.99
Connecticut	6.75	0.18	(34)	-3.71	New York	6.62 ^b	0.08	(43)	-5.55
Delaware	6.80	0.37	(33)	-3.08	North Carolina	6.89	0.12	(30)	-1.75
D.C.	5.67	0.45	(50)	-19.14 ^c	North Dakota	7.58	0.44	(10)	8.12
Florida	6.63 ^b	0.07	(40)	-5.41	Ohio	7.44 ^b	0.11	(18)	6.13
Georgia	6.72	0.13	(35)	-4.20	Oklahoma	7.61	0.20	(09)	8.52
Hawaii	5.49 ^b	0.27	(51)	-21.70 ^c	Oregon	6.97	0.18	(28)	-0.67
Idaho	7.12	0.31	(25)	1.46	Pennsylvania	7.48 ^b	0.10	(16)	6.61
Illinois	7.32	0.11	(20)	4.39	Rhode Island	7.03	0.34	(26)	0.23
Indiana	7.77 ^b	0.15	(06)	10.82	South Carolina	6.62	0.17	(44)	-5.58
Iowa	7.15	0.20	(23)	1.91	South Dakota	7.56	0.41	(11)	7.79
Kansas	7.92 ^b	0.23	(05)	12.91	Tennessee	7.53	0.15	(13)	7.33
Kentucky	7.49	0.18	(15)	6.72	Texas	6.84	0.08	(31)	-2.48
Louisiana	7.40	0.18	(19)	5.47	Utah	6.50	0.26	(46)	-7.30
Maine	7.14	0.30	(24)	1.78	Vermont	6.63	0.44	(41)	-5.44
Maryland	6.57	0.15	(45)	-6.32	Virginia	6.71	0.13	(36)	-4.34
Massachusetts	6.64	0.14	(38)	-5.35	Washington	7.29	0.15	(21)	3.97
Michigan	7.45 ^b	0.12	(17)	6.23	West Virginia	7.94	0.27	(04)	13.24
Minnesota	7.96 ^b	0.17	(03)	13.43	Wisconsin	7.97 ^b	0.16	(02)	13.61
Mississippi	7.66	0.23	(07)	9.24	Wyoming	8.39	0.55	(01)	19.63 ^c
Missouri	7.50	0.15	(14)	6.90					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.27
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	9.43	0.04		
High Five States				
Wisconsin	11.01 ^b	0.29	(01)	16.79 ^c
Wyoming	10.87	0.95	(02)	15.34 ^c
Minnesota	10.83 ^b	0.31	(03)	14.95
West Virginia	10.55	0.48	(04)	11.91
Indiana	10.53 ^b	0.28	(05)	11.77
Low Five States				
New Hampshire	8.54	0.54	(47)	-9.40
D.C.	8.39	0.88	(48)	-11.02
Arizona	8.35 ^b	0.24	(49)	-11.45
New Mexico	7.82 ^b	0.42	(50)	-16.99 ^c
Hawaii	7.01 ^b	0.46	(51)	-25.68 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.51	0.33	(07)	11.50	Montana	9.51	0.62	(28)	0.91
Alaska	9.81	1.10	(23)	4.07	Nebraska	9.15	0.47	(34)	-2.92
Arizona	8.35 ^b	0.24	(49)	-11.45	Nevada	8.88	0.42	(41)	-5.81
Arkansas	10.45	0.40	(08)	10.85	New Hampshire	8.54	0.54	(47)	-9.40
California	8.59 ^b	0.11	(46)	-8.87	New Jersey	9.07	0.22	(36)	-3.76
Colorado	8.81	0.31	(43)	-6.48	New Mexico	7.82 ^b	0.42	(50)	-16.99 ^c
Connecticut	9.21	0.33	(32)	-2.32	New York	9.00	0.14	(39)	-4.48
Delaware	9.62	0.68	(25)	2.05	North Carolina	9.59	0.23	(26)	1.76
D.C.	8.39	0.88	(48)	-11.02	North Dakota	10.53	0.79	(06)	11.69
Florida	8.81 ^b	0.13	(44)	-6.51	Ohio	10.06	0.20	(18)	6.75
Georgia	9.26	0.25	(31)	-1.77	Oklahoma	10.38	0.36	(10)	10.14
Hawaii	7.01 ^b	0.46	(51)	-25.68 ^c	Oregon	9.16	0.32	(33)	-2.80
Idaho	9.53	0.54	(27)	1.11	Pennsylvania	10.12 ^b	0.18	(16)	7.41
Illinois	9.93	0.19	(21)	5.31	Rhode Island	9.45	0.61	(30)	0.22
Indiana	10.53 ^b	0.28	(05)	11.77	South Carolina	9.03	0.31	(37)	-4.15
Iowa	9.73	0.36	(24)	3.19	South Dakota	10.26	0.72	(14)	8.88
Kansas	10.33	0.40	(13)	9.59	Tennessee	10.33	0.28	(12)	9.61
Kentucky	10.41	0.34	(09)	10.41	Texas	9.00	0.15	(38)	-4.47
Louisiana	10.16	0.34	(15)	7.75	Utah	8.84	0.45	(42)	-6.19
Maine	9.98	0.54	(20)	5.86	Vermont	9.46	0.80	(29)	0.37
Maryland	8.76	0.28	(45)	-7.08	Virginia	9.08	0.24	(35)	-3.63
Massachusetts	8.96	0.25	(40)	-4.98	Washington	9.90	0.27	(22)	5.09
Michigan	10.01	0.21	(19)	6.19	West Virginia	10.55	0.48	(04)	11.91
Minnesota	10.83 ^b	0.31	(03)	14.95	Wisconsin	11.01 ^b	0.29	(01)	16.79 ^c
Mississippi	10.08	0.42	(17)	7.00	Wyoming	10.87	0.95	(02)	15.34 ^c
Missouri	10.33	0.28	(11)	9.65					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.28
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.27	0.02		
High Five States				
Wyoming	6.41	0.66	(01)	21.54 ^c
Kansas	6.12	0.27	(02)	16.08 ^c
West Virginia	6.10	0.31	(03)	15.79 ^c
Mississippi	5.96	0.26	(04)	13.06
Indiana	5.92 ^b	0.18	(05)	12.39
Low Five States				
Delaware	4.56	0.40	(47)	-13.54
Vermont	4.53	0.49	(48)	-13.95
Nevada	4.53	0.26	(49)	-14.00
Hawaii	4.28	0.33	(50)	-18.83 ^c
D.C.	3.86	0.48	(51)	-26.76 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.53	0.20	(16)	4.98	Montana	5.45	0.43	(20)	3.37
Alaska	4.74	0.67	(45)	-10.10	Nebraska	5.12	0.31	(31)	-2.93
Arizona	4.90	0.17	(43)	-7.06	Nevada	4.53	0.26	(49)	-14.00
Arkansas	5.61	0.25	(09)	6.38	New Hampshire	5.78	0.39	(07)	9.58
California	5.14	0.07	(29)	-2.54	New Jersey	5.02	0.14	(34)	-4.67
Colorado	4.98	0.20	(39)	-5.60	New Mexico	4.84	0.29	(44)	-8.15
Connecticut	5.01	0.21	(36)	-5.00	New York	4.97	0.09	(40)	-5.72
Delaware	4.56	0.40	(47)	-13.54	North Carolina	5.08	0.14	(32)	-3.68
D.C.	3.86	0.48	(51)	-26.76 ^c	North Dakota	5.39	0.51	(23)	2.20
Florida	4.95 ^b	0.09	(41)	-6.16	Ohio	5.59	0.12	(11)	6.12
Georgia	5.02	0.15	(35)	-4.82	Oklahoma	5.60	0.23	(10)	6.19
Hawaii	4.28	0.33	(50)	-18.83 ^c	Oregon	5.34	0.22	(26)	1.24
Idaho	5.17	0.36	(28)	-1.83	Pennsylvania	5.58	0.11	(13)	5.86
Illinois	5.54	0.12	(15)	5.11	Rhode Island	5.41	0.40	(22)	2.70
Indiana	5.92 ^b	0.18	(05)	12.39	South Carolina	4.93	0.19	(42)	-6.45
Iowa	5.36	0.23	(24)	1.64	South Dakota	5.45	0.46	(18)	3.47
Kansas	6.12	0.27	(02)	16.08 ^c	Tennessee	5.58	0.17	(14)	5.86
Kentucky	5.42	0.21	(21)	2.85	Texas	5.26	0.09	(27)	-0.18
Louisiana	5.48	0.21	(17)	3.91	Utah	4.66	0.29	(46)	-11.61
Maine	4.98	0.33	(38)	-5.52	Vermont	4.53	0.49	(48)	-13.95
Maryland	5.05	0.18	(33)	-4.13	Virginia	4.99	0.15	(37)	-5.26
Massachusetts	5.13	0.16	(30)	-2.59	Washington	5.35	0.17	(25)	1.53
Michigan	5.59	0.14	(12)	6.12	West Virginia	6.10	0.31	(03)	15.79 ^c
Minnesota	5.83	0.20	(06)	10.68	Wisconsin	5.72	0.18	(08)	8.55
Mississippi	5.96	0.26	(04)	13.06	Wyoming	6.41	0.66	(01)	21.54 ^c
Missouri	5.45	0.17	(19)	3.42					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.18
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.76	0.02		
High Five States				
D.C.	8.70 ^b	0.55	(01)	51.03 ^c
Louisiana	7.58 ^b	0.18	(02)	31.45 ^c
Texas	7.52 ^b	0.08	(03)	30.40 ^c
Hawaii	7.28 ^b	0.31	(04)	26.25 ^c
California	7.03 ^b	0.06	(05)	22.01 ^c
Low Five States				
South Dakota	4.27 ^b	0.30	(47)	-25.93 ^c
Utah	4.13 ^b	0.20	(48)	-28.37 ^c
Montana	4.07 ^b	0.26	(49)	-29.31 ^c
Idaho	4.01 ^b	0.23	(50)	-30.44 ^c
North Dakota	3.74 ^b	0.31	(51)	-35.15 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.94	0.15	(12)	2.98	Montana	4.07 ^b	0.26	(49)	-29.31 ^c
Alaska	5.71	0.49	(18)	-0.97	Nebraska	4.41 ^b	0.21	(46)	-23.52 ^c
Arizona	5.57	0.13	(22)	-3.40	Nevada	5.84	0.21	(14)	1.34
Arkansas	6.20	0.19	(08)	7.61	New Hampshire	4.53 ^b	0.25	(44)	-21.44 ^c
California	7.03 ^b	0.06	(05)	22.01 ^c	New Jersey	5.34 ^b	0.10	(26)	-7.39
Colorado	5.04 ^b	0.15	(35)	-12.51	New Mexico	6.45	0.24	(07)	11.86
Connecticut	5.04 ^b	0.16	(37)	-12.56	New York	5.78	0.07	(16)	0.29
Delaware	6.06	0.34	(11)	5.13	North Carolina	5.22 ^b	0.10	(31)	-9.38
D.C.	8.70 ^b	0.55	(01)	51.03 ^c	North Dakota	3.74 ^b	0.31	(51)	-35.15 ^c
Florida	5.53 ^b	0.07	(24)	-4.04	Ohio	5.06 ^b	0.09	(34)	-12.12
Georgia	5.22 ^b	0.11	(32)	-9.49	Oklahoma	5.69	0.17	(19)	-1.24
Hawaii	7.28 ^b	0.31	(04)	26.25 ^c	Oregon	5.62	0.16	(20)	-2.45
Idaho	4.01 ^b	0.23	(50)	-30.44 ^c	Pennsylvania	5.32 ^b	0.08	(27)	-7.65
Illinois	5.46	0.09	(25)	-5.18	Rhode Island	5.92	0.31	(13)	2.67
Indiana	4.74 ^b	0.12	(40)	-17.75 ^c	South Carolina	5.61	0.15	(21)	-2.65
Iowa	4.55 ^b	0.16	(43)	-20.99 ^c	South Dakota	4.27 ^b	0.30	(47)	-25.93 ^c
Kansas	4.51 ^b	0.17	(45)	-21.73 ^c	Tennessee	5.84	0.13	(15)	1.34
Kentucky	5.22 ^b	0.15	(33)	-9.49	Texas	7.52 ^b	0.08	(03)	30.40 ^c
Louisiana	7.58 ^b	0.18	(02)	31.45 ^c	Utah	4.13 ^b	0.20	(48)	-28.37 ^c
Maine	4.70 ^b	0.24	(41)	-18.46 ^c	Vermont	5.31	0.39	(28)	-7.79
Maryland	5.76	0.14	(17)	0.02	Virginia	5.31 ^b	0.11	(29)	-7.95
Massachusetts	6.13	0.13	(10)	6.34	Washington	6.17	0.13	(09)	7.08
Michigan	5.30 ^b	0.10	(30)	-8.11	West Virginia	4.92 ^b	0.21	(39)	-14.71
Minnesota	4.93 ^b	0.13	(38)	-14.46	Wisconsin	5.04 ^b	0.13	(36)	-12.56
Mississippi	6.84 ^b	0.21	(06)	18.61 ^c	Wyoming	4.61	0.40	(42)	-19.93 ^c
Missouri	5.54	0.13	(23)	-3.86					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.19
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.54	0.03		
High Five States				
D.C.	13.97 ^b	1.05	(01)	63.50 ^c
Louisiana	11.57 ^b	0.34	(02)	35.42 ^c
Texas	11.00 ^b	0.15	(03)	28.73 ^c
Hawaii	10.47 ^b	0.54	(04)	22.54 ^c
Mississippi	10.27 ^b	0.40	(05)	20.24 ^c
Low Five States				
Utah	5.68 ^b	0.34	(47)	-33.56 ^c
Idaho	5.53 ^b	0.39	(48)	-35.27 ^c
Montana	5.47 ^b	0.44	(49)	-35.92 ^c
North Dakota	5.43 ^b	0.55	(50)	-36.49 ^c
South Dakota	5.19 ^b	0.49	(51)	-39.24 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.86	0.28	(14)	3.74	Montana	5.47 ^b	0.44	(49)	-35.92 ^c
Alaska	7.97	0.86	(29)	-6.73	Nebraska	6.49 ^b	0.38	(45)	-24.06 ^c
Arizona	7.81	0.22	(30)	-8.60	Nevada	8.40	0.38	(18)	-1.64
Arkansas	9.20	0.35	(10)	7.69	New Hampshire	6.59 ^b	0.45	(42)	-22.81 ^c
California	10.26 ^b	0.11	(06)	20.08 ^c	New Jersey	7.99	0.20	(28)	-6.51
Colorado	7.40 ^b	0.27	(36)	-13.41	New Mexico	9.34	0.43	(09)	9.34
Connecticut	7.63	0.29	(35)	-10.67	New York	8.75	0.14	(16)	2.39
Delaware	9.50	0.64	(08)	11.23	North Carolina	8.00	0.19	(27)	-6.39
D.C.	13.97 ^b	1.05	(01)	63.50 ^c	North Dakota	5.43 ^b	0.55	(50)	-36.49 ^c
Florida	8.37	0.12	(19)	-1.98	Ohio	7.65 ^b	0.16	(34)	-10.47
Georgia	7.70 ^b	0.21	(32)	-9.82	Oklahoma	8.21	0.30	(21)	-3.87
Hawaii	10.47 ^b	0.54	(04)	22.54 ^c	Oregon	8.16	0.28	(22)	-4.49
Idaho	5.53 ^b	0.39	(48)	-35.27 ^c	Pennsylvania	8.00 ^b	0.15	(26)	-6.32
Illinois	8.07	0.17	(25)	-5.53	Rhode Island	8.88	0.57	(13)	3.92
Indiana	6.89 ^b	0.21	(40)	-19.35 ^c	South Carolina	8.47	0.28	(17)	-0.80
Iowa	6.35 ^b	0.28	(46)	-25.62 ^c	South Dakota	5.19 ^b	0.49	(51)	-39.24 ^c
Kansas	6.56 ^b	0.30	(43)	-23.23 ^c	Tennessee	8.99	0.25	(11)	5.27
Kentucky	7.78	0.28	(31)	-8.94	Texas	11.00 ^b	0.15	(03)	28.73 ^c
Louisiana	11.57 ^b	0.34	(02)	35.42 ^c	Utah	5.68 ^b	0.34	(47)	-33.56 ^c
Maine	6.94 ^b	0.43	(39)	-18.81 ^c	Vermont	8.34	0.72	(20)	-2.39
Maryland	8.83	0.26	(15)	3.34	Virginia	8.12	0.22	(24)	-5.00
Massachusetts	9.73 ^b	0.25	(07)	13.86	Washington	8.99	0.24	(12)	5.21
Michigan	7.70 ^b	0.18	(33)	-9.83	West Virginia	6.88 ^b	0.37	(41)	-19.49 ^c
Minnesota	7.16 ^b	0.24	(38)	-16.22 ^c	Wisconsin	7.35 ^b	0.23	(37)	-14.02
Mississippi	10.27 ^b	0.40	(05)	20.24 ^c	Wyoming	6.51	0.71	(44)	-23.75 ^c
Missouri	8.16	0.23	(23)	-4.54					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.20
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.43	0.02		
High Five States				
Texas	4.53 ^b	0.09	(01)	32.02 ^c
D.C.	4.47	0.53	(02)	30.29 ^c
Louisiana	4.33 ^b	0.18	(03)	26.35 ^c
Hawaii	4.33	0.32	(04)	26.31 ^c
California	4.27 ^b	0.07	(05)	24.62 ^c
Low Five States				
New Hampshire	2.72	0.27	(47)	-20.72 ^c
Kansas	2.70 ^b	0.18	(48)	-21.20 ^c
Idaho	2.68	0.26	(49)	-21.97 ^c
Nebraska	2.66	0.22	(50)	-22.46 ^c
North Dakota	2.22 ^b	0.32	(51)	-35.27 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.61	0.16	(10)	5.14	Montana	2.76	0.30	(45)	-19.50 ^c
Alaska	3.51	0.55	(14)	2.35	Nebraska	2.66	0.22	(50)	-22.46 ^c
Arizona	3.55	0.14	(12)	3.57	Nevada	3.51	0.23	(15)	2.34
Arkansas	3.69	0.20	(08)	7.56	New Hampshire	2.72	0.27	(47)	-20.72 ^c
California	4.27 ^b	0.07	(05)	24.62 ^c	New Jersey	3.23	0.11	(26)	-5.88
Colorado	2.99	0.16	(37)	-12.89	New Mexico	3.91	0.26	(07)	14.07
Connecticut	2.92	0.16	(42)	-14.79	New York	3.38	0.07	(17)	-1.46
Delaware	3.19	0.34	(29)	-7.11	North Carolina	2.94 ^b	0.10	(41)	-14.22
D.C.	4.47	0.53	(02)	26.31 ^c	North Dakota	2.22 ^b	0.32	(51)	-35.27 ^c
Florida	3.08 ^b	0.07	(32)	-10.28	Ohio	2.97 ^b	0.09	(38)	-13.40
Georgia	3.21	0.12	(28)	-6.36	Oklahoma	3.52	0.18	(13)	2.60
Hawaii	4.33	0.32	(04)	26.31 ^c	Oregon	3.35	0.17	(19)	-2.30
Idaho	2.68	0.26	(49)	-21.97 ^c	Pennsylvania	3.11 ^b	0.08	(31)	-9.24
Illinois	3.38	0.10	(18)	-1.55	Rhode Island	3.58	0.33	(11)	4.47
Indiana	3.02	0.13	(35)	-11.96	South Carolina	3.23	0.16	(27)	-5.89
Iowa	2.97	0.17	(39)	-13.49	South Dakota	3.41	0.37	(16)	-0.64
Kansas	2.70 ^b	0.18	(48)	-21.20 ^c	Tennessee	3.31	0.13	(21)	-3.40
Kentucky	3.12	0.15	(30)	-9.14	Texas	4.53 ^b	0.09	(01)	32.02 ^c
Louisiana	4.33 ^b	0.18	(03)	26.35 ^c	Utah	2.75	0.22	(46)	-19.94 ^c
Maine	2.89	0.25	(43)	-15.81 ^c	Vermont	2.78	0.38	(44)	-19.00 ^c
Maryland	3.30	0.14	(22)	-3.70	Virginia	3.00 ^b	0.12	(36)	-12.48
Massachusetts	3.27	0.13	(24)	-4.71	Washington	3.69	0.14	(09)	7.45
Michigan	3.29	0.10	(23)	-4.19	West Virginia	3.26	0.23	(25)	-5.06
Minnesota	3.04	0.14	(34)	-11.39	Wisconsin	3.07	0.13	(33)	-10.56
Mississippi	4.10	0.22	(06)	19.68 ^c	Wyoming	2.96	0.45	(40)	-13.76
Missouri	3.31	0.13	(20)	-3.39					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.24
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	48.37	0.05		
High Five States				
Kentucky	72.03 ^b	0.56	(01)	48.91 ^c
Arkansas	63.79 ^b	0.62	(02)	31.88 ^c
West Virginia	63.77 ^b	0.74	(03)	31.82 ^c
Tennessee	62.83 ^b	0.43	(04)	29.88 ^c
Mississippi	62.63 ^b	0.64	(05)	29.48 ^c
Low Five States				
California	37.48 ^b	0.15	(47)	-22.52 ^c
Colorado	35.64 ^b	0.41	(48)	-26.33 ^c
Hawaii	34.84 ^b	0.68	(49)	-27.98 ^c
New Mexico	34.25 ^b	0.57	(50)	-29.20 ^c
Utah	20.87 ^b	0.46	(51)	-56.87 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	59.65 ^b	0.48	(07)	23.31 ^c	Montana	45.10 ^b	0.89	(36)	-6.77 ^c
Alaska	53.13	1.61	(17)	9.84 ^c	Nebraska	45.22 ^b	0.68	(35)	-6.52 ^c
Arizona	39.20 ^b	0.34	(46)	-18.97 ^c	Nevada	50.92 ^b	0.65	(22)	5.27 ^c
Arkansas	63.79 ^b	0.62	(02)	31.88 ^c	New Hampshire	49.88	0.84	(24)	3.11
California	37.48 ^b	0.15	(47)	-22.52 ^c	New Jersey	43.60 ^b	0.30	(39)	-9.87 ^c
Colorado	35.64 ^b	0.41	(48)	-26.33 ^c	New Mexico	34.25 ^b	0.57	(50)	-29.20 ^c
Connecticut	43.02 ^b	0.46	(40)	-11.06 ^c	New York	42.78 ^b	0.20	(42)	-11.57 ^c
Delaware	55.08 ^b	1.05	(12)	13.87 ^c	North Carolina	54.41 ^b	0.33	(13)	12.47 ^c
D.C.	44.38	1.27	(38)	-8.25 ^c	North Dakota	40.35 ^b	1.03	(44)	-16.58 ^c
Florida	47.55 ^b	0.20	(31)	-1.70	Ohio	56.15 ^b	0.29	(11)	16.08 ^c
Georgia	51.33 ^b	0.35	(19)	6.11 ^c	Oklahoma	60.04 ^b	0.55	(06)	24.11 ^c
Hawaii	34.84 ^b	0.68	(49)	-27.98 ^c	Oregon	48.11	0.48	(28)	-0.54
Idaho	40.16 ^b	0.73	(45)	-16.98 ^c	Pennsylvania	49.55 ^b	0.25	(26)	2.42
Illinois	50.94 ^b	0.28	(21)	5.31 ^c	Rhode Island	51.05	0.92	(20)	5.53 ^c
Indiana	59.00 ^b	0.42	(08)	21.96 ^c	South Carolina	54.21 ^b	0.47	(15)	12.06 ^c
Iowa	48.86	0.53	(27)	1.00	South Dakota	45.69	1.00	(34)	-5.54 ^c
Kansas	49.82	0.58	(25)	2.99	Tennessee	62.83 ^b	0.43	(04)	29.88 ^c
Kentucky	72.03 ^b	0.56	(01)	48.91 ^c	Texas	44.87 ^b	0.21	(37)	-7.24 ^c
Louisiana	58.40 ^b	0.51	(09)	20.74 ^c	Utah	20.87 ^b	0.46	(51)	-56.87 ^c
Maine	54.33 ^b	0.81	(14)	12.31 ^c	Vermont	52.24	1.21	(18)	7.99 ^c
Maryland	47.73	0.41	(30)	-1.33	Virginia	49.97 ^b	0.36	(23)	3.29
Massachusetts	48.04	0.37	(29)	-0.70	Washington	47.28	0.38	(32)	-2.26
Michigan	53.27 ^b	0.31	(16)	10.12 ^c	West Virginia	63.77 ^b	0.74	(03)	31.82 ^c
Minnesota	42.80 ^b	0.40	(41)	-11.52 ^c	Wisconsin	46.82 ^b	0.39	(33)	-3.21
Mississippi	62.63 ^b	0.64	(05)	29.48 ^c	Wyoming	41.11 ^b	1.22	(43)	-15.02 ^c
Missouri	57.69 ^b	0.42	(10)	19.27 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.25
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	61.58	0.09		
High Five States				
Kentucky	94.49 ^b	0.98	(01)	53.44 ^c
Mississippi	92.02 ^b	1.22	(02)	49.42 ^c
Arkansas	88.13 ^b	1.12	(03)	43.10 ^c
Tennessee	86.50 ^b	0.79	(04)	40.45 ^c
Alabama	85.52 ^b	0.89	(05)	38.87 ^c
Low Five States				
Hawaii	46.68 ^b	1.18	(47)	-24.19 ^c
California	45.54 ^b	0.25	(48)	-26.06 ^c
Colorado	42.78 ^b	0.68	(49)	-30.54 ^c
New Mexico	42.25 ^b	0.96	(50)	-31.39 ^c
Utah	26.51 ^b	0.77	(51)	-56.95 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	85.52 ^b	0.89	(05)	38.87 ^c	Montana	52.26 ^b	1.42	(41)	-15.14 ^c
Alaska	62.84	2.66	(24)	2.04	Nebraska	57.92	1.16	(33)	-5.95 ^c
Arizona	47.85 ^b	0.56	(46)	-22.30 ^c	Nevada	57.97	1.03	(32)	-5.87 ^c
Arkansas	88.13 ^b	1.12	(03)	43.10 ^c	New Hampshire	59.22	1.38	(28)	-3.84
California	45.54 ^b	0.25	(48)	-26.06 ^c	New Jersey	53.88 ^b	0.52	(38)	-12.51 ^c
Colorado	42.78 ^b	0.68	(49)	-30.54 ^c	New Mexico	42.25 ^b	0.96	(50)	-31.39 ^c
Connecticut	51.30 ^b	0.78	(43)	-16.71 ^c	New York	52.87 ^b	0.35	(39)	-14.15 ^c
Delaware	67.27	1.76	(16)	9.23 ^c	North Carolina	74.42 ^b	0.61	(11)	20.84 ^c
D.C.	57.07	2.26	(35)	-7.33 ^c	North Dakota	52.70 ^b	1.76	(40)	-14.43 ^c
Florida	59.25 ^b	0.33	(27)	-3.80	Ohio	72.79 ^b	0.51	(13)	18.20 ^c
Georgia	70.08 ^b	0.65	(14)	13.80 ^c	Oklahoma	78.45 ^b	0.95	(08)	27.39 ^c
Hawaii	46.68 ^b	1.18	(47)	-24.19 ^c	Oregon	56.38 ^b	0.78	(36)	-8.45 ^c
Idaho	48.00 ^b	1.19	(45)	-22.06 ^c	Pennsylvania	63.92 ^b	0.44	(22)	3.79
Illinois	64.18 ^b	0.49	(20)	4.21	Rhode Island	63.93	1.59	(21)	3.81
Indiana	77.29 ^b	0.74	(09)	25.50 ^c	South Carolina	74.87 ^b	0.86	(10)	21.58 ^c
Iowa	63.17	0.90	(23)	2.58	South Dakota	60.03	1.72	(26)	-2.53
Kansas	64.32	0.99	(19)	4.44	Tennessee	86.50 ^b	0.79	(04)	40.45 ^c
Kentucky	94.49 ^b	0.98	(01)	53.44 ^c	Texas	58.64 ^b	0.37	(30)	-4.77
Louisiana	79.07 ^b	0.92	(07)	28.40 ^c	Utah	26.51 ^b	0.77	(51)	-56.95 ^c
Maine	67.65 ^b	1.37	(15)	9.85 ^c	Vermont	61.31	1.98	(25)	-0.45
Maryland	58.88 ^b	0.70	(29)	-4.39	Virginia	64.73 ^b	0.63	(18)	5.11 ^c
Massachusetts	58.02 ^b	0.62	(31)	-5.79 ^c	Washington	55.58 ^b	0.62	(37)	-9.75 ^c
Michigan	66.53 ^b	0.53	(17)	8.03 ^c	West Virginia	82.31 ^b	1.28	(06)	33.65 ^c
Minnesota	51.85 ^b	0.66	(42)	-15.80 ^c	Wisconsin	57.85 ^b	0.65	(34)	-6.06 ^c
Mississippi	92.02 ^b	1.22	(02)	49.42 ^c	Wyoming	48.93 ^b	1.99	(44)	-20.55 ^c
Missouri	74.17 ^b	0.72	(12)	20.44 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.26
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	38.53	0.07		
High Five States				
Kentucky	55.53 ^b	0.65	(01)	44.13 ^c
West Virginia	49.82 ^b	0.88	(02)	29.32 ^c
Oklahoma	46.16 ^b	0.65	(03)	19.81 ^c
Delaware	45.89 ^b	1.28	(04)	19.12 ^c
Tennessee	45.65 ^b	0.49	(05)	18.49 ^c
Low Five States				
North Dakota	31.36 ^b	1.24	(47)	-18.59 ^c
Colorado	30.34 ^b	0.50	(48)	-21.25 ^c
New Mexico	27.90 ^b	0.69	(49)	-27.59 ^c
Hawaii	25.54 ^b	0.78	(50)	-33.70 ^c
Utah	16.24 ^b	0.55	(51)	-57.84 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	40.70 ^b	0.53	(23)	5.64 ^c	Montana	39.43	1.14	(26)	2.35
Alaska	45.33	2.02	(09)	17.65 ^c	Nebraska	35.49 ^b	0.81	(39)	-7.89 ^c
Arizona	32.24 ^b	0.42	(45)	-16.32 ^c	Nevada	45.23 ^b	0.83	(11)	17.40 ^c
Arkansas	45.27 ^b	0.71	(10)	17.50 ^c	New Hampshire	43.09 ^b	1.05	(16)	11.85 ^c
California	31.45 ^b	0.18	(46)	-18.37 ^c	New Jersey	36.55 ^b	0.37	(36)	-5.14 ^c
Colorado	30.34 ^b	0.50	(48)	-21.25 ^c	New Mexico	27.90 ^b	0.69	(49)	-27.59 ^c
Connecticut	37.22	0.58	(35)	-3.40	New York	35.77 ^b	0.25	(38)	-7.16 ^c
Delaware	45.89 ^b	1.28	(04)	19.12 ^c	North Carolina	39.94 ^b	0.38	(24)	3.66
D.C.	35.42	1.48	(40)	-8.07 ^c	North Dakota	31.36 ^b	1.24	(47)	-18.59 ^c
Florida	38.12	0.24	(33)	-1.06	Ohio	43.94 ^b	0.35	(13)	14.04 ^c
Georgia	38.06	0.40	(34)	-1.22	Oklahoma	46.16 ^b	0.65	(03)	19.81 ^c
Hawaii	25.54 ^b	0.78	(50)	-33.70 ^c	Oregon	41.76 ^b	0.61	(18)	8.39 ^c
Idaho	33.87 ^b	0.91	(44)	-12.08 ^c	Pennsylvania	39.25	0.30	(27)	1.88
Illinois	41.56 ^b	0.34	(19)	7.87 ^c	Rhode Island	42.58 ^b	1.13	(17)	10.52 ^c
Indiana	45.54 ^b	0.49	(06)	18.21 ^c	South Carolina	39.02	0.53	(29)	1.28
Iowa	38.13	0.63	(32)	-1.04	South Dakota	34.98	1.19	(41)	-9.21 ^c
Kansas	38.76	0.69	(30)	0.60	Tennessee	45.65 ^b	0.49	(05)	18.49 ^c
Kentucky	55.53 ^b	0.65	(01)	44.13 ^c	Texas	34.48 ^b	0.24	(43)	-10.50 ^c
Louisiana	43.15 ^b	0.58	(15)	11.99 ^c	Utah	16.24 ^b	0.55	(51)	-57.84 ^c
Maine	44.46 ^b	0.99	(12)	15.40 ^c	Vermont	45.37 ^b	1.52	(08)	17.75 ^c
Maryland	39.66	0.50	(25)	2.95	Virginia	39.15	0.42	(28)	1.61
Massachusetts	41.16 ^b	0.45	(21)	6.83 ^c	Washington	41.04 ^b	0.48	(22)	6.52 ^c
Michigan	43.55 ^b	0.38	(14)	13.03 ^c	West Virginia	49.82 ^b	0.88	(02)	29.32 ^c
Minnesota	36.14 ^b	0.49	(37)	-6.20 ^c	Wisconsin	38.62	0.48	(31)	0.23
Mississippi	41.34 ^b	0.69	(20)	7.31 ^c	Wyoming	34.61	1.53	(42)	-10.16 ^c
Missouri	45.53 ^b	0.50	(07)	18.17 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 16.18
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.72	0.01		
High Five States				
Utah	3.51 ^b	0.18	(01)	28.91 ^c
Colorado	3.49 ^b	0.12	(02)	27.98 ^c
Idaho	3.39	0.21	(03)	24.44 ^c
Kentucky	3.37 ^b	0.12	(04)	23.75 ^c
Oklahoma	3.31 ^b	0.13	(05)	21.68 ^c
Low Five States				
Louisiana	2.17 ^b	0.10	(47)	-20.45 ^c
North Dakota	2.16	0.24	(48)	-20.70 ^c
Alaska	2.11	0.30	(49)	-22.37 ^c
Hawaii	1.39 ^b	0.13	(50)	-49.10 ^c
D.C.	1.11 ^b	0.20	(51)	-59.20 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.87	0.11	(26)	5.30	Montana	2.85	0.23	(27)	4.67
Alaska	2.11	0.30	(49)	-22.37 ^c	Nebraska	2.87	0.17	(25)	5.52
Arizona	2.92	0.09	(23)	7.26	Nevada	3.06	0.16	(11)	12.41
Arkansas	2.80	0.13	(28)	2.88	New Hampshire	2.79	0.20	(29)	2.54
California	2.56 ^b	0.04	(36)	-6.04	New Jersey	2.55	0.07	(37)	-6.38
Colorado	3.49 ^b	0.12	(02)	27.98 ^c	New Mexico	2.70	0.16	(32)	-0.86
Connecticut	2.52	0.11	(38)	-7.60	New York	2.19 ^b	0.05	(46)	-19.63 ^c
Delaware	2.94	0.24	(20)	8.00	North Carolina	2.94	0.08	(19)	8.04
D.C.	1.11 ^b	0.20	(51)	-59.20 ^c	North Dakota	2.16	0.24	(48)	-20.70 ^c
Florida	2.94 ^b	0.05	(18)	8.06	Ohio	2.88	0.07	(24)	5.64
Georgia	2.45 ^b	0.08	(40)	-9.89	Oklahoma	3.31 ^b	0.13	(05)	21.68 ^c
Hawaii	1.39 ^b	0.13	(50)	-49.10 ^c	Oregon	3.23 ^b	0.12	(07)	18.69 ^c
Idaho	3.39	0.21	(03)	24.44 ^c	Pennsylvania	2.93	0.06	(22)	7.46
Illinois	2.45 ^b	0.06	(41)	-10.21	Rhode Island	2.29	0.19	(45)	-15.88 ^c
Indiana	3.11 ^b	0.10	(09)	14.31	South Carolina	2.69	0.11	(33)	-1.05
Iowa	2.79	0.13	(30)	2.33	South Dakota	2.65	0.24	(34)	-2.57
Kansas	3.05	0.14	(12)	12.07	Tennessee	3.02	0.10	(15)	10.95
Kentucky	3.37 ^b	0.12	(04)	23.75 ^c	Texas	2.50 ^b	0.05	(39)	-8.05
Louisiana	2.17 ^b	0.10	(47)	-20.45 ^c	Utah	3.51 ^b	0.18	(01)	28.91 ^c
Maine	2.95	0.19	(17)	8.26	Vermont	3.07	0.30	(10)	12.68
Maryland	2.61	0.10	(35)	-4.19	Virginia	3.00	0.09	(16)	10.29
Massachusetts	2.93	0.09	(21)	7.66	Washington	3.02	0.09	(14)	10.96
Michigan	2.44 ^b	0.07	(42)	-10.42	West Virginia	3.22	0.17	(08)	18.33 ^c
Minnesota	2.42	0.09	(43)	-11.15	Wisconsin	2.72	0.09	(31)	-0.31
Mississippi	2.34	0.13	(44)	-13.93	Wyoming	3.31	0.35	(06)	21.65 ^c
Missouri	3.03	0.10	(13)	11.13					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.19
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.08	0.02		
High Five States				
Utah	5.46 ^b	0.34	(01)	33.81 ^c
Colorado	5.03 ^b	0.23	(02)	23.37 ^c
Idaho	4.96	0.38	(03)	21.69 ^c
Oklahoma	4.91	0.24	(04)	20.27 ^c
Kentucky	4.88 ^b	0.22	(05)	19.62 ^c
Low Five States				
South Dakota	3.32	0.40	(46)	-18.62 ^c
New York	3.24 ^b	0.09	(47)	-20.67 ^c
North Dakota	3.13	0.43	(48)	-23.26 ^c
Alaska	2.79	0.54	(49)	-31.61 ^c
Hawaii	2.08 ^b	0.24	(50)	-48.92 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.51	0.21	(15)	10.60	Montana	3.61	0.37	(41)	-11.47
Alaska	2.79	0.54	(49)	-31.61 ^c	Nebraska	3.95	0.30	(32)	-3.18
Arizona	4.38	0.17	(21)	7.50	Nevada	4.76	0.29	(09)	16.59 ^c
Arkansas	4.40	0.25	(20)	7.87	New Hampshire	3.96	0.36	(31)	-2.85
California	3.85	0.07	(35)	-5.63	New Jersey	3.74	0.14	(37)	-8.32
Colorado	5.03 ^b	0.23	(02)	23.37 ^c	New Mexico	3.89	0.29	(33)	-4.55
Connecticut	3.49	0.20	(43)	-14.48	New York	3.24 ^b	0.09	(47)	-20.67 ^c
Delaware	4.42	0.46	(19)	8.27	North Carolina	4.52	0.15	(12)	10.94
D.C.	-	-	(-)	-	North Dakota	3.13	0.43	(48)	-23.26 ^c
Florida	4.43 ^b	0.09	(18)	8.66	Ohio	4.30	0.13	(26)	5.34
Georgia	3.89	0.16	(34)	-4.70	Oklahoma	4.91	0.24	(04)	20.27 ^c
Hawaii	2.08 ^b	0.24	(50)	-48.92 ^c	Oregon	4.34	0.21	(23)	6.49
Idaho	4.96	0.38	(03)	21.69 ^c	Pennsylvania	4.33	0.12	(24)	6.17
Illinois	3.66 ^b	0.12	(39)	-10.25	Rhode Island	3.48	0.36	(44)	-14.79
Indiana	4.68	0.18	(10)	14.66	South Carolina	4.30	0.21	(25)	5.55
Iowa	4.15	0.23	(28)	1.69	South Dakota	3.32	0.40	(46)	-18.62 ^c
Kansas	4.52	0.26	(13)	10.74	Tennessee	4.77 ^b	0.19	(08)	16.97 ^c
Kentucky	4.88 ^b	0.22	(05)	19.62 ^c	Texas	3.82	0.09	(36)	-6.45
Louisiana	3.39 ^b	0.19	(45)	-16.87 ^c	Utah	5.46 ^b	0.34	(01)	33.81 ^c
Maine	4.45	0.36	(17)	9.01	Vermont	4.56	0.56	(11)	11.76
Maryland	4.08	0.19	(29)	0.06	Virginia	4.50	0.17	(16)	10.30
Massachusetts	4.26	0.17	(27)	4.37	Washington	4.37	0.17	(22)	7.14
Michigan	3.63 ^b	0.12	(40)	-11.09	West Virginia	4.81	0.32	(06)	17.91 ^c
Minnesota	3.56	0.17	(42)	-12.76	Wisconsin	4.07	0.17	(30)	-0.18
Mississippi	3.73	0.25	(38)	-8.57	Wyoming	4.79	0.62	(07)	17.43 ^c
Missouri	4.51	0.18	(14)	10.64					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.20
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.70	0.01		
High Five States				
Oregon	2.36 ^b	0.15	(01)	38.80 ^c
Colorado	2.29 ^b	0.14	(02)	34.46 ^c
Kentucky	2.23 ^b	0.14	(03)	31.12 ^c
Montana	2.19	0.28	(04)	28.62 ^c
South Dakota	2.09	0.30	(05)	22.85 ^c
Low Five States				
Mississippi	1.42	0.13	(47)	-16.27 ^c
North Dakota	1.31	0.25	(48)	-22.84 ^c
Louisiana	1.28 ^b	0.10	(49)	-24.62 ^c
D.C.	1.14	0.27	(50)	-32.79 ^c
Hawaii	0.78 ^b	0.13	(51)	-54.07 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.68	0.11	(31)	-1.48	Montana	2.19	0.28	(04)	28.62 ^c
Alaska	1.58	0.35	(37)	-7.14	Nebraska	2.02	0.20	(09)	18.89 ^c
Arizona	1.71	0.10	(30)	0.74	Nevada	1.62	0.16	(34)	-4.50
Arkansas	1.57	0.14	(38)	-7.85	New Hampshire	1.89	0.22	(17)	11.16
California	1.55 ^b	0.04	(39)	-8.71	New Jersey	1.67	0.08	(32)	-1.75
Colorado	2.29 ^b	0.14	(02)	34.46 ^c	New Mexico	1.75	0.18	(29)	2.64
Connecticut	1.85	0.13	(25)	8.54	New York	1.44 ^b	0.05	(46)	-15.60 ^c
Delaware	1.88	0.27	(19)	10.37	North Carolina	1.85	0.08	(22)	8.85
D.C.	1.14	0.27	(50)	-32.79 ^c	North Dakota	1.31	0.25	(48)	-22.84 ^c
Florida	1.76	0.05	(27)	3.36	Ohio	1.85	0.07	(24)	8.54
Georgia	1.46	0.08	(45)	-14.10	Oklahoma	2.04	0.14	(07)	20.10 ^c
Hawaii	0.78 ^b	0.13	(51)	-54.07 ^c	Oregon	2.36 ^b	0.15	(01)	38.80 ^c
Idaho	2.05	0.23	(06)	20.38 ^c	Pennsylvania	1.87	0.07	(21)	10.13
Illinois	1.59	0.07	(36)	-6.65	Rhode Island	1.52	0.21	(43)	-10.62
Indiana	1.95	0.10	(13)	14.88	South Carolina	1.53	0.11	(41)	-10.07
Iowa	1.75	0.14	(28)	2.96	South Dakota	2.09	0.30	(05)	22.85 ^c
Kansas	1.92	0.15	(15)	12.91	Tennessee	1.76	0.10	(26)	3.45
Kentucky	2.23 ^b	0.14	(03)	31.12 ^c	Texas	1.50 ^b	0.05	(44)	-12.07
Louisiana	1.28 ^b	0.10	(49)	-24.62 ^c	Utah	1.89	0.18	(18)	11.02
Maine	1.85	0.21	(23)	8.84	Vermont	1.99	0.32	(11)	16.94 ^c
Maryland	1.60	0.10	(35)	-5.86	Virginia	1.87	0.09	(20)	10.26
Massachusetts	1.95	0.10	(14)	14.84	Washington	1.96	0.10	(12)	15.13 ^c
Michigan	1.53	0.07	(42)	-10.16	West Virginia	2.02	0.19	(10)	18.77 ^c
Minnesota	1.54	0.10	(40)	-9.50	Wisconsin	1.65	0.10	(33)	-2.70
Mississippi	1.42	0.13	(47)	-16.27 ^c	Wyoming	2.04	0.37	(08)	19.81 ^c
Missouri	1.92	0.11	(16)	12.86					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.18
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.36	0.01		
High Five States				
D.C.	5.45 ^b	0.45	(01)	62.30 ^c
Alabama	3.97 ^b	0.12	(02)	18.29 ^c
Louisiana	3.83 ^b	0.13	(03)	14.08
North Carolina	3.83 ^b	0.09	(04)	13.94
Maryland	3.80 ^b	0.12	(05)	13.18
Low Five States				
Rhode Island	2.86	0.22	(47)	-14.82
Connecticut	2.86 ^b	0.12	(48)	-14.83
New Mexico	2.84	0.16	(49)	-15.46 ^c
Nevada	2.77 ^b	0.15	(50)	-17.68 ^c
Hawaii	2.46 ^b	0.18	(51)	-26.71 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.97 ^b	0.12	(02)	18.29 ^c	Montana	3.26	0.24	(35)	-2.89
Alaska	2.91	0.39	(46)	-13.26	Nebraska	3.32	0.18	(32)	-1.30
Arizona	2.96 ^b	0.09	(45)	-11.90	Nevada	2.77 ^b	0.15	(50)	-17.68 ^c
Arkansas	3.50	0.15	(22)	4.30	New Hampshire	3.34	0.22	(28)	-0.57
California	3.14 ^b	0.04	(39)	-6.58	New Jersey	3.22	0.08	(38)	-4.05
Colorado	3.09	0.12	(41)	-8.12	New Mexico	2.84	0.16	(49)	-15.46 ^c
Connecticut	2.86 ^b	0.12	(48)	-14.83	New York	3.11 ^b	0.05	(40)	-7.28
Delaware	3.71	0.27	(09)	10.39	North Carolina	3.83 ^b	0.09	(04)	13.94
D.C.	5.45 ^b	0.45	(01)	62.30 ^c	North Dakota	3.58	0.30	(13)	6.51
Florida	2.98 ^b	0.05	(44)	-11.40	Ohio	3.55	0.07	(16)	5.79
Georgia	3.57	0.09	(15)	6.25	Oklahoma	3.53	0.13	(19)	5.20
Hawaii	2.46 ^b	0.18	(51)	-26.71 ^c	Oregon	3.55	0.13	(18)	5.59
Idaho	3.27	0.21	(33)	-2.54	Pennsylvania	3.36	0.07	(27)	0.01
Illinois	3.41	0.07	(26)	1.55	Rhode Island	2.86	0.22	(47)	-14.82
Indiana	3.53	0.10	(20)	5.19	South Carolina	3.75	0.13	(08)	11.78
Iowa	3.32	0.14	(30)	-1.18	South Dakota	3.33	0.27	(29)	-0.95
Kansas	3.66	0.16	(11)	8.91	Tennessee	3.78 ^b	0.11	(07)	12.49
Kentucky	3.42	0.12	(24)	1.79	Texas	3.23	0.06	(37)	-3.82
Louisiana	3.83 ^b	0.13	(03)	14.08	Utah	3.42	0.19	(25)	1.79
Maine	3.06	0.19	(42)	-9.02	Vermont	2.98	0.29	(43)	-11.35
Maryland	3.80 ^b	0.12	(05)	13.18	Virginia	3.80 ^b	0.10	(06)	13.07
Massachusetts	3.25	0.09	(36)	-3.28	Washington	3.32	0.10	(31)	-1.29
Michigan	3.70 ^b	0.08	(10)	10.13	West Virginia	3.51	0.17	(21)	4.58
Minnesota	3.49	0.11	(23)	4.05	Wisconsin	3.64	0.11	(12)	8.32
Mississippi	3.55	0.15	(17)	5.59	Wyoming	3.27	0.35	(34)	-2.74
Missouri	3.58	0.10	(14)	6.48					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.19
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.27	0.03		
High Five States				
D.C.	6.86	0.79	(01)	60.59 ^c
Tennessee	5.10 ^b	0.20	(02)	19.37 ^c
Louisiana	5.02	0.24	(03)	17.36 ^c
Alabama	5.00	0.22	(04)	17.00 ^c
Oregon	4.84	0.23	(05)	13.33
Low Five States				
New Mexico	3.61	0.28	(47)	-15.51 ^c
Nevada	3.44	0.26	(48)	-19.49 ^c
Rhode Island	3.33	0.36	(49)	-22.19 ^c
Hawaii	3.25	0.31	(50)	-23.93 ^c
Vermont	3.14	0.45	(51)	-26.51 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.00	0.22	(04)	17.00 ^c	Montana	3.94	0.39	(41)	-7.90
Alaska	4.59	0.76	(15)	7.44	Nebraska	4.14	0.31	(33)	-3.24
Arizona	3.81	0.16	(43)	-10.92	Nevada	3.44	0.26	(48)	-19.49 ^c
Arkansas	4.61	0.26	(13)	7.94	New Hampshire	4.50	0.39	(20)	5.34
California	4.05	0.07	(38)	-5.23	New Jersey	4.10	0.14	(36)	-4.15
Colorado	4.06	0.21	(37)	-5.11	New Mexico	3.61	0.28	(47)	-15.51 ^c
Connecticut	3.78	0.21	(44)	-11.63	New York	3.86 ^b	0.09	(42)	-9.69
Delaware	4.57	0.46	(16)	6.96	North Carolina	4.83 ^b	0.16	(06)	13.10
D.C.	6.86	0.79	(01)	60.59 ^c	North Dakota	4.43	0.52	(25)	3.55
Florida	3.72 ^b	0.08	(46)	-12.86	Ohio	4.48	0.13	(22)	4.89
Georgia	4.46	0.17	(23)	4.35	Oklahoma	4.52	0.23	(18)	5.77
Hawaii	3.25	0.31	(50)	-23.93 ^c	Oregon	4.84	0.23	(05)	13.33
Idaho	4.11	0.35	(35)	-3.91	Pennsylvania	4.21	0.11	(31)	-1.48
Illinois	4.40	0.13	(26)	2.91	Rhode Island	3.33	0.36	(49)	-22.19 ^c
Indiana	4.71	0.19	(11)	10.27	South Carolina	4.73	0.22	(09)	10.63
Iowa	4.35	0.24	(28)	1.67	South Dakota	3.94	0.44	(40)	-7.71
Kansas	4.72	0.27	(10)	10.37	Tennessee	5.10 ^b	0.20	(02)	19.37 ^c
Kentucky	4.46	0.22	(24)	4.25	Texas	4.13	0.10	(34)	-3.33
Louisiana	5.02	0.24	(03)	17.36 ^c	Utah	4.60	0.33	(14)	7.63
Maine	3.73	0.33	(45)	-12.66	Vermont	3.14	0.45	(51)	-26.51 ^c
Maryland	4.57	0.20	(17)	6.91	Virginia	4.77	0.18	(07)	11.56
Massachusetts	4.24	0.17	(30)	-0.69	Washington	4.31	0.18	(29)	0.77
Michigan	4.68	0.14	(12)	9.44	West Virginia	4.39	0.30	(27)	2.80
Minnesota	4.51	0.20	(19)	5.62	Wisconsin	4.73	0.19	(08)	10.71
Mississippi	4.19	0.27	(32)	-2.02	Wyoming	4.01	0.58	(39)	-6.23
Missouri	4.50	0.18	(21)	5.21					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.20
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.70	0.02		
High Five States				
D.C.	4.42	0.52	(01)	64.03 ^c
Maryland	3.28 ^b	0.14	(02)	21.58 ^c
Alabama	3.23 ^b	0.15	(03)	19.98 ^c
Mississippi	3.15	0.19	(04)	16.82 ^c
Virginia	3.15 ^b	0.12	(05)	16.80 ^c
Low Five States				
New Mexico	2.24	0.20	(47)	-17.04 ^c
Nevada	2.21	0.19	(48)	-18.00 ^c
Connecticut	2.21 ^b	0.14	(49)	-18.03 ^c
Hawaii	1.86 ^b	0.21	(50)	-30.84 ^c
Alaska	1.59	0.39	(51)	-41.17 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.23 ^b	0.15	(03)	19.98 ^c	Montana	2.72	0.30	(27)	0.92
Alaska	1.59	0.39	(51)	-41.17 ^c	Nebraska	2.73	0.22	(24)	1.36
Arizona	2.28 ^b	0.11	(46)	-15.26 ^c	Nevada	2.21	0.19	(48)	-18.00 ^c
Arkansas	2.70	0.17	(29)	0.08	New Hampshire	2.53	0.25	(40)	-6.13
California	2.47 ^b	0.05	(42)	-8.42	New Jersey	2.61	0.10	(32)	-3.04
Colorado	2.35	0.14	(45)	-12.71	New Mexico	2.24	0.20	(47)	-17.04 ^c
Connecticut	2.21 ^b	0.14	(49)	-18.03 ^c	New York	2.60	0.07	(33)	-3.55
Delaware	3.07	0.33	(08)	14.05	North Carolina	3.15 ^b	0.11	(06)	16.78 ^c
D.C.	4.42	0.52	(01)	64.03 ^c	North Dakota	2.95	0.37	(12)	9.53
Florida	2.39 ^b	0.06	(44)	-11.28	Ohio	2.90	0.09	(14)	7.65
Georgia	2.95	0.11	(11)	9.57	Oklahoma	2.77	0.16	(22)	2.87
Hawaii	1.86 ^b	0.21	(50)	-30.84 ^c	Oregon	2.53	0.15	(39)	-6.04
Idaho	2.61	0.25	(31)	-3.03	Pennsylvania	2.79	0.08	(21)	3.37
Illinois	2.72	0.09	(25)	1.01	Rhode Island	2.60	0.27	(34)	-3.64
Indiana	2.75	0.12	(23)	2.18	South Carolina	3.09	0.15	(07)	14.50
Iowa	2.50	0.15	(41)	-7.38	South Dakota	2.83	0.32	(18)	4.79
Kansas	2.91	0.19	(13)	7.91	Tennessee	2.82	0.12	(19)	4.79
Kentucky	2.72	0.14	(26)	0.97	Texas	2.56	0.07	(35)	-4.93
Louisiana	3.01	0.15	(09)	11.66	Utah	2.46	0.21	(43)	-8.70
Maine	2.56	0.23	(36)	-5.13	Vermont	2.83	0.37	(17)	4.87
Maryland	3.28 ^b	0.14	(02)	21.58 ^c	Virginia	3.15 ^b	0.12	(05)	16.80 ^c
Massachusetts	2.55	0.11	(37)	-5.26	Washington	2.54	0.12	(38)	-5.90
Michigan	2.96	0.10	(10)	9.93	West Virginia	2.82	0.20	(20)	4.64
Minnesota	2.71	0.13	(28)	0.57	Wisconsin	2.86	0.13	(16)	6.14
Mississippi	3.15	0.19	(04)	16.82 ^c	Wyoming	2.69	0.43	(30)	-0.08
Missouri	2.87	0.12	(15)	6.48					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.22
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	6.29	0.02		
High Five States				
West Virginia	7.17 ^b	0.25	(01)	14.07
Ohio	7.08 ^b	0.10	(02)	12.66
Minnesota	7.04 ^b	0.16	(03)	11.98
Kentucky	6.99 ^b	0.18	(04)	11.12
Michigan	6.98 ^b	0.11	(05)	11.05
Low Five States				
Hawaii	5.62	0.27	(47)	-10.56
Nevada	5.58	0.22	(48)	-11.19
North Dakota	5.54	0.38	(49)	-11.89
New Mexico	5.19 ^b	0.22	(50)	-17.38 ^c
D.C.	5.03	0.42	(51)	-20.01 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.43	0.16	(22)	2.31	Montana	6.00	0.32	(31)	-4.62
Alaska	6.30	0.57	(23)	0.18	Nebraska	6.74	0.26	(15)	7.16
Arizona	5.93	0.13	(37)	-5.67	Nevada	5.58	0.22	(48)	-11.19
Arkansas	6.78	0.21	(13)	7.80	New Hampshire	5.78	0.28	(42)	-8.00
California	6.03 ^b	0.06	(28)	-4.15	New Jersey	6.00	0.11	(32)	-4.64
Colorado	5.68 ^b	0.16	(45)	-9.69	New Mexico	5.19 ^b	0.22	(50)	-17.38 ^c
Connecticut	5.85	0.17	(40)	-6.89	New York	5.99 ^b	0.08	(33)	-4.78
Delaware	6.00	0.35	(30)	-4.52	North Carolina	5.95	0.11	(36)	-5.40
D.C.	5.03	0.42	(51)	-20.01 ^c	North Dakota	5.54	0.38	(49)	-11.89
Florida	5.95 ^b	0.07	(35)	-5.30	Ohio	7.08 ^b	0.10	(02)	12.66
Georgia	5.63 ^b	0.12	(46)	-10.54	Oklahoma	6.96 ^b	0.19	(06)	10.77
Hawaii	5.62	0.27	(47)	-10.56	Oregon	6.48	0.18	(20)	3.07
Idaho	6.52	0.30	(19)	3.70	Pennsylvania	6.86 ^b	0.09	(10)	9.15
Illinois	6.45	0.10	(21)	2.51	Rhode Island	5.70	0.31	(44)	-9.38
Indiana	6.95 ^b	0.14	(07)	10.51	South Carolina	5.87	0.16	(38)	-6.71
Iowa	6.81	0.19	(12)	8.23	South Dakota	6.24	0.36	(25)	-0.71
Kansas	6.93	0.21	(08)	10.29	Tennessee	6.76	0.14	(14)	7.51
Kentucky	6.99 ^b	0.18	(04)	11.12	Texas	6.02 ^b	0.08	(29)	-4.21
Louisiana	6.57	0.17	(17)	4.56	Utah	5.81	0.24	(41)	-7.66
Maine	6.89	0.29	(09)	9.54	Vermont	6.28	0.42	(24)	-0.18
Maryland	5.75 ^b	0.14	(43)	-8.60	Virginia	6.16	0.13	(26)	-2.05
Massachusetts	5.86	0.13	(39)	-6.87	Washington	6.63	0.14	(16)	5.52
Michigan	6.98 ^b	0.11	(05)	11.05	West Virginia	7.17 ^b	0.25	(01)	14.07
Minnesota	7.04 ^b	0.16	(03)	11.98	Wisconsin	6.82 ^b	0.15	(11)	8.46
Mississippi	5.99	0.20	(34)	-4.80	Wyoming	6.11	0.48	(27)	-2.85
Missouri	6.55	0.14	(18)	4.10					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.23
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.07	0.03		
High Five States				
Minnesota	9.39 ^b	0.29	(01)	16.38 ^c
Ohio	9.19 ^b	0.19	(02)	13.92
Kansas	9.18	0.38	(03)	13.72
Maine	9.11	0.52	(04)	12.91
Indiana	9.07 ^b	0.26	(05)	12.36
Low Five States				
Wyoming	7.31	0.81	(47)	-9.35
Nevada	6.91	0.36	(48)	-14.35
North Dakota	6.70	0.63	(49)	-16.98 ^c
D.C.	6.49	0.76	(50)	-19.59 ^c
New Mexico	6.21 ^b	0.37	(51)	-23.06 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.21	0.29	(22)	1.79	Montana	7.78	0.55	(28)	-3.54
Alaska	8.18	0.98	(24)	1.31	Nebraska	8.30	0.44	(19)	2.91
Arizona	7.32	0.22	(46)	-9.33	Nevada	6.91	0.36	(48)	-14.35
Arkansas	8.34	0.35	(17)	3.41	New Hampshire	7.40	0.50	(43)	-8.30
California	7.73	0.10	(30)	-4.26	New Jersey	7.48	0.20	(39)	-7.27
Colorado	7.41	0.29	(42)	-8.15	New Mexico	6.21 ^b	0.37	(51)	-23.06 ^c
Connecticut	7.32	0.29	(45)	-9.31	New York	7.71	0.13	(31)	-4.45
Delaware	7.45	0.60	(40)	-7.70	North Carolina	7.48	0.20	(38)	-7.24
D.C.	6.49	0.76	(50)	-19.59 ^c	North Dakota	6.70	0.63	(49)	-16.98 ^c
Florida	7.57 ^b	0.12	(35)	-6.23	Ohio	9.19 ^b	0.19	(02)	13.92
Georgia	7.51	0.22	(36)	-6.95	Oklahoma	8.83	0.33	(10)	9.41
Hawaii	7.42	0.47	(41)	-8.01	Oregon	8.38	0.30	(16)	3.79
Idaho	7.84	0.49	(26)	-2.83	Pennsylvania	8.85 ^b	0.17	(08)	9.71
Illinois	8.30	0.18	(20)	2.84	Rhode Island	7.75	0.56	(29)	-4.00
Indiana	9.07 ^b	0.26	(05)	12.36	South Carolina	7.70	0.29	(32)	-4.63
Iowa	8.85	0.34	(09)	9.66	South Dakota	7.63	0.61	(34)	-5.39
Kansas	9.18	0.38	(03)	13.72	Tennessee	8.92	0.26	(06)	10.52
Kentucky	8.60	0.31	(13)	6.54	Texas	7.79	0.14	(27)	-3.45
Louisiana	8.64	0.31	(12)	7.12	Utah	7.32	0.41	(44)	-9.29
Maine	9.11	0.52	(04)	12.91	Vermont	8.34	0.75	(18)	3.30
Maryland	7.49	0.25	(37)	-7.23	Virginia	8.19	0.23	(23)	1.49
Massachusetts	7.68	0.23	(33)	-4.85	Washington	8.56	0.25	(14)	6.04
Michigan	8.89 ^b	0.20	(07)	10.19	West Virginia	8.43	0.42	(15)	4.48
Minnesota	9.39 ^b	0.29	(01)	16.38 ^c	Wisconsin	8.73	0.26	(11)	8.25
Mississippi	7.96	0.37	(25)	-1.35	Wyoming	7.31	0.81	(47)	-9.35
Missouri	8.27	0.25	(21)	2.52					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 19.24
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.95	0.02		
High Five States				
West Virginia	6.19 ^b	0.31	(01)	25.05 ^c
Kentucky	5.79 ^b	0.21	(02)	16.88 ^c
Arkansas	5.62	0.25	(03)	13.45
Oklahoma	5.56	0.22	(04)	12.28
Michigan	5.56 ^b	0.13	(05)	12.23
Low Five States				
Georgia	4.31 ^b	0.14	(47)	-13.00
New Mexico	4.30	0.27	(48)	-13.06
Hawaii	4.27	0.32	(49)	-13.81
Rhode Island	4.23	0.34	(50)	-14.59
D.C.	3.94	0.49	(51)	-20.48 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.15	0.19	(18)	4.06	Montana	4.51	0.38	(43)	-8.88
Alaska	4.70	0.67	(32)	-5.13	Nebraska	5.53	0.31	(06)	11.78
Arizona	4.80	0.16	(26)	-3.02	Nevada	4.42	0.27	(45)	-10.61
Arkansas	5.62	0.25	(03)	13.45	New Hampshire	4.56	0.34	(41)	-7.84
California	4.73	0.07	(31)	-4.50	New Jersey	4.97	0.13	(24)	0.31
Colorado	4.33	0.19	(46)	-12.58	New Mexico	4.30	0.27	(48)	-13.06
Connecticut	4.78	0.20	(29)	-3.50	New York	4.76	0.09	(30)	-3.76
Delaware	4.79	0.42	(28)	-3.23	North Carolina	4.80	0.13	(27)	-3.11
D.C.	3.94	0.49	(51)	-20.48 ^c	North Dakota	4.61	0.46	(37)	-6.79
Florida	4.63 ^b	0.08	(36)	-6.36	Ohio	5.53 ^b	0.12	(07)	11.71
Georgia	4.31 ^b	0.14	(47)	-13.00	Oklahoma	5.56	0.22	(04)	12.28
Hawaii	4.27	0.32	(49)	-13.81	Oregon	5.03	0.21	(23)	1.68
Idaho	5.44	0.37	(08)	9.83	Pennsylvania	5.39 ^b	0.11	(11)	8.89
Illinois	5.15	0.12	(19)	4.01	Rhode Island	4.23	0.34	(50)	-14.59
Indiana	5.41	0.17	(09)	9.19	South Carolina	4.59	0.18	(39)	-7.34
Iowa	5.30	0.23	(13)	6.99	South Dakota	5.14	0.44	(20)	3.75
Kansas	5.21	0.25	(17)	5.25	Tennessee	5.13	0.16	(21)	3.63
Kentucky	5.79 ^b	0.21	(02)	16.88 ^c	Texas	4.69	0.09	(33)	-5.30
Louisiana	4.96	0.20	(25)	0.26	Utah	4.57	0.29	(40)	-7.64
Maine	5.25	0.34	(15)	6.06	Vermont	4.66	0.48	(34)	-5.78
Maryland	4.46	0.17	(44)	-9.85	Virginia	4.66	0.14	(35)	-5.83
Massachusetts	4.60	0.15	(38)	-7.16	Washington	5.13	0.17	(22)	3.57
Michigan	5.56 ^b	0.13	(05)	12.23	West Virginia	6.19 ^b	0.31	(01)	25.05 ^c
Minnesota	5.24	0.18	(16)	5.87	Wisconsin	5.40	0.17	(10)	9.16
Mississippi	4.55	0.23	(42)	-8.07	Wyoming	5.32	0.60	(12)	7.56
Missouri	5.26	0.17	(14)	6.33					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 20.24
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.46	0.01		
High Five States				
Mississippi	3.44 ^b	0.15	(01)	40.19 ^c
D.C.	3.42	0.35	(02)	39.28 ^c
Louisiana	3.29 ^b	0.12	(03)	33.98 ^c
Arkansas	3.06 ^b	0.14	(04)	24.71 ^c
Tennessee	3.06 ^b	0.09	(05)	24.47 ^c
Low Five States				
Kansas	1.93 ^b	0.11	(47)	-21.26 ^c
Colorado	1.91 ^b	0.09	(48)	-22.18 ^c
South Dakota	1.86	0.20	(49)	-24.42 ^c
Alaska	1.73	0.26	(50)	-29.76 ^c
Utah	1.48 ^b	0.12	(51)	-39.69 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.82	0.10	(08)	14.64	Montana	2.01	0.19	(45)	-18.30 ^c
Alaska	1.73	0.26	(50)	-29.76 ^c	Nebraska	2.43	0.16	(28)	-1.08
Arizona	2.01 ^b	0.08	(44)	-18.04 ^c	Nevada	2.31	0.14	(33)	-6.05
Arkansas	3.06 ^b	0.14	(04)	24.71 ^c	New Hampshire	2.38	0.18	(30)	-3.23
California	2.47	0.04	(24)	0.59	New Jersey	2.07 ^b	0.07	(42)	-15.87 ^c
Colorado	1.91 ^b	0.09	(48)	-22.18 ^c	New Mexico	2.24	0.14	(36)	-8.92
Connecticut	2.05 ^b	0.10	(43)	-16.60 ^c	New York	2.17 ^b	0.04	(39)	-11.81
Delaware	2.59	0.23	(16)	5.48	North Carolina	2.55	0.07	(19)	3.92
D.C.	3.42	0.35	(02)	39.28 ^c	North Dakota	2.12	0.24	(40)	-13.67
Florida	2.73 ^b	0.05	(10)	10.94	Ohio	2.58	0.06	(17)	5.15
Georgia	2.56	0.08	(18)	4.25	Oklahoma	2.69	0.12	(12)	9.57
Hawaii	2.64	0.19	(13)	7.52	Oregon	2.30	0.10	(34)	-6.35
Idaho	2.59	0.18	(15)	5.51	Pennsylvania	2.22 ^b	0.05	(38)	-9.84
Illinois	2.39	0.06	(29)	-2.71	Rhode Island	2.09	0.19	(41)	-15.12 ^c
Indiana	2.32	0.08	(32)	-5.56	South Carolina	3.01 ^b	0.11	(06)	22.47 ^c
Iowa	2.23	0.11	(37)	-9.17	South Dakota	1.86	0.20	(49)	-24.42 ^c
Kansas	1.93 ^b	0.11	(47)	-21.26 ^c	Tennessee	3.06 ^b	0.09	(05)	24.47 ^c
Kentucky	2.70	0.11	(11)	10.01	Texas	2.52	0.05	(23)	2.59
Louisiana	3.29 ^b	0.12	(03)	33.98 ^c	Utah	1.48 ^b	0.12	(51)	-39.69 ^c
Maine	2.53	0.17	(20)	3.02	Vermont	2.45	0.25	(25)	-0.26
Maryland	2.44	0.09	(27)	-0.87	Virginia	2.28	0.08	(35)	-7.03
Massachusetts	2.52	0.08	(21)	2.76	Washington	2.97 ^b	0.09	(07)	20.78 ^c
Michigan	2.52	0.07	(22)	2.65	West Virginia	2.60	0.15	(14)	5.88
Minnesota	1.98 ^b	0.08	(46)	-19.57 ^c	Wisconsin	2.45	0.09	(26)	-0.34
Mississippi	3.44 ^b	0.15	(01)	40.19 ^c	Wyoming	2.77	0.31	(09)	12.61
Missouri	2.35	0.08	(31)	-4.18					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.25
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.77	0.02		
High Five States				
Mississippi	5.69 ^b	0.29	(01)	51.04 ^c
Louisiana	5.46 ^b	0.23	(02)	44.81 ^c
D.C.	5.44	0.67	(03)	44.27 ^c
South Carolina	5.03 ^b	0.22	(04)	33.41 ^c
Tennessee	4.86 ^b	0.18	(05)	28.98 ^c
Low Five States				
South Dakota	2.73	0.36	(47)	-27.64 ^c
Montana	2.72	0.32	(48)	-27.74 ^c
Colorado	2.67 ^b	0.16	(49)	-29.11 ^c
Alaska	2.61	0.48	(50)	-30.85 ^c
Utah	1.81 ^b	0.20	(51)	-51.87 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.52 ^b	0.20	(07)	19.89 ^c	Montana	2.72	0.32	(48)	-27.74 ^c
Alaska	2.61	0.48	(50)	-30.85 ^c	Nebraska	3.34	0.28	(40)	-11.28
Arizona	3.14 ^b	0.14	(43)	-16.66 ^c	Nevada	3.42	0.24	(38)	-9.15
Arkansas	4.81 ^b	0.26	(06)	27.72 ^c	New Hampshire	3.68	0.33	(26)	-2.47
California	3.67	0.07	(27)	-2.57	New Jersey	3.12 ^b	0.12	(44)	-17.29 ^c
Colorado	2.67 ^b	0.16	(49)	-29.11 ^c	New Mexico	3.28	0.26	(41)	-13.04
Connecticut	3.21	0.19	(42)	-14.88	New York	3.39 ^b	0.09	(39)	-10.14
Delaware	4.07	0.43	(15)	7.98	North Carolina	4.10	0.14	(13)	8.71
D.C.	5.44	0.67	(03)	44.27 ^c	North Dakota	3.70	0.46	(24)	-1.97
Florida	4.21 ^b	0.09	(09)	11.67	Ohio	3.98	0.12	(18)	5.48
Georgia	4.12	0.15	(12)	9.33	Oklahoma	4.04	0.21	(17)	7.11
Hawaii	4.09	0.34	(14)	8.42	Oregon	3.47	0.19	(34)	-8.06
Idaho	3.69	0.32	(25)	-2.11	Pennsylvania	3.44	0.10	(36)	-8.85
Illinois	3.59	0.11	(29)	-4.80	Rhode Island	3.47	0.36	(33)	-8.02
Indiana	3.60	0.15	(28)	-4.39	South Carolina	5.03 ^b	0.22	(04)	33.41 ^c
Iowa	3.43	0.21	(37)	-9.00	South Dakota	2.73	0.36	(47)	-27.64 ^c
Kansas	2.83 ^b	0.20	(45)	-24.80 ^c	Tennessee	4.86 ^b	0.18	(05)	28.98 ^c
Kentucky	4.28	0.21	(08)	13.54	Texas	3.92	0.09	(20)	4.05
Louisiana	5.46 ^b	0.23	(02)	44.81 ^c	Utah	1.81 ^b	0.20	(51)	-51.87 ^c
Maine	3.81	0.32	(22)	1.05	Vermont	3.44	0.46	(35)	-8.78
Maryland	3.70	0.17	(23)	-1.94	Virginia	3.58	0.15	(30)	-4.92
Massachusetts	3.94	0.16	(19)	4.63	Washington	4.13	0.16	(11)	9.52
Michigan	3.90	0.13	(21)	3.45	West Virginia	4.18	0.28	(10)	10.93
Minnesota	2.83 ^b	0.15	(46)	-24.93 ^c	Wisconsin	3.51	0.16	(31)	-6.90
Mississippi	5.69 ^b	0.29	(01)	51.04 ^c	Wyoming	4.05	0.56	(16)	7.37
Missouri	3.49	0.15	(32)	-7.30					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.26
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.37	0.01		
High Five States				
Washington	1.97 ^b	0.10	(01)	43.85 ^c
D.C.	1.86	0.34	(02)	36.37 ^c
Wyoming	1.67	0.33	(03)	22.00 ^c
Mississippi	1.65	0.14	(04)	21.03 ^c
Vermont	1.63	0.27	(05)	19.36 ^c
Low Five States				
Connecticut	1.12	0.10	(46)	-17.85 ^c
South Dakota	1.07	0.20	(47)	-21.70 ^c
Arizona	1.04 ^b	0.08	(48)	-23.64 ^c
Rhode Island	1.03	0.18	(49)	-24.52 ^c
North Dakota	0.83	0.21	(50)	-39.01 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.44	0.10	(16)	5.51	Montana	1.34	0.21	(28)	-2.10
Alaska	-	-	(-)	-	Nebraska	1.62	0.17	(07)	18.90 ^c
Arizona	1.04 ^b	0.08	(48)	-23.64 ^c	Nevada	1.31	0.14	(32)	-3.83
Arkansas	1.63	0.14	(06)	19.31 ^c	New Hampshire	1.26	0.18	(34)	-7.48
California	1.47	0.04	(13)	7.43	New Jersey	1.24	0.07	(40)	-9.13
Colorado	1.25	0.10	(37)	-8.46	New Mexico	1.31	0.15	(33)	-3.92
Connecticut	1.12	0.10	(46)	-17.85 ^c	New York	1.21 ^b	0.04	(42)	-11.63
Delaware	1.35	0.22	(27)	-1.18	North Carolina	1.32	0.07	(31)	-3.30
D.C.	1.86	0.34	(02)	36.37 ^c	North Dakota	0.83	0.21	(50)	-39.01 ^c
Florida	1.45	0.05	(14)	6.36	Ohio	1.42	0.06	(18)	3.83
Georgia	1.33	0.07	(29)	-2.44	Oklahoma	1.53	0.12	(11)	12.26
Hawaii	1.38	0.18	(21)	1.06	Oregon	1.32	0.11	(30)	-3.27
Idaho	1.55	0.20	(10)	13.22	Pennsylvania	1.25	0.05	(38)	-8.80
Illinois	1.41	0.06	(20)	2.87	Rhode Island	1.03	0.18	(49)	-24.52 ^c
Indiana	1.24	0.08	(39)	-9.11	South Carolina	1.36	0.10	(25)	-0.85
Iowa	1.23	0.11	(41)	-10.01	South Dakota	1.07	0.20	(47)	-21.70 ^c
Kansas	1.18	0.12	(44)	-13.89	Tennessee	1.56	0.09	(08)	14.13
Kentucky	1.42	0.10	(17)	4.02	Texas	1.35	0.05	(26)	-1.16
Louisiana	1.50	0.11	(12)	9.88	Utah	1.16	0.14	(45)	-14.79
Maine	1.44	0.17	(15)	5.51	Vermont	1.63	0.27	(05)	19.36 ^c
Maryland	1.41	0.09	(19)	3.32	Virginia	1.26	0.08	(35)	-7.68
Massachusetts	1.38	0.08	(22)	0.64	Washington	1.97 ^b	0.10	(01)	43.85 ^c
Michigan	1.37	0.07	(23)	-0.07	West Virginia	1.20	0.14	(43)	-11.89
Minnesota	1.26	0.09	(36)	-8.17	Wisconsin	1.55	0.10	(09)	13.30
Mississippi	1.65	0.14	(04)	21.03 ^c	Wyoming	1.67	0.33	(03)	22.00 ^c
Missouri	1.36	0.09	(24)	-0.66					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 21.14
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.88	0.03		
High Five States				
Washington	9.17 ^b	0.22	(01)	16.38 ^c
Oregon	9.10 ^b	0.28	(02)	15.54 ^c
Iowa	8.96 ^b	0.31	(03)	13.82
Alabama	8.94 ^b	0.25	(04)	13.52
Montana	8.85	0.53	(05)	12.34
Low Five States				
Missouri	7.07 ^b	0.20	(47)	-10.21
Nevada	6.86	0.32	(48)	-12.96
Alaska	6.85	0.75	(49)	-12.98
Rhode Island	6.75	0.44	(50)	-14.34
Hawaii	5.81 ^b	0.38	(51)	-26.25 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.94 ^b	0.25	(04)	13.52	Montana	8.85	0.53	(05)	12.34
Alaska	6.85	0.75	(49)	-12.98	Nebraska	7.11	0.36	(45)	-9.69
Arizona	7.62	0.21	(36)	-3.26	Nevada	6.86	0.32	(48)	-12.96
Arkansas	8.39	0.31	(11)	6.49	New Hampshire	7.67	0.44	(33)	-2.64
California	7.85	0.09	(23)	-0.29	New Jersey	8.53 ^b	0.18	(07)	8.24
Colorado	8.48	0.26	(09)	7.71	New Mexico	7.77	0.37	(28)	-1.35
Connecticut	7.73	0.26	(29)	-1.85	New York	7.83	0.11	(26)	-0.54
Delaware	8.18	0.54	(14)	3.84	North Carolina	7.46	0.16	(38)	-5.25
D.C.	7.36	0.68	(41)	-6.58	North Dakota	7.68	0.61	(32)	-2.55
Florida	7.17 ^b	0.11	(44)	-8.94	Ohio	8.01	0.15	(18)	1.69
Georgia	8.26	0.19	(12)	4.91	Oklahoma	7.85	0.27	(22)	-0.27
Hawaii	5.81 ^b	0.38	(51)	-26.25 ^c	Oregon	9.10 ^b	0.28	(02)	15.54 ^c
Idaho	7.51	0.43	(37)	-4.60	Pennsylvania	8.48 ^b	0.14	(08)	7.71
Illinois	7.93	0.15	(20)	0.74	Rhode Island	6.75	0.44	(50)	-14.34
Indiana	8.14	0.21	(15)	3.37	South Carolina	8.00	0.24	(19)	1.54
Iowa	8.96 ^b	0.31	(03)	13.82	South Dakota	7.42	0.54	(39)	-5.77
Kansas	8.12	0.32	(16)	3.05	Tennessee	7.62	0.20	(35)	-3.21
Kentucky	7.71	0.24	(30)	-2.12	Texas	7.38 ^b	0.11	(40)	-6.26
Louisiana	7.32	0.24	(43)	-7.07	Utah	7.33	0.37	(42)	-6.95
Maine	7.89	0.41	(21)	0.18	Vermont	7.64	0.63	(34)	-2.97
Maryland	7.70	0.22	(31)	-2.26	Virginia	8.21	0.19	(13)	4.21
Massachusetts	7.84	0.20	(25)	-0.46	Washington	9.17 ^b	0.22	(01)	16.38 ^c
Michigan	8.11	0.16	(17)	2.97	West Virginia	8.40	0.36	(10)	6.68
Minnesota	7.85	0.23	(24)	-0.37	Wisconsin	8.71 ^b	0.23	(06)	10.63
Mississippi	7.11	0.29	(46)	-9.78	Wyoming	7.78	0.72	(27)	-1.25
Missouri	7.07 ^b	0.20	(47)	-10.21					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.18
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	10.90	0.03		
High Five States				
D.C.	14.19 ^b	0.71	(01)	30.17 ^c
Louisiana	12.73 ^b	0.24	(02)	16.81 ^c
Mississippi	11.73	0.28	(03)	7.57
Michigan	11.72 ^b	0.15	(04)	7.52
New Jersey	11.65 ^b	0.16	(05)	6.90
Low Five States				
Arizona	9.86 ^b	0.17	(47)	-9.51
Montana	9.83	0.41	(48)	-9.83
West Virginia	9.65 ^b	0.29	(49)	-11.49
New Mexico	9.51 ^b	0.30	(50)	-12.79
Utah	9.05 ^b	0.30	(51)	-16.98 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	11.39	0.21	(12)	4.52	Montana	9.83	0.41	(48)	-9.83
Alaska	11.24	0.74	(19)	3.12	Nebraska	10.42	0.32	(38)	-4.47
Arizona	9.86 ^b	0.17	(47)	-9.51	Nevada	11.06	0.30	(25)	1.47
Arkansas	11.23	0.26	(21)	3.01	New Hampshire	11.50	0.40	(09)	5.47
California	10.42 ^b	0.08	(36)	-4.43	New Jersey	11.65 ^b	0.16	(05)	6.90
Colorado	9.89 ^b	0.21	(46)	-9.28	New Mexico	9.51 ^b	0.30	(50)	-12.79
Connecticut	11.63	0.24	(06)	6.70	New York	11.22	0.10	(22)	2.89
Delaware	11.31	0.47	(15)	3.72	North Carolina	10.50	0.15	(33)	-3.72
D.C.	14.19 ^b	0.71	(01)	30.17 ^c	North Dakota	10.27	0.51	(41)	-5.81
Florida	10.34 ^b	0.09	(40)	-5.17	Ohio	11.50 ^b	0.13	(08)	5.52
Georgia	10.47	0.16	(35)	-3.98	Oklahoma	10.63	0.23	(32)	-2.46
Hawaii	11.17	0.38	(23)	2.47	Oregon	10.94	0.23	(30)	0.31
Idaho	10.38	0.37	(39)	-4.74	Pennsylvania	11.54 ^b	0.12	(07)	5.89
Illinois	11.35	0.13	(13)	4.14	Rhode Island	10.26	0.41	(42)	-5.89
Indiana	10.96	0.18	(29)	0.54	South Carolina	11.34	0.22	(14)	4.06
Iowa	10.42	0.24	(37)	-4.43	South Dakota	10.00	0.46	(45)	-8.26
Kansas	10.77	0.27	(31)	-1.18	Tennessee	11.10	0.18	(24)	1.77
Kentucky	11.00	0.22	(26)	0.91	Texas	10.18 ^b	0.10	(43)	-6.63
Louisiana	12.73 ^b	0.24	(02)	16.81 ^c	Utah	9.05 ^b	0.30	(51)	-16.98 ^c
Maine	11.00	0.36	(27)	0.87	Vermont	11.25	0.55	(18)	3.22
Maryland	11.48	0.20	(10)	5.29	Virginia	10.99	0.17	(28)	0.76
Massachusetts	11.45	0.18	(11)	5.06	Washington	11.30	0.18	(16)	3.62
Michigan	11.72 ^b	0.15	(04)	7.52	West Virginia	9.65 ^b	0.29	(49)	-11.49
Minnesota	10.48	0.19	(34)	-3.84	Wisconsin	11.27	0.19	(17)	3.39
Mississippi	11.73	0.28	(03)	7.57	Wyoming	10.08	0.60	(44)	-7.51
Missouri	11.24	0.18	(20)	3.09					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.19
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	12.54	0.04		
High Five States				
D.C.	16.43	1.21	(01)	30.99 ^c
Louisiana	14.31 ^b	0.39	(02)	14.13 ^c
Mississippi	13.83	0.47	(03)	10.26 ^c
Connecticut	13.81	0.40	(04)	10.13 ^c
Michigan	13.78 ^b	0.24	(05)	9.88
Low Five States				
Nebraska	11.26	0.51	(47)	-10.17 ^c
New Mexico	11.02	0.48	(48)	-12.16 ^c
Colorado	10.77 ^b	0.33	(49)	-14.08 ^c
South Dakota	10.41	0.70	(50)	-16.95 ^c
Utah	10.31 ^b	0.48	(51)	-17.76 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	13.27	0.35	(11)	5.85	Montana	11.91	0.68	(40)	-5.01
Alaska	13.36	1.19	(08)	6.57	Nebraska	11.26	0.51	(47)	-10.17 ^c
Arizona	11.35 ^b	0.27	(46)	-9.52	Nevada	12.76	0.48	(27)	1.74
Arkansas	13.26	0.43	(12)	5.72	New Hampshire	13.54	0.66	(07)	7.97
California	11.71 ^b	0.13	(45)	-6.65	New Jersey	13.62 ^b	0.26	(06)	8.59
Colorado	10.77 ^b	0.33	(49)	-14.08 ^c	New Mexico	11.02	0.48	(48)	-12.16 ^c
Connecticut	13.81	0.40	(04)	10.13 ^c	New York	12.91	0.17	(21)	2.97
Delaware	12.89	0.77	(22)	2.76	North Carolina	11.78	0.24	(43)	-6.06
D.C.	16.43	1.21	(01)	30.99 ^c	North Dakota	12.78	0.86	(25)	1.89
Florida	12.01 ^b	0.15	(39)	-4.25	Ohio	13.31 ^b	0.22	(10)	6.14
Georgia	12.30	0.27	(32)	-1.94	Oklahoma	12.32	0.38	(31)	-1.74
Hawaii	12.84	0.61	(23)	2.37	Oregon	12.25	0.36	(35)	-2.28
Idaho	12.12	0.59	(36)	-3.36	Pennsylvania	13.32 ^b	0.20	(09)	6.26
Illinois	12.98	0.22	(17)	3.50	Rhode Island	12.26	0.69	(34)	-2.23
Indiana	12.82	0.30	(24)	2.23	South Carolina	13.11	0.36	(14)	4.56
Iowa	12.27	0.40	(33)	-2.13	South Dakota	10.41	0.70	(50)	-16.95 ^c
Kansas	12.52	0.44	(30)	-0.17	Tennessee	13.08	0.31	(15)	4.31
Kentucky	12.93	0.36	(18)	3.09	Texas	11.76 ^b	0.16	(44)	-6.24
Louisiana	14.31 ^b	0.39	(02)	14.13 ^c	Utah	10.31 ^b	0.48	(51)	-17.76 ^c
Maine	12.08	0.57	(37)	-3.66	Vermont	12.91	0.90	(20)	2.99
Maryland	13.22	0.33	(13)	5.40	Virginia	12.67	0.28	(29)	1.03
Massachusetts	12.72	0.29	(28)	1.47	Washington	12.77	0.30	(26)	1.87
Michigan	13.78 ^b	0.24	(05)	9.88	West Virginia	11.87	0.49	(42)	-5.31
Minnesota	12.01	0.32	(38)	-4.23	Wisconsin	13.03	0.31	(16)	3.88
Mississippi	13.83	0.47	(03)	10.26 ^c	Wyoming	11.89	0.97	(41)	-5.21
Missouri	12.92	0.30	(19)	3.01					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.20
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	9.56	0.03		
High Five States				
D.C.	12.42	0.87	(01)	29.93 ^c
Louisiana	11.39 ^b	0.30	(02)	19.23 ^c
Massachusetts	10.41 ^b	0.22	(03)	8.92
Maryland	10.13	0.25	(04)	6.05
Delaware	10.12	0.60	(05)	5.91
Low Five States				
New Mexico	8.26	0.38	(47)	-13.59
North Dakota	8.11	0.61	(48)	-15.10 ^c
Montana	8.03	0.51	(49)	-15.97 ^c
Utah	8.01 ^b	0.38	(50)	-16.17 ^c
West Virginia	7.83 ^b	0.35	(51)	-18.06 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.84	0.26	(18)	2.92	Montana	8.03	0.51	(49)	-15.97 ^c
Alaska	9.15	0.90	(37)	-4.25	Nebraska	9.62	0.42	(26)	0.62
Arizona	8.58 ^b	0.22	(45)	-10.19	Nevada	9.59	0.38	(28)	0.37
Arkansas	9.47	0.32	(30)	-0.89	New Hampshire	9.88	0.50	(17)	3.36
California	9.35	0.10	(33)	-2.15	New Jersey	10.10	0.19	(06)	5.72
Colorado	9.07	0.27	(39)	-5.07	New Mexico	8.26	0.38	(47)	-13.59
Connecticut	9.93	0.29	(14)	3.91	New York	9.90	0.13	(16)	3.59
Delaware	10.12	0.60	(05)	5.91	North Carolina	9.41	0.18	(32)	-1.53
D.C.	12.42	0.87	(01)	29.93 ^c	North Dakota	8.11	0.61	(48)	-15.10 ^c
Florida	8.93 ^b	0.11	(40)	-6.58	Ohio	10.06	0.17	(09)	5.26
Georgia	9.08	0.20	(38)	-4.94	Oklahoma	9.16	0.29	(36)	-4.18
Hawaii	9.69	0.48	(24)	1.40	Oregon	9.80	0.29	(21)	2.52
Idaho	8.84	0.47	(43)	-7.54	Pennsylvania	10.09 ^b	0.15	(07)	5.53
Illinois	10.03	0.16	(11)	4.90	Rhode Island	8.70	0.50	(44)	-8.92
Indiana	9.47	0.22	(31)	-0.92	South Carolina	9.95	0.27	(13)	4.13
Iowa	8.90	0.30	(41)	-6.92	South Dakota	9.71	0.62	(23)	1.58
Kansas	9.27	0.33	(34)	-3.05	Tennessee	9.60	0.22	(27)	0.41
Kentucky	9.48	0.27	(29)	-0.82	Texas	8.89 ^b	0.12	(42)	-6.99
Louisiana	11.39 ^b	0.30	(02)	19.23 ^c	Utah	8.01 ^b	0.38	(50)	-16.17 ^c
Maine	9.91	0.46	(15)	3.72	Vermont	9.72	0.69	(22)	1.72
Maryland	10.13	0.25	(04)	6.05	Virginia	9.62	0.21	(25)	0.69
Massachusetts	10.41 ^b	0.22	(03)	8.92	Washington	10.08	0.24	(08)	5.44
Michigan	10.04	0.18	(10)	5.09	West Virginia	7.83 ^b	0.35	(51)	-18.06 ^c
Minnesota	9.18	0.24	(35)	-3.97	Wisconsin	9.83	0.24	(19)	2.83
Mississippi	9.97	0.34	(12)	4.28	Wyoming	8.49	0.75	(46)	-11.16
Missouri	9.82	0.23	(20)	2.80					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 23.14
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	22.35	0.06		
High Five States				
D.C.	37.59 ^b	1.93	(01)	68.21 ^c
Mississippi	29.57 ^b	0.76	(02)	32.32 ^c
Alabama	28.17 ^b	0.56	(03)	26.07 ^c
South Carolina	26.29 ^b	0.56	(04)	17.66 ^c
Georgia	25.97 ^b	0.44	(05)	16.20 ^c
Low Five States				
Texas	20.43 ^b	0.23	(47)	-8.56
Kansas	20.43	0.57	(48)	-8.58
Arizona	20.07 ^b	0.38	(49)	-10.18 ^c
Florida	19.52 ^b	0.19	(50)	-12.66 ^c
Hawaii	15.56 ^b	0.68	(51)	-30.36 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	28.17 ^b	0.56	(03)	26.07 ^c	Montana	24.78	1.02	(08)	10.89 ^c
Alaska	22.62	1.80	(29)	1.23	Nebraska	22.63	0.74	(28)	1.26
Arizona	20.07 ^b	0.38	(49)	-10.18 ^c	Nevada	22.77	0.71	(25)	1.89
Arkansas	24.31	0.63	(12)	8.78	New Hampshire	21.39	0.88	(40)	-4.30
California	21.88	0.18	(35)	-2.09	New Jersey	21.20	0.34	(43)	-5.12
Colorado	22.83	0.53	(24)	2.17	New Mexico	22.95	0.75	(23)	2.71
Connecticut	21.46	0.51	(37)	-3.97	New York	21.44 ^b	0.23	(38)	-4.07
Delaware	23.56	1.11	(17)	5.43	North Carolina	24.63 ^b	0.38	(09)	10.22 ^c
D.C.	37.59 ^b	1.93	(01)	68.21 ^c	North Dakota	22.65	1.16	(27)	1.35
Florida	19.52 ^b	0.19	(50)	-12.66 ^c	Ohio	23.23	0.30	(21)	3.97
Georgia	25.97 ^b	0.44	(05)	16.20 ^c	Oklahoma	23.11	0.55	(22)	3.41
Hawaii	15.56 ^b	0.68	(51)	-30.36 ^c	Oregon	23.94	0.53	(14)	7.14
Idaho	25.78 ^b	0.92	(06)	15.35 ^c	Pennsylvania	22.05	0.26	(34)	-1.34
Illinois	23.31	0.31	(20)	4.32	Rhode Island	21.32	0.93	(41)	-4.62
Indiana	22.71	0.42	(26)	1.64	South Carolina	26.29 ^b	0.56	(04)	17.66 ^c
Iowa	21.66	0.54	(36)	-3.08	South Dakota	22.06	1.05	(32)	-1.27
Kansas	20.43	0.57	(48)	-8.58	Tennessee	24.00 ^b	0.45	(13)	7.39
Kentucky	22.33	0.53	(31)	-0.08	Texas	20.43 ^b	0.23	(47)	-8.56
Louisiana	25.06 ^b	0.56	(07)	12.13 ^c	Utah	24.62	0.79	(10)	10.16 ^c
Maine	22.05	0.83	(33)	-1.32	Vermont	22.42	1.26	(30)	0.33
Maryland	23.74	0.47	(16)	6.23	Virginia	23.91 ^b	0.41	(15)	6.98
Massachusetts	21.40	0.38	(39)	-4.25	Washington	23.42	0.42	(19)	4.82
Michigan	21.11 ^b	0.32	(44)	-5.53	West Virginia	20.74	0.69	(45)	-7.21
Minnesota	23.44	0.46	(18)	4.89	Wisconsin	24.32 ^b	0.44	(11)	8.83
Mississippi	29.57 ^b	0.76	(02)	32.32 ^c	Wyoming	21.29	1.40	(42)	-4.72
Missouri	20.69 ^b	0.40	(46)	-7.43					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.18
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.46	0.01		
High Five States				
Hawaii	5.61 ^b	0.27	(01)	62.51 ^c
Louisiana	4.72 ^b	0.15	(02)	36.52 ^c
D.C.	4.60	0.41	(03)	33.26 ^c
California	4.35 ^b	0.05	(04)	25.95 ^c
New Mexico	4.19 ^b	0.20	(05)	21.23 ^c
Low Five States				
Vermont	2.50 ^b	0.26	(47)	-27.53 ^c
Iowa	2.47 ^b	0.12	(48)	-28.65 ^c
Montana	2.41 ^b	0.21	(49)	-30.23 ^c
Wyoming	2.13 ^b	0.28	(50)	-38.30 ^c
Utah	2.10 ^b	0.15	(51)	-39.11 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.57	0.12	(17)	3.22	Montana	2.41 ^b	0.21	(49)	-30.23 ^c
Alaska	3.93	0.43	(11)	13.62	Nebraska	2.54 ^b	0.16	(45)	-26.63 ^c
Arizona	2.96 ^b	0.10	(34)	-14.30	Nevada	3.08	0.16	(29)	-10.85
Arkansas	3.12	0.14	(26)	-9.81	New Hampshire	2.60 ^b	0.19	(43)	-24.87 ^c
California	4.35 ^b	0.05	(04)	25.95 ^c	New Jersey	3.95 ^b	0.09	(10)	14.34
Colorado	2.74 ^b	0.11	(39)	-20.73 ^c	New Mexico	4.19 ^b	0.20	(05)	21.23 ^c
Connecticut	3.74	0.14	(13)	8.39	New York	3.99 ^b	0.06	(08)	15.59 ^c
Delaware	4.09	0.29	(06)	18.37 ^c	North Carolina	3.24	0.08	(21)	-6.20
D.C.	4.60	0.41	(03)	33.26 ^c	North Dakota	3.00	0.27	(33)	-13.31
Florida	3.06 ^b	0.05	(31)	-11.52	Ohio	3.09 ^b	0.07	(28)	-10.68
Georgia	3.67	0.09	(15)	6.23	Oklahoma	3.12	0.13	(25)	-9.58
Hawaii	5.61 ^b	0.27	(01)	62.51 ^c	Oregon	2.55 ^b	0.11	(44)	-26.05 ^c
Idaho	2.68 ^b	0.19	(41)	-22.38 ^c	Pennsylvania	3.21 ^b	0.06	(24)	-7.13
Illinois	3.75 ^b	0.08	(12)	8.67	Rhode Island	3.43	0.23	(18)	-0.69
Indiana	2.73 ^b	0.09	(40)	-21.04 ^c	South Carolina	4.02 ^b	0.13	(07)	16.30 ^c
Iowa	2.47 ^b	0.12	(48)	-28.65 ^c	South Dakota	3.09	0.26	(27)	-10.55
Kansas	2.51 ^b	0.13	(46)	-27.36 ^c	Tennessee	3.35	0.10	(20)	-3.17
Kentucky	3.04 ^b	0.12	(32)	-12.09	Texas	3.73 ^b	0.06	(14)	8.03
Louisiana	4.72 ^b	0.15	(02)	36.52 ^c	Utah	2.10 ^b	0.15	(51)	-39.11 ^c
Maine	2.65 ^b	0.18	(42)	-23.17 ^c	Vermont	2.50 ^b	0.26	(47)	-27.53 ^c
Maryland	3.60	0.11	(16)	4.34	Virginia	3.21	0.09	(23)	-7.05
Massachusetts	3.42	0.10	(19)	-1.06	Washington	3.07 ^b	0.10	(30)	-11.13
Michigan	3.24	0.08	(22)	-6.21	West Virginia	2.89 ^b	0.16	(37)	-16.30 ^c
Minnesota	2.90 ^b	0.10	(36)	-16.18 ^c	Wisconsin	2.92 ^b	0.10	(35)	-15.44 ^c
Mississippi	3.98	0.16	(09)	15.22 ^c	Wyoming	2.13 ^b	0.28	(50)	-38.30 ^c
Missouri	2.75 ^b	0.09	(38)	-20.38 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.19
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.73	0.03		
High Five States				
Hawaii	7.70 ^b	0.48	(01)	62.90 ^c
Louisiana	6.62 ^b	0.27	(02)	40.05 ^c
D.C.	6.53	0.76	(03)	38.18 ^c
Delaware	6.04	0.53	(04)	27.63 ^c
South Carolina	5.91 ^b	0.24	(05)	24.99 ^c
Low Five States				
Iowa	3.45 ^b	0.21	(47)	-26.97 ^c
Kansas	3.40 ^b	0.23	(48)	-28.13 ^c
Montana	3.14 ^b	0.35	(49)	-33.56 ^c
Wyoming	2.90 ^b	0.48	(50)	-38.58 ^c
Utah	2.85 ^b	0.25	(51)	-39.77 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.86	0.22	(19)	2.78	Montana	3.14 ^b	0.35	(49)	-33.56 ^c
Alaska	5.02	0.74	(15)	6.17	Nebraska	3.53 ^b	0.29	(44)	-25.41 ^c
Arizona	3.95 ^b	0.16	(38)	-16.54 ^c	Nevada	4.00	0.28	(33)	-15.36 ^c
Arkansas	4.18	0.25	(29)	-11.55	New Hampshire	3.48 ^b	0.33	(46)	-26.41 ^c
California	5.72 ^b	0.09	(07)	20.88 ^c	New Jersey	5.49 ^b	0.17	(09)	16.10 ^c
Colorado	3.77 ^b	0.20	(40)	-20.26 ^c	New Mexico	5.72	0.35	(06)	20.88 ^c
Connecticut	5.38	0.25	(10)	13.72	New York	5.54 ^b	0.11	(08)	17.08 ^c
Delaware	6.04	0.53	(04)	27.63 ^c	North Carolina	4.54	0.15	(24)	-4.05
D.C.	6.53	0.76	(03)	38.18 ^c	North Dakota	3.97	0.48	(35)	-16.07 ^c
Florida	4.04 ^b	0.09	(32)	-14.60	Ohio	4.34	0.13	(26)	-8.23
Georgia	4.91	0.17	(17)	3.77	Oklahoma	4.56	0.23	(23)	-3.55
Hawaii	7.70 ^b	0.48	(01)	62.90 ^c	Oregon	3.55 ^b	0.20	(43)	-24.85 ^c
Idaho	3.52 ^b	0.33	(45)	-25.47 ^c	Pennsylvania	4.56	0.12	(22)	-3.49
Illinois	5.20	0.14	(12)	10.02	Rhode Island	5.13	0.44	(13)	8.40
Indiana	3.96 ^b	0.17	(36)	-16.30 ^c	South Carolina	5.91 ^b	0.24	(05)	24.99 ^c
Iowa	3.45 ^b	0.21	(47)	-26.97 ^c	South Dakota	3.98	0.44	(34)	-15.95 ^c
Kansas	3.40 ^b	0.23	(48)	-28.13 ^c	Tennessee	4.87	0.19	(18)	3.01
Kentucky	4.22	0.21	(27)	-10.75	Texas	4.95	0.10	(16)	4.70
Louisiana	6.62 ^b	0.27	(02)	40.05 ^c	Utah	2.85 ^b	0.25	(51)	-39.77 ^c
Maine	3.88	0.33	(39)	-17.86 ^c	Vermont	3.61	0.49	(42)	-23.59 ^c
Maryland	5.05	0.21	(14)	6.83	Virginia	4.37	0.16	(25)	-7.68
Massachusetts	4.77	0.18	(20)	0.85	Washington	4.22	0.17	(28)	-10.82
Michigan	4.59	0.14	(21)	-2.93	West Virginia	3.95	0.28	(37)	-16.38 ^c
Minnesota	4.06 ^b	0.18	(31)	-14.15	Wisconsin	4.16	0.18	(30)	-12.10
Mississippi	5.37	0.30	(11)	13.50	Wyoming	2.90 ^b	0.48	(50)	-38.58 ^c
Missouri	3.67 ^b	0.16	(41)	-22.35 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 24.20
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.48	0.02		
High Five States				
Hawaii	3.98 ^b	0.30	(01)	60.54 ^c
California	3.30 ^b	0.06	(02)	32.90 ^c
Louisiana	3.29 ^b	0.16	(03)	32.49 ^c
D.C.	3.22	0.45	(04)	29.95 ^c
Mississippi	3.04	0.19	(05)	22.47 ^c
Low Five States				
Nebraska	1.72 ^b	0.18	(47)	-30.67 ^c
Iowa	1.70 ^b	0.13	(48)	-31.34 ^c
Maine	1.68 ^b	0.19	(49)	-32.12 ^c
Utah	1.51 ^b	0.17	(50)	-39.01 ^c
Wyoming	1.43	0.31	(51)	-42.43 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.60	0.13	(13)	4.80	Montana	1.78	0.24	(44)	-28.09 ^c
Alaska	2.96	0.50	(06)	19.37 ^c	Nebraska	1.72 ^b	0.18	(47)	-30.67 ^c
Arizona	2.15	0.11	(31)	-13.12	Nevada	2.33	0.19	(21)	-6.11
Arkansas	2.28	0.16	(23)	-7.93	New Hampshire	1.85	0.21	(42)	-25.57 ^c
California	3.30 ^b	0.06	(02)	32.90 ^c	New Jersey	2.83	0.10	(09)	14.17
Colorado	1.92 ^b	0.13	(40)	-22.64 ^c	New Mexico	2.93	0.23	(07)	18.14 ^c
Connecticut	2.49	0.14	(16)	0.46	New York	2.88 ^b	0.07	(08)	16.01 ^c
Delaware	2.40	0.30	(19)	-3.05	North Carolina	2.27	0.09	(24)	-8.56
D.C.	3.22	0.45	(04)	29.95 ^c	North Dakota	2.24	0.32	(26)	-9.82
Florida	2.26 ^b	0.06	(25)	-8.85	Ohio	2.14 ^b	0.08	(32)	-13.59
Georgia	2.76	0.11	(11)	11.26	Oklahoma	2.01	0.14	(37)	-19.03 ^c
Hawaii	3.98 ^b	0.30	(01)	60.54 ^c	Oregon	1.76 ^b	0.13	(45)	-28.95 ^c
Idaho	1.93	0.22	(39)	-22.14 ^c	Pennsylvania	2.18 ^b	0.07	(29)	-12.23
Illinois	2.69	0.09	(12)	8.37	Rhode Island	2.33	0.26	(20)	-5.92
Indiana	1.81 ^b	0.10	(43)	-27.06 ^c	South Carolina	2.51	0.14	(15)	1.28
Iowa	1.70 ^b	0.13	(48)	-31.34 ^c	South Dakota	2.45	0.31	(17)	-1.33
Kansas	1.88 ^b	0.15	(41)	-24.16 ^c	Tennessee	2.23	0.11	(27)	-10.05
Kentucky	2.12	0.13	(34)	-14.70	Texas	2.79 ^b	0.07	(10)	12.29
Louisiana	3.29 ^b	0.16	(03)	32.49 ^c	Utah	1.51 ^b	0.17	(50)	-39.01 ^c
Maine	1.68 ^b	0.19	(49)	-32.12 ^c	Vermont	1.72	0.29	(46)	-30.50 ^c
Maryland	2.56	0.13	(14)	3.40	Virginia	2.33	0.10	(22)	-6.19
Massachusetts	2.42	0.11	(18)	-2.33	Washington	2.16	0.11	(30)	-12.81
Michigan	2.21	0.08	(28)	-10.74	West Virginia	2.13	0.19	(33)	-14.04
Minnesota	2.01 ^b	0.12	(36)	-18.99 ^c	Wisconsin	1.99 ^b	0.11	(38)	-19.82 ^c
Mississippi	3.04	0.19	(05)	22.47 ^c	Wyoming	1.43	0.31	(51)	-42.43 ^c
Missouri	2.01 ^b	0.11	(35)	-18.78 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 25.14
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.25	0.01		
High Five States				
Oklahoma	0.37	0.07	(01)	48.28 ^c
Arizona	0.35	0.05	(02)	42.96 ^c
Missouri	0.35	0.05	(03)	41.23 ^c
Pennsylvania	0.32	0.03	(04)	28.87 ^c
California	0.31 ^b	0.02	(05)	25.65 ^c
Low Five States				
Georgia	0.19	0.03	(24)	-24.53 ^c
South Carolina	0.17	0.04	(25)	-31.44 ^c
Massachusetts	0.14	0.03	(26)	-42.36 ^c
Virginia	0.14 ^b	0.03	(27)	-43.04 ^c
Minnesota	0.13 ^b	0.03	(28)	-47.69 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.30	0.05	(06)	19.95 ^c	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.35	0.05	(02)	42.96 ^c	Nevada	-	-	(-)	-
Arkansas	0.28	0.07	(08)	14.38	New Hampshire	-	-	(-)	-
California	0.31 ^b	0.02	(05)	25.65 ^c	New Jersey	0.25	0.03	(12)	1.78
Colorado	0.24	0.04	(16)	-4.22	New Mexico	-	-	(-)	-
Connecticut	-	-	(-)	-	New York	0.20	0.02	(22)	-17.73 ^c
Delaware	-	-	(-)	-	North Carolina	0.28	0.04	(09)	13.64
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.26	0.02	(11)	3.78	Ohio	0.23	0.03	(18)	-7.54
Georgia	0.19	0.03	(24)	-24.53 ^c	Oklahoma	0.37	0.07	(01)	48.28 ^c
Hawaii	-	-	(-)	-	Oregon	-	-	(-)	-
Idaho	-	-	(-)	-	Pennsylvania	0.32	0.03	(04)	28.87 ^c
Illinois	0.24	0.03	(17)	-4.42	Rhode Island	-	-	(-)	-
Indiana	0.26	0.04	(10)	4.89	South Carolina	0.17	0.04	(25)	-31.44 ^c
Iowa	-	-	(-)	-	South Dakota	-	-	(-)	-
Kansas	-	-	(-)	-	Tennessee	0.23	0.04	(19)	-8.00
Kentucky	0.24	0.05	(15)	-3.06	Texas	0.29	0.02	(07)	16.60 ^c
Louisiana	0.19	0.04	(23)	-24.24 ^c	Utah	-	-	(-)	-
Maine	-	-	(-)	-	Vermont	-	-	(-)	-
Maryland	0.22	0.04	(20)	-9.37	Virginia	0.14 ^b	0.03	(27)	-43.04 ^c
Massachusetts	0.14	0.03	(26)	-42.36 ^c	Washington	0.22	0.04	(21)	-9.61
Michigan	0.25	0.03	(13)	1.37	West Virginia	-	-	(-)	-
Minnesota	0.13 ^b	0.03	(28)	-47.69 ^c	Wisconsin	0.25	0.04	(14)	-0.09
Mississippi	-	-	(-)	-	Wyoming	-	-	(-)	-
Missouri	0.35	0.05	(03)	41.23 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 26.18
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Alaska	0.80	0.19	(01)	57.62 ^c
Vermont	0.76	0.14	(02)	48.16 ^c
Idaho	0.66	0.09	(03)	29.84 ^c
Hawaii	0.66	0.09	(04)	29.12 ^c
D.C.	0.65	0.15	(05)	28.23 ^c
Low Five States				
Maryland	0.42	0.04	(46)	-17.44 ^c
Mississippi	0.40	0.05	(47)	-20.87 ^c
South Carolina	0.40	0.04	(48)	-21.42 ^c
Georgia	0.39 ^b	0.03	(49)	-22.76 ^c
Montana	0.38	0.08	(50)	-24.91 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.45	0.04	(44)	-11.62	Montana	0.38	0.08	(50)	-24.91 ^c
Alaska	0.80	0.19	(01)	57.62 ^c	Nebraska	0.48	0.07	(36)	-5.72
Arizona	0.49	0.04	(28)	-3.65	Nevada	0.46	0.06	(42)	-9.78
Arkansas	0.49	0.06	(26)	-3.60	New Hampshire	0.54	0.09	(14)	6.21
California	0.61 ^b	0.02	(07)	19.20 ^c	New Jersey	0.49	0.03	(33)	-4.60
Colorado	0.57	0.05	(09)	11.09	New Mexico	0.51	0.07	(25)	-0.77
Connecticut	0.45	0.05	(43)	-11.45	New York	0.53	0.02	(17)	4.00
Delaware	0.62	0.11	(06)	20.79 ^c	North Carolina	0.49	0.03	(31)	-4.26
D.C.	0.65	0.15	(05)	28.23 ^c	North Dakota	0.49	0.11	(27)	-3.64
Florida	0.43 ^b	0.02	(45)	-14.91	Ohio	0.52	0.03	(20)	2.02
Georgia	0.39 ^b	0.03	(49)	-22.76 ^c	Oklahoma	0.49	0.05	(34)	-4.65
Hawaii	0.66	0.09	(04)	29.12 ^c	Oregon	0.52	0.05	(19)	2.70
Idaho	0.66	0.09	(03)	29.84 ^c	Pennsylvania	0.52	0.03	(21)	1.91
Illinois	0.48	0.03	(35)	-5.61	Rhode Island	0.49	0.09	(32)	-4.32
Indiana	0.49	0.04	(29)	-4.16	South Carolina	0.40	0.04	(48)	-21.42 ^c
Iowa	0.47	0.05	(41)	-8.45	South Dakota	0.56	0.11	(10)	9.61
Kansas	0.53	0.06	(16)	4.09	Tennessee	0.48	0.04	(37)	-6.41
Kentucky	0.51	0.05	(23)	-0.27	Texas	0.55	0.02	(13)	8.18
Louisiana	0.47	0.05	(40)	-8.40	Utah	0.53	0.07	(18)	3.60
Maine	0.47	0.07	(39)	-8.09	Vermont	0.76	0.14	(02)	48.16 ^c
Maryland	0.42	0.04	(46)	-17.44 ^c	Virginia	0.49	0.04	(30)	-4.22
Massachusetts	0.55	0.04	(12)	8.37	Washington	0.57	0.04	(08)	11.11
Michigan	0.47	0.03	(38)	-7.52	West Virginia	0.52	0.07	(22)	0.94
Minnesota	0.51	0.04	(24)	-0.76	Wisconsin	0.56	0.04	(11)	9.55
Mississippi	0.40	0.05	(47)	-20.87 ^c	Wyoming	-	-	(-)	-
Missouri	0.53	0.04	(15)	4.21					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.19
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Utah	0.68	0.12	(01)	33.05 ^c
New Hampshire	0.66	0.15	(02)	30.34 ^c
Kansas	0.64	0.10	(03)	25.79 ^c
Colorado	0.63	0.08	(04)	22.85 ^c
Idaho	0.61	0.14	(05)	20.45 ^c
Low Five States				
Arkansas	0.41	0.08	(36)	-19.86 ^c
Illinois	0.40	0.04	(37)	-20.63 ^c
South Carolina	0.40	0.06	(38)	-22.13 ^c
New Mexico	0.39	0.09	(39)	-22.56 ^c
Mississippi	0.35	0.08	(40)	-30.87 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.51	0.07	(21)	-0.55	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.55	0.11	(14)	7.27
Arizona	0.53	0.06	(17)	4.97	Nevada	0.44	0.08	(31)	-14.01
Arkansas	0.41	0.08	(36)	-19.86 ^c	New Hampshire	0.66	0.15	(02)	30.34 ^c
California	0.55	0.03	(13)	7.74	New Jersey	0.53	0.05	(18)	4.01
Colorado	0.63	0.08	(04)	22.85 ^c	New Mexico	0.39	0.09	(39)	-22.56 ^c
Connecticut	0.47	0.07	(29)	-8.20	New York	0.54	0.04	(15)	5.72
Delaware	-	-	(-)	-	North Carolina	0.49	0.05	(23)	-3.07
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.44	0.03	(30)	-13.18	Ohio	0.57	0.05	(07)	12.81
Georgia	0.42	0.05	(35)	-17.30 ^c	Oklahoma	0.57	0.08	(10)	11.02
Hawaii	-	-	(-)	-	Oregon	0.52	0.08	(20)	2.64
Idaho	0.61	0.14	(05)	20.45 ^c	Pennsylvania	0.50	0.04	(22)	-2.56
Illinois	0.40	0.04	(37)	-20.63 ^c	Rhode Island	-	-	(-)	-
Indiana	0.47	0.06	(28)	-7.57	South Carolina	0.40	0.06	(38)	-22.13 ^c
Iowa	0.56	0.09	(12)	10.46	South Dakota	-	-	(-)	-
Kansas	0.64	0.10	(03)	25.79 ^c	Tennessee	0.48	0.06	(26)	-5.41
Kentucky	0.49	0.07	(25)	-4.46	Texas	0.56	0.04	(11)	10.85
Louisiana	0.42	0.07	(34)	-16.73 ^c	Utah	0.68	0.12	(01)	33.05 ^c
Maine	-	-	(-)	-	Vermont	-	-	(-)	-
Maryland	0.43	0.06	(32)	-15.95 ^c	Virginia	0.57	0.06	(08)	12.34
Massachusetts	0.58	0.06	(06)	13.84	Washington	0.57	0.06	(09)	11.54
Michigan	0.47	0.05	(27)	-7.28	West Virginia	0.43	0.09	(33)	-16.41 ^c
Minnesota	0.54	0.07	(16)	5.12	Wisconsin	0.49	0.06	(24)	-3.89
Mississippi	0.35	0.08	(40)	-30.87 ^c	Wyoming	-	-	(-)	-
Missouri	0.53	0.06	(19)	3.28					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 26.20
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Vermont	1.02	0.23	(01)	101.48 ^c
Hawaii	0.89	0.15	(02)	75.47 ^c
Idaho	0.71	0.13	(03)	40.59 ^c
North Dakota	0.65	0.18	(04)	28.67 ^c
California	0.65 ^b	0.03	(05)	28.39 ^c
Low Five States				
Iowa	0.40	0.06	(42)	-20.48 ^c
South Carolina	0.40	0.05	(43)	-21.12 ^c
Utah	0.39	0.09	(44)	-22.19 ^c
Nebraska	0.38	0.08	(45)	-24.35 ^c
Georgia	0.36 ^b	0.04	(46)	-28.45 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.41	0.05	(40)	-18.50 ^c	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.38	0.08	(45)	-24.35 ^c
Arizona	0.45	0.05	(33)	-10.23	Nevada	0.47	0.08	(28)	-7.44
Arkansas	0.55	0.08	(10)	7.87	New Hampshire	0.45	0.11	(32)	-10.18
California	0.65 ^b	0.03	(05)	28.39 ^c	New Jersey	0.46	0.04	(31)	-9.57
Colorado	0.50	0.06	(21)	-0.89	New Mexico	0.58	0.10	(08)	13.69
Connecticut	0.44	0.06	(34)	-12.88	New York	0.52	0.03	(19)	3.49
Delaware	0.61	0.14	(07)	19.92 ^c	North Carolina	0.48	0.04	(24)	-4.56
D.C.	-	-	(-)	-	North Dakota	0.65	0.18	(04)	28.67 ^c
Florida	0.42	0.02	(38)	-16.12 ^c	Ohio	0.47	0.04	(26)	-6.80
Georgia	0.36 ^b	0.04	(46)	-28.45 ^c	Oklahoma	0.44	0.06	(35)	-13.37
Hawaii	0.89	0.15	(02)	75.47 ^c	Oregon	0.53	0.07	(15)	5.05
Idaho	0.71	0.13	(03)	40.59 ^c	Pennsylvania	0.53	0.03	(14)	5.14
Illinois	0.53	0.04	(18)	3.94	Rhode Island	0.40	0.10	(41)	-20.43 ^c
Indiana	0.48	0.05	(23)	-4.49	South Carolina	0.40	0.05	(43)	-21.12 ^c
Iowa	0.40	0.06	(42)	-20.48 ^c	South Dakota	-	-	(-)	-
Kansas	0.46	0.07	(30)	-9.10	Tennessee	0.47	0.05	(27)	-7.09
Kentucky	0.53	0.06	(17)	4.32	Texas	0.54	0.03	(13)	6.31
Louisiana	0.50	0.06	(22)	-2.24	Utah	0.39	0.09	(44)	-22.19 ^c
Maine	0.54	0.11	(12)	7.40	Vermont	1.02	0.23	(01)	101.48 ^c
Maryland	0.42	0.05	(39)	-17.67 ^c	Virginia	0.43	0.04	(37)	-14.77
Massachusetts	0.52	0.05	(20)	3.40	Washington	0.56	0.06	(09)	10.04
Michigan	0.46	0.04	(29)	-8.53	West Virginia	0.55	0.09	(11)	7.63
Minnesota	0.48	0.06	(25)	-5.87	Wisconsin	0.61	0.06	(06)	21.00 ^c
Mississippi	0.43	0.07	(36)	-14.31	Wyoming	-	-	(-)	-
Missouri	0.53	0.05	(16)	4.72					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.18
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.42	0.02		
High Five States				
Maine	6.19 ^b	0.27	(01)	40.17 ^c
Vermont	5.60	0.40	(02)	26.88 ^c
Nevada	5.35 ^b	0.22	(03)	21.06 ^c
Massachusetts	5.18 ^b	0.12	(04)	17.35 ^c
Ohio	5.06 ^b	0.09	(05)	14.70
Low Five States				
South Dakota	3.71	0.28	(47)	-16.07 ^c
New Mexico	3.68 ^b	0.19	(48)	-16.74 ^c
Mississippi	3.44 ^b	0.15	(49)	-21.99 ^c
Alaska	3.42	0.43	(50)	-22.61 ^c
Hawaii	2.55 ^b	0.18	(51)	-42.14 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.80 ^b	0.12	(43)	-13.95	Montana	4.75	0.28	(17)	7.62
Alaska	3.42	0.43	(50)	-22.61 ^c	Nebraska	4.09	0.20	(39)	-7.43
Arizona	4.15	0.11	(36)	-6.04	Nevada	5.35 ^b	0.22	(03)	21.06 ^c
Arkansas	4.54	0.17	(23)	2.92	New Hampshire	4.94	0.26	(06)	11.89
California	3.97 ^b	0.05	(42)	-10.16	New Jersey	4.89 ^b	0.10	(08)	10.75
Colorado	3.78 ^b	0.13	(45)	-14.47	New Mexico	3.68 ^b	0.19	(48)	-16.74 ^c
Connecticut	4.84	0.15	(10)	9.58	New York	4.49	0.06	(25)	1.73
Delaware	4.74	0.31	(18)	7.34	North Carolina	4.20	0.09	(34)	-4.93
D.C.	4.08	0.38	(40)	-7.71	North Dakota	4.85	0.34	(09)	9.89
Florida	4.49	0.06	(24)	1.73	Ohio	5.06 ^b	0.09	(05)	14.70
Georgia	4.37	0.11	(29)	-0.93	Oklahoma	4.48	0.15	(27)	1.38
Hawaii	2.55 ^b	0.18	(51)	-42.14 ^c	Oregon	4.81	0.15	(15)	8.98
Idaho	4.30	0.24	(31)	-2.59	Pennsylvania	4.84 ^b	0.08	(11)	9.54
Illinois	4.44	0.08	(28)	0.51	Rhode Island	4.94	0.28	(07)	11.79
Indiana	4.70	0.12	(19)	6.34	South Carolina	4.14	0.13	(37)	-6.32
Iowa	4.58	0.16	(22)	3.81	South Dakota	3.71	0.28	(47)	-16.07 ^c
Kansas	4.07	0.16	(41)	-7.76	Tennessee	4.23	0.11	(33)	-4.15
Kentucky	4.77	0.15	(16)	8.04	Texas	3.74 ^b	0.06	(46)	-15.32 ^c
Louisiana	4.15	0.14	(35)	-6.02	Utah	3.79	0.20	(44)	-14.13
Maine	6.19 ^b	0.27	(01)	40.17 ^c	Vermont	5.60	0.40	(02)	26.88 ^c
Maryland	4.83	0.13	(13)	9.34	Virginia	4.27	0.11	(32)	-3.34
Massachusetts	5.18 ^b	0.12	(04)	17.35 ^c	Washington	4.83	0.12	(12)	9.47
Michigan	4.81 ^b	0.09	(14)	9.04	West Virginia	4.61	0.20	(21)	4.32
Minnesota	4.10	0.12	(38)	-7.10	Wisconsin	4.66	0.12	(20)	5.47
Mississippi	3.44 ^b	0.15	(49)	-21.99 ^c	Wyoming	4.49	0.40	(26)	1.72
Missouri	4.36	0.11	(30)	-1.36					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.19
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.67	0.03		
High Five States				
Maine	10.86 ^b	0.56	(01)	41.53 ^c
Vermont	9.69	0.82	(02)	26.25 ^c
Massachusetts	9.01 ^b	0.25	(03)	17.46 ^c
Ohio	8.88 ^b	0.19	(04)	15.73 ^c
New Hampshire	8.66	0.55	(05)	12.82
Low Five States				
New Mexico	6.47	0.39	(47)	-15.63 ^c
Mississippi	6.25 ^b	0.34	(48)	-18.55 ^c
D.C.	6.07	0.76	(49)	-20.95 ^c
Alaska	5.49	0.83	(50)	-28.48 ^c
Hawaii	4.41 ^b	0.36	(51)	-42.56 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.74	0.27	(41)	-12.12	Montana	8.61	0.59	(06)	12.18
Alaska	5.49	0.83	(50)	-28.48 ^c	Nebraska	7.04	0.41	(38)	-8.26
Arizona	7.12	0.22	(36)	-7.18	Nevada	8.49	0.43	(12)	10.61
Arkansas	7.56	0.34	(29)	-1.44	New Hampshire	8.66	0.55	(05)	12.82
California	6.73 ^b	0.10	(42)	-12.28	New Jersey	8.59 ^b	0.21	(07)	11.99
Colorado	6.48 ^b	0.27	(46)	-15.51 ^c	New Mexico	6.47	0.39	(47)	-15.63 ^c
Connecticut	8.48	0.32	(14)	10.48	New York	7.96	0.14	(22)	3.74
Delaware	8.11	0.64	(21)	5.70	North Carolina	7.40	0.21	(31)	-3.55
D.C.	6.07	0.76	(49)	-20.95 ^c	North Dakota	8.57	0.70	(08)	11.70
Florida	7.62	0.12	(28)	-0.74	Ohio	8.88 ^b	0.19	(04)	15.73 ^c
Georgia	7.63	0.23	(27)	-0.53	Oklahoma	7.83	0.32	(23)	2.10
Hawaii	4.41 ^b	0.36	(51)	-42.56 ^c	Oregon	8.44	0.31	(17)	9.99
Idaho	7.40	0.48	(32)	-3.59	Pennsylvania	8.55 ^b	0.16	(09)	11.47
Illinois	7.76	0.17	(25)	1.16	Rhode Island	8.45	0.58	(16)	10.10
Indiana	8.24	0.25	(20)	7.42	South Carolina	6.78	0.28	(40)	-11.66
Iowa	8.50	0.34	(10)	10.77	South Dakota	6.56	0.57	(43)	-14.52
Kansas	6.89	0.33	(39)	-10.25	Tennessee	7.64	0.25	(26)	-0.42
Kentucky	8.50	0.32	(11)	10.76	Texas	6.51 ^b	0.13	(45)	-15.18 ^c
Louisiana	7.05	0.29	(37)	-8.08	Utah	6.56	0.40	(44)	-14.53
Maine	10.86 ^b	0.56	(01)	41.53 ^c	Vermont	9.69	0.82	(02)	26.25 ^c
Maryland	8.49	0.28	(13)	10.60	Virginia	7.36	0.22	(33)	-4.14
Massachusetts	9.01 ^b	0.25	(03)	17.46 ^c	Washington	8.33	0.25	(19)	8.51
Michigan	8.48 ^b	0.20	(15)	10.47	West Virginia	7.76	0.41	(24)	1.17
Minnesota	7.27	0.25	(35)	-5.19	Wisconsin	8.36	0.26	(18)	9.01
Mississippi	6.25 ^b	0.34	(48)	-18.55 ^c	Wyoming	7.28	0.78	(34)	-5.18
Missouri	7.51	0.24	(30)	-2.15					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.20
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.22	0.02		
High Five States				
D.C.	2.96	0.43	(01)	33.31 ^c
Nevada	2.91	0.22	(02)	31.41 ^c
Maine	2.90	0.25	(03)	30.93 ^c
Massachusetts	2.72 ^b	0.11	(04)	22.52 ^c
Vermont	2.71	0.36	(05)	22.39 ^c
Low Five States				
South Dakota	1.69	0.24	(47)	-23.64 ^c
Mississippi	1.69 ^b	0.14	(48)	-23.73 ^c
Utah	1.66	0.18	(49)	-24.99 ^c
New Mexico	1.60 ^b	0.17	(50)	-28.01 ^c
Hawaii	1.20 ^b	0.16	(51)	-45.87 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.87	0.11	(43)	-15.61 ^c	Montana	1.88	0.24	(42)	-15.13 ^c
Alaska	1.74	0.42	(46)	-21.48 ^c	Nebraska	2.09	0.19	(35)	-5.77
Arizona	1.89	0.10	(41)	-14.73	Nevada	2.91	0.22	(02)	31.41 ^c
Arkansas	2.47	0.16	(10)	11.59	New Hampshire	2.39	0.24	(16)	7.79
California	2.04 ^b	0.05	(36)	-7.94	New Jersey	2.54 ^b	0.09	(08)	14.76
Colorado	1.82	0.12	(45)	-17.98 ^c	New Mexico	1.60 ^b	0.17	(50)	-28.01 ^c
Connecticut	2.45	0.14	(12)	10.36	New York	2.28	0.06	(24)	2.97
Delaware	2.46	0.29	(11)	10.85	North Carolina	2.21	0.09	(29)	-0.12
D.C.	2.96	0.43	(01)	33.31 ^c	North Dakota	2.23	0.31	(26)	0.71
Florida	2.16	0.06	(34)	-2.69	Ohio	2.54 ^b	0.08	(07)	14.77
Georgia	2.35	0.10	(20)	5.78	Oklahoma	2.18	0.14	(32)	-1.61
Hawaii	1.20 ^b	0.16	(51)	-45.87 ^c	Oregon	2.18	0.13	(33)	-1.71
Idaho	1.96	0.22	(38)	-11.54	Pennsylvania	2.41	0.07	(13)	8.85
Illinois	2.28	0.08	(25)	2.75	Rhode Island	2.70	0.27	(06)	21.79 ^c
Indiana	2.36	0.11	(19)	6.33	South Carolina	2.41	0.13	(14)	8.58
Iowa	1.95	0.13	(39)	-12.04	South Dakota	1.69	0.24	(47)	-23.64 ^c
Kansas	2.19	0.16	(31)	-1.01	Tennessee	2.03	0.10	(37)	-8.46
Kentucky	2.33	0.13	(21)	5.10	Texas	1.85 ^b	0.06	(44)	-16.72 ^c
Louisiana	2.29	0.13	(23)	3.23	Utah	1.66	0.18	(49)	-24.99 ^c
Maine	2.90	0.25	(03)	30.93 ^c	Vermont	2.71	0.36	(05)	22.39 ^c
Maryland	2.54	0.12	(09)	14.51	Virginia	2.23	0.10	(28)	0.41
Massachusetts	2.72 ^b	0.11	(04)	22.52 ^c	Washington	2.37	0.11	(18)	6.83
Michigan	2.37	0.09	(17)	6.99	West Virginia	2.41	0.19	(15)	8.49
Minnesota	1.92	0.11	(40)	-13.29	Wisconsin	2.20	0.11	(30)	-0.64
Mississippi	1.69 ^b	0.14	(48)	-23.73 ^c	Wyoming	2.32	0.39	(22)	4.48
Missouri	2.23	0.11	(27)	0.58					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.