

Table 1.10
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 2003-2012

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.1*	-2.1*	-0.2	-2.1*	-2.6*	-1.6*
Oral Cavity & Pharynx:	-2.3*	-2.4*	-2.3*	-3.4*	-3.8*	-2.3*
Lip	-	-	-	-	-	-
Tongue	-1.1	-0.8	-1.8	-3.5*	-4.4*	-1.8
Salivary gland	-1.8*	-2.7	-1.1	1.6	2.7	0.0
Floor of mouth	-3.8*	-3.3	-5.0*	-	-	-
Gum & other oral cavity	-3.0*	-4.0*	-2.1*	-5.2*	-5.8*	-4.3*
Nasopharynx	-2.6	-2.9	-1.8	-3.2*	-3.7*	-2.1
Tonsil	-0.5	-0.2	-1.6	-1.3	-1.2*	-1.7
Oropharynx	-2.2	-2.6	-	-0.1	-1.1	2.0
Hypopharynx	-5.1*	-5.0*	-5.9*	-3.8*	-5.3*	-
Other oral cavity & pharynx	-4.0	-3.9	-	-5.1*	-4.8*	-6.5*
Digestive System:	-1.6*	-1.7*	-1.6*	-1.6*	-1.5*	-1.9*
Esophagus	-4.8*	-5.0*	-4.5*	-4.5*	-4.5*	-4.8*
Stomach	-2.2*	-3.5*	-0.9	-3.0*	-3.3*	-2.8*
Small intestine	2.5*	3.1*	1.9	0.4	0.3	0.8
Colon & Rectum:	-2.8*	-3.0*	-2.8*	-3.0*	-2.6*	-3.4*
Colon	-3.2*	-3.5*	-3.1*	-	-	-
Rectum	-1.7*	-1.5*	-2.0*	-	-	-
Anus, anal canal & anorectum	1.7*	2.4	0.9	3.8*	6.9*	0.8
Liver & intrahepatic bile duct	3.2*	3.4*	2.3*	2.5*	2.7*	1.6*
Gallbladder	0.6	2.4	0.2	1.5	3.5	0.8
Other biliary	1.8	2.0	1.7	-0.9	-0.9	-0.7
Pancreas	-0.3	-0.3	-0.3	-0.4*	-0.5	-0.2
Retroperitoneum	0.9	-	0.8	-7.0*	-	-
Peritoneum, omentum & mesentery	1.6	-	2.9	-0.4	-	-0.3
Other digestive system	4.7*	6.8	2.6	1.0	0.5	1.3
Respiratory System:	-2.2*	-3.0*	-1.2*	-2.7*	-3.4*	-1.8*
Nose, nasal cavity & middle ear	-0.9	-1.7	0.6	-1.4	-3.6	1.8
Larynx	-2.9*	-3.3*	-1.2	-3.8*	-4.2*	-2.7*
Lung & bronchus	-2.1*	-2.9*	-1.3*	-2.6*	-3.4*	-1.8*
Pleura	-	-	-	-	-	-
Trachea & other respiratory organs	-2.4	-	-	-0.3	-	-
Bones & joints	-1.1	1.7	-3.1	-0.7	-0.4	-1.5
Soft tissue (including heart)	-0.1	-0.9	0.6	0.7	0.1	1.1
Skin (excl. basal & squamous):	-0.9	-0.1	-1.7*	-1.8*	-1.7	-1.9
Melanoma of the skin	0.0	-0.4	0.0	-1.5*	-1.6	-1.6
Other non-epithelial skin	-1.8	0.2	-3.5*	-2.2	-1.7	-2.4
Breast	0.5	2.2	0.7*	-1.5*	-0.4	-1.4*
Breast (<i>in situ</i>)	1.8*	-	2.0*	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

* The APC is significantly different from zero ($p<.05$).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.10 - continued
 SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
 Blacks, 2003-2012

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:						
Cervix uteri	0.3	-	0.5	-0.6*	-	-0.4
Corpus uteri	-3.2*	-	-3.1*	-2.4*	-	-2.2*
Uterus, NOS	2.5*	-	2.7*	-0.5	-	-0.2
Ovary ^c	4.6*	-	4.9*	2.7*	-	2.9*
Vagina	-1.6*	-	-1.4*	-1.7*	-	-1.6*
Vulva	0.0	-	-0.2	-3.3*	-	-3.0*
Other female genital system	0.1	-	0.2	1.8	-	2.1
	5.1*	-	4.9*	2.1	-	2.0
Male Genital System:	-2.4*	-2.9*	-	-3.3*	-3.9*	-
Prostate	-2.4*	-2.9*	-	-3.4*	-3.9*	-
Testis	3.2*	3.2*	-	-3.7	-3.9	-
Penis	1.1	1.0	-	-3.1	-4.0	-
Other male genital system	-	-	-	-	-	-
Urinary System:	1.0	0.8	1.1	-1.1*	-1.0*	-1.4*
Urinary bladder	-0.6	-0.3	-1.5	-0.9	-0.7	-1.5*
Kidney & renal pelvis	2.3*	1.7*	2.8*	-1.2*	-1.3*	-1.1
Ureter	-3.1*	-	-	0.2	-	-
Other urinary system	1.1	-	-	-2.4	-	-4.3
Eye & Orbit	0.1	-	-	-	-	-
Brain & Nervous System: ^d	-0.4	0.2	-0.8	0.1	0.0	0.4
Brain	0.0	0.3	-0.1	-	-	-
Cranial nerves & other nervous system	-3.8	-2.1	-5.1*	-	-	-
Endocrine System:	4.4*	2.6	5.0*	1.2	3.0*	0.1
Thyroid	4.9*	3.2*	5.3*	1.2	2.6	0.5
Other endocrine & thymus	1.0	0.5	1.3	1.3	3.3	-0.7
Lymphoma:	-0.4	-0.6	-0.1	-1.7*	-1.1	-2.3*
Hodgkin lymphoma	0.1	0.2	-0.5	-2.2	-2.0	-2.6*
Non-Hodgkin lymphoma	-0.4	-0.8	-0.1	-1.7*	-1.0	-2.3*
Myeloma	0.7*	0.1	1.1	-1.4*	-1.4*	-1.5*
Leukemia:	-0.3	-0.6	0.1	-1.4*	-1.6*	-1.3*
Lymphocytic:	-0.5	-0.7	-0.2	-2.1*	-2.5*	-1.6*
Acute lymphocytic	1.5	1.2	1.7	-2.1*	-3.1*	-0.4
Chronic lymphocytic	-1.1	-1.2	-0.9	-2.0*	-2.1	-2.2*
Other lymphocytic	-0.3	-	-	-3.3	-7.7	-
Myeloid & Monocytic:	0.7	0.1	1.2	-1.0*	-1.0	-1.2
Acute myeloid	1.7	1.5	2.1	-0.4	-0.2	-0.5
Chronic myeloid	-0.5	-2.7	1.3	-4.7*	-6.3*	-3.0
Acute monocytic	-3.1*	-	-	-	-	-
Other myeloid & monocytic	-	-	-	-0.7	2.7	-5.5
Other leukemia:	-5.1*	-4.6	-5.5	-1.4*	-1.5	-1.3
Other acute leukemia	-3.9	-	-	-4.0*	-4.8*	-3.5*
Aleukemic, subleukemic & NOS	-5.6	-6.0*	-4.8	-0.2	0.1	-0.4
Kaposi Sarcoma ^e	-0.5	-0.4	-	-	-	-
Mesothelioma ^e	-2.5	-3.3	-	-	-	-
Ill-defined & unspecified	-2.9*	-3.5*	-2.2*	-3.2*	-3.1*	-3.2*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population
 (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero ($p<.05$).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.