

Table 1.22
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2013^a
 Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			38-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 38 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 38 Years ^d
All Sites	Both Sexes	4,683,633	4,777,436	5,281,162	13,793,147	14,017,809
	Male	2,371,817	2,408,477	2,654,461	6,558,326	6,635,084
	Female	2,311,816	2,368,959	2,626,701	7,234,821	7,382,725
Oral Cavity & Pharynx	Both Sexes	111,858	129,718	134,909	292,631	324,216
	Male	78,943	90,724	93,801	195,801	215,214
	Female	32,915	38,994	41,108	96,830	109,002
Esophagus	Both Sexes	21,605	26,812	26,884	36,756	43,489
	Male	17,134	21,074	21,110	28,826	33,848
	Female	4,471	5,738	5,774	7,930	9,641
Stomach	Both Sexes	38,885	46,951	47,520	78,761	91,003
	Male	22,739	27,768	28,102	44,833	52,171
	Female	16,146	19,183	19,418	33,928	38,832
Colon & Rectum	Both Sexes	393,064	452,807	462,351	1,160,780	1,291,277
	Male	201,444	232,173	236,571	579,395	642,194
	Female	191,620	220,634	225,780	581,385	649,083
Liver & Intrahepatic Bile Duct	Both Sexes	38,940	44,690	44,772	54,464	61,346
	Male	28,968	32,749	32,797	39,263	43,716
	Female	9,972	11,941	11,975	15,201	17,630
Pancreas	Both Sexes	36,831	44,775	44,811	49,363	58,920
	Male	18,821	22,954	22,966	24,251	29,201
	Female	18,010	21,821	21,845	25,112	29,719
Larynx	Both Sexes	32,033	37,848	38,327	87,910	98,553
	Male	25,889	30,586	30,982	70,533	78,905
	Female	6,144	7,262	7,345	17,377	19,648
Lung & Bronchus	Both Sexes	235,891	305,893	316,680	409,799	509,857
	Male	107,952	142,292	146,705	182,649	228,690
	Female	127,939	163,601	169,975	227,150	281,167
Melanoma of the Skin	Both Sexes	293,419	335,454	352,673	1,001,852	1,085,564
	Male	158,125	185,809	196,879	498,832	548,924
	Female	135,294	149,645	155,794	503,020	536,640
Breast	Female	888,374	963,711	1,026,553	3,011,973	3,194,990
Cervix	Female	39,026	41,272	41,346	216,287	222,528
Corpus & Uterus, NOS	Female	190,409	213,555	213,718	615,772	673,630
Ovary ^f	Female	61,283	70,222	70,275	181,920	204,129

^a U.S. 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER 9 registries and 1/1/2013 U.S. population estimates based on the average of 2012 and 2013 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 38 years (1975-2012)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2008-2012)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2008; Melanoma in 2009.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 38-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2013.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 38-year limited duration prevalence. The 2008 breast cancer is counted for the breast 5-year and 38-year limited duration prevalence.

In method (e) the 2008 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2009 melanoma is counted for 5-year limited duration prevalence for melanoma.

^f Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.22 - continued
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2013^a
 Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			38-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 38 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 38 Years ^d
Prostate	Male	996,735	1,077,771	1,077,866	2,849,303	3,033,987
Testis	Male	42,726	43,748	44,378	226,015	229,281
Urinary Bladder	Both Sexes	203,937	256,726	262,276	578,563	677,871
	Male	154,968	196,098	200,600	432,356	507,228
	Female	48,969	60,628	61,676	146,207	170,643
Kidney & Renal Pelvis	Both Sexes	158,086	192,740	195,379	384,082	448,864
	Male	97,484	120,269	122,119	228,920	270,059
	Female	60,602	72,471	73,260	155,162	178,805
Brain & Nervous System	Both Sexes	46,359	49,525	49,948	140,209	144,923
	Male	24,907	26,640	26,879	74,171	76,649
	Female	21,452	22,885	23,069	66,038	68,274
Thyroid	Both Sexes	194,792	218,107	218,918	600,285	645,059
	Male	43,408	51,452	51,619	130,022	143,922
	Female	151,384	166,655	167,299	470,263	501,137
Hodgkin Lymphoma	Both Sexes	37,358	39,844	39,912	181,370	186,607
	Male	20,235	21,627	21,671	94,047	96,687
	Female	17,123	18,217	18,241	87,323	89,920
Non-Hodgkin Lymphoma	Both Sexes	208,024	245,167	249,216	561,810	630,027
	Male	112,213	133,122	135,245	296,343	333,061
	Female	95,811	112,045	113,971	265,467	296,966
Myeloma	Both Sexes	60,408	71,647	71,915	95,457	110,345
	Male	32,121	39,215	39,380	51,385	60,395
	Female	28,287	32,432	32,535	44,072	49,950
Leukemia	Both Sexes	124,326	145,980	146,454	328,477	363,794
	Male	72,087	85,110	85,378	186,605	207,382
	Female	52,239	60,870	61,076	141,872	156,412
Acute Lymphocytic Leukemia	Both Sexes	17,063	17,597	17,597	74,508	75,300
	Male	9,316	9,590	9,590	40,643	41,001
	Female	7,747	8,007	8,007	33,865	34,299
Childhood (Ages 0-19)	Both Sexes	66,771	66,880	67,473	337,686	338,228
	Male	34,885	34,966	35,300	173,483	173,747
	Female	31,886	31,914	32,173	164,203	164,481
Kaposi Sarcoma	Both Sexes	7,288	7,954	7,954	26,999	28,478
	Male	6,806	7,404	7,404	25,587	26,891
	Female	482	550	550	1,412	1,587
Mesothelioma	Both Sexes	3,073	4,014	4,014	4,738	5,803
	Male	2,172	2,842	2,842	2,854	3,587
	Female	901	1,172	1,172	1,884	2,216

^a U.S. 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER 9 registries and 1/1/2013 U.S. population estimates based on the average of 2012 and 2013 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 38 years (1975-2012)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2008-2012)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2008; Melanoma in 2009.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 38-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2013.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 38-year limited duration prevalence. The 2008 breast cancer is counted for the breast 5-year and 38-year limited duration prevalence.

In method (e) the 2008 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2009 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.23
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2013^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	6,630,144	18,977	42,515	89,810	160,506	352,035	920,363	1,777,405	3,268,534
Females	7,510,110	17,057	36,959	90,895	235,457	635,354	1,378,596	1,900,156	3,215,637
Oral Cavity & Pharynx									
Males	200,280	28	471	1,653	3,431	13,626	48,036	65,705	67,330
Females	100,402	96	546	1,517	3,868	8,513	19,303	25,928	40,633
Esophagus									
Males	28,878	0	0	6	138	1,080	4,812	10,234	12,607
Females	7,979	0	0	22	34	189	1,353	2,058	4,324
Stomach									
Males	45,442	0	40	177	555	2,588	7,433	12,737	21,912
Females	34,401	0	27	231	757	2,445	4,965	7,644	18,332
Colon & Rectum									
Males	585,136	11	94	1,316	6,064	26,227	85,943	144,658	320,823
Females	592,420	0	230	1,444	6,369	23,982	73,478	117,870	369,046
Liver & Intrahep									
Males	39,334	612	484	604	641	1,590	11,144	15,911	8,350
Females	15,620	413	570	434	455	1,044	3,383	4,681	4,640
Pancreas									
Males	24,363	34	23	153	365	1,584	5,088	7,791	9,324
Females	25,257	0	12	288	536	1,593	4,671	7,156	11,001
Larynx									
Males	71,488	0	0	89	292	1,713	10,143	21,196	38,055
Females	17,593	0	0	28	144	847	3,543	5,051	7,979
Lung & Bronchus									
Males	186,030	34	75	517	1,120	4,955	23,975	55,413	99,941
Females	229,677	34	104	399	1,489	6,572	31,088	61,774	128,218
Melanoma of the Skin									
Males	509,094	56	696	4,938	17,619	45,039	99,914	144,398	196,432
Females	525,366	113	834	10,310	34,836	72,511	121,614	127,402	157,747

^a U.S. 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2013 U.S. population estimates based on the average of 2012 and 2013 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2013^a
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	15,781	0	0	11	115	703	1,911	4,527	8,514
Females	3,053,450	11	47	2,984	36,611	211,630	567,039	845,870	1,389,258
Cervix									
Females	248,920	0	68	1,821	14,820	40,574	59,368	60,139	72,130
Corpus & Uterus, NOS									
Females	635,437	11	23	629	6,040	25,247	93,125	182,278	328,082
Ovary ^d									
Females	195,767	105	1,012	3,796	6,939	18,071	41,987	52,895	70,961
Prostate									
Males	2,850,139	34	46	104	343	18,486	237,236	815,954	1,777,936
Urinary Bladder									
Males	438,068	68	105	551	2,315	9,819	42,325	105,894	276,992
Females	149,358	68	34	241	968	3,908	13,820	31,806	98,514
Kidney & Renal Pelvis									
Males	234,706	1,375	2,285	2,963	5,082	19,921	46,181	68,291	88,610
Females	159,630	1,455	2,581	3,065	5,082	12,617	28,144	40,702	65,983
Hodgkin Lymphoma									
Males	100,096	203	2,405	9,526	16,482	23,489	23,526	16,032	8,431
Females	93,449	34	2,104	9,344	16,640	22,518	21,509	13,067	8,232
Non-Hodgkin Lymphoma									
Males	300,417	927	3,738	8,182	14,080	29,312	57,496	77,491	109,191
Females	269,119	452	1,681	4,846	9,401	21,417	43,880	66,307	121,135
Myeloma									
Males	51,465	0	6	110	539	3,457	9,101	16,940	21,313
Females	44,223	0	0	45	377	2,237	7,664	12,935	20,965
Leukemia									
Males	189,019	6,519	13,341	13,964	12,067	15,409	26,626	39,493	61,600
Females	144,956	6,039	10,458	12,073	11,027	11,191	17,766	26,051	50,352
Acute Lymphocytic Leuk									
Males	42,395	5,394	11,202	10,799	6,965	4,601	1,951	925	559
Females	35,460	5,139	8,796	8,782	6,216	3,365	1,545	1,069	549

^a U.S. 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2013 U.S. population estimates based on the average of 2012 and 2013 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.21
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	4,683,633	3,302,218	2,388,027	1,512,405	914,197	534,563	12,499,657	13,793,147	347,107	14,140,254
	Males	2,371,817	1,687,487	1,160,300	679,900	350,628	174,108	6,148,336	6,558,326	71,818	6,630,144
	Females	2,311,816	1,614,731	1,227,727	832,505	563,569	360,455	6,351,321	7,234,821	275,289	7,510,110
White ^b	Both Sexes	3,918,402	2,837,504	2,087,003	1,335,669	824,524	485,175	10,730,652	11,904,805	302,738	12,207,543
	Males	1,978,263	1,442,623	1,007,901	598,480	319,935	159,759	5,253,660	5,630,885	67,228	5,698,113
	Females	1,940,139	1,394,881	1,079,102	737,189	504,589	325,416	5,476,992	6,273,920	235,510	6,509,430
Black ^b	Both Sexes	496,816	308,718	202,127	116,659	57,326	30,618	1,163,805	1,238,540	21,389	1,259,929
	Males	262,272	169,178	109,086	57,293	20,443	8,443	612,963	632,563	1,749	634,312
	Females	234,544	139,540	93,041	59,366	36,883	22,175	550,842	605,977	19,640	625,617
Asian/ Pacific Islander ^c	Both Sexes	152,791	96,072	63,136	+	+	+	354,019	+	+	+
	Males	63,892	39,982	24,230	+	+	+	142,568	+	+	+
	Females	88,899	56,090	38,906	+	+	+	211,451	+	+	+
Hispanic ^d	Both Sexes	340,558	215,426	141,713	+	+	+	793,696	+	+	+
	Males	159,609	101,176	63,795	+	+	+	363,689	+	+	+
	Females	180,949	114,250	77,918	+	+	+	430,007	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	3.4482%	0.0866%	0.1883%	0.3439%	0.7993%	1.9479%	4.4448%	9.5130%	16.2143%	18.3787%	3.2687%
	Males	3.4122%	0.0900%	0.1984%	0.3357%	0.6326%	1.3471%	3.5996%	10.0279%	20.4610%	26.0822%	3.6128%
	Females	3.4833%	0.0830%	0.1777%	0.3525%	0.9668%	2.5434%	5.2551%	9.0480%	12.7542%	13.7760%	3.0751%
White ^c	Both Sexes	3.8037%	0.0938%	0.2073%	0.3904%	0.8983%	2.0934%	4.6502%	9.9348%	17.1170%	19.1670%	3.4481%
	Males	3.7336%	0.0966%	0.2203%	0.3830%	0.7191%	1.4735%	3.7359%	10.2671%	21.1825%	26.6695%	3.7488%
	Females	3.8735%	0.0910%	0.1936%	0.3982%	1.0869%	2.7343%	5.5560%	9.6241%	13.7170%	14.6278%	3.2916%
Black ^c	Both Sexes	2.6037%	0.0659%	0.1323%	0.2221%	0.5426%	1.5976%	4.2293%	9.9567%	16.2183%	16.3911%	3.1028%
	Males	2.8675%	0.0683%	0.1304%	0.2199%	0.3999%	1.1222%	4.0876%	12.8552%	24.2461%	27.8880%	4.0796%
	Females	2.3635%	0.0634%	0.1343%	0.2243%	0.6678%	2.0155%	4.3528%	7.6916%	10.6142%	10.7379%	2.5029%
Asian/ Pacific Islander ^c	Both Sexes	2.1998%	0.0665%	0.1395%	0.2080%	0.4921%	1.3929%	3.1810%	6.0037%	9.8741%	12.8225%	2.1496%
	Males	1.8903%	0.0731%	0.1446%	0.1907%	0.3314%	0.7223%	1.9710%	5.2688%	11.4823%	17.8667%	2.1040%
	Females	2.4836%	0.0595%	0.1342%	0.2256%	0.6373%	1.9859%	4.2231%	6.6081%	8.6238%	9.7323%	2.2421%
Hispanic ^d	Both Sexes	1.5228%	0.0812%	0.1766%	0.2786%	0.5914%	1.3797%	3.1675%	6.7103%	11.6348%	13.7425%	2.3775%
	Males	1.3722%	0.0850%	0.1866%	0.2807%	0.4606%	0.8730%	2.2715%	6.6874%	15.1864%	20.3684%	2.6257%
	Females	1.6761%	0.0774%	0.1662%	0.2763%	0.7304%	1.9070%	4.0297%	6.7300%	9.0024%	9.7529%	2.2775%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 3.21
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	46,359	26,407	21,630	16,369	12,681	8,623	119,068	140,209	12,542	152,751
	Males	24,907	14,055	10,678	8,929	6,913	4,615	63,225	74,171	5,653	79,824
	Females	21,452	12,352	10,952	7,440	5,768	4,008	55,843	66,038	6,889	72,927
White ^b	Both Sexes	39,980	22,783	19,086	14,228	11,101	7,740	103,303	122,191	3,207	125,398
	Males	21,552	12,361	9,483	7,819	6,030	4,058	55,233	64,997	1,817 ⁱ	66,814
	Females	18,428	10,422	9,603	6,409	5,071	3,682	48,070	57,194	1,390 ⁱ	58,584
Black ^b	Both Sexes	3,743	2,197	1,645	1,302	1,104	610	9,608	11,278	+	+
	Males	1,883	953	794	663	619	387	4,730	5,589	+	+
	Females	1,860	1,244	851	639	485	223	4,878	5,689	+	+
Asian/ Pacific Islander ^c	Both Sexes	1,542	855	568	+	+	+	3,519	+	+	+
	Males	812	396	282	+	+	+	1,761	+	+	+
	Females	730	459	286	+	+	+	1,758	+	+	+
Hispanic ^d	Both Sexes	4,886	3,112	2,463	+	+	+	12,273	+	+	+
	Males	2,718	1,710	1,265	+	+	+	6,579	+	+	+
	Females	2,168	1,402	1,198	+	+	+	5,694	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0329%	0.0145%	0.0329%	0.0346%	0.0335%	0.0371%	0.0416%	0.0409%	0.0328%	0.0184%	0.0323%
	Males	0.0355%	0.0151%	0.0344%	0.0363%	0.0367%	0.0410%	0.0456%	0.0459%	0.0342%	0.0214%	0.0349%
	Females	0.0302%	0.0138%	0.0314%	0.0329%	0.0303%	0.0333%	0.0377%	0.0363%	0.0317%	0.0167%	0.0298%
White ^c	Both Sexes	0.0373%	0.0163%	0.0368%	0.0398%	0.0385%	0.0427%	0.0477%	0.0450%	0.0371%	0.0189%	0.0367%
	Males	0.0404%	0.0169%	0.0390%	0.0418%	0.0417%	0.0472%	0.0522%	0.0498%	0.0393%	0.0218%	0.0396%
	Females	0.0342%	0.0155%	0.0344%	0.0378%	0.0352%	0.0380%	0.0432%	0.0405%	0.0353%	0.0171%	0.0338%
Black ^c	Both Sexes	0.0210%	0.0104%	0.0243%	0.0229%	0.0221%	0.0223%	0.0227%	0.0293%	0.0131%	0.0156%	0.0207%
	Males	0.0212%	0.0105%	0.0228%	0.0245%	0.0219%	0.0227%	0.0227%	0.0321%	0.0105%	0.0178%	0.0208%
	Females	0.0209%	0.0104%	0.0258%	0.0214%	0.0222%	0.0218%	0.0228%	0.0272%	0.0149%	0.0145%	0.0206%
Asian/ Pacific Islander ^c	Both Sexes	0.0191%	0.0089%	0.0203%	0.0195%	0.0188%	0.0205%	0.0222%	0.0245%	0.0209%	0.0148%	0.0188%
	Males	0.0200%	0.0090%	0.0197%	0.0191%	0.0221%	0.0212%	0.0237%	0.0292%	0.0177%	0.0171%	0.0197%
	Females	0.0182%	0.0088%	0.0209%	0.0198%	0.0159%	0.0199%	0.0210%	0.0205%	0.0233%	0.0135%	0.0181%
Hispanic ^d	Both Sexes	0.0231%	0.0108%	0.0248%	0.0217%	0.0212%	0.0281%	0.0355%	0.0332%	0.0261%	0.0239%	0.0243%
	Males	0.0245%	0.0113%	0.0259%	0.0213%	0.0213%	0.0322%	0.0390%	0.0409%	0.0236%	0.0303%	0.0261%
	Females	0.0217%	0.0104%	0.0236%	0.0222%	0.0210%	0.0239%	0.0320%	0.0265%	0.0280%	0.0200%	0.0225%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 4.24
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	894,709	687,678	571,451	381,567	246,862	141,227	2,697,125	3,027,655	41,576	3,069,231
	Males	6,335	3,856	2,472	1,476	706	450	14,650	15,682	99	15,781
	Females	888,374	683,822	568,979	380,091	246,156	140,777	2,682,475	3,011,973	41,477	3,053,450
White ^b	Both Sexes	746,789	592,304	503,572	338,960	221,949	127,864	2,326,391	2,625,985	36,580	2,662,565
	Males	5,157	3,344	2,071	1,305	603	392	12,323	13,236	0 ¹	13,236
	Females	741,632	588,960	501,501	337,655	221,346	127,472	2,314,068	2,612,749	36,580	2,649,329
Black ^b	Both Sexes	96,917	62,687	43,796	28,425	16,751	9,164	243,131	264,386	+	+
	Males	917	413	292	110	97	49	1,792	1,902	+	+
	Females	96,000	62,274	43,504	28,315	16,654	9,115	241,339	262,484	3,261	265,745
Asian/ Pacific Islander ^c	Both Sexes	37,519	25,527	18,619	+	+	+	94,100	+	+	+
	Males	151	83	57	+	+	+	334	+	+	+
	Females	37,368	25,444	18,562	+	+	+	93,766	+	+	+
Hispanic ^d	Both Sexes	65,003	44,545	31,261	+	+	+	162,074	+	+	+
	Males	320	167	90	+	+	+	613	+	+	+
	Females	64,683	44,378	31,171	+	+	+	161,461	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.7517%	-	0.0001%	0.0067%	0.0861%	0.4766%	1.2197%	2.2556%	3.2003%	3.6644%	0.6971%
	Males	0.0081%	-	-	-	0.0005%	0.0028%	0.0087%	0.0255%	0.0496%	0.0683%	0.0086%
	Females	1.4763%	-	0.0003%	0.0135%	0.1720%	0.9462%	2.3807%	4.2695%	5.7673%	5.8130%	1.2814%
White ^c	Both Sexes	0.8239%	-	0.0002%	0.0064%	0.0847%	0.4758%	1.2487%	2.3649%	3.4267%	3.9136%	0.7291%
	Males	0.0088%	-	-	-	0.0004%	0.0025%	0.0078%	0.0266%	0.0512%	0.0740%	0.0088%
	Females	1.6356%	-	0.0004%	0.0132%	0.1734%	0.9651%	2.4779%	4.5509%	6.2495%	6.2367%	1.3575%
Black ^c	Both Sexes	0.5507%	-	-	0.0081%	0.0982%	0.4861%	1.1440%	2.0443%	2.7618%	3.1188%	0.6352%
	Males	0.0082%	-	-	-	-	0.0059%	0.0169%	0.0329%	0.0610%	0.0623%	0.0110%
	Females	1.0449%	-	-	0.0159%	0.1835%	0.9083%	2.1276%	3.6162%	4.6472%	4.6217%	1.0993%
Asian/ Pacific Islander ^c	Both Sexes	0.5778%	-	-	0.0064%	0.0826%	0.4779%	1.1319%	1.7675%	2.2007%	2.3789%	0.5500%
	Males	0.0044%	-	-	-	-	-	0.0065%	0.0127%	0.0304%	0.0323%	0.0049%
	Females	1.1036%	-	-	0.0129%	0.1565%	0.8994%	2.1010%	3.2106%	3.8879%	3.8165%	0.9867%
Hispanic ^d	Both Sexes	0.3141%	-	-	0.0046%	0.0648%	0.3299%	0.8846%	1.6052%	2.1663%	2.3799%	0.4860%
	Males	0.0023%	-	-	-	-	0.0015%	0.0041%	0.0161%	0.0235%	0.0341%	0.0046%
	Females	0.6313%	-	-	0.0095%	0.1334%	0.6715%	1.7320%	2.9752%	3.7545%	3.7924%	0.8851%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 5.15
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>										
All Races ^b	39,026	32,190	33,728	33,192	29,376	21,967	156,248	216,287	32,633	248,920
White ^b	30,672	25,506	27,004	27,137	24,572	18,472	125,354	175,330	27,483	202,813
Black ^b	5,129	4,385	4,428	3,673	3,068	2,439	19,606	26,717	+	+
Asian/Pacific Islander ^c	2,178	1,751	1,666	+	+	+	7,418	+	+	+
Hispanic ^d	8,002	7,064	6,973	+	+	+	28,867	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races ^c	0.0974%	-	0.0002%	0.0078%	0.0686%	0.1842%	0.2094%	0.1923%	0.1657%	0.1080%	0.0908%	
White ^c	0.1018%	-	-	0.0093%	0.0784%	0.2055%	0.2178%	0.1858%	0.1502%	0.0885%	0.0947%	
Black ^c	0.0814%	-	-	0.0037%	0.0540%	0.1451%	0.1867%	0.2037%	0.1884%	0.1565%	0.0830%	
Asian/Pacific Islander ^c	0.0848%	-	-	0.0039%	0.0317%	0.1050%	0.1732%	0.2099%	0.2339%	0.1994%	0.0765%	
Hispanic ^d	0.1125%	-	-	0.0093%	0.0753%	0.2282%	0.3428%	0.3554%	0.3527%	0.2662%	0.1418%	

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

^e (d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry. Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 6.28
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	393,064	281,906	206,320	121,413	79,266	46,430	1,054,865	1,160,780	16,776	1,177,556
	Males	201,444	141,586	103,205	60,694	38,518	21,344	532,958	579,395	5,741	585,136
	Females	191,620	140,320	103,115	60,719	40,748	25,086	521,907	581,385	11,035	592,420
White ^b	Both Sexes	318,724	235,878	175,895	103,853	69,080	40,904	879,946	973,061	14,682	987,743
	Males	164,730	120,728	89,034	52,292	34,129	19,056	449,833	491,231	5,097	496,328
	Females	153,994	115,150	86,861	51,561	34,951	21,848	430,113	481,830	9,585	491,415
Black ^b	Both Sexes	47,070	29,885	19,890	11,665	6,934	3,916	112,879	121,718	1,275	122,993
	Males	22,212	12,561	8,611	5,214	2,882	1,392	50,552	53,636	393	54,029
	Females	24,858	17,324	11,279	6,451	4,052	2,524	62,327	68,082	882	68,964
Asian/ Pacific Islander ^c	Both Sexes	18,561	11,624	7,604	+	+	+	42,739	+	+	+
	Males	9,770	5,985	3,819	+	+	+	22,100	+	+	+
	Females	8,791	5,639	3,785	+	+	+	20,639	+	+	+
Hispanic ^d	Both Sexes	32,255	18,902	10,733	+	+	+	68,895	+	+	+
	Males	17,364	9,764	5,528	+	+	+	36,267	+	+	+
	Females	14,891	9,138	5,205	+	+	+	32,628	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted ^f		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.3002%	-	0.0005%	0.0052%	0.0274%	0.1109%	0.3521%	0.7582%	1.5052%	2.3794%	0.2841%
	Males	0.3084%	-	0.0003%	0.0050%	0.0265%	0.1180%	0.3875%	0.8884%	1.7808%	2.6789%	0.3259%
	Females	0.2922%	-	0.0007%	0.0054%	0.0282%	0.1038%	0.3181%	0.6407%	1.2806%	2.2005%	0.2506%
White ^c	Both Sexes	0.3125%	-	0.0006%	0.0056%	0.0287%	0.1094%	0.3359%	0.7253%	1.4977%	2.4090%	0.2803%
	Males	0.3214%	-	0.0004%	0.0054%	0.0273%	0.1155%	0.3724%	0.8543%	1.7810%	2.7000%	0.3221%
	Females	0.3036%	-	0.0008%	0.0058%	0.0303%	0.1031%	0.2998%	0.6047%	1.2607%	2.2329%	0.2459%
Black ^c	Both Sexes	0.2556%	-	-	0.0042%	0.0220%	0.1200%	0.4171%	0.9561%	1.6767%	2.2572%	0.3138%
	Males	0.2416%	-	-	0.0044%	0.0215%	0.1230%	0.4250%	1.0706%	1.8217%	2.4139%	0.3387%
	Females	0.2683%	-	-	0.0040%	0.0224%	0.1173%	0.4103%	0.8666%	1.5755%	2.1801%	0.2966%
Asian/ Pacific Islander ^c	Both Sexes	0.2754%	-	-	0.0026%	0.0230%	0.1028%	0.3610%	0.7600%	1.3920%	2.2152%	0.2712%
	Males	0.2937%	-	-	0.0027%	0.0239%	0.1187%	0.4128%	0.9069%	1.6873%	2.5942%	0.3215%
	Females	0.2586%	-	-	0.0025%	0.0223%	0.0887%	0.3164%	0.6393%	1.1624%	1.9831%	0.2329%
Hispanic ^d	Both Sexes	0.1326%	-	-	0.0037%	0.0216%	0.0814%	0.2803%	0.6468%	1.1988%	1.8001%	0.2268%
	Males	0.1374%	-	-	0.0032%	0.0214%	0.0860%	0.3031%	0.7643%	1.5176%	2.2005%	0.2716%
	Females	0.1278%	-	-	0.0042%	0.0218%	0.0766%	0.2583%	0.5455%	0.9624%	1.5591%	0.1934%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 7.15
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>										
All Races ^b	190,409	131,077	102,712	73,340	48,841	32,618	529,671	615,772	19,665	635,437
White ^b	161,701	116,009	92,657	66,999	45,395	30,608	467,264	548,026	16,348	564,374
Black ^b	17,049	8,164	5,300	3,088	1,771	1,058	34,703	37,414	672	38,086
Asian/Pacific Islander ^c	7,713	4,660	3,176	+	+	+	17,931	+	+	+
Hispanic ^d	15,050	8,909	5,628	+	+	+	33,874	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.2867%	-	-	0.0034%	0.0314%	0.1217%	0.3988%	0.9484%	1.2559%	1.0819%	0.2471%
White ^c	0.3261%	-	-	0.0033%	0.0314%	0.1211%	0.4156%	1.0303%	1.4062%	1.2172%	0.2681%
Black ^c	0.1507%	-	-	0.0036%	0.0236%	0.0882%	0.2262%	0.6325%	0.8881%	0.5960%	0.1618%
Asian/Pacific Islander ^c	0.2089%	-	-	0.0034%	0.0316%	0.1414%	0.4355%	0.7171%	0.6716%	0.5085%	0.1835%
Hispanic ^d	0.1324%	-	-	0.0047%	0.0418%	0.1318%	0.3325%	0.6877%	0.8177%	0.6328%	0.1841%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

^e (d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry. Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 8.21
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	21,605	8,307	4,234	1,640	581	254	36,292	36,756	101	36,857
	Males	17,134	6,323	3,311	1,332	415	201	28,455	28,826	52	28,878
	Females	4,471	1,984	923	308	166	53	7,837	7,930	49	7,979
White ^b	Both Sexes	19,425	7,539	3,836	1,439	512	224	32,691	33,108	80	33,188
	Males	15,719	5,807	3,095	1,187	368	175	26,127	26,459	47	26,506
	Females	3,706	1,732	741	252	144	49	6,564	6,649	33	6,682
Black ^b	Both Sexes	1,496	556	265	161	50	26	2,515	2,554	23	2,577
	Males	916	354	134	112	37	26	1,540	1,579	6	1,585
	Females	580	202	131	49	13	0	975	975	17	992
Asian/ Pacific Islander ^c	Both Sexes	464	115	88	+	+	+	710	+	+	+
	Males	337	89	54	+	+	+	512	+	+	+
	Females	127	26	34	+	+	+	198	+	+	+
Hispanic ^d	Both Sexes	1,065	302	121	+	+	+	1,584	+	+	+
	Males	874	250	99	+	+	+	1,301	+	+	+
	Females	191	52	22	+	+	+	283	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	Age Specific (Crude)				80+	Age-Adjusted ^f	
						30-39	40-49	50-59	60-69	70-79	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0094%	-	-	-	0.0004%	0.0026%	0.0118%	0.0323%	0.0533%	0.0451%	0.0088%
	Males	0.0148%	-	-	-	0.0005%	0.0044%	0.0190%	0.0559%	0.0914%	0.0824%	0.0151%
	Females	0.0041%	-	-	-	0.0002%	0.0008%	0.0049%	0.0110%	0.0223%	0.0229%	0.0036%
White ^c	Both Sexes	0.0110%	-	-	-	0.0005%	0.0030%	0.0131%	0.0370%	0.0591%	0.0476%	0.0098%
	Males	0.0176%	-	-	-	0.0007%	0.0052%	0.0213%	0.0641%	0.1030%	0.0868%	0.0169%
	Females	0.0045%	-	-	-	0.0002%	0.0008%	0.0051%	0.0117%	0.0224%	0.0239%	0.0037%
Black ^c	Both Sexes	0.0056%	-	-	-	-	0.0022%	0.0103%	0.0226%	0.0399%	0.0333%	0.0066%
	Males	0.0072%	-	-	-	-	0.0034%	0.0137%	0.0329%	0.0558%	0.0625%	0.0098%
	Females	0.0041%	-	-	-	-	-	0.0074%	0.0146%	0.0288%	0.0189%	0.0044%
Asian/ Pacific Islander ^c	Both Sexes	0.0046%	-	-	-	-	0.0012%	0.0060%	0.0120%	0.0290%	0.0334%	0.0046%
	Males	0.0068%	-	-	-	-	0.0014%	0.0102%	0.0215%	0.0449%	0.0585%	0.0076%
	Females	0.0025%	-	-	-	-	0.0011%	0.0024%	0.0042%	0.0166%	0.0180%	0.0024%
Hispanic ^d	Both Sexes	0.0030%	-	-	-	0.0003%	0.0021%	0.0068%	0.0165%	0.0259%	0.0347%	0.0050%
	Males	0.0049%	-	-	-	-	0.0036%	0.0117%	0.0295%	0.0514%	0.0684%	0.0095%
	Females	0.0011%	-	-	-	-	-	0.0021%	0.0053%	0.0070%	0.0143%	0.0016%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 9.21
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	37,358	34,292	28,727	25,325	22,323	17,430	139,536	181,370	12,175	193,545
	Males	20,235	18,129	14,510	12,916	11,125	8,871	72,770	94,047	6,049	100,096
	Females	17,123	16,163	14,217	12,409	11,198	8,559	66,766	87,323	6,126	93,449
White ^b	Both Sexes	30,903	28,953	24,933	22,403	20,422	16,164	119,721	158,840	13,473	172,313
	Males	16,904	15,439	12,599	11,460	10,227	8,058	62,752	82,343	6,989	89,332
	Females	13,999	13,514	12,334	10,943	10,195	8,106	56,969	76,497	6,484	82,981
Black ^b	Both Sexes	4,743	4,127	2,817	2,469	1,501	1,031	15,221	17,329	957	18,286
	Males	2,438	2,028	1,369	1,240	698	695	7,586	8,960	667	9,627
	Females	2,305	2,099	1,448	1,229	803	336	7,635	8,369	290	8,659
Asian/ Pacific Islander ^c	Both Sexes	999	838	567	+	+	+	2,742	+	+	+
	Males	503	434	304	+	+	+	1,382	+	+	+
	Females	496	404	263	+	+	+	1,360	+	+	+
Hispanic ^d	Both Sexes	4,295	3,436	2,580	+	+	+	12,418	+	+	+
	Males	2,290	1,890	1,438	+	+	+	6,757	+	+	+
	Females	2,005	1,546	1,142	+	+	+	5,661	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0377%	0.0006%	0.0104%	0.0393%	0.0655%	0.0655%	0.0469%	0.0369%	0.0338%	0.0251%	0.0376%
	Males	0.0400%	0.0009%	0.0115%	0.0392%	0.0634%	0.0694%	0.0534%	0.0449%	0.0404%	0.0273%	0.0398%
	Females	0.0354%	0.0003%	0.0092%	0.0393%	0.0676%	0.0616%	0.0407%	0.0295%	0.0284%	0.0237%	0.0356%
White ^c	Both Sexes	0.0429%	0.0006%	0.0115%	0.0444%	0.0754%	0.0753%	0.0529%	0.0415%	0.0383%	0.0285%	0.0429%
	Males	0.0454%	0.0008%	0.0127%	0.0440%	0.0721%	0.0791%	0.0594%	0.0508%	0.0451%	0.0310%	0.0450%
	Females	0.0404%	0.0005%	0.0102%	0.0449%	0.0789%	0.0714%	0.0464%	0.0329%	0.0326%	0.0270%	0.0410%
Black ^c	Both Sexes	0.0340%	0.0007%	0.0095%	0.0314%	0.0613%	0.0650%	0.0467%	0.0348%	0.0296%	0.0179%	0.0351%
	Males	0.0352%	0.0013%	0.0099%	0.0294%	0.0606%	0.0681%	0.0571%	0.0365%	0.0383%	0.0218%	0.0371%
	Females	0.0329%	-	0.0092%	0.0333%	0.0620%	0.0623%	0.0376%	0.0336%	0.0236%	0.0160%	0.0335%
Asian/ Pacific Islander ^c	Both Sexes	0.0146%	-	0.0054%	0.0220%	0.0263%	0.0194%	0.0148%	0.0100%	0.0116%	0.0066%	0.0139%
	Males	0.0154%	-	0.0067%	0.0256%	0.0246%	0.0196%	0.0147%	0.0115%	0.0159%	0.0076%	0.0148%
	Females	0.0139%	-	0.0040%	0.0184%	0.0279%	0.0192%	0.0148%	0.0088%	0.0083%	0.0060%	0.0131%
Hispanic ^d	Both Sexes	0.0235%	0.0009%	0.0097%	0.0275%	0.0378%	0.0366%	0.0322%	0.0363%	0.0368%	0.0241%	0.0257%
	Males	0.0253%	0.0010%	0.0125%	0.0288%	0.0355%	0.0395%	0.0399%	0.0408%	0.0411%	0.0350%	0.0282%
	Females	0.0217%	0.0008%	0.0068%	0.0261%	0.0402%	0.0335%	0.0248%	0.0324%	0.0336%	0.0176%	0.0235%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 10.15
Kaposi SarcomaEstimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	7,288	5,797	4,453	6,636	1,913	685	25,480	26,999	+	+
	Males	6,806	5,465	4,217	6,471	1,792	672	24,145	25,587	+	+
	Females	482	332	236	165	121	13	1,335	1,412	+	+
White ^b	Both Sexes	4,389	4,194	3,136	5,357	1,534	597	18,146	19,400	+	+
	Males	4,085	3,950	2,947	5,250	1,435	584	17,203	18,392	+	+
	Females	304	244	189	107	99	13	943	1,008	+	+
Black ^b	Both Sexes	2,382	1,395	1,035	897	191	37	5,842	5,949	+	+
	Males	2,265	1,326	1,000	862	181	37	5,575	5,682	+	+
	Females	117	69	35	35	10	0	267	267	+	+
Asian/ Pacific Islander ^c	Both Sexes	153	89	79	+	+	+	407	+	+	+
	Males	151	86	79	+	+	+	402	+	+	+
	Females	2	3	0	+	+	+	5	+	+	+
Hispanic ^d	Both Sexes	1,181	1,130	914	+	+	+	4,475	+	+	+
	Males	1,040	1,057	846	+	+	+	4,171	+	+	+
	Females	141	73	68	+	+	+	304	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0087%	-	-	0.0012%	0.0060%	0.0180%	0.0221%	0.0126%	0.0090%	0.0156%	0.0083%
	Males	0.0169%	-	-	0.0022%	0.0118%	0.0355%	0.0444%	0.0257%	0.0171%	0.0292%	0.0164%
	Females	0.0008%	-	-	-	-	0.0007%	0.0008%	0.0008%	0.0024%	0.0075%	0.0007%
White ^c	Both Sexes	0.0087%	-	-	0.0005%	0.0048%	0.0169%	0.0225%	0.0132%	0.0100%	0.0166%	0.0080%
	Males	0.0165%	-	-	0.0010%	0.0093%	0.0326%	0.0445%	0.0264%	0.0186%	0.0310%	0.0156%
	Females	0.0008%	-	-	-	-	0.0007%	0.0006%	0.0009%	0.0027%	0.0079%	0.0007%
Black ^c	Both Sexes	0.0136%	-	-	0.0052%	0.0176%	0.0350%	0.0307%	0.0146%	0.0053%	0.0127%	0.0140%
	Males	0.0274%	-	-	0.0100%	0.0371%	0.0740%	0.0634%	0.0322%	0.0106%	0.0207%	0.0291%
	Females	0.0009%	-	-	-	-	-	0.0023%	-	-	0.0087%	0.0010%
Asian/ Pacific Islander ^c	Both Sexes	0.0023%	-	-	-	0.0015%	0.0040%	0.0062%	0.0037%	0.0028%	0.0053%	0.0022%
	Males	0.0048%	-	-	-	0.0032%	0.0083%	0.0135%	0.0082%	0.0063%	0.0125%	0.0047%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic ^d	Both Sexes	0.0086%	-	-	0.0008%	0.0069%	0.0214%	0.0267%	0.0178%	0.0158%	0.0415%	0.0111%
	Males	0.0158%	-	-	0.0016%	0.0132%	0.0409%	0.0530%	0.0369%	0.0262%	0.0587%	0.0207%
	Females	0.0012%	-	-	-	-	0.0011%	0.0015%	-	0.0081%	0.0312%	0.0020%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 11.21
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	158,086	101,181	55,073	32,357	19,160	10,131	359,192	384,082	10,254	394,336
	Males	97,484	59,911	32,420	18,537	10,668	5,615	215,575	228,920	5,786	234,706
	Females	60,602	41,270	22,653	13,820	8,492	4,516	143,617	155,162	4,468	159,630
White ^b	Both Sexes	130,651	85,726	47,204	27,843	17,085	8,965	302,534	324,605	11,448	336,053
	Males	81,177	50,894	28,124	16,410	9,692	5,062	183,103	195,243	5,022	200,265
	Females	49,474	34,832	19,080	11,433	7,393	3,903	119,431	129,362	6,426	135,788
Black ^b	Both Sexes	19,351	10,830	5,803	3,473	1,619	920	40,475	42,789	2,042	44,831
	Males	11,221	6,376	3,035	1,559	747	402	22,727	23,687	709	24,396
	Females	8,130	4,454	2,768	1,914	872	518	17,748	19,102	1,333	20,435
Asian/ Pacific Islander ^c	Both Sexes	4,279	2,367	1,215	+	+	+	8,524	+	+	+
	Males	2,678	1,429	721	+	+	+	5,237	+	+	+
	Females	1,601	938	494	+	+	+	3,287	+	+	+
Hispanic ^d	Both Sexes	15,748	8,692	4,575	+	+	+	31,601	+	+	+
	Males	8,908	4,829	2,573	+	+	+	17,684	+	+	+
	Females	6,840	3,863	2,002	+	+	+	13,917	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0992%	0.0070%	0.0104%	0.0072%	0.0156%	0.0591%	0.1479%	0.2967%	0.4455%	0.4096%	0.0931%
	Males	0.1215%	0.0065%	0.0091%	0.0066%	0.0167%	0.0745%	0.1915%	0.3985%	0.6070%	0.5702%	0.1223%
	Females	0.0774%	0.0077%	0.0117%	0.0078%	0.0145%	0.0439%	0.1061%	0.2047%	0.3139%	0.3136%	0.0686%
White ^c	Both Sexes	0.1094%	0.0072%	0.0111%	0.0079%	0.0165%	0.0628%	0.1557%	0.3120%	0.4750%	0.4291%	0.0983%
	Males	0.1336%	0.0063%	0.0105%	0.0072%	0.0169%	0.0784%	0.1989%	0.4132%	0.6467%	0.6014%	0.1284%
	Females	0.0852%	0.0081%	0.0117%	0.0087%	0.0161%	0.0467%	0.1129%	0.2173%	0.3314%	0.3248%	0.0725%
Black ^c	Both Sexes	0.0884%	0.0090%	0.0126%	0.0093%	0.0165%	0.0626%	0.1684%	0.3401%	0.4605%	0.3917%	0.1005%
	Males	0.1051%	0.0103%	0.0084%	0.0078%	0.0181%	0.0781%	0.2255%	0.4714%	0.6068%	0.4948%	0.1300%
	Females	0.0731%	0.0077%	0.0170%	0.0107%	0.0151%	0.0490%	0.1186%	0.2375%	0.3585%	0.3410%	0.0782%
Asian/ Pacific Islander ^c	Both Sexes	0.0528%	0.0040%	0.0039%	0.0020%	0.0081%	0.0332%	0.0780%	0.1598%	0.2381%	0.2775%	0.0516%
	Males	0.0674%	0.0024%	0.0025%	0.0022%	0.0099%	0.0445%	0.1097%	0.2310%	0.3268%	0.3917%	0.0709%
	Females	0.0394%	0.0056%	0.0054%	0.0018%	0.0065%	0.0231%	0.0507%	0.1012%	0.1691%	0.2075%	0.0365%
Hispanic ^d	Both Sexes	0.0606%	0.0055%	0.0086%	0.0047%	0.0161%	0.0563%	0.1403%	0.2941%	0.4565%	0.4616%	0.0938%
	Males	0.0669%	0.0044%	0.0070%	0.0036%	0.0154%	0.0642%	0.1648%	0.3740%	0.6266%	0.5710%	0.1171%
	Females	0.0542%	0.0067%	0.0104%	0.0059%	0.0169%	0.0480%	0.1168%	0.2252%	0.3304%	0.3957%	0.0756%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

^e (d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry. Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 12.21
Cancer of the Larynx (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	32,033	19,496	14,224	9,251	6,278	3,797	79,147	87,910	1,171	89,081
	Males	25,889	15,426	11,526	7,418	4,999	3,094	63,539	70,533	955	71,488
	Females	6,144	4,070	2,698	1,833	1,279	703	15,608	17,377	216	17,593
White ^b	Both Sexes	26,843	16,515	12,110	8,088	5,454	3,285	67,159	74,794	910	75,704
	Males	21,712	13,117	9,861	6,465	4,412	2,738	54,015	60,217	739	60,956
	Females	5,131	3,398	2,249	1,623	1,042	547	13,144	14,577	171	14,748
Black ^b	Both Sexes	4,140	2,465	1,668	954	673	411	9,688	10,547	80	10,627
	Males	3,271	1,866	1,280	793	495	295	7,568	8,188	69	8,257
	Females	869	599	388	161	178	116	2,120	2,359	11	2,370
Asian/ Pacific Islander ^c	Both Sexes	568	346	251	+	+	+	1,330	+	+	+
	Males	484	298	214	+	+	+	1,143	+	+	+
	Females	84	48	37	+	+	+	187	+	+	+
Hispanic ^d	Both Sexes	1,894	1,093	882	+	+	+	4,362	+	+	+
	Males	1,650	910	718	+	+	+	3,692	+	+	+
	Females	244	183	164	+	+	+	670	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0206%	-	-	0.0003%	0.0009%	0.0051%	0.0259%	0.0683%	0.1205%	0.0998%	0.0193%
	Males	0.0339%	-	-	0.0004%	0.0011%	0.0070%	0.0405%	0.1184%	0.2253%	0.2238%	0.0353%
	Females	0.0075%	-	-	-	0.0008%	0.0032%	0.0120%	0.0230%	0.0351%	0.0257%	0.0066%
White ^c	Both Sexes	0.0229%	-	-	0.0003%	0.0010%	0.0059%	0.0278%	0.0721%	0.1263%	0.1037%	0.0204%
	Males	0.0373%	-	-	0.0005%	0.0012%	0.0078%	0.0429%	0.1229%	0.2331%	0.2290%	0.0366%
	Females	0.0085%	-	-	-	0.0009%	0.0038%	0.0128%	0.0246%	0.0369%	0.0278%	0.0071%
Black ^c	Both Sexes	0.0214%	-	-	-	0.0014%	0.0046%	0.0361%	0.0939%	0.1728%	0.1094%	0.0257%
	Males	0.0352%	-	-	-	0.0021%	0.0068%	0.0550%	0.1698%	0.3448%	0.2863%	0.0500%
	Females	0.0087%	-	-	-	-	0.0027%	0.0196%	0.0347%	0.0528%	0.0224%	0.0091%
Asian/ Pacific Islander ^c	Both Sexes	0.0086%	-	-	-	-	0.0017%	0.0072%	0.0278%	0.0537%	0.0676%	0.0087%
	Males	0.0155%	-	-	-	-	0.0027%	0.0135%	0.0535%	0.1057%	0.1563%	0.0175%
	Females	0.0023%	-	-	-	-	-	0.0017%	0.0067%	0.0133%	0.0133%	0.0022%
Hispanic ^d	Both Sexes	0.0083%	-	-	0.0004%	0.0007%	0.0033%	0.0151%	0.0429%	0.1000%	0.0984%	0.0148%
	Males	0.0140%	-	-	0.0007%	0.0008%	0.0048%	0.0250%	0.0790%	0.2061%	0.2268%	0.0294%
	Females	0.0026%	-	-	-	-	0.0017%	0.0057%	0.0118%	0.0213%	0.0211%	0.0039%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 13.29
LeukemiaEstimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	124,326	80,785	49,405	31,735	19,435	12,358	298,813	328,477	5,498	333,975
	Males	72,087	46,052	27,849	17,875	10,823	6,755	170,703	186,605	2,414	189,019
	Females	52,239	34,733	21,556	13,860	8,612	5,603	128,110	141,872	3,084	144,956
White ^b	Both Sexes	106,763	71,231	44,054	28,247	17,433	11,168	261,586	288,470	4,601	293,071
	Males	62,036	40,744	24,844	15,958	9,861	6,173	149,845	164,453	1,683	166,136
	Females	44,727	30,487	19,210	12,289	7,572	4,995	111,741	124,017	2,918	126,935
Black ^b	Both Sexes	9,870	5,670	2,937	1,913	1,095	603	21,121	22,581	813	23,394
	Males	5,543	3,137	1,528	995	531	291	11,531	12,182	384	12,566
	Females	4,327	2,533	1,409	918	564	312	9,590	10,399	429	10,828
Asian/ Pacific Islander ^c	Both Sexes	3,092	2,081	1,379	+	+	+	7,675	+	+	+
	Males	1,826	1,143	791	+	+	+	4,354	+	+	+
	Females	1,266	938	588	+	+	+	3,321	+	+	+
Hispanic ^d	Both Sexes	11,855	7,822	5,310	+	+	+	29,226	+	+	+
	Males	6,686	4,562	2,816	+	+	+	16,563	+	+	+
	Females	5,169	3,260	2,494	+	+	+	12,663	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted ^f		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0839%	0.0322%	0.0608%	0.0475%	0.0300%	0.0452%	0.0850%	0.1711%	0.2882%	0.3378%	0.0815%
	Males	0.0979%	0.0333%	0.0665%	0.0527%	0.0338%	0.0540%	0.1052%	0.2217%	0.3813%	0.4509%	0.1003%
	Females	0.0702%	0.0311%	0.0550%	0.0420%	0.0262%	0.0366%	0.0656%	0.1254%	0.2123%	0.2702%	0.0656%
White ^c	Both Sexes	0.0961%	0.0361%	0.0692%	0.0542%	0.0333%	0.0480%	0.0927%	0.1895%	0.3239%	0.3766%	0.0906%
	Males	0.1114%	0.0363%	0.0753%	0.0596%	0.0376%	0.0567%	0.1139%	0.2431%	0.4256%	0.5017%	0.1108%
	Females	0.0808%	0.0359%	0.0627%	0.0484%	0.0289%	0.0390%	0.0718%	0.1394%	0.2389%	0.3009%	0.0733%
Black ^c	Both Sexes	0.0470%	0.0166%	0.0273%	0.0287%	0.0211%	0.0383%	0.0586%	0.1165%	0.1826%	0.2128%	0.0523%
	Males	0.0540%	0.0182%	0.0311%	0.0320%	0.0228%	0.0462%	0.0750%	0.1518%	0.2389%	0.2579%	0.0643%
	Females	0.0407%	0.0150%	0.0233%	0.0255%	0.0197%	0.0313%	0.0442%	0.0890%	0.1433%	0.1907%	0.0432%
Asian/ Pacific Islander ^c	Both Sexes	0.0431%	0.0262%	0.0511%	0.0314%	0.0212%	0.0343%	0.0465%	0.0665%	0.1013%	0.1096%	0.0436%
	Males	0.0512%	0.0308%	0.0551%	0.0364%	0.0234%	0.0421%	0.0579%	0.0832%	0.1309%	0.1678%	0.0531%
	Females	0.0356%	0.0214%	0.0469%	0.0264%	0.0191%	0.0273%	0.0368%	0.0527%	0.0783%	0.0740%	0.0357%
Hispanic ^d	Both Sexes	0.0548%	0.0363%	0.0695%	0.0527%	0.0314%	0.0416%	0.0560%	0.0982%	0.1585%	0.1623%	0.0616%
	Males	0.0613%	0.0378%	0.0756%	0.0603%	0.0396%	0.0481%	0.0629%	0.1216%	0.1751%	0.2272%	0.0717%
	Females	0.0482%	0.0347%	0.0632%	0.0445%	0.0227%	0.0349%	0.0494%	0.0780%	0.1463%	0.1233%	0.0528%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 13.30
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	17,063	13,626	11,827	10,122	8,389	6,522	57,873	74,508	3,347	77,855
	Males	9,316	7,652	6,474	5,589	4,617	3,505	31,895	40,643	1,752	42,395
	Females	7,747	5,974	5,353	4,533	3,772	3,017	25,978	33,865	1,595	35,460
White ^b	Both Sexes	14,355	11,393	10,366	8,768	7,380	5,787	49,494	64,481	+	+
	Males	7,757	6,370	5,649	4,848	4,126	3,130	27,213	35,154	+	+
	Females	6,598	5,023	4,717	3,920	3,254	2,657	22,281	29,327	+	+
Black ^b	Both Sexes	1,492	1,151	701	679	508	360	4,346	5,150	+	+
	Males	904	700	363	351	247	187	2,455	2,846	+	+
	Females	588	451	338	328	261	173	1,891	2,304	+	+
Asian/ Pacific Islander ^c	Both Sexes	762	712	544	+	+	+	2,570	+	+	+
	Males	430	384	334	+	+	+	1,451	+	+	+
	Females	332	328	210	+	+	+	1,119	+	+	+
Hispanic ^d	Both Sexes	5,285	3,777	2,870	+	+	+	14,642	+	+	+
	Males	2,989	2,253	1,484	+	+	+	8,356	+	+	+
	Females	2,296	1,524	1,386	+	+	+	6,286	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0186%	0.0272%	0.0515%	0.0343%	0.0094%	0.0055%	0.0043%	0.0046%	0.0041%	0.0019%	0.0192%
	Males	0.0212%	0.0278%	0.0563%	0.0393%	0.0117%	0.0061%	0.0049%	0.0045%	0.0044%	0.0026%	0.0212%
	Females	0.0162%	0.0265%	0.0464%	0.0291%	0.0071%	0.0049%	0.0037%	0.0047%	0.0038%	0.0014%	0.0172%
White ^c	Both Sexes	0.0211%	0.0310%	0.0591%	0.0404%	0.0117%	0.0059%	0.0047%	0.0049%	0.0040%	0.0017%	0.0222%
	Males	0.0237%	0.0311%	0.0642%	0.0459%	0.0146%	0.0067%	0.0053%	0.0047%	0.0048%	0.0019%	0.0243%
	Females	0.0185%	0.0310%	0.0539%	0.0346%	0.0088%	0.0051%	0.0042%	0.0051%	0.0033%	0.0016%	0.0200%
Black ^c	Both Sexes	0.0096%	0.0120%	0.0220%	0.0177%	0.0031%	0.0033%	0.0018%	0.0041%	-	-	0.0089%
	Males	0.0113%	0.0134%	0.0258%	0.0202%	0.0037%	0.0031%	0.0021%	0.0035%	-	-	0.0100%
	Females	0.0080%	0.0104%	0.0181%	0.0152%	0.0025%	0.0036%	0.0015%	0.0046%	-	-	0.0077%
Asian/ Pacific Islander ^c	Both Sexes	0.0131%	0.0212%	0.0411%	0.0207%	0.0044%	0.0049%	0.0037%	0.0028%	0.0045%	0.0026%	0.0142%
	Males	0.0155%	0.0236%	0.0451%	0.0253%	0.0049%	0.0054%	0.0050%	0.0033%	-	0.0069%	0.0160%
	Females	0.0108%	0.0186%	0.0370%	0.0161%	0.0039%	0.0045%	0.0027%	0.0024%	0.0073%	-	0.0125%
Hispanic ^d	Both Sexes	0.0270%	0.0317%	0.0604%	0.0401%	0.0124%	0.0077%	0.0063%	0.0070%	0.0072%	-	0.0233%
	Males	0.0305%	0.0328%	0.0652%	0.0470%	0.0163%	0.0092%	0.0077%	0.0050%	-	-	0.0258%
	Females	0.0235%	0.0305%	0.0554%	0.0325%	0.0084%	0.0062%	0.0049%	0.0087%	0.0091%	-	0.0205%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 14.21
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	38,940	9,558	3,239	1,214	716	527	53,480	54,464	490	54,954
	Males	28,968	6,762	2,047	722	410	213	38,824	39,263	71	39,334
	Females	9,972	2,796	1,192	492	306	314	14,656	15,201	419	15,620
White ^b	Both Sexes	28,607	6,931	2,307	860	559	438	39,117	39,907	401	40,308
	Males	21,328	5,056	1,433	495	304	159	28,559	28,887	264 ⁱ	29,151
	Females	7,279	1,875	874	365	255	279	10,558	11,020	137 ⁱ	11,157
Black ^b	Both Sexes	5,786	983	231	94	71	23	7,143	7,237	+	+
	Males	4,360	559	132	47	35	13	5,122	5,171	+	+
	Females	1,426	424	99	47	36	10	2,021	2,066	+	+
Asian/ Pacific Islander ^c	Both Sexes	3,850	1,443	626	+	+	+	6,121	+	+	+
	Males	2,716	1,039	422	+	+	+	4,323	+	+	+
	Females	1,134	404	204	+	+	+	1,798	+	+	+
Hispanic ^d	Both Sexes	6,209	1,333	497	+	+	+	8,291	+	+	+
	Males	4,545	948	284	+	+	+	5,953	+	+	+
	Females	1,664	385	213	+	+	+	2,338	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0191%	0.0029%	0.0027%	0.0013%	0.0019%	0.0069%	0.0358%	0.0731%	0.0703%	0.0447%	0.0171%
	Males	0.0278%	0.0033%	0.0026%	0.0015%	0.0023%	0.0096%	0.0571%	0.1173%	0.1004%	0.0729%	0.0259%
	Females	0.0107%	0.0024%	0.0028%	0.0011%	0.0015%	0.0042%	0.0154%	0.0331%	0.0457%	0.0279%	0.0095%
White ^c	Both Sexes	0.0161%	0.0032%	0.0028%	0.0015%	0.0014%	0.0054%	0.0309%	0.0609%	0.0488%	0.0283%	0.0137%
	Males	0.0234%	0.0036%	0.0027%	0.0016%	0.0013%	0.0072%	0.0491%	0.0977%	0.0702%	0.0462%	0.0204%
	Females	0.0087%	0.0029%	0.0030%	0.0013%	0.0014%	0.0035%	0.0128%	0.0265%	0.0309%	0.0175%	0.0075%
Black ^c	Both Sexes	0.0158%	0.0009%	0.0007%	0.0014%	0.0024%	0.0046%	0.0338%	0.0959%	0.0554%	0.0313%	0.0163%
	Males	0.0232%	-	-	0.0017%	0.0036%	0.0057%	0.0513%	0.1628%	0.0820%	0.0560%	0.0258%
	Females	0.0091%	-	-	-	-	0.0036%	0.0185%	0.0437%	0.0368%	0.0191%	0.0090%
Asian/ Pacific Islander ^c	Both Sexes	0.0377%	0.0027%	0.0037%	0.0007%	0.0039%	0.0151%	0.0630%	0.1235%	0.1978%	0.1499%	0.0366%
	Males	0.0552%	0.0037%	0.0036%	-	0.0059%	0.0240%	0.1079%	0.2002%	0.2856%	0.2405%	0.0570%
	Females	0.0217%	0.0016%	0.0039%	-	0.0021%	0.0072%	0.0244%	0.0605%	0.1296%	0.0945%	0.0203%
Hispanic ^d	Both Sexes	0.0159%	0.0038%	0.0026%	0.0012%	0.0010%	0.0073%	0.0508%	0.0980%	0.0935%	0.0683%	0.0229%
	Males	0.0225%	0.0046%	0.0027%	0.0012%	0.0009%	0.0116%	0.0852%	0.1560%	0.1269%	0.1039%	0.0349%
	Females	0.0091%	0.0030%	0.0025%	0.0011%	0.0011%	0.0027%	0.0177%	0.0481%	0.0688%	0.0468%	0.0127%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 15.27
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	235,891	83,695	42,860	23,477	12,767	6,435	394,265	409,799	5,908	415,707
	Males	107,952	35,821	18,093	10,288	5,514	2,756	175,769	182,649	3,381	186,030
	Females	127,939	47,874	24,767	13,189	7,253	3,679	218,496	227,150	2,527	229,677
White ^b	Both Sexes	196,844	72,476	37,590	20,395	11,351	5,731	334,693	348,422	2,220	350,642
	Males	89,694	30,463	15,697	8,669	4,861	2,435	147,734	153,689	1,017	154,706
	Females	107,150	42,013	21,893	11,726	6,490	3,296	186,959	194,733	1,203	195,936
Black ^b	Both Sexes	28,016	7,976	3,825	2,212	1,075	525	42,780	44,120	207	44,327
	Males	12,881	3,775	1,693	1,103	502	224	19,767	20,453	101	20,554
	Females	15,135	4,201	2,132	1,109	573	301	23,013	23,667	106	23,773
Asian/ Pacific Islander ^c	Both Sexes	8,262	2,441	1,070	+	+	+	12,423	+	+	+
	Males	4,052	1,216	498	+	+	+	6,131	+	+	+
	Females	4,210	1,225	572	+	+	+	6,292	+	+	+
Hispanic ^d	Both Sexes	9,250	2,945	1,508	+	+	+	14,573	+	+	+
	Males	4,177	1,333	636	+	+	+	6,555	+	+	+
	Females	5,073	1,612	872	+	+	+	8,018	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1047%	0.0002%	0.0004%	0.0019%	0.0054%	0.0229%	0.1066%	0.3138%	0.6574%	0.6428%	0.1003%
	Males	0.0952%	0.0002%	0.0003%	0.0019%	0.0047%	0.0202%	0.0965%	0.3166%	0.6615%	0.7013%	0.1009%
	Females	0.1140%	-	0.0004%	0.0019%	0.0062%	0.0257%	0.1164%	0.3113%	0.6541%	0.6078%	0.1005%
White ^c	Both Sexes	0.1126%	0.0001%	0.0005%	0.0021%	0.0057%	0.0228%	0.1043%	0.3159%	0.6832%	0.6588%	0.1023%
	Males	0.0996%	-	0.0004%	0.0020%	0.0046%	0.0200%	0.0924%	0.3075%	0.6664%	0.7001%	0.0999%
	Females	0.1255%	-	0.0006%	0.0021%	0.0069%	0.0258%	0.1162%	0.3238%	0.6973%	0.6339%	0.1050%
Black ^c	Both Sexes	0.0930%	-	-	0.0021%	0.0056%	0.0278%	0.1567%	0.4080%	0.6850%	0.5465%	0.1120%
	Males	0.0896%	-	-	0.0023%	0.0060%	0.0261%	0.1501%	0.4680%	0.7491%	0.6264%	0.1221%
	Females	0.0962%	-	-	0.0020%	0.0052%	0.0293%	0.1625%	0.3612%	0.6402%	0.5073%	0.1054%
Asian/ Pacific Islander ^c	Both Sexes	0.0802%	-	-	0.0010%	0.0041%	0.0202%	0.0814%	0.2372%	0.5070%	0.5967%	0.0806%
	Males	0.0823%	-	-	0.0012%	0.0040%	0.0179%	0.0813%	0.2702%	0.5869%	0.7415%	0.0923%
	Females	0.0783%	-	-	-	0.0043%	0.0221%	0.0815%	0.2101%	0.4449%	0.5080%	0.0723%
Hispanic ^d	Both Sexes	0.0280%	-	-	0.0016%	0.0031%	0.0122%	0.0478%	0.1410%	0.3292%	0.3561%	0.0500%
	Males	0.0248%	-	-	0.0014%	0.0028%	0.0108%	0.0435%	0.1428%	0.3387%	0.4009%	0.0516%
	Females	0.0313%	-	-	0.0017%	0.0035%	0.0136%	0.0519%	0.1395%	0.3221%	0.3291%	0.0492%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 16.21
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	293,419	223,601	167,410	118,534	84,082	60,557	857,129	1,001,852	32,608	1,034,460
	Males	158,125	114,854	83,196	57,801	38,344	25,391	439,591	498,832	10,262	509,094
	Females	135,294	108,747	84,214	60,733	45,738	35,166	417,538	503,020	22,346	525,366
White ^b	Both Sexes	280,033	216,671	161,996	113,664	81,097	57,703	824,652	963,585	30,683	994,268
	Males	151,569	111,682	80,554	55,443	37,276	24,174	424,105	481,074	9,757	490,831
	Females	128,464	104,989	81,442	58,221	43,821	33,529	400,547	482,511	20,926	503,437
Black ^b	Both Sexes	981	730	611	418	327	143	2,929	3,368	89	3,457
	Males	436	226	297	242	117	12	1,284	1,392	35	1,427
	Females	545	504	314	176	210	131	1,645	1,976	54	2,030
Asian/ Pacific Islander ^c	Both Sexes	780	549	415	+	+	+	2,039	+	+	+
	Males	392	246	144	+	+	+	914	+	+	+
	Females	388	303	271	+	+	+	1,125	+	+	+
Hispanic ^d	Both Sexes	5,309	3,915	3,107	+	+	+	14,650	+	+	+
	Males	2,185	1,408	1,062	+	+	+	5,442	+	+	+
	Females	3,124	2,507	2,045	+	+	+	9,208	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.2125%	0.0004%	0.0031%	0.0267%	0.0977%	0.2066%	0.3602%	0.5505%	0.7222%	0.7923%	0.2000%
	Males	0.2236%	0.0002%	0.0030%	0.0175%	0.0683%	0.1655%	0.3552%	0.6679%	1.0015%	1.2134%	0.2268%
	Females	0.2017%	0.0006%	0.0032%	0.0363%	0.1272%	0.2473%	0.3650%	0.4444%	0.4946%	0.5408%	0.1828%
White ^c	Both Sexes	0.2774%	0.0004%	0.0039%	0.0352%	0.1298%	0.2685%	0.4567%	0.6876%	0.8979%	0.9545%	0.2521%
	Males	0.2895%	0.0002%	0.0039%	0.0230%	0.0885%	0.2095%	0.4439%	0.8206%	1.2298%	1.4571%	0.2799%
	Females	0.2652%	0.0005%	0.0039%	0.0483%	0.1734%	0.3294%	0.4693%	0.5633%	0.6203%	0.6505%	0.2354%
Black ^c	Both Sexes	0.0064%	-	-	-	0.0031%	0.0075%	0.0091%	0.0173%	0.0333%	0.0500%	0.0077%
	Males	0.0062%	-	-	-	0.0021%	0.0052%	0.0092%	0.0212%	0.0442%	0.0795%	0.0091%
	Females	0.0065%	-	-	-	0.0041%	0.0095%	0.0090%	0.0142%	0.0256%	0.0355%	0.0070%
Asian/ Pacific Islander ^c	Both Sexes	0.0122%	0.0008%	0.0010%	0.0025%	0.0063%	0.0142%	0.0186%	0.0272%	0.0403%	0.0496%	0.0118%
	Males	0.0115%	-	-	0.0022%	0.0055%	0.0114%	0.0157%	0.0320%	0.0512%	0.0567%	0.0120%
	Females	0.0128%	-	-	0.0027%	0.0071%	0.0167%	0.0211%	0.0234%	0.0318%	0.0453%	0.0118%
Hispanic ^d	Both Sexes	0.0282%	0.0004%	0.0013%	0.0043%	0.0172%	0.0421%	0.0715%	0.1112%	0.1381%	0.1840%	0.0404%
	Males	0.0206%	-	0.0011%	0.0024%	0.0102%	0.0256%	0.0490%	0.0959%	0.1506%	0.2281%	0.0352%
	Females	0.0359%	0.0006%	0.0014%	0.0063%	0.0247%	0.0593%	0.0931%	0.1243%	0.1288%	0.1575%	0.0464%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 17.13
MesotheliomaEstimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	3,073	635	368	190	202	132	4,394	4,738	+	+
	Males	2,172	354	157	24	52	48	2,744	2,854	+	+
	Females	901	281	211	166	150	84	1,650	1,884	+	+
White ^b	Both Sexes	2,928	589	286	166	186	120	4,084	4,389	+	+
	Males	2,072	331	122	24	48	36	2,585	2,667	+	+
	Females	856	258	164	142	138	84	1,499	1,722	+	+
Black ^b	Both Sexes	82	26	58	13	12	12	192	228	+	+
	Males	47	15	35	0	0	12	97	122	+	+
	Females	35	11	23	13	12	0	95	106	+	+
Asian/ Pacific Islander ^c	Both Sexes	38	19	5	+	+	+	69	+	+	+
	Males	24	8	2	+	+	+	41	+	+	+
	Females	14	11	3	+	+	+	28	+	+	+
Hispanic ^d	Both Sexes	223	64	52	+	+	+	339	+	+	+
	Males	142	31	41	+	+	+	214	+	+	+
	Females	81	33	11	+	+	+	125	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	Age Specific (Crude)					Age-Adjusted ^f	
						30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0012%	-	-	-	0.0003%	0.0006%	0.0013%	0.0037%	0.0056%	0.0071%	0.0011%
	Males	0.0015%	-	-	-	0.0002%	0.0005%	0.0014%	0.0053%	0.0083%	0.0139%	0.0016%
	Females	0.0009%	-	-	-	0.0004%	0.0006%	0.0012%	0.0023%	0.0033%	0.0030%	0.0008%
White ^c	Both Sexes	0.0015%	-	-	-	0.0003%	0.0007%	0.0014%	0.0044%	0.0065%	0.0080%	0.0013%
	Males	0.0019%	-	-	-	0.0003%	0.0007%	0.0015%	0.0060%	0.0099%	0.0152%	0.0019%
	Females	0.0010%	-	-	-	0.0004%	0.0008%	0.0014%	0.0028%	0.0037%	0.0037%	0.0009%
Black ^c	Both Sexes	0.0005%	-	-	-	-	-	0.0010%	0.0017%	0.0038%	-	0.0006%
	Males	0.0006%	-	-	-	-	-	-	0.0033%	-	-	0.0008%
	Females	0.0004%	-	-	-	-	-	-	-	-	-	0.0005%
Asian/ Pacific Islander ^c	Both Sexes	0.0004%	-	-	-	-	-	0.0007%	0.0013%	-	0.0026%	0.0004%
	Males	0.0006%	-	-	-	-	-	-	0.0025%	-	0.0069%	0.0006%
	Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0003%
Hispanic ^d	Both Sexes	0.0007%	-	-	-	-	0.0006%	0.0017%	0.0038%	0.0024%	0.0077%	0.0010%
	Males	0.0008%	-	-	-	-	-	0.0020%	0.0057%	0.0047%	0.0150%	0.0015%
	Females	0.0005%	-	-	-	-	-	0.0014%	0.0021%	-	-	0.0006%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 18.21
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	60,408	21,641	8,022	3,226	1,463	481	94,421	95,457	231	95,688	
	Males	32,121	11,898	4,491	1,669	841	270	50,817	51,385	80	51,465	
	Females	28,287	9,743	3,531	1,557	622	211	43,604	44,072	151	44,223	
White ^b	Both Sexes	44,457	16,678	6,179	2,568	1,076	397	70,751	71,492	0	71,492	
	Males	24,387	9,669	3,654	1,336	624	216	39,527	39,955	0 ⁱ	39,955	
	Females	20,070	7,009	2,525	1,232	452	181	31,224	31,537	0 ⁱ	31,537	
Black ^b	Both Sexes	12,858	4,251	1,497	531	308	63	19,335	19,581	+	+	
	Males	6,057	1,911	648	253	185	49	8,993	9,127	+	+	
	Females	6,801	2,340	849	278	123	14	10,342	10,454	+	+	
Asian/ Pacific Islander ^c	Both Sexes	1,615	445	222	+	+	+	2,384	+	+	+	
	Males	887	221	109	+	+	+	1,265	+	+	+	
	Females	728	224	113	+	+	+	1,119	+	+	+	
Hispanic ^d	Both Sexes	4,600	1,510	631	+	+	+	6,994	+	+	+	
	Males	2,613	842	332	+	+	+	3,944	+	+	+	
	Females	1,987	668	299	+	+	+	3,050	+	+	+	

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races ^c	Both Sexes	0.0264%	-	-	0.0004%	0.0022%	0.0122%	0.0369%	0.0860%	0.1393%	0.1162%	0.0248%
	Males	0.0290%	-	-	0.0006%	0.0027%	0.0150%	0.0417%	0.1013%	0.1645%	0.1476%	0.0295%
	Females	0.0239%	-	-	-	0.0017%	0.0094%	0.0323%	0.0721%	0.1187%	0.0974%	0.0209%
White ^c	Both Sexes	0.0259%	-	-	0.0004%	0.0021%	0.0109%	0.0335%	0.0801%	0.1316%	0.1104%	0.0231%
	Males	0.0293%	-	-	0.0006%	0.0027%	0.0135%	0.0392%	0.0972%	0.1587%	0.1476%	0.0283%
	Females	0.0224%	-	-	-	0.0016%	0.0082%	0.0278%	0.0641%	0.1089%	0.0878%	0.0187%
Black ^c	Both Sexes	0.0429%	-	-	-	0.0036%	0.0282%	0.0782%	0.1774%	0.2730%	0.2301%	0.0506%
	Males	0.0417%	-	-	-	0.0043%	0.0336%	0.0757%	0.1948%	0.3122%	0.2370%	0.0549%
	Females	0.0439%	-	-	-	0.0030%	0.0234%	0.0804%	0.1638%	0.2457%	0.2267%	0.0474%
Asian/ Pacific Islander ^c	Both Sexes	0.0150%	-	-	-	0.0014%	0.0056%	0.0204%	0.0488%	0.0876%	0.0753%	0.0146%
	Males	0.0164%	-	-	-	0.0018%	0.0075%	0.0260%	0.0559%	0.1001%	0.0903%	0.0174%
	Females	0.0137%	-	-	-	0.0010%	0.0039%	0.0156%	0.0431%	0.0778%	0.0661%	0.0125%
Hispanic ^d	Both Sexes	0.0134%	-	-	0.0004%	0.0028%	0.0101%	0.0329%	0.0692%	0.1316%	0.0974%	0.0218%
	Males	0.0149%	-	-	0.0007%	0.0036%	0.0132%	0.0397%	0.0854%	0.1498%	0.1256%	0.0265%
	Females	0.0119%	-	-	-	0.0019%	0.0069%	0.0263%	0.0552%	0.1181%	0.0804%	0.0180%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 19.25
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	208,024	147,198	93,446	53,791	29,696	17,350	522,304	561,810	7,726	569,536	
	Males	112,213	75,991	48,589	28,402	15,226	9,437	275,343	296,343	4,074	300,417	
	Females	95,811	71,207	44,857	25,389	14,470	7,913	246,961	265,467	3,652	269,119	
White ^b	Both Sexes	177,772	129,114	82,376	47,190	26,689	15,706	454,266	490,081	8,097	498,178	
	Males	96,367	67,062	42,958	24,852	13,724	8,505	240,356	259,429	4,601	264,030	
	Females	81,405	62,052	39,418	22,338	12,965	7,201	213,910	230,652	3,496	234,148	
Black ^b	Both Sexes	17,916	11,753	7,007	4,000	1,905	1,053	41,985	44,379	657	45,036	
	Males	9,210	5,685	3,495	2,144	951	608	21,173	22,442	275	22,717	
	Females	8,706	6,068	3,512	1,856	954	445	20,812	21,937	382	22,319	
Asian/ Pacific Islander ^c	Both Sexes	6,610	4,003	2,560	+	+	+	14,811	+	+	+	
	Males	3,504	2,073	1,331	+	+	+	7,733	+	+	+	
	Females	3,106	1,930	1,229	+	+	+	7,078	+	+	+	
Hispanic ^d	Both Sexes	17,016	10,193	6,238	+	+	+	37,304	+	+	+	
	Males	8,761	5,202	3,274	+	+	+	19,291	+	+	+	
	Females	8,255	4,991	2,964	+	+	+	18,013	+	+	+	

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races ^c	Both Sexes	0.1453%	0.0032%	0.0122%	0.0261%	0.0470%	0.1000%	0.2003%	0.3794%	0.6058%	0.7099%	0.1377%
	Males	0.1555%	0.0042%	0.0166%	0.0316%	0.0550%	0.1164%	0.2326%	0.4283%	0.6806%	0.8179%	0.1581%
	Females	0.1354%	0.0021%	0.0077%	0.0203%	0.0390%	0.0837%	0.1694%	0.3353%	0.5448%	0.6453%	0.1199%
White ^c	Both Sexes	0.1630%	0.0031%	0.0128%	0.0274%	0.0484%	0.1067%	0.2145%	0.4077%	0.6658%	0.7634%	0.1481%
	Males	0.1736%	0.0043%	0.0178%	0.0333%	0.0561%	0.1247%	0.2487%	0.4554%	0.7442%	0.8736%	0.1693%
	Females	0.1524%	0.0020%	0.0077%	0.0211%	0.0402%	0.0881%	0.1806%	0.3632%	0.6002%	0.6968%	0.1290%
Black ^c	Both Sexes	0.0927%	0.0036%	0.0112%	0.0258%	0.0541%	0.0949%	0.1718%	0.2952%	0.3768%	0.3583%	0.1037%
	Males	0.0985%	0.0034%	0.0139%	0.0327%	0.0678%	0.1074%	0.1989%	0.3454%	0.3815%	0.3716%	0.1161%
	Females	0.0875%	0.0038%	0.0083%	0.0191%	0.0421%	0.0839%	0.1482%	0.2560%	0.3735%	0.3517%	0.0932%
Asian/ Pacific Islander ^c	Both Sexes	0.0908%	0.0032%	0.0106%	0.0191%	0.0317%	0.0609%	0.1261%	0.2405%	0.3636%	0.5004%	0.0885%
	Males	0.0983%	0.0046%	0.0132%	0.0216%	0.0355%	0.0679%	0.1475%	0.2748%	0.4244%	0.6218%	0.1037%
	Females	0.0841%	0.0017%	0.0078%	0.0165%	0.0283%	0.0547%	0.1077%	0.2123%	0.3164%	0.4260%	0.0765%
Hispanic ^d	Both Sexes	0.0716%	0.0030%	0.0101%	0.0207%	0.0327%	0.0790%	0.1516%	0.2929%	0.4728%	0.5580%	0.1070%
	Males	0.0729%	0.0034%	0.0131%	0.0248%	0.0385%	0.0890%	0.1702%	0.3041%	0.4833%	0.6015%	0.1155%
	Females	0.0702%	0.0025%	0.0069%	0.0163%	0.0266%	0.0685%	0.1338%	0.2832%	0.4651%	0.5319%	0.0993%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 20.27
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	111,858	68,930	43,767	28,840	17,144	11,802	264,846	292,631	8,051	300,682	
	Males	78,943	47,378	28,716	18,000	10,233	6,795	180,085	195,801	4,479	200,280	
	Females	32,915	21,552	15,051	10,840	6,911	5,007	84,761	96,830	3,572	100,402	
White ^b	Both Sexes	96,634	59,621	38,015	25,179	14,923	10,459	229,456	253,694	6,920	260,614	
	Males	68,960	41,709	25,365	15,910	9,147	6,142	158,279	172,373	3,858	176,231	
	Females	27,674	17,912	12,650	9,269	5,776	4,317	71,177	81,321	3,062	84,383	
Black ^b	Both Sexes	8,360	4,855	3,182	1,846	1,189	819	19,035	21,048	382	21,430	
	Males	5,684	2,951	1,859	971	555	366	11,815	12,610	153	12,763	
	Females	2,676	1,904	1,323	875	634	453	7,220	8,438	229	8,667	
Asian/ Pacific Islander ^c	Both Sexes	4,170	2,698	1,776	+	+	+	10,054	+	+	+	
	Males	2,666	1,638	1,037	+	+	+	6,169	+	+	+	
	Females	1,504	1,060	739	+	+	+	3,885	+	+	+	
Hispanic ^d	Both Sexes	6,441	3,466	2,136	+	+	+	13,569	+	+	+	
	Males	4,283	2,047	1,336	+	+	+	8,447	+	+	+	
	Females	2,158	1,419	800	+	+	+	5,122	+	+	+	

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races ^c	Both Sexes	0.0735%	0.0004%	0.0025%	0.0065%	0.0161%	0.0476%	0.1335%	0.2388%	0.2701%	0.2447%	0.0672%
	Males	0.1005%	0.0003%	0.0023%	0.0063%	0.0155%	0.0594%	0.1969%	0.3680%	0.4081%	0.3542%	0.0969%
	Females	0.0471%	0.0004%	0.0027%	0.0066%	0.0167%	0.0359%	0.0728%	0.1222%	0.1576%	0.1793%	0.0416%
White ^c	Both Sexes	0.0805%	0.0004%	0.0024%	0.0064%	0.0149%	0.0465%	0.1396%	0.2572%	0.2914%	0.2586%	0.0705%
	Males	0.1114%	0.0003%	0.0021%	0.0062%	0.0136%	0.0596%	0.2084%	0.3962%	0.4393%	0.3750%	0.1023%
	Females	0.0498%	0.0005%	0.0028%	0.0066%	0.0162%	0.0329%	0.0714%	0.1273%	0.1678%	0.1881%	0.0422%
Black ^c	Both Sexes	0.0435%	-	0.0033%	0.0058%	0.0143%	0.0342%	0.1033%	0.1709%	0.1785%	0.1197%	0.0472%
	Males	0.0571%	-	0.0033%	0.0071%	0.0139%	0.0365%	0.1447%	0.2698%	0.2809%	0.1798%	0.0673%
	Females	0.0311%	-	0.0034%	0.0045%	0.0147%	0.0321%	0.0671%	0.0936%	0.1071%	0.0901%	0.0320%
Asian/ Pacific Islander ^c	Both Sexes	0.0603%	-	0.0020%	0.0072%	0.0225%	0.0601%	0.1159%	0.1665%	0.1979%	0.2016%	0.0572%
	Males	0.0771%	-	0.0022%	0.0065%	0.0259%	0.0724%	0.1640%	0.2408%	0.2845%	0.2754%	0.0778%
	Females	0.0448%	-	0.0017%	0.0080%	0.0194%	0.0493%	0.0744%	0.1054%	0.1307%	0.1564%	0.0406%
Hispanic ^d	Both Sexes	0.0260%	0.0004%	0.0015%	0.0048%	0.0096%	0.0253%	0.0683%	0.1303%	0.1623%	0.1704%	0.0388%
	Males	0.0320%	-	0.0014%	0.0045%	0.0090%	0.0301%	0.0934%	0.1852%	0.2467%	0.2267%	0.0532%
	Females	0.0200%	-	0.0016%	0.0051%	0.0101%	0.0203%	0.0442%	0.0830%	0.0998%	0.1365%	0.0271%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 21.15
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>										
All Races ^b	61,283	32,091	24,676	19,688	16,709	13,475	148,318	181,920	13,847	195,767
White ^b	52,691	27,607	21,380	17,388	15,009	12,143	128,618	158,579	11,701	170,280
Black ^b	4,766	2,346	1,667	1,225	1,052	836	10,618	13,009	1,431	14,440
Asian/Pacific Islander ^c	2,784	1,688	1,236	+	+	+	6,622	+	+	+
Hispanic ^d	6,393	2,771	2,133	+	+	+	12,965	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races ^c	0.0851%	0.0004%	0.0061%	0.0194%	0.0339%	0.0733%	0.1512%	0.2307%	0.2586%	0.2080%	0.0756%	
White ^c	0.0940%	0.0004%	0.0061%	0.0206%	0.0354%	0.0744%	0.1614%	0.2501%	0.2837%	0.2236%	0.0806%	
Black ^c	0.0470%	-	0.0051%	0.0142%	0.0265%	0.0551%	0.0867%	0.1326%	0.1606%	0.1414%	0.0486%	
Asian/Pacific Islander ^c	0.0747%	-	0.0067%	0.0184%	0.0339%	0.0803%	0.1499%	0.1922%	0.1837%	0.1445%	0.0671%	
Hispanic ^d	0.0504%	-	0.0084%	0.0226%	0.0338%	0.0610%	0.1188%	0.1896%	0.2121%	0.1798%	0.0644%	

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 22.21
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	36,831	6,610	2,782	1,466	856	442	48,327	49,363	257	49,620	
	Males	18,821	2,927	1,360	497	399	108	23,890	24,251	112	24,363	
	Females	18,010	3,683	1,422	969	457	334	24,437	25,112	145	25,257	
White ^b	Both Sexes	30,057	5,611	2,283	1,123	617	320	39,581	40,320	208	40,528	
	Males	15,874	2,509	1,228	399	313	66	20,246	20,507	101	20,608	
	Females	14,183	3,102	1,055	724	304	254	19,335	19,813	107	19,920	
Black ^b	Both Sexes	4,965	651	322	170	155	89	6,176	6,408	123	6,531	
	Males	2,104	292	71	36	39	38	2,503	2,591	44	2,635	
	Females	2,861	359	251	134	116	51	3,673	3,817	79	3,896	
Asian/ Pacific Islander ^c	Both Sexes	1,429	258	137	+	+	+	1,893	+	+	+	
	Males	640	112	46	+	+	+	822	+	+	+	
	Females	789	146	91	+	+	+	1,071	+	+	+	
Hispanic ^d	Both Sexes	2,668	524	199	+	+	+	3,576	+	+	+	
	Males	1,335	169	90	+	+	+	1,628	+	+	+	
	Females	1,333	355	109	+	+	+	1,948	+	+	+	

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races ^c	Both Sexes	0.0139%	-	0.0001%	0.0009%	0.0021%	0.0072%	0.0205%	0.0429%	0.0648%	0.0657%	0.0130%
	Males	0.0138%	-	-	0.0004%	0.0019%	0.0072%	0.0219%	0.0468%	0.0736%	0.0672%	0.0139%
	Females	0.0140%	-	-	0.0013%	0.0024%	0.0072%	0.0192%	0.0394%	0.0576%	0.0647%	0.0123%
White ^c	Both Sexes	0.0145%	-	0.0001%	0.0008%	0.0021%	0.0069%	0.0199%	0.0430%	0.0659%	0.0638%	0.0129%
	Males	0.0148%	-	-	0.0005%	0.0018%	0.0070%	0.0215%	0.0482%	0.0774%	0.0684%	0.0142%
	Females	0.0141%	-	-	0.0011%	0.0024%	0.0068%	0.0184%	0.0382%	0.0563%	0.0611%	0.0118%
Black ^c	Both Sexes	0.0137%	-	-	0.0013%	0.0025%	0.0098%	0.0281%	0.0551%	0.0688%	0.0702%	0.0157%
	Males	0.0119%	-	-	-	0.0022%	0.0110%	0.0289%	0.0532%	0.0612%	0.0533%	0.0144%
	Females	0.0153%	-	-	0.0020%	0.0028%	0.0088%	0.0275%	0.0566%	0.0741%	0.0786%	0.0164%
Asian/ Pacific Islander ^c	Both Sexes	0.0119%	-	-	0.0008%	0.0021%	0.0066%	0.0179%	0.0338%	0.0550%	0.0721%	0.0116%
	Males	0.0106%	-	-	-	0.0023%	0.0053%	0.0184%	0.0340%	0.0585%	0.0620%	0.0112%
	Females	0.0130%	-	-	0.0014%	0.0019%	0.0078%	0.0175%	0.0336%	0.0523%	0.0783%	0.0118%
Hispanic ^d	Both Sexes	0.0069%	-	-	0.0008%	0.0016%	0.0058%	0.0169%	0.0357%	0.0500%	0.0628%	0.0109%
	Males	0.0062%	-	-	-	0.0016%	0.0046%	0.0154%	0.0374%	0.0653%	0.0483%	0.0110%
	Females	0.0076%	-	-	0.0012%	0.0016%	0.0071%	0.0183%	0.0343%	0.0386%	0.0715%	0.0107%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 23.15
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>										
All Races ^b	996,735	835,034	588,312	302,931	105,849	16,054	2,810,743	2,849,303	836	2,850,139
White ^b	793,823	686,303	496,168	260,108	95,674	14,647	2,315,571	2,350,701	988	2,351,689
Black ^b	148,024	111,939	73,675	34,907	8,142	1,133	375,436	378,155	84	378,239
Asian/Pacific Islander ^c	21,381	16,882	10,000	+	+	+	52,881	+	+	+
Hispanic ^d	60,772	45,886	29,420	+	+	+	150,385	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	1.5451%	-	-	0.0003%	0.0014%	0.0774%	1.0088%	4.8974%	11.8301%	15.0969%	1.6948%
White ^c	1.6325%	-	-	0.0003%	0.0012%	0.0666%	0.9540%	4.8256%	11.9166%	15.0297%	1.6837%
Black ^c	1.7495%	-	-	-	0.0034%	0.2055%	2.0295%	8.4702%	17.8005%	20.8709%	2.6527%
Asian/Pacific Islander ^c	0.7380%	-	-	-	-	0.0193%	0.3384%	2.0299%	5.8994%	9.8450%	0.8715%
Hispanic ^d	0.5682%	-	-	-	-	0.0406%	0.6020%	3.3836%	9.3451%	12.9804%	1.3247%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 24.21
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	38,885	17,420	9,999	5,516	3,718	2,018	74,128	78,761	1,082	79,843
	Males	22,739	10,092	5,701	2,948	1,720	1,001	42,588	44,833	609	45,442
	Females	16,146	7,328	4,298	2,568	1,998	1,017	31,540	33,928	473	34,401
White ^b	Both Sexes	28,731	12,921	7,188	3,979	2,717	1,466	54,465	57,927	859	58,786
	Males	17,480	7,789	4,224	2,081	1,215	751	32,321	34,062	470	34,532
	Females	11,251	5,132	2,964	1,898	1,502	715	22,144	23,865	389	24,254
Black ^b	Both Sexes	6,058	2,556	1,684	845	495	322	11,453	12,084	153	12,237
	Males	2,902	1,231	830	461	277	140	5,616	5,903	85	5,988
	Females	3,156	1,325	854	384	218	182	5,837	6,181	68	6,249
Asian/ Pacific Islander ^c	Both Sexes	3,411	1,920	1,071	+	+	+	7,166	+	+	+
	Males	1,963	1,055	605	+	+	+	4,060	+	+	+
	Females	1,448	865	466	+	+	+	3,106	+	+	+
Hispanic ^d	Both Sexes	6,116	2,476	1,348	+	+	+	10,818	+	+	+
	Males	3,298	1,377	721	+	+	+	5,769	+	+	+
	Females	2,818	1,099	627	+	+	+	5,049	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0261%	-	0.0001%	0.0009%	0.0035%	0.0127%	0.0317%	0.0683%	0.1314%	0.1784%	0.0249%
	Males	0.0301%	-	-	0.0008%	0.0032%	0.0131%	0.0385%	0.0888%	0.1764%	0.2343%	0.0317%
	Females	0.0223%	-	-	0.0011%	0.0037%	0.0122%	0.0251%	0.0498%	0.0947%	0.1450%	0.0195%
White ^c	Both Sexes	0.0225%	-	-	0.0008%	0.0033%	0.0115%	0.0270%	0.0572%	0.1059%	0.1333%	0.0203%
	Males	0.0264%	-	-	0.0007%	0.0031%	0.0121%	0.0333%	0.0759%	0.1449%	0.1744%	0.0261%
	Females	0.0186%	-	-	0.0010%	0.0035%	0.0110%	0.0208%	0.0397%	0.0733%	0.1085%	0.0155%
Black ^c	Both Sexes	0.0249%	-	-	0.0009%	0.0030%	0.0147%	0.0442%	0.0875%	0.1596%	0.1963%	0.0301%
	Males	0.0254%	-	-	-	0.0025%	0.0149%	0.0464%	0.1083%	0.1956%	0.2201%	0.0348%
	Females	0.0245%	-	-	-	0.0035%	0.0146%	0.0422%	0.0714%	0.1345%	0.1846%	0.0269%
Asian/ Pacific Islander ^c	Both Sexes	0.0469%	-	-	0.0015%	0.0040%	0.0170%	0.0468%	0.1149%	0.2537%	0.4492%	0.0470%
	Males	0.0551%	-	-	0.0011%	0.0039%	0.0176%	0.0627%	0.1485%	0.3420%	0.6116%	0.0621%
	Females	0.0394%	-	-	0.0018%	0.0041%	0.0165%	0.0331%	0.0873%	0.1851%	0.3496%	0.0360%
Hispanic ^d	Both Sexes	0.0208%	-	-	0.0009%	0.0055%	0.0165%	0.0434%	0.0915%	0.1944%	0.2512%	0.0351%
	Males	0.0218%	-	-	0.0006%	0.0053%	0.0164%	0.0490%	0.1073%	0.2666%	0.2914%	0.0427%
	Females	0.0198%	-	-	0.0012%	0.0057%	0.0166%	0.0380%	0.0779%	0.1409%	0.2270%	0.0295%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 25.15
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>										
All Races ^b	42,726	39,615	37,513	32,672	28,999	22,670	170,263	226,015	14,357	240,372
White ^b	39,092	37,146	35,108	31,004	27,916	21,895	159,376	213,123	13,723	226,846
Black ^b	1,303	971	1,089	663	368	298	4,245	5,135	150	5,285
Asian/Pacific Islander ^c	936	708	648	+	+	+	2,868	+	+	+
Hispanic ^d	6,848	5,262	3,793	+	+	+	19,147	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.0966%	0.0025%	0.0070%	0.0709%	0.1733%	0.2172%	0.1591%	0.0754%	0.0291%	0.0148%	0.0961%
White ^c	0.1192%	0.0024%	0.0083%	0.0884%	0.2171%	0.2692%	0.1921%	0.0900%	0.0345%	0.0174%	0.1185%
Black ^c	0.0209%	0.0019%	0.0023%	0.0140%	0.0312%	0.0508%	0.0424%	0.0206%	-	-	0.0216%
Asian/Pacific Islander ^c	0.0324%	0.0040%	0.0047%	0.0298%	0.0519%	0.0646%	0.0518%	0.0235%	0.0087%	-	0.0311%
Hispanic ^d	0.0717%	0.0029%	0.0089%	0.0785%	0.1583%	0.1462%	0.0931%	0.0364%	0.0183%	-	0.0728%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 26.21
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	194,792	136,396	88,785	61,311	44,208	34,113	509,772	600,285	36,830	637,115
	Males	43,408	29,562	19,306	12,908	9,224	6,950	111,234	130,022	8,385	138,407
	Females	151,384	106,834	69,479	48,403	34,984	27,163	398,538	470,263	28,445	498,708
White ^b	Both Sexes	165,975	119,118	77,847	54,387	39,765	30,621	442,887	524,546	33,329	557,875
	Males	38,349	26,465	17,462	11,709	8,449	6,305	99,514	116,744	7,036	123,780
	Females	127,626	92,653	60,385	42,678	31,316	24,316	343,373	407,802	26,293	434,095
Black ^b	Both Sexes	13,424	8,515	5,086	3,143	2,008	1,561	31,536	35,626	1,594	37,220
	Males	2,134	1,435	849	558	459	261	5,273	5,935	172	6,107
	Females	11,290	7,080	4,237	2,585	1,549	1,300	26,263	29,691	1,422	31,113
Asian/ Pacific Islander ^c	Both Sexes	9,955	6,328	4,268	+	+	+	23,756	+	+	+
	Males	1,909	1,068	791	+	+	+	4,272	+	+	+
	Females	8,046	5,260	3,477	+	+	+	19,484	+	+	+
Hispanic ^d	Both Sexes	20,560	13,418	8,413	+	+	+	48,775	+	+	+
	Males	3,550	2,187	1,456	+	+	+	8,170	+	+	+
	Females	17,010	11,231	6,957	+	+	+	40,605	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1471%	0.0002%	0.0055%	0.0417%	0.1373%	0.2382%	0.2910%	0.2966%	0.2770%	0.1837%	0.1405%
	Males	0.0637%	-	0.0021%	0.0132%	0.0437%	0.0905%	0.1258%	0.1587%	0.1701%	0.1210%	0.0623%
	Females	0.2284%	0.0002%	0.0090%	0.0715%	0.2314%	0.3846%	0.4492%	0.4211%	0.3642%	0.2211%	0.2151%
White ^c	Both Sexes	0.1611%	0.0002%	0.0060%	0.0480%	0.1532%	0.2607%	0.3152%	0.3091%	0.2858%	0.1817%	0.1514%
	Males	0.0718%	-	0.0023%	0.0159%	0.0476%	0.1004%	0.1394%	0.1720%	0.1824%	0.1237%	0.0681%
	Females	0.2499%	0.0003%	0.0099%	0.0823%	0.2644%	0.4264%	0.4893%	0.4372%	0.3723%	0.2167%	0.2342%
Black ^c	Both Sexes	0.0720%	-	0.0018%	0.0135%	0.0575%	0.1258%	0.1507%	0.1960%	0.1992%	0.1344%	0.0775%
	Males	0.0250%	-	0.0013%	0.0041%	0.0147%	0.0417%	0.0531%	0.0792%	0.0948%	0.0869%	0.0296%
	Females	0.1148%	-	0.0024%	0.0226%	0.0950%	0.1996%	0.2359%	0.2873%	0.2720%	0.1578%	0.1172%
Asian/ Pacific Islander ^c	Both Sexes	0.1361%	-	0.0055%	0.0327%	0.1192%	0.2119%	0.2645%	0.2837%	0.2687%	0.2155%	0.1290%
	Males	0.0516%	-	0.0017%	0.0068%	0.0429%	0.0757%	0.1004%	0.1263%	0.1398%	0.1125%	0.0517%
	Females	0.2136%	-	0.0096%	0.0588%	0.1882%	0.3324%	0.4058%	0.4131%	0.3689%	0.2785%	0.1952%
Hispanic ^d	Both Sexes	0.0937%	-	0.0055%	0.0341%	0.1081%	0.1733%	0.2425%	0.2690%	0.2645%	0.1954%	0.1175%
	Males	0.0309%	-	0.0019%	0.0081%	0.0277%	0.0523%	0.0832%	0.1154%	0.1498%	0.1050%	0.0438%
	Females	0.1576%	-	0.0093%	0.0625%	0.1936%	0.2992%	0.3959%	0.4014%	0.3495%	0.2499%	0.1879%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 27.21
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	203,937	140,089	94,287	62,385	37,347	22,853	525,220	578,563	8,863	587,426
	Males	154,968	105,240	70,014	45,589	27,601	16,956	393,949	432,356	5,712	438,068
	Females	48,969	34,849	24,273	16,796	9,746	5,897	131,271	146,207	3,151	149,358
White ^b	Both Sexes	185,397	129,466	88,061	58,758	35,511	21,630	484,832	535,590	6,980	542,570
	Males	141,682	97,696	65,571	42,906	26,240	15,989	365,004	401,493	4,407	405,900
	Females	43,715	31,770	22,490	15,852	9,271	5,641	119,828	134,097	2,573	136,670
Black ^b	Both Sexes	10,121	5,389	3,603	2,124	1,104	792	22,057	23,605	158	23,763
	Males	7,173	3,625	2,368	1,599	791	658	15,307	16,483	129	16,612
	Females	2,948	1,764	1,235	525	313	134	6,750	7,122	29	7,151
Asian/ Pacific Islander ^c	Both Sexes	3,918	2,595	1,376	+	+	+	8,879	+	+	+
	Males	2,973	1,998	1,038	+	+	+	6,739	+	+	+
	Females	945	597	338	+	+	+	2,140	+	+	+
Hispanic ^d	Both Sexes	8,143	5,179	3,082	+	+	+	18,583	+	+	+
	Males	6,039	3,950	2,274	+	+	+	13,843	+	+	+
	Females	2,104	1,229	808	+	+	+	4,740	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1372%	0.0003%	0.0004%	0.0015%	0.0068%	0.0266%	0.1080%	0.3510%	0.8108%	1.2231%	0.1320%
	Males	0.2098%	0.0003%	0.0005%	0.0019%	0.0096%	0.0397%	0.1691%	0.5737%	1.3877%	2.3637%	0.2306%
	Females	0.0665%	0.0002%	0.0002%	0.0010%	0.0040%	0.0136%	0.0494%	0.1499%	0.3408%	0.5416%	0.0574%
White ^c	Both Sexes	0.1657%	0.0002%	0.0004%	0.0016%	0.0082%	0.0315%	0.1223%	0.4019%	0.9379%	1.3573%	0.1502%
	Males	0.2514%	-	0.0006%	0.0022%	0.0113%	0.0455%	0.1885%	0.6467%	1.5860%	2.6237%	0.2596%
	Females	0.0803%	0.0003%	-	0.0009%	0.0048%	0.0169%	0.0568%	0.1730%	0.3958%	0.5910%	0.0654%
Black ^c	Both Sexes	0.0516%	-	-	0.0011%	0.0034%	0.0152%	0.0661%	0.1892%	0.3745%	0.5758%	0.0659%
	Males	0.0760%	-	-	-	0.0053%	0.0258%	0.1071%	0.3243%	0.6406%	1.0403%	0.1141%
	Females	0.0293%	-	-	0.0015%	0.0016%	0.0060%	0.0303%	0.0837%	0.1887%	0.3473%	0.0336%
Asian/ Pacific Islander ^c	Both Sexes	0.0593%	-	-	-	0.0026%	0.0090%	0.0495%	0.1439%	0.3314%	0.6678%	0.0598%
	Males	0.0932%	-	-	-	0.0034%	0.0150%	0.0830%	0.2477%	0.5933%	1.2399%	0.1073%
	Females	0.0282%	-	-	-	0.0018%	0.0037%	0.0206%	0.0586%	0.1278%	0.3174%	0.0255%
Hispanic ^d	Both Sexes	0.0357%	-	-	0.0012%	0.0037%	0.0144%	0.0521%	0.1622%	0.3861%	0.6519%	0.0658%
	Males	0.0525%	-	-	0.0016%	0.0050%	0.0192%	0.0807%	0.2599%	0.6959%	1.2539%	0.1166%
	Females	0.0186%	-	-	0.0007%	0.0024%	0.0094%	0.0247%	0.0779%	0.1565%	0.2895%	0.0294%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person.

Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data

(excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997,

Table 28.12
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^f	Complete ^g
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	66,771	56,113	52,196	45,861	39,730	33,795	245,597	337,686	+	+
	Males	34,885	29,802	26,799	23,976	20,735	16,999	128,572	173,483	+	+
	Females	31,886	26,311	25,397	21,885	18,995	16,796	117,025	164,203	+	+
White ^b	Both Sexes	53,357	46,327	44,082	38,388	33,831	29,635	203,123	283,768	+	+
	Males	28,049	24,840	22,579	20,048	17,788	14,965	106,787	146,395	+	+
	Females	25,308	21,487	21,503	18,340	16,043	14,670	96,336	137,373	+	+
Black ^b	Both Sexes	8,264	6,170	5,071	4,649	3,813	2,673	26,476	34,244	+	+
	Males	4,181	3,169	2,583	2,372	1,909	1,283	13,479	16,982	+	+
	Females	4,083	3,001	2,488	2,277	1,904	1,390	12,997	17,262	+	+
Asian/ Pacific Islander ^c	Both Sexes	2,933	2,281	1,967	+	+	+	9,192	+	+	+
	Males	1,518	1,179	1,053	+	+	+	4,778	+	+	+
	Females	1,415	1,102	914	+	+	+	4,414	+	+	+
Hispanic ^d	Both Sexes	14,118	10,853	8,919	+	+	+	42,077	+	+	+
	Males	7,461	6,089	4,733	+	+	+	22,698	+	+	+
	Females	6,657	4,764	4,186	+	+	+	19,379	+	+	+

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry (d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^{f g} (f) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <38 and >=38. Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
+ Not available.

Table 29.7
U.S. Childhood Cancer Survivors at January 1, 2013
Number of People Previously Diagnosed with Cancer as Children (Ages 0-19 Years) in the United States and Alive January 1, 2013
By International Classification of Childhood Cancer (ICCC)^a Selected Group and Subgroup, Age Group, and Sex

Site	Sex	Complete Prevalence Counts by Age at Prevalence							All Ages		
		0-19	20-29	30-39	40-49	50-59	60+	All Ages	Diagnosed between 1975-2012	Diagnosed before 1975	% ^b
									Count	Count	
All Sites	Both	116,243	101,788	68,311	55,703	43,495	22,743	408,283	337,415	70,868	17
	Male	61,956	53,753	33,848	25,551	18,225	11,361	204,694	173,596	31,098	15
	Female ^c	54,287	48,035	34,463	30,152	25,270	11,382	203,589	163,819	39,770	20
I Leukemia	Both	36,587	22,644	13,340	6,296	2,239	344	81,450	77,160	4,290	5
	Male	19,999	12,220	6,702	3,312	1,060	146	43,439	41,538	1,901	4
	Female	16,588	10,424	6,638	2,984	1,179	198	38,011	35,622	2,389	6
Acute lymphocytic leukemia ^d	Both	30,608	18,643	11,279	5,559	1,097	17	67,203	64,006	3,197	5
	Male	16,636	10,245	5,795	3,154	595	9	36,434	34,769	1,665	5
	Female	13,972	8,398	5,484	2,405	502	8	30,769	29,237	1,532	5
I(b) Acute myeloid leukemia	Both	4,292	2,704	1,270	963	823	455	10,507	8,912	1,595	15
	Male	2,282	1,282	540	441	417	239	5,201	4,397	804	15
	Female	2,010	1,422	730	522	406	216	5,306	4,515	791	15
II(a) Hodgkin lymphoma	Both	4,774	9,382	8,295	7,623	5,649	1,885	37,608	32,457	5,151	14
	Male	2,624	4,830	4,124	3,814	2,749	971	19,112	16,363	2,749	14
	Female	2,150	4,552	4,171	3,809	2,900	914	18,496	16,094	2,402	13
II(b,c,e) Non-Hodgkin lymphoma	Both	6,877	7,834	5,035	3,426	2,042	1,400	26,614	23,175	3,439	13
	Male	4,714	5,103	3,422	2,218	1,078	801	17,336	15,477	1,859	11
	Female	2,163	2,731	1,613	1,208	964	599	9,278	7,698	1,580	17
III CNS and misc intracranial & intraspinal neoplasms	Both	21,219	15,694	9,853	8,337	5,871	2,285	63,259	52,621	10,638	17
	Male	11,405	8,694	5,393	4,169	2,651	1,002	33,314	28,499	4,815	14
	Female	9,814	7,000	4,460	4,168	3,220	1,283	29,945	24,122	5,823	19
IV Neuroblastoma & other peripheral nervous cell tumor	Both	9,783	4,201	2,475	1,870	1,586	1,082	20,997	16,752	4,245	20
	Male	4,996	2,235	1,147	607	551	245	9,781	8,476	1,305	13
	Female	4,787	1,966	1,328	1,263	1,035	837	11,216	8,276	2,940	26
VI Renal tumors	Both	7,533	5,053	3,717	4,450	2,981	1,145	24,879	16,944	7,935	32
	Male	3,595	2,434	1,630	2,506	1,619	662	12,446	7,910	4,536	36
	Female	3,938	2,619	2,087	1,944	1,362	483	12,433	9,034	3,399	27
VIII Malignant bone tumors	Both	3,657	4,481	3,333	2,508	2,624	1,183	17,786	14,327	3,459	19
	Male	2,194	2,606	1,773	1,337	1,146	620	9,676	8,087	1,589	16
	Female	1,463	1,875	1,560	1,171	1,478	563	8,110	6,240	1,870	23
VIII(a) Osteosarcoma	Both	1,966	2,435	1,754	1,364	1,280	897	9,696	7,729	1,967	20
	Male	1,077	1,343	968	714	565	396	5,063	4,196	867	17
	Female	889	1,092	786	650	715	501	4,633	3,533	1,100	24
VIII(c) Ewing tumor & related sarcomas of bone	Both	1,339	1,476	1,053	714	435	220	5,237	4,648	589	11
	Male	891	896	499	352	206	90	2,934	2,693	241	8
	Female	448	580	554	362	229	130	2,303	1,955	348	15
IX Soft tissue & other extraosseous sarcomas	Both	7,120	7,147	4,601	4,616	4,649	5,214	33,347	23,009	10,338	31
	Male	3,886	4,019	2,240	2,347	2,244	2,914	17,650	12,245	5,405	31
	Female	3,234	3,128	2,361	2,269	2,405	2,300	15,697	10,764	4,933	31
X Germ cell & trophoblastic tumors & neoplasms of gonads	Both	5,185	8,402	5,693	6,084	6,790	7,575	39,729	26,146	13,583	34
	Male	2,664	5,501	3,523	3,297	3,362	3,526	21,873	15,467	6,406	29
	Female	2,521	2,901	2,170	2,787	3,428	4,049	17,856	10,679	7,177	40

^a International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.

^b Percent diagnosed before 1975 of all ages complete prevalence counts.

^c All sites females excludes breast cancer.

^d SEER site recode Acute Lymphocytic Leukemia was used instead of the ICCC classification lymphocytic leukemia (Ia) because of statistical modeling requirements.

For more information regarding the modeling of the prevalence estimates, see Mariotto et al. Long-Term Survivors of Childhood Cancers in the United States. Cancer Epidemiol Biomarkers Prev 2009, page 1034.