

Table 23.1
Cancer of the Prostate (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2017 With up to Five Joinpoints, 2000-2017 With up to Three Joinpoints,
All Ages by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	2008-17	2013-17								
SEER 9 Delay-Adjusted Incidence^a, 1975-2017														
All Races	1975-88	2.6*	1988-92	16.1*	1992-95	-9.5	1995-08	-0.2	2008-14	-7.9*	2014-17	4.4	-4.0*	1.2
White	1975-88	2.8*	1988-92	16.0*	1992-95	-10.6	1995-07	0.1	2007-14	-7.0*	2014-17	3.2	-3.7*	0.5
Black	1975-89	2.0*	1989-92	22.0*	1992-95	-4.6	1995-09	-1.1*	2009-14	-7.9*	2014-17	4.3*	-3.2*	1.1
SEER 21 Delay-Adjusted Incidence^b, 2000-2017														
All Races	2000-10	-1.7*	2010-14	-8.8*	2014-17	4.3							-3.0	0.9
White	2000-10	-1.9*	2010-14	-9.1*	2014-17	4.2							-3.2	0.7
White NH ^{ef}	2000-10	-1.8*	2010-14	-9.0*	2014-17	5.0							-2.9	1.3
Black	2000-11	-1.9*	2011-14	-9.0	2014-17	3.2							-2.7	0.0
Black NH ^{ef}	2000-11	-1.8*	2011-14	-9.0	2014-17	3.5							-2.6	0.2
API ^e	2000-11	-2.9*	2011-14	-10.8	2014-17	5.4							-3.0	1.1
AI/AN ^{eg}	2000-03	6.8	2003-17	-3.8*									-3.8*	-3.8*
Hispanic ^f	2000-10	-2.0*	2010-14	-8.7*	2014-17	0.3							-4.3*	-2.1
U.S. Cancer Mortality^c, 1975-2017														
All Races	1975-87	0.9*	1987-91	3.0*	1991-94	-0.5	1994-98	-4.2*	1998-13	-3.5*	2013-17	-0.3	-2.1*	-0.3
White	1975-87	0.8*	1987-91	3.1*	1991-94	-0.7	1994-99	-4.3*	1999-13	-3.3*	2013-17	0.0	-1.9*	0.0
Black	1975-88	1.9*	1988-93	3.2*	1993-00	-2.5*	2000-10	-3.7*	2010-13	-6.2*	2013-17	-1.4	-3.5*	-1.4
U.S. Cancer Mortality^c, 2000-2017														
All Races	2000-13	-3.5*	2013-17	-0.3									-2.1*	-0.3
White	2000-13	-3.3*	2013-17	0.0									-1.9*	0.0
White NH ^{ef}	2000-13	-3.3*	2013-17	0.1									-1.8*	0.1
Black	2000-14	-4.2*	2014-17	-1.7									-3.3*	-2.3
Black NH ^{ef}	2000-17	-3.8*											-3.8*	-3.8*
API ^e	2000-17	-2.3*											-2.3*	-2.3*
AI/AN ^{eg}	2000-17	-1.7*											-1.7*	-1.7*
Hispanic ^f	2000-14	-3.0*	2014-17	-0.1									-2.0*	-0.8

Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>). The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 21 areas (SEER 9 areas, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 23.2
Cancer of the Prostate (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2017 With up to Five Joinpoints, 2000-2017 With up to Three Joinpoints,
Ages <65 by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	2008-17	2013-17								
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2017</u>														
All Races	1975-88	3.1*	1988-92	23.4*	1992-01	3.3*	2001-09	0.2	2009-14	-8.2*	2014-17	3.0	-3.7*	0.1
White	1975-88	3.5*	1988-92	23.0*	1992-01	3.1*	2001-09	0.1	2009-14	-8.6*	2014-17	2.1	-4.2*	-0.7
Black	1975-89	2.0*	1989-93	26.4*	1993-01	2.7*	2001-10	-0.5	2010-14	-7.5*	2014-17	4.4	-2.1	1.3
<u>SEER 21 Delay-Adjusted Incidence^b, 2000-2017</u>														
All Races	2000-10	-0.1	2010-14	-8.7*	2014-17	3.0							-3.1	-0.1
White	2000-09	0.2	2009-14	-8.0*	2014-17	1.9							-3.9*	-0.7
White NH ^{ef}	2000-09	0.3	2009-14	-7.9*	2014-17	2.7							-3.6	0.0
Black	2000-11	-0.3	2011-14	-8.1	2014-17	2.5							-2.1	-0.3
Black NH ^{ef}	2000-11	-0.3	2011-14	-8.1	2014-17	2.7							-2.0	-0.1
API ^e	2000-11	0.0	2011-14	-10.1	2014-17	3.7							-2.3	0.0
AI/AN ^{eg}	2000-17	-2.9*											-2.9*	-2.9*
Hispanic ^f	2000-09	-0.3	2009-17	-4.7*									-4.2*	-4.7*
<u>U.S. Cancer Mortality^c, 1975-2017</u>														
All Races	1975-87	0.5*	1987-90	3.1	1990-96	-3.2*	1996-00	-5.8*	2000-12	-1.4*	2012-17	-0.1	-0.7	-0.1
White	1975-87	0.7*	1987-90	3.5	1990-02	-4.3*	2002-17	-0.8*					-0.8*	-0.8*
Black	1975-92	0.5*	1992-12	-2.9*	2012-17	0.0							-1.3*	0.0
<u>U.S. Cancer Mortality^c, 2000-2017</u>														
All Races	2000-17	-1.1*											-1.1*	-1.1*
White	2000-17	-1.0*											-1.0*	-1.0*
White NH ^{ef}	2000-17	-1.0*											-1.0*	-1.0*
Black	2000-17	-2.0*											-2.0*	-2.0*
Black NH ^{ef}	2000-17	-1.9*											-1.9*	-1.9*
API ^e	2000-17	0.2											0.2	0.2
AI/AN ^{eg}	2000-17	-0.1											-0.1	-0.1
Hispanic ^f	2000-17	-0.9*											-0.9*	-0.9*

Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>). The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 21 areas (SEER 9 areas, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

* The APC/AAPC is significantly different from zero ($p<.05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 23.3
Cancer of the Prostate (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2017 With up to Five Joinpoints, 2000-2017 With up to Three Joinpoints,
Ages 65+ by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2008-17	2013-17
SEER 9 Delay-Adjusted Incidence^a, 1975-2017														
All Races	1975-88	2.5*	1988-92	14.9*	1992-95	-13.3*	1995-07	-0.8*	2007-14	-7.3*	2014-17	4.7	-3.4*	1.6
White	1975-87	2.3*	1987-92	12.6*	1992-95	-13.5*	1995-07	-0.7	2007-14	-7.3*	2014-17	4.7	-3.5*	1.5
Black	1975-89	2.0*	1989-92	20.3*	1992-95	-9.0*	1995-09	-2.1*	2009-14	-8.7*	2014-17	4.5	-3.7*	1.1
SEER 21 Delay-Adjusted Incidence^b, 2000-2017														
All Races	2000-10	-2.5*	2010-14	-8.9*	2014-17	5.0							-3.0	1.4
White	2000-10	-2.6*	2010-14	-8.9*	2014-17	4.9							-3.1	1.3
White NH ^{ef}	2000-10	-2.6*	2010-14	-8.8*	2014-17	5.6							-2.8	1.8
Black	2000-11	-2.8*	2011-14	-9.6*	2014-17	3.8							-3.0*	0.3
Black NH ^{ef}	2000-11	-2.7*	2011-14	-9.7*	2014-17	4.0							-3.0	0.4
API ^e	2000-11	-3.7*	2011-14	-11.2	2014-17	6.1							-3.2	1.5
AI/AN ^{eg}	2000-17	-3.0*											-3.0*	-3.0*
Hispanic ^f	2000-10	-2.6*	2010-14	-9.3*	2014-17	0.6							-4.6*	-2.0
U.S. Cancer Mortality^c, 1975-2017														
All Races	1975-87	0.9*	1987-90	3.3*	1990-93	1.3	1993-13	-3.6*	2013-17	-0.3			-2.2*	-0.3
White	1975-87	0.8*	1987-91	3.2*	1991-94	-0.5	1994-98	-4.4*	1998-13	-3.5*	2013-17	0.0	-1.9*	0.0
Black	1975-88	2.1*	1988-93	3.5*	1993-01	-2.6*	2001-10	-3.9*	2010-13	-6.4*	2013-17	-1.5	-3.7*	-1.5
U.S. Cancer Mortality^c, 2000-2017														
All Races	2000-13	-3.6*	2013-17	-0.4									-2.2*	-0.4
White	2000-13	-3.5*	2013-17	0.0									-1.9*	0.0
White NH ^{ef}	2000-13	-3.5*	2013-17	0.1									-1.9*	0.1
Black	2000-17	-4.1*											-4.1*	-4.1*
Black NH ^{ef}	2000-17	-4.0*											-4.0*	-4.0*
API ^e	2000-17	-2.5*											-2.5*	-2.5*
AI/AN ^{eg}	2000-17	-1.8*											-1.8*	-1.8*
Hispanic ^f	2000-14	-3.1*	2014-17	0.0									-2.1*	-0.8

Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>). The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 21 areas (SEER 9 areas, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 23.4
Cancer of the Prostate (Invasive)

Delay-adjusted SEER Incidence^a Rates by Year, Race and Age

Year of Diagnosis:	All Races, Males			White Males			Black Males		
	All	<65	65+	All	<65	65+	All	<65	65+
1975-1979	99.14	14.71	682.70	97.69	14.16	675.05	150.71	27.73	1,000.84
1980	106.05	15.52	731.77	105.00	14.69	729.23	161.14	33.79	1,041.42
1981	108.88	16.84	745.11	107.87	15.96	743.20	162.01	34.72	1,041.86
1982	108.25	16.36	743.41	107.33	15.81	739.99	167.75	31.46	1,109.83
1983	111.65	17.18	764.62	110.85	16.64	762.09	171.07	34.02	1,118.41
1984	111.70	17.27	764.40	110.11	16.22	759.14	178.74	37.05	1,158.14
1985	115.52	17.95	790.00	114.65	17.69	784.82	170.41	31.59	1,129.99
1986	119.10	18.55	814.14	119.05	18.30	815.47	168.59	34.08	1,098.34
1987	133.78	21.53	909.65	134.64	21.15	919.13	189.42	36.06	1,249.49
1988	137.65	22.19	935.77	138.75	22.27	943.93	190.20	34.88	1,263.80
1989	145.43	24.21	983.30	146.23	24.11	990.33	191.83	37.41	1,259.25
1990	171.14	28.91	1,154.25	172.93	28.99	1,167.93	222.53	44.83	1,450.91
1991	214.89	38.94	1,431.05	216.61	39.45	1,441.16	289.85	57.64	1,894.97
1992	237.60	49.52	1,537.72	238.75	49.55	1,546.53	331.08	78.77	2,075.15
1993	209.67	50.50	1,309.86	204.74	48.93	1,281.77	347.90	95.71	2,091.16
1994	180.43	48.70	1,090.98	174.50	46.91	1,056.48	316.05	96.20	1,835.74
1995	169.55	50.19	994.60	164.81	48.19	970.90	286.13	100.99	1,565.91
1996	169.68	53.63	971.85	165.68	52.16	950.38	286.38	101.44	1,564.74
1997	173.83	55.38	992.64	170.54	54.06	975.69	288.43	98.97	1,598.03
1998	171.25	55.10	974.13	167.84	53.19	960.38	292.51	104.98	1,588.75
1999	183.72	60.92	1,032.56	179.64	58.88	1,014.44	297.09	114.90	1,556.49
2000	183.37	61.90	1,023.03	179.83	59.81	1,009.46	297.30	117.66	1,539.01
2001	185.41	64.75	1,019.48	183.36	62.77	1,016.91	280.68	117.15	1,411.11
2002	182.74	65.67	991.99	180.21	64.06	983.06	293.69	118.98	1,501.35
2003	170.19	62.14	917.06	166.99	59.86	907.48	266.11	118.13	1,289.03
2004	166.36	62.17	886.56	163.09	60.65	871.22	263.05	108.06	1,334.40
2005	157.10	58.14	841.15	152.64	55.57	823.59	256.94	108.38	1,283.80
2006	172.69	64.92	917.60	169.24	62.76	905.29	260.24	111.31	1,289.73
2007	175.79	66.97	928.05	171.46	64.08	913.67	272.32	121.47	1,315.08
2008	158.65	62.84	820.95	154.27	60.15	804.86	257.22	112.75	1,255.87
2009	156.02	62.74	800.84	152.38	60.52	787.34	248.80	111.92	1,194.94
2010	148.77	59.31	767.18	145.88	57.00	760.26	235.78	107.70	1,121.17
2011	142.84	57.83	730.52	139.29	55.45	718.79	225.76	106.67	1,048.92
2012	116.95	47.08	599.93	112.33	43.95	585.00	196.09	94.05	901.43
2013	111.86	44.92	574.56	108.44	42.53	564.02	189.84	88.85	887.88
2014	102.81	41.86	524.14	100.34	39.69	519.62	165.16	79.33	758.41
2015	109.14	42.96	566.58	106.38	40.01	565.22	175.15	87.20	783.06
2016	112.02	43.66	584.57	107.80	40.40	573.70	185.53	87.16	865.55
2017	119.00	45.89	624.40	114.35	42.06	614.01	192.49	92.86	881.21

Reference: [Cancer Incidence Rates Adjusted for Reporting Delay](#).

Individual rates for diagnosis years 1975-1979 available in web browser version of [Cancer Statistics Review](#).

SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.

Table 23.5
Cancer of the Prostate (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

	All Races, Males			White Males			Black Males		
	All	<65	65+	All	<65	65+	All	<65	65+
Year of Diagnosis:									
1975-2017	144.47	43.96	839.26	141.41	41.63	831.11	227.55	83.86	1,220.78
1975-1979	99.14	14.71	682.70	97.69	14.16	675.05	150.71	27.73	1,000.84
1980	106.05	15.52	731.77	105.00	14.69	729.23	161.14	33.79	1,041.42
1981	108.88	16.84	745.11	107.87	15.96	743.20	162.01	34.72	1,041.86
1982	108.25	16.36	743.41	107.33	15.81	739.99	167.75	31.46	1,109.83
1983	111.65	17.18	764.62	110.85	16.64	762.09	171.07	34.02	1,118.41
1984	111.70	17.27	764.40	110.11	16.22	759.14	178.74	37.05	1,158.14
1985	115.52	17.95	790.00	114.65	17.69	784.82	170.41	31.59	1,129.99
1986	119.10	18.55	814.14	119.05	18.30	815.47	168.59	34.08	1,098.34
1987	133.78	21.53	909.65	134.64	21.15	919.13	189.42	36.06	1,249.49
1988	137.65	22.19	935.77	138.75	22.27	943.93	190.20	34.88	1,263.80
1989	145.43	24.21	983.30	146.23	24.11	990.33	191.83	37.41	1,259.25
1990	171.14	28.91	1,154.25	172.93	28.99	1,167.93	222.53	44.83	1,450.91
1991	214.89	38.94	1,431.05	216.61	39.45	1,441.16	289.85	57.64	1,894.97
1992	237.60	49.52	1,537.72	238.75	49.55	1,546.53	331.08	78.77	2,075.15
1993	209.67	50.50	1,309.86	204.74	48.93	1,281.77	347.90	95.71	2,091.16
1994	180.43	48.70	1,090.98	174.50	46.91	1,056.48	316.05	96.20	1,835.74
1995	169.55	50.19	994.60	164.81	48.19	970.90	286.13	100.99	1,565.91
1996	169.68	53.63	971.85	165.68	52.16	950.38	286.38	101.44	1,564.74
1997	173.83	55.38	992.64	170.54	54.06	975.69	288.43	98.97	1,598.03
1998	171.25	55.10	974.13	167.84	53.19	960.38	292.51	104.98	1,588.75
1999	183.72	60.92	1,032.56	179.64	58.88	1,014.44	297.09	114.90	1,556.49
2000	183.37	61.90	1,023.03	179.83	59.81	1,009.46	297.30	117.66	1,539.01
2001	185.41	64.75	1,019.48	183.36	62.77	1,016.91	280.68	117.15	1,411.11
2002	182.74	65.67	991.99	180.21	64.06	983.06	293.69	118.98	1,501.35
2003	170.19	62.14	917.06	166.99	59.86	907.48	266.11	118.13	1,289.03
2004	166.36	62.17	886.56	163.09	60.65	871.22	263.05	108.06	1,334.40
2005	157.10	58.14	841.15	152.64	55.57	823.59	256.94	108.38	1,283.80
2006	172.69	64.92	917.60	169.24	62.76	905.29	260.24	111.31	1,289.73
2007	175.79	66.97	928.05	171.46	64.08	913.67	272.32	121.47	1,315.08
2008	158.65	62.84	820.95	154.27	60.15	804.86	257.22	112.75	1,255.87
2009	155.82	62.68	799.62	151.97	60.37	785.14	247.79	111.48	1,190.01
2010	148.40	59.21	764.93	145.11	56.72	756.07	233.83	106.84	1,111.68
2011	142.31	57.68	727.32	138.19	55.05	712.84	223.05	105.41	1,036.24
2012	116.37	46.92	596.42	111.16	43.54	578.58	192.87	92.57	886.19
2013	111.15	44.73	570.32	107.02	42.03	556.27	185.87	87.07	868.83
2014	102.02	41.64	519.42	98.75	39.12	510.96	161.19	77.47	739.91
2015	107.88	42.61	559.03	104.03	39.20	552.13	169.92	84.70	759.04
2016	110.01	43.08	572.69	104.46	39.26	555.11	178.39	83.91	831.46
2017	115.24	44.71	602.76	108.69	40.18	582.26	180.92	87.51	826.57

^a Individual rates for diagnosis years 1975-1979 available in web browser version of [Cancer Statistics Review](#). SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 23.6
Cancer of the Prostate (Invasive)

Age-adjusted U.S. Death^a Rates by Year, Race and Age

	All Races, Males			White Males			Black Males		
	All	<65	65+	All	<65	65+	All	<65	65+
Year of Death:									
1975-2017	28.52	2.17	210.63	26.49	1.87	196.62	58.70	5.41	427.02
1975-1979	32.04	2.62	235.43	30.08	2.25	222.44	57.10	6.75	405.12
1980	33.05	2.67	243.08	30.77	2.33	227.39	61.99	6.57	445.03
1981	33.17	2.68	243.95	30.76	2.30	227.52	63.61	7.06	454.45
1982	33.36	2.68	245.47	31.09	2.34	229.85	62.36	6.62	447.67
1983	33.92	2.76	249.30	31.57	2.40	233.18	64.55	6.99	462.46
1984	34.06	2.71	250.77	31.58	2.37	233.49	65.93	6.74	475.13
1985	33.91	2.73	249.43	31.28	2.34	231.34	67.26	7.16	482.68
1986	34.93	2.80	256.99	32.46	2.47	239.72	67.39	6.63	487.36
1987	35.11	2.71	259.06	32.57	2.38	241.27	68.88	6.66	499.03
1988	35.88	2.83	264.40	33.38	2.47	247.08	69.57	6.98	502.28
1989	37.10	2.97	273.01	34.54	2.62	255.20	71.71	7.06	518.54
1990	38.56	3.02	284.23	35.67	2.65	263.98	77.98	7.43	565.65
1991	39.31	2.89	291.08	36.50	2.52	271.39	78.24	7.06	570.21
1992	39.22	2.86	290.58	36.29	2.44	270.28	79.84	7.66	578.73
1993	39.34	2.72	292.48	36.26	2.33	270.82	81.86	7.04	599.08
1994	38.54	2.67	286.48	35.64	2.31	266.04	79.96	6.76	585.93
1995	37.29	2.58	277.23	34.41	2.22	256.93	78.20	6.70	572.47
1996	36.00	2.50	267.61	33.00	2.13	246.38	78.84	6.45	579.20
1997	34.15	2.31	254.26	31.35	1.94	234.58	74.30	6.32	544.22
1998	32.63	2.27	242.49	29.89	1.90	223.36	72.77	6.28	532.41
1999	31.56	2.06	235.48	28.91	1.72	216.88	70.13	5.80	514.78
2000	30.39	1.98	226.82	27.78	1.66	208.29	68.90	5.40	507.82
2001	29.52	1.94	220.14	26.97	1.63	202.08	67.82	5.33	499.72
2002	28.71	1.92	213.87	26.36	1.60	197.48	64.98	5.26	477.81
2003	27.19	1.90	201.97	25.02	1.59	186.95	60.37	5.30	441.02
2004	26.19	1.82	194.64	24.04	1.55	179.54	59.00	4.79	433.72
2005	25.40	1.80	188.57	23.33	1.51	174.20	57.02	4.83	417.76
2006	24.24	1.84	179.14	22.34	1.58	165.81	53.73	4.57	393.54
2007	24.23	1.75	179.68	22.23	1.49	165.58	54.88	4.51	403.03
2008	23.01	1.78	169.73	21.34	1.52	158.30	49.11	4.48	357.62
2009	22.12	1.70	163.30	20.34	1.43	151.06	49.35	4.45	359.70
2010	21.81	1.78	160.30	20.10	1.51	148.62	48.08	4.40	350.03
2011	20.79	1.66	153.03	19.26	1.39	142.76	43.98	4.25	318.62
2012	19.57	1.66	143.39	18.14	1.44	133.56	41.65	3.86	302.84
2013	19.28	1.67	141.01	18.01	1.44	132.55	39.15	3.95	282.49
2014	19.12	1.66	139.83	17.91	1.42	131.93	38.02	3.95	273.54
2015	18.96	1.65	138.61	17.74	1.41	130.60	37.56	3.84	270.66
2016	19.40	1.72	141.60	18.18	1.45	133.83	38.41	4.21	274.84
2017	18.85	1.61	138.00	17.75	1.38	130.95	36.37	3.83	261.31

^a Individual rates for years of death 1975-1979 available in web browser version of [Cancer Statistics Review](#). US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 23.7
Cancer of the Prostate (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2013-2017</u>						
Age at Diagnosis/Death:						
All ages	109.8	102.3	175.2	19.1	17.9	37.9
Under 65	43.9	39.4	82.0	1.7	1.4	4.0
65 and over	565.3	536.8	819.7	139.8	132.0	272.2
All ages (WHO world std) ^c	83.6	77.4	136.9	8.5	7.8	17.6
<u>Age-Specific Rates, 2013-2017</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	0.1	-	-	-	-	-
35-39	0.6	0.4	1.3	-	-	-
40-44	7.1	5.8	17.5	0.1	0.1	0.3
45-49	31.4	26.0	77.2	0.8	0.7	1.6
50-54	114.1	100.1	229.1	3.1	2.6	6.9
55-59	249.1	223.9	458.7	8.5	7.1	20.5
60-64	418.3	384.0	719.4	20.0	17.1	48.5
65-69	631.1	588.5	1,014.0	38.7	34.2	92.3
70-74	652.8	620.7	946.9	69.3	63.1	157.3
75-79	577.3	547.2	797.3	123.2	114.2	261.0
80-84	436.6	422.0	550.1	216.6	205.0	422.7
85+	368.1	364.3	478.7	447.8	435.9	751.9

^a SEER 21 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 23.8
Cancer of the Prostate (Invasive)

5-Year Relative and Period Survival (Percent) by Race, Diagnosis Year, Stage and Age

	All Races, Males			White Males			Black Males		
	All	<65	65+	All	<65	65+	All	<65	65+
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1960-1963 ^a	-	-	-	50	-	-	35	-	-
1970-1973 ^a	-	-	-	63	-	-	55	-	-
1975-1977 ^b	67.8	73.4	66.1	68.5	75.2	66.6	60.7	62.0	60.1
1978-1980 ^b	70.1	75.2	68.6	71.2	76.7	69.6	62.1	65.0	60.6
1981-1983 ^b	72.0	76.0	70.9	73.1	77.6	71.9	62.8	65.0	61.9
1984-1986 ^b	75.1	77.2	74.5	76.3	78.9	75.6	65.6	67.2	64.9
1987-1989 ^b	82.8	83.2	82.7	84.4	84.7	84.3	71.2	72.8	70.7
1990-1992 ^b	92.9	92.9	92.9	94.1	94.0	94.1	84.5	86.8	83.6
1993-1995 ^b	95.2	95.9	94.9	96.0	96.5	95.7	91.9	93.4	91.0
1996-1998 ^b	97.3	98.5	96.7	97.9	98.9	97.3	95.0	96.7	93.7
1999-2001 ^b	99.3	99.1	99.3	99.7	99.5	99.6	97.4	98.1	96.4
2002-2004 ^b	99.6	99.5	99.7	99.8	99.6	99.7	98.0	99.2	96.3
2005-2009 ^b	99.5	99.6	99.4	99.8 ^f	99.9	99.6	98.3	98.6	97.8
2010-2016 ^b	98.3 ^f	99.0 ^f	97.7 ^f	98.7 ^f	99.4 ^f	98.1 ^f	96.3 ^f	97.5 ^f	94.4 ^f
5-Year Period Survival (Percent)^{cd}									
2016	97.3	97.9	96.8	97.2	98.0	96.6	95.9	97.2	94.2
Stage Distribution (%) 2010-2016^{ce}									
All Stages									
Number of cases	332,075	145,886	186,189	248,870	104,945	143,925	53,305	29,514	23,791
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	76	77	75	76	77	76	78	79	78
Regional	13	16	11	14	17	12	11	14	8
Distant	6	4	8	6	4	8	7	5	9
Unstaged/Unknown	5	3	6	4	2	5	4	3	5
5-Year Relative Survival (Percent), 2010-2016^c									
Age at Diagnosis:									
Ages <45	96.1	-	-	95.0	-	-	97.7	-	-
Ages 45-54	97.3	-	-	96.9	-	-	97.9	-	-
Ages 55-64	98.5	-	-	98.7	-	-	97.2	-	-
Ages 65-74	100.0	-	-	100.0	-	-	97.6	-	-
Ages 75+	90.9	-	-	90.4	-	-	84.7	-	-
Ages <65	98.2	-	-	98.3	-	-	97.5	-	-
Ages 65+	97.5	-	-	97.4	-	-	94.5	-	-
Stage^e:									
All Stages	97.8	98.2	97.5	97.8	98.3	97.4	96.2	97.5	94.5
Localized	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Regional	100.0	99.7	100.0	100.0	99.6	100.0	100.0	100.0	99.7
Distant	30.2	33.5	28.8	29.5	32.7	28.3	29.9	33.6	27.0
Unstaged/Unknown	83.3	92.4	79.8	76.8	89.7	72.5	74.4	86.9	66.7

Based on End Results data from a series of hospital registries and one population-based registry.

SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Based on follow-up of patients into 2017. [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Expected survival rates are derived from life tables by socio-economic status, geography and race developed by the SEER program.

Based on follow-up of patients into 2017.

Period survival provides a 2016 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2010-2012: 0-1 year survival), (2009-2011: 1-2 year survival), (2008-2010: 2-3 year survival), (2007-2009: 3-4 year survival), (2006-2008: 4-5 years survival).

Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding.

The difference between 1975-1977 and 2010-2016 is statistically significant ($p<.05$).

The standard error is between 5 and 10 percentage points.

The standard error is greater than 10 percentage points.

Statistic could not be calculated due to fewer than 25 cases during the time period.

a

b

c

d

e

f

g

h

-

Table 23.9
Cancer of the Prostate (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975-	1980-	1985-	1990-	1979	1984	1989	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																														
1-year	91.4	93.4	95.5	98.8	98.7	99.1	99.2	99.5	99.6	99.7	99.9	99.8	99.7	99.8	99.8	99.8	99.8	99.8	99.7	99.6	99.6	99.7	99.3	99.4	98.7	99.2	99.3			
2-year	84.6	87.4	91.0	97.3	97.7	98.0	98.8	98.9	99.6	99.5	99.9	99.8	99.6	99.7	99.6	99.8	99.8	99.6	99.4	99.5	99.4	98.7	98.8	98.0	98.2					
3-year	78.4	81.5	86.9	96.0	97.0	97.4	98.4	98.5	99.6	99.1	99.9	99.7	99.4	99.7	99.5	99.7	99.8	99.3	99.3	99.5	99.3	98.4	98.3	97.5						
4-year	73.3	76.5	83.6	94.9	96.4	96.9	97.9	98.2	99.2	98.9	99.8	99.7	99.4	99.7	99.5	99.6	99.8	99.2	99.3	99.5	99.3	98.2	97.9							
5-year	68.8	71.9	80.4	93.7	95.7	96.4	97.4	98.1	99.2	98.8	99.8	99.7	99.1	99.7	99.1	99.6	99.8	99.2	99.3	99.5	99.2	98.1								
6-year	65.3	68.5	77.9	92.7	95.4	96.0	97.1	97.9	99.2	98.5	99.8	99.7	99.1	99.7	99.1	99.6	99.8	99.2	99.3	99.5	99.2									
7-year	61.8	65.0	75.4	91.6	95.0	95.4	97.1	97.7	99.1	98.5	99.8	99.7	99.1	99.7	99.1	99.6	99.8	99.2	99.3	99.5										
8-year	58.6	62.0	73.4	90.6	94.6	95.0	97.0	97.7	98.7	98.2	99.8	99.7	99.0	99.7	99.1	99.6	99.8	99.2	99.3											
9-year	55.7	59.7	71.4	89.4	94.3	94.4	96.6	97.0	98.5	98.2	99.8	99.7	98.5	99.6	99.1	99.6	99.8	99.2												
10-year	53.2	57.7	69.6	88.5	93.5	94.3	96.5	96.6	98.1	98.2	99.8	99.7	98.5	99.4	99.1	99.6	99.8													
11-year	50.7	55.5	67.9	87.5	92.8	93.9	96.4	96.2	97.9	97.3	99.8	99.7	98.3	99.2	99.1	99.6														
12-year	48.5	53.6	66.1	86.6	92.0	93.5	96.1	95.5	97.4	97.2	99.8	99.7	97.9	98.8	99.1															
13-year	46.5	51.6	64.3	85.9	91.0	93.0	95.3	95.3	97.2	96.6	99.8	99.7	97.9	97.9	98.8															
14-year	44.8	50.0	63.0	85.1	90.1	92.8	94.7	95.0	96.6	96.1	99.8	99.7	98.5	99.4	99.1	99.6	99.8													
15-year	43.6	48.7	61.5	84.2	89.4	92.1	94.7	94.9	96.0	95.4	99.8	99.7																		
16-year	41.9	46.9	60.0	83.3	88.9	91.2	94.1	94.4	95.6	95.4	99.3																			
17-year	40.1	45.8	59.4	82.4	88.0	90.6	93.6	94.1	95.6	95.4	95.4																			
18-year	38.8	44.5	58.3	81.8	87.7	90.4	92.9	94.1	95.2																					
19-year	38.0	43.1	57.9	80.8	87.3	89.7	92.6	94.1																						
20-year	37.0	41.7	56.9	80.4	85.8	88.8	92.6																							
21-year	35.4	40.8	56.1	79.5	84.9	88.8																								
22-year	34.0	39.5	54.8	78.7	84.3																									
23-year	33.2	38.3	53.6	77.7																										
24-year	32.4	37.4	52.5																											
25-year	31.0	35.9	51.4																											
26-year	30.4	34.7	50.4																											
27-year	29.7	34.4	49.5																											
28-year	29.1	34.0	48.8																											
29-year	28.5	33.3																												
30-year	27.0	32.4																												

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 23.10
Cancer of the Prostate (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2015-2017 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer						Risk of Being Diagnosed with Cancer						Risk of Dying from Cancer								
		+10 yrs			+20 yrs			+30 yrs			Ever			+10 yrs			+20 yrs			+30 yrs		
All Races	0	0.00	0.00	0.00	12.12	2.43		Asian/	0	0.00	0.00	0.00	7.94	2.21								
	10	0.00	0.00	0.01	12.25	2.46		Pacific	10	0.00	0.00	0.00	8.00	2.23								
	20	0.00	0.01	0.22	12.30	2.47		Islander	20	0.00	0.00	0.06	8.02	2.24								
	30	0.01	0.23	1.94	12.46	2.50			30	0.00	0.06	0.73	8.06	2.25								
	40	0.23	1.97	6.36	12.67	2.56			40	0.06	0.74	3.15	8.12	2.26								
	50	1.80	6.34	10.70	12.86	2.64			50	0.68	3.13	6.07	8.18	2.29								
	60	4.95	9.71	11.65	12.07	2.78			60	2.56	5.62	7.23	7.82	2.35								
	70	5.80	8.15	-	8.67	2.94			70	3.43	5.22	-	5.88	2.43								
	80	3.48	-	-	4.23	3.01			80	2.28	-	-	3.12	2.53								
White	0	0.00	0.00	0.00	11.28	2.28		American	0	0.00	0.00	0.00	5.62	2.04								
	10	0.00	0.00	0.01	11.38	2.31		Indian/	10	0.00	0.00	0.00	5.68	2.07								
	20	0.00	0.01	0.19	11.43	2.32		Alaska	20	0.00	0.00	0.05	5.72	2.09								
	30	0.01	0.19	1.71	11.58	2.35		Native ^a	30	0.00	0.06	0.67	5.85	2.14								
	40	0.19	1.74	5.80	11.78	2.40			40	0.06	0.69	2.47	6.06	2.23								
	50	1.60	5.80	9.92	11.97	2.47			50	0.67	2.54	4.66	6.31	2.38								
	60	4.57	9.06	10.91	11.29	2.61			60	2.06	4.40	5.87	6.22	2.60								
	70	5.44	7.68	-	8.14	2.76			70	2.79	4.55	-	4.96	2.89								
	80	3.30	-	-	3.99	2.85			80	2.52	-	-	3.12	2.98								
Black	0	0.00	0.00	0.00	16.29	3.87		Hispanic ^b	0	0.00	0.00	0.00	11.03	2.80								
	10	0.00	0.00	0.02	16.61	3.96			10	0.00	0.00	0.00	11.13	2.82								
	20	0.00	0.02	0.52	16.72	3.99			20	0.00	0.00	0.13	11.17	2.84								
	30	0.02	0.53	3.61	17.03	4.07			30	0.00	0.13	1.35	11.29	2.87								
	40	0.53	3.69	10.25	17.46	4.19			40	0.13	1.36	4.98	11.45	2.91								
	50	3.33	10.22	15.46	17.79	4.38			50	1.27	4.98	8.99	11.61	2.97								
	60	7.90	13.91	15.95	16.58	4.74			60	3.97	8.27	10.35	11.07	3.10								
	70	8.17	10.94	-	11.79	5.19			70	5.03	7.47	-	8.31	3.24								
	80	4.59	-	-	6.00	5.48			80	3.34	-	-	4.51	3.27								

Devcan 6.7.8, April 2020, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the PRCD(A(Purchased/Referred Care Delivery Areas) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 23.11
Cancer of the Prostate (Invasive)
SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity

Race/Ethnicity	SEER 21 Areas ^b		SEER 21 Areas ^b		SEER 9 Areas ^b	
	Rate 2013-2017		Trend 2008-2017 ^c		Trend 2008-2017 ^d	
	Rate per 100,000 persons	Males	AAPC (%)	Males	AAPC (%)	Males
SEER Incidence						
All Races	109.8		-3.4*		-4.3*	
White	102.3		-4.1*		-4.2*	
White Hispanic ^e	85.2		-6.2*		-	
White Non-Hispanic ^e	105.2		-3.8*		-	
Black	175.2		-3.4*		-3.8*	
Asian/Pacific Islander	56.7		-4.3*		-	
Amer Ind/Alaska Nat ^f	54.6		-6.0*		-	
Hispanic ^e	92.0		-5.1*		-	
U.S. Mortality^g						
	Rate 2013-2017		Trend 2008-2017 ^h			
	Rate per 100,000 persons		AAPC (%)			
Race/Ethnicity						
All Races	19.1		-2.1*			
White	17.9		-1.9*			
White Hispanic ^e	16.4		-1.9*			
White Non-Hispanic ^e	18.0		-1.8*			
Black	37.9		-3.3*			
Asian/Pacific Islander	8.6		-2.3*			
Amer Ind/Alaska Nat						
Total U.S.	13.6		-2.9*			
PRCDA Counties	18.7		-1.7*			
Non-PRCDA Counties	8.0		-4.8*			
Hispanic ^e	15.8		-2.0*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

a Trend based on less than 10 cases for at least one year within the time interval.

b Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

c The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

d The SEER 21 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts.

e The 2008-2017 AAPC estimates are based on a Joinpoint analysis with up to 3 Joinpoints over diagnosis years 2000-2017.

f The 2008-2017 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2017.

g Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

h Incidence data for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

i US Mortality Files, National Center for Health Statistics, CDC.

j The 2008-2017 mortality AAPCs are based on a Joinpoint analysis using years of death 2000-2017.

* The APC is significantly different from zero ($p < .05$).

Table 23.12
Cancer of the Prostate (Invasive)

Age-Adjusted SEER Incidence Rates^a
By Registry, Race and Age

	All Races			Whites			Blacks		
	All Ages	Ages <65	Ages 65+	All Ages	Ages <65	Ages 65+	All Ages	Ages <65	Ages 65+
<u>SEER Incidence Rates^a, 2013-2017</u>									
California	92.96	34.18	499.23	89.69	32.43	485.54	137.94	66.82	629.56
Greater Bay Area	92.24	35.09	487.27	94.43	36.02	498.18	142.59	70.71	639.46
San Francisco-Oakland	93.54	36.24	489.60	94.21	36.14	495.66	145.14	73.30	641.68
San Jose-Monterey	89.81	32.98	482.66	94.77	35.81	502.33	130.36	58.84	624.73
Los Angeles	90.32	33.32	484.30	84.30	29.77	461.20	134.70	65.63	612.14
Greater California	94.33	34.24	509.70	90.35	32.44	490.65	138.95	66.22	641.68
Connecticut	111.30	45.34	567.20	104.40	41.08	542.09	164.26	76.49	770.99
Detroit	128.17	56.33	624.80	116.25	47.33	592.66	178.25	93.46	764.35
Georgia	124.19	51.78	624.77	103.41	38.69	550.76	193.32	87.38	925.56
Atlanta	139.81	57.37	709.61	115.89	42.04	626.36	200.36	88.78	971.67
Rural Georgia	133.10	64.31	608.65	114.36	53.50	535.11	177.35	83.32	827.34
Greater Georgia	117.26	48.54	592.29	99.20	37.06	528.74	189.00	86.37	898.40
Hawaii	88.22	27.93	504.92	86.25	29.31	479.91	162.54	103.38	571.45
Idaho	105.28	37.10	576.58	102.34	35.77	562.49	139.68	57.71	-
Iowa	107.70	42.87	555.84	106.13	41.91	550.02	174.64	85.26	792.46
Kentucky	104.08	43.34	523.93	92.94	37.13	478.67	144.86	65.65	692.38
Louisiana	131.22	56.71	646.28	115.24	46.89	587.69	179.97	82.85	851.26
Massachusetts	102.42	44.97	499.53	95.92	41.41	472.69	158.25	77.74	714.75
New Jersey	131.25	53.74	667.04	120.50	47.74	623.41	187.07	84.33	897.24
New Mexico	82.78	31.55	436.88	80.60	30.83	424.61	113.12	61.73	468.35
New York	124.95	52.38	626.60	114.67	46.19	588.01	193.51	90.12	908.18
Seattle-Puget Sound	108.70	39.32	588.28	108.96	38.61	595.24	160.13	67.73	798.86
Utah	112.76	41.27	606.92	112.70	41.56	604.45	154.68	63.29	786.42
SEER 9 Areas ^b	109.37	43.36	565.68	104.67	39.96	551.97	175.27	84.14	805.20
SEER 13 Areas ^b	104.09	40.51	543.56	100.09	37.62	531.95	165.17	79.71	755.90
SEER 18 Areas ^b	107.03	42.02	556.38	100.06	37.80	530.42	169.94	79.67	793.88
SEER 21 Areas ^b	109.80	43.91	565.32	102.26	39.40	536.76	175.22	81.99	819.68

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.
The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

The SEER 21 areas comprise the SEER 18 areas plus Idaho, New York and Massachusetts.

^b Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 23.13
Cancer of the Prostate (Invasive)

Age-Adjusted SEER Death Rates^a
By Registry, Race and Age

	All Races			Whites			Blacks		
	All Ages	Ages <65	Ages 65+	All Ages	Ages <65	Ages 65+	All Ages	Ages <65	Ages 65+
SEER Death Rates^a, 2013-2017									
California	19.76	1.59	145.34	20.45	1.53	151.21	41.31	4.57	295.23
Greater Bay Area	16.87	1.23	125.05	18.44	1.19	137.69	39.73	4.43	283.74
San Francisco-Oakland	17.07	1.27	126.28	17.75	1.13	132.64	41.46	4.45	297.26
San Jose-Monterey	16.50	1.15	122.60	19.62	1.28	146.40	28.35	-	195.54
Los Angeles	19.63	1.60	144.28	19.57	1.37	145.37	45.04	5.03	321.66
Greater California	20.85	1.72	153.04	21.34	1.70	157.17	38.37	4.26	274.10
Connecticut	17.91	1.40	132.06	17.25	1.21	128.12	30.77	3.85	216.80
Detroit	19.41	1.91	140.38	16.05	1.31	117.91	35.21	4.30	248.93
Georgia	22.01	1.99	160.36	17.43	1.44	127.98	42.91	3.60	314.67
Atlanta	24.37	2.02	178.84	18.12	1.37	133.87	44.40	3.41	327.74
Rural Georgia	31.07	-	231.85	22.32	-	170.06	58.00	-	432.50
Greater Georgia	20.83	1.98	151.10	17.04	1.47	124.65	41.47	3.74	302.23
Hawaii	14.04	1.27	102.32	19.82	1.30	147.89	42.06	-	-
Idaho	22.94	1.35	172.13	23.08	1.32	173.52	-	-	-
Iowa	19.54	1.45	144.52	19.53	1.45	144.53	26.61	-	193.86
Kentucky	19.55	1.45	144.67	18.85	1.37	139.66	35.24	2.77	259.65
Louisiana	20.78	2.04	150.30	17.04	1.35	125.50	34.79	3.90	248.28
Massachusetts	18.42	1.37	136.31	18.24	1.37	134.89	32.57	2.22	242.37
New Jersey	17.78	1.37	131.19	16.31	1.17	121.02	38.92	3.37	284.64
New Mexico	19.59	1.44	145.02	19.56	1.50	144.43	24.07	-	186.28
New York	18.03	1.56	131.90	16.86	1.35	124.13	32.61	3.23	235.68
Seattle-Puget Sound	19.74	1.58	145.28	20.17	1.55	148.89	28.76	3.56	202.99
Utah	20.05	1.34	149.33	20.35	1.36	151.59	-	-	-
SEER 9 Areas ^b	19.01	1.54	139.76	18.59	1.36	137.73	37.23	3.74	268.72
SEER 13 Areas ^b	19.02	1.53	139.92	18.84	1.36	139.63	39.35	4.03	283.49
SEER 18 Areas ^b	19.57	1.61	143.69	18.98	1.43	140.32	38.73	3.87	279.66
SEER 21 Areas ^b	19.27	1.59	141.50	18.62	1.41	137.61	36.92	3.67	266.82
Total U.S.	19.12	1.66	139.78	17.92	1.42	131.96	37.86	3.96	272.24

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

The SEER 21 areas comprise the SEER 18 areas plus Idaho, New York and Massachusetts.


- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 23.14
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2013-2017

Males									
State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	19.12	0.05							
High Five States									
D.C.	28.07	1.54	01(01,03)	46.83 ^c					
Mississippi	24.67	0.63	02(01,04)	29.03 ^c					
Idaho	22.94	0.78	03(02,13)	19.98 ^c					
Montana	22.69	0.90	04(02,18)	18.69 ^c					
Georgia	22.01	0.35	05(03,10)	15.11 ^c					
Low Five States									
West Virginia	17.39	0.59	47(28,50)	-9.03					
Florida	16.71	0.16	48(45,50)	-12.58 ^c					
Delaware	16.59	0.83	49(27,51)	-13.23 ^c					
Wyoming	15.94	1.09	50(28,51)	-16.62 ^c					
Hawaii	14.04	0.59	51(49,51)	-26.58 ^c					

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	21.15	0.44	08(04,21)	10.62 ^c	Montana	22.69	0.90	04(02,18)	18.69 ^c
Alaska	18.01	1.38	40(04,51)	-5.79	Nebraska	18.35	0.63	38(13,49)	-4.02
Arizona	17.47	0.31	46(35,49)	-8.60	Nevada	19.77	0.57	22(07,39)	3.43
Arkansas	18.55	0.51	36(16,48)	-2.98	New Hampshire	18.89	0.75	31(08,48)	-1.20
California	19.76	0.15	23(13,30)	3.37	New Jersey	17.78	0.29	43(32,48)	-7.00
Colorado	21.47	0.45	07(03,18)	12.33 ^c	New Mexico	19.59	0.62	24(07,43)	2.47
Connecticut	17.91	0.44	41(27,49)	-6.31	New York	18.03	0.20	39(32,46)	-5.67
Delaware	16.59	0.83	49(27,51)	-13.23 ^c	North Carolina	19.87	0.30	19(10,33)	3.93
D.C.	28.07	1.54	01(01,03)	46.83 ^c	North Dakota	17.76	0.98	44(11,50)	-7.11
Florida	16.71	0.16	48(45,50)	-12.58 ^c	Ohio	19.08	0.26	29(19,38)	-0.18
Georgia	22.01	0.35	05(03,10)	15.11 ^c	Oklahoma	20.35	0.48	13(06,32)	6.44
Hawaii	14.04	0.59	51(49,51)	-26.58 ^c	Oregon	20.68	0.45	11(05,28)	8.15
Idaho	22.94	0.78	03(02,13)	19.98 ^c	Pennsylvania	18.75	0.23	32(24,40)	-1.90
Illinois	20.14	0.27	15(09,28)	5.33	Rhode Island	18.70	0.84	33(08,50)	-2.20
Indiana	19.46	0.37	27(12,37)	1.80	South Carolina	21.81	0.45	06(03,13)	14.09 ^c
Iowa	19.54	0.49	26(09,40)	2.19	South Dakota	18.96	0.93	30(05,50)	-0.84
Kansas	18.60	0.52	35(15,47)	-2.72	Tennessee	19.83	0.37	21(09,34)	3.71
Kentucky	19.55	0.45	25(09,39)	2.25	Texas	17.90	0.19	42(34,47)	-6.36
Louisiana	20.78	0.46	10(04,27)	8.70	Utah	20.05	0.64	16(05,39)	4.85
Maine	20.41	0.73	12(04,38)	6.75	Vermont	19.10	1.08	28(05,50)	-0.10
Maryland	20.02	0.40	17(08,33)	4.69	Virginia	19.83	0.34	20(10,33)	3.71
Massachusetts	18.42	0.34	37(25,46)	-3.64	Washington	20.19	0.36	14(08,30)	5.59
Michigan	18.69	0.28	34(24,42)	-2.24	West Virginia	17.39	0.59	47(28,50)	-9.03
Minnesota	19.91	0.39	18(08,34)	4.14	Wisconsin	20.84	0.38	09(05,24)	8.98
Mississippi	24.67	0.63	02(01,04)	29.03 ^c	Wyoming	15.94	1.09	50(28,51)	-16.62 ^c
Missouri	17.50	0.34	45(34,49)	-8.45					



Age-Adjusted Death Rates per 100,000 Quantile Interval

■	20.85 - 28.07
■	20.03 - 20.84
■	19.56 - 20.02
■	18.70 - 19.55
■	17.79 - 18.69
■	14.04 - 17.78
■	Suppressed
U.S. rate: 19.12	

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 23.15
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2017
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to 5	> 5 to 10	> 10 to 15	> 15 to 20	> 20 to 25	<=25 ^b	> 25 ^d	Complete ^e
<u>Race</u>								
All Races	900,214	953,463	678,436	418,862	184,721	3,135,696	34,643	3,170,339
White	701,239	769,223	563,050	354,061	159,744	2,547,317	36,576	2,583,893
Black	138,010	131,943	86,907	50,644	20,399	427,902	4,086	431,988
Asian/Pacific Islander	21,319	20,629	13,351	6,827	2,333	64,458	+	+
Hispanic	59,273	57,004	38,768	20,529	7,678	183,252	+	+

Estimated prevalence percent^a on January 1, 2017, of the SEER population diagnosed in the previous 25 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)									Age-Adjusted ^c All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	
<u>Race</u>										
All Races	1.9646	0.0001	0.0002	0.0003	0.0007	0.0628	0.9938	4.9473	12.7506	17.7721
White	2.0311	0.0002	0.0003	0.0003	0.0005	0.0480	0.8646	4.5811	12.2431	17.3251
Black	1.9574	-	-	0.0002	0.0017	0.1674	1.9500	8.2063	18.4635	22.7226
Asian/Pacific Islander	0.6447	-	-	-	-	0.0144	0.2968	1.9438	5.7590	9.4841
Hispanic	0.6240	0.0002	0.0002	0.0001	0.0005	0.0286	0.5448	3.1347	9.2044	13.7068

^a US 2017 cancer prevalence counts are based on 2017 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2017 US population estimates based on the average of 2016 and 2017 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 25 years. Maximum limited-duration prevalence is 25 years for 1992-2017 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 25 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).


^e Complete prevalence is obtained by summing <=25 and > 25.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Figure 23.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Prostate, by Race



^a Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 23.2

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Prostate, by Age and Race



^a Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.


* The APC is significantly different from zero ($p < 0.05$).

Figure 23.3

SEER Incidence and US Death Rates^a

Cancer of the Prostate

Joinpoint Analyses for Whites and Blacks from 1975-2017
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 2000-2017



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 21 areas (SEER 9 areas, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2017 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 2000-2017 allow a maximum of 3 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.


^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 23.4

Cancer of the Prostate

Delay-Adjusted SEER Incidence & US Mortality


1975-2017



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).
 Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines are calculated using the Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute.

Figure 23.5


Delay-Adjusted Incidence Rates for Cancer of the Prostate and Cancer of the Lung and Bronchus All Races, Males 1975-2017



Source: SEER 9 areas. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.8, April 2020, National Cancer Institute.

Figure 23.6

Cancer of the Prostate
5-Year SEER Conditional Relative Survival and
95% Confidence Intervals
 Probability of surviving the next 5 years given the cohort
 has already survived 0, 1, or 3 years
 2000-2015 by stage at diagnosis



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Percent surviving is not shown if based on less than 25 cases.

Confidence intervals are not shown if length of the confidence interval is greater than 5 times the standard error.