

Table 1.22  
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2010<sup>a</sup>  
Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			35-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 35 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 35 Years <sup>d</sup>
All Sites	Both Sexes	4,511,266	4,598,344	5,044,168	12,600,627	12,810,577
	Male	2,332,050	2,366,640	2,584,260	5,990,343	6,060,178
	Female	2,179,216	2,231,704	2,459,908	6,610,284	6,750,399
Oral Cavity & Pharynx	Both Sexes	102,019	117,112	121,258	264,620	291,343
	Male	71,123	80,929	83,544	175,045	190,951
	Female	30,896	36,183	37,714	89,575	100,392
Esophagus	Both Sexes	21,268	25,973	26,092	33,702	39,594
	Male	16,808	20,399	20,447	26,317	30,715
	Female	4,460	5,574	5,645	7,385	8,879
Stomach	Both Sexes	35,420	42,280	42,609	70,821	81,142
	Male	21,286	25,765	25,901	40,560	46,844
	Female	14,134	16,515	16,708	30,261	34,298
Colon & Rectum	Both Sexes	407,833	471,251	480,046	1,129,666	1,258,620
	Male	207,302	239,824	244,132	561,245	622,211
	Female	200,531	231,427	235,914	568,421	636,409
Liver & Intrahepatic Bile Duct	Both Sexes	30,577	34,834	34,867	40,940	45,832
	Male	22,266	25,232	25,265	28,891	32,221
	Female	8,311	9,602	9,602	12,049	13,611
Pancreas	Both Sexes	31,442	38,275	38,275	41,284	49,174
	Male	15,733	19,467	19,467	20,155	24,367
	Female	15,709	18,808	18,808	21,129	24,807
Larynx	Both Sexes	30,947	37,152	37,518	87,135	98,063
	Male	25,107	30,036	30,356	70,117	78,586
	Female	5,840	7,116	7,162	17,018	19,477
Lung & Bronchus	Both Sexes	231,996	296,283	304,764	391,564	480,133
	Male	108,211	140,119	143,487	178,912	220,472
	Female	123,785	156,164	161,277	212,652	259,661
Melanoma of the Skin	Both Sexes	276,312	312,334	325,566	884,259	953,581
	Male	148,749	171,783	180,200	440,298	480,917
	Female	127,563	140,551	145,366	443,961	472,664
Breast	Female	842,193	909,424	965,835	2,770,060	2,934,536
Cervix	Female	38,963	40,783	40,907	208,065	213,819
Corpus & Uterus, NOS	Female	169,126	189,983	190,106	569,448	620,803
Ovary <sup>f</sup>	Female	57,207	66,063	66,139	169,398	190,511

<sup>a</sup> U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 35 years (1975-2009)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2005-2009)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2005; Melanoma in 2006.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 35-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2010.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 35-year limited duration prevalence. The 2005 breast cancer is counted for the breast 5-year and 35-year limited duration prevalence.

In method (e) the 2005 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2006 melanoma is counted for 5-year limited duration prevalence for melanoma.

<sup>f</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.22 - continued  
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2010<sup>a</sup>  
 Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			35-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 35 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 35 Years <sup>d</sup>
Prostate	Male	1,026,391	1,107,903	1,107,977	2,616,190	2,787,286
Testis	Male	42,302	43,012	43,616	205,130	207,844
Urinary Bladder	Both Sexes	198,998	248,244	253,143	551,278	641,106
	Male	151,489	190,075	194,009	411,638	479,012
	Female	47,509	58,169	59,134	139,640	162,094
Kidney & Renal Pelvis	Both Sexes	145,766	177,066	179,051	330,450	385,370
	Male	88,557	108,930	110,317	196,335	230,734
	Female	57,209	68,136	68,734	134,115	154,636
Brain & Nervous System	Both Sexes	43,995	46,883	47,345	128,063	132,164
	Male	24,044	25,790	26,033	68,158	70,466
	Female	19,951	21,093	21,312	59,905	61,698
Thyroid	Both Sexes	165,253	183,524	184,093	494,969	528,131
	Male	37,076	42,905	43,058	107,747	117,569
	Female	128,177	140,619	141,035	387,222	410,562
Hodgkin Lymphoma	Both Sexes	39,048	41,589	41,611	168,309	172,937
	Male	21,122	22,557	22,568	87,084	89,522
	Female	17,926	19,032	19,043	81,225	83,415
Non-Hodgkin Lymphoma	Both Sexes	198,946	233,585	235,457	499,712	558,340
	Male	105,553	125,451	126,442	261,560	292,961
	Female	93,393	108,134	109,015	238,152	265,379
Myeloma	Both Sexes	49,451	58,292	58,328	77,360	88,490
	Male	26,622	32,063	32,087	42,092	48,802
	Female	22,829	26,229	26,241	35,268	39,688
Leukemia	Both Sexes	112,120	129,298	129,528	282,150	310,046
	Male	64,915	75,612	75,711	160,039	176,801
	Female	47,205	53,686	53,817	122,111	133,245
Acute Lymphocytic Leukemia	Both Sexes	16,205	16,686	16,686	66,030	66,759
	Male	9,215	9,443	9,443	36,418	36,708
	Female	6,990	7,243	7,243	29,612	30,051
Childhood (Ages 0-19)	Both Sexes	63,904	63,977	64,503	304,247	304,716
	Male	33,948	33,981	34,250	156,819	157,032
	Female	29,956	29,996	30,253	147,428	147,684
Kaposi Sarcoma	Both Sexes	7,319	7,750	7,750	24,815	25,977
	Male	6,865	7,208	7,208	23,533	24,518
	Female	454	542	542	1,282	1,459
Mesothelioma	Both Sexes	2,968	3,926	3,938	4,538	5,658
	Male	1,998	2,761	2,773	2,601	3,467
	Female	970	1,165	1,165	1,937	2,191

<sup>a</sup> U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever  
<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 35 years (1975-2009)  
<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2005-2009)  
 For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2005; Melanoma in 2006.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 35-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2010. In method (d) the 1981 melanoma is counted for the melanoma and all sites 35-year limited duration prevalence. The 2005 breast cancer is counted for the breast 5-year and 35-year limited duration prevalence. In method (e) the 2005 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2006 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.23  
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2010<sup>a</sup>  
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	6,078,974	18,390	42,087	83,873	155,608	354,801	860,143	1,578,142	2,985,931
Females	6,948,940	16,117	36,351	86,197	226,859	636,726	1,276,932	1,684,808	2,984,950
Oral Cavity & Pharynx									
Males	181,084	68	468	1,480	3,518	14,224	43,847	54,509	62,971
Females	94,109	90	553	1,470	3,635	8,681	17,691	22,821	39,167
Esophagus									
Males	26,388	0	0	22	219	1,177	4,653	9,419	10,897
Females	7,451	0	0	0	28	256	1,263	1,790	4,114
Stomach									
Males	41,386	0	18	64	607	2,589	7,095	10,242	20,773
Females	30,883	4	12	165	568	2,307	4,398	6,312	17,118
Colon & Rectum									
Males	569,930	11	117	1,190	5,632	25,029	80,675	136,002	321,273
Females	584,551	0	75	1,251	5,467	23,433	67,578	110,244	376,504
Liver & Intrahep									
Males	28,967	484	535	554	454	1,582	10,000	9,083	6,274
Females	12,437	471	470	312	411	895	2,803	3,028	4,048
Pancreas									
Males	20,293	11	12	127	376	1,616	4,320	6,183	7,649
Females	21,316	0	66	155	413	1,376	3,843	5,483	9,980
Larynx									
Males	71,659	0	0	78	243	1,915	10,556	20,739	38,129
Females	17,370	0	0	25	134	1,011	3,256	4,677	8,267
Lung & Bronchus									
Males	183,410	45	90	444	1,184	5,545	24,951	54,888	96,262
Females	216,021	31	71	396	1,435	7,883	28,898	59,047	118,261
Melanoma of the Skin									
Males	452,540	78	656	5,014	18,294	46,766	95,157	122,709	163,865
Females	469,240	113	993	10,894	33,972	73,514	109,961	106,450	133,343

<sup>a</sup> U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued  
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2010<sup>a</sup>  
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	14,588	0	0	11	89	586	2,018	4,133	7,751
Females	2,829,041	0	53	2,740	36,291	216,912	539,229	767,846	1,265,969
Cervix									
Females	249,496	0	45	2,239	16,405	42,748	59,766	55,248	73,044
Corpus & Uterus, NOS									
Females	600,346	0	47	530	5,666	24,621	87,167	157,833	324,482
Ovary <sup>d</sup>									
Females	186,138	84	1,018	3,453	6,883	18,736	40,700	47,632	67,632
Prostate									
Males	2,617,682	34	59	101	324	21,033	228,561	740,428	1,627,144
Urinary Bladder									
Males	419,648	45	105	611	2,438	11,535	42,898	99,644	262,372
Females	143,992	23	57	217	964	4,266	13,698	30,030	94,737
Kidney & Renal Pelvis									
Males	202,516	1,454	2,505	2,659	5,172	17,901	40,060	57,511	75,253
Females	138,989	1,343	2,681	2,851	4,901	12,071	23,973	33,716	57,452
Hodgkin Lymphoma									
Males	93,890	181	2,315	9,272	16,487	24,007	21,438	13,274	6,916
Females	88,038	101	1,901	9,617	16,530	23,444	18,638	10,651	7,157
Non-Hodgkin Lymphoma									
Males	266,478	773	3,513	7,582	13,366	29,224	52,496	67,493	92,032
Females	242,587	430	1,847	4,348	8,879	20,902	41,030	56,723	108,427
Myeloma									
Males	42,185	0	6	84	442	2,859	8,141	13,608	17,045
Females	35,432	0	0	39	281	2,266	6,273	10,676	15,896
Leukemia									
Males	162,651	6,545	12,947	12,388	10,928	13,500	22,064	32,827	51,452
Females	125,312	5,438	10,680	10,750	9,514	9,728	14,680	20,623	43,899
Acute Lymphocytic Leuk									
Males	38,239	5,461	11,081	9,597	6,075	3,568	1,238	775	444
Females	31,328	4,527	9,014	7,859	5,163	2,755	884	717	408

<sup>a</sup> U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.21  
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	4,511,266	3,100,895	2,115,695	1,341,528	776,939	454,693	11,069,385	12,600,627	427,287	13,027,914
	Males	2,332,050	1,572,925	998,128	593,178	268,188	144,398	5,496,281	5,990,343	88,631	6,078,974
	Females	2,179,216	1,527,970	1,117,567	748,350	508,751	310,295	5,573,104	6,610,284	338,656	6,948,940
White <sup>b</sup>	Both Sexes	3,822,952	2,699,375	1,859,648	1,200,350	705,111	415,002	9,582,325	10,975,277	375,017	11,350,294
	Males	1,968,442	1,361,458	872,314	532,143	245,842	133,749	4,734,357	5,189,363	82,898	5,272,261
	Females	1,854,510	1,337,917	987,334	668,207	459,269	281,253	4,847,968	5,785,914	292,119	6,078,033
Black <sup>b</sup>	Both Sexes	454,240	274,223	172,669	94,496	45,620	26,053	995,627	1,084,133	25,176	1,109,309
	Males	244,991	152,451	89,843	43,171	14,121	6,671	530,455	554,560	2,277	556,837
	Females	209,249	121,772	82,826	51,325	31,499	19,382	465,172	529,573	22,899	552,472
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	131,655	80,549	49,631	+	+	+	280,924	+	+	+
	Males	56,813	33,143	18,552	+	+	+	115,176	+	+	+
	Females	74,842	47,406	31,079	+	+	+	165,748	+	+	+
Hispanic <sup>d</sup>	Both Sexes	298,868	183,359	113,698	+	+	+	595,926	+	+	+
	Males	143,051	86,170	50,138	+	+	+	279,359	+	+	+
	Females	155,817	97,189	63,560	+	+	+	316,567	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	2.8371%	0.0848%	0.1635%	0.2848%	0.7167%	1.7097%	3.8957%	8.4802%	13.8744%	14.4502%	2.8019%
	Males	2.8615%	0.0897%	0.1723%	0.2731%	0.5580%	1.1719%	3.2482%	9.4103%	18.3650%	20.6809%	3.1737%
	Females	2.8134%	0.0796%	0.1543%	0.2970%	0.8759%	2.2425%	4.5132%	7.6391%	10.2898%	10.8412%	2.5744%
White <sup>c</sup>	Both Sexes	3.1221%	0.0930%	0.1823%	0.3226%	0.8024%	1.8281%	4.0711%	8.8786%	14.5767%	14.8617%	2.9449%
	Males	3.1171%	0.0984%	0.1936%	0.3096%	0.6357%	1.2672%	3.3618%	9.6546%	18.9306%	20.8081%	3.2756%
	Females	3.1271%	0.0874%	0.1705%	0.3365%	0.9776%	2.4052%	4.7703%	8.1555%	11.0145%	11.3939%	2.7498%
Black <sup>c</sup>	Both Sexes	2.1287%	0.0592%	0.1108%	0.1776%	0.4923%	1.4068%	3.7539%	8.8626%	13.8728%	12.9297%	2.6643%
	Males	2.3998%	0.0631%	0.1105%	0.1696%	0.3486%	0.9964%	3.7865%	12.0610%	21.6295%	21.9777%	3.5887%
	Females	1.8825%	0.0551%	0.1111%	0.1851%	0.6172%	1.7701%	3.7258%	6.3343%	8.5405%	8.6493%	2.0848%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1.8066%	0.0648%	0.1125%	0.1714%	0.4444%	1.2280%	2.7556%	5.1339%	8.6692%	10.6803%	1.8554%
	Males	1.6079%	0.0694%	0.1175%	0.1589%	0.2824%	0.6460%	1.7332%	4.8329%	10.7492%	15.4330%	1.8966%
	Females	1.9885%	0.0600%	0.1072%	0.1836%	0.5898%	1.7450%	3.6315%	5.3847%	7.0857%	7.7430%	1.8755%
Hispanic <sup>d</sup>	Both Sexes	1.2175%	0.0795%	0.1532%	0.2142%	0.5087%	1.1942%	2.7535%	5.9179%	10.0925%	11.2607%	2.0455%
	Males	1.1171%	0.0848%	0.1630%	0.2108%	0.3928%	0.7214%	2.0261%	6.3201%	14.0475%	16.7861%	2.3340%
	Females	1.3203%	0.0740%	0.1430%	0.2180%	0.6319%	1.6852%	3.4420%	5.5772%	7.1880%	7.9935%	1.9028%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 3.21  
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	43,995	26,358	19,018	15,886	11,066	6,626	105,257	128,063	13,490	141,553
	Males	24,044	13,517	10,015	8,797	5,700	3,526	56,373	68,158	6,195	74,353
	Females	19,951	12,841	9,003	7,089	5,366	3,100	48,884	59,905	7,295	67,200
White <sup>b</sup>	Both Sexes	38,023	23,231	16,619	13,897	9,867	5,933	91,769	112,206	4,836	117,042
	Males	21,245	12,032	8,807	7,635	5,000	3,185	49,718	60,252	2,663 <sup>i</sup>	62,915
	Females	16,778	11,199	7,812	6,262	4,867	2,748	42,051	51,954	2,173 <sup>i</sup>	54,127
Black <sup>b</sup>	Both Sexes	3,719	2,011	1,493	1,299	859	499	8,522	10,240	+	+
	Males	1,683	934	801	738	489	245	4,156	5,042	+	+
	Females	2,036	1,077	692	561	370	254	4,366	5,198	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1,249	723	556	+	+	+	2,779	+	+	+
	Males	610	356	253	+	+	+	1,363	+	+	+
	Females	639	367	303	+	+	+	1,416	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,461	2,908	1,988	+	+	+	9,357	+	+	+
	Males	2,460	1,551	989	+	+	+	5,000	+	+	+
	Females	2,001	1,357	999	+	+	+	4,357	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0266%	0.0141%	0.0295%	0.0242%	0.0263%	0.0316%	0.0326%	0.0321%	0.0263%	0.0165%	0.0264%
	Males	0.0289%	0.0140%	0.0312%	0.0253%	0.0284%	0.0349%	0.0366%	0.0367%	0.0293%	0.0186%	0.0286%
	Females	0.0244%	0.0142%	0.0277%	0.0231%	0.0242%	0.0282%	0.0288%	0.0279%	0.0239%	0.0153%	0.0243%
White <sup>c</sup>	Both Sexes	0.0302%	0.0158%	0.0334%	0.0276%	0.0302%	0.0369%	0.0368%	0.0357%	0.0291%	0.0170%	0.0299%
	Males	0.0331%	0.0159%	0.0361%	0.0288%	0.0324%	0.0413%	0.0412%	0.0409%	0.0330%	0.0199%	0.0327%
	Females	0.0273%	0.0157%	0.0305%	0.0263%	0.0279%	0.0324%	0.0324%	0.0308%	0.0260%	0.0154%	0.0273%
Black <sup>c</sup>	Both Sexes	0.0169%	0.0091%	0.0205%	0.0169%	0.0165%	0.0162%	0.0207%	0.0221%	0.0160%	0.0146%	0.0169%
	Males	0.0166%	0.0091%	0.0195%	0.0163%	0.0169%	0.0147%	0.0223%	0.0242%	0.0118%	-	0.0164%
	Females	0.0173%	0.0091%	0.0215%	0.0176%	0.0162%	0.0175%	0.0195%	0.0204%	0.0189%	0.0168%	0.0172%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0150%	0.0097%	0.0171%	0.0140%	0.0143%	0.0161%	0.0188%	0.0174%	0.0141%	0.0114%	0.0149%
	Males	0.0151%	0.0090%	0.0160%	0.0149%	0.0157%	0.0154%	0.0197%	0.0175%	0.0140%	0.0114%	0.0150%
	Females	0.0149%	0.0104%	0.0183%	0.0131%	0.0129%	0.0167%	0.0180%	0.0174%	0.0141%	0.0115%	0.0149%
Hispanic <sup>d</sup>	Both Sexes	0.0189%	0.0103%	0.0223%	0.0140%	0.0177%	0.0249%	0.0279%	0.0245%	0.0279%	0.0165%	0.0202%
	Males	0.0199%	0.0097%	0.0245%	0.0136%	0.0185%	0.0276%	0.0308%	0.0283%	0.0277%	0.0155%	0.0215%
	Females	0.0179%	0.0109%	0.0199%	0.0144%	0.0167%	0.0222%	0.0251%	0.0213%	0.0280%	0.0171%	0.0189%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 4.25  
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	847,990	682,322	520,032	338,523	215,193	111,063	2,388,868	2,784,496	59,133	2,843,629
	Males	5,797	3,879	2,093	1,249	771	417	13,018	14,436	152	14,588
	Females	842,193	678,443	517,939	337,274	214,422	110,646	2,375,850	2,770,060	58,981	2,829,041
White <sup>b</sup>	Both Sexes	719,295	597,566	460,928	303,258	195,605	100,654	2,081,047	2,440,716	52,282	2,492,998
	Males	5,005	3,241	1,846	1,091	678	401	11,183	12,478	+ <sup>i</sup>	12,471
	Females	714,290	594,325	459,082	302,167	194,927	100,253	2,069,864	2,428,238	52,289	2,480,527
Black <sup>b</sup>	Both Sexes	85,890	56,331	39,352	24,087	13,669	7,561	205,661	230,970	+	+
	Males	610	506	160	97	85	12	1,374	1,483	+	+
	Females	85,280	55,825	39,192	23,990	13,584	7,549	204,287	229,487	4,397	233,884
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	31,091	21,837	14,564	+	+	+	72,975	+	+	+
	Males	137	66	43	+	+	+	280	+	+	+
	Females	30,954	21,771	14,521	+	+	+	72,695	+	+	+
Hispanic <sup>d</sup>	Both Sexes	57,058	38,748	25,302	+	+	+	121,107	+	+	+
	Males	259	140	52	+	+	+	451	+	+	+
	Females	56,799	38,608	25,250	+	+	+	120,656	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.6089%	-	0.0001%	0.0061%	0.0863%	0.4609%	1.1010%	1.9094%	2.5026%	2.8320%	0.5873%
	Males	0.0068%	-	-	-	0.0004%	0.0027%	0.0083%	0.0239%	0.0408%	0.0588%	0.0075%
	Females	1.1951%	-	0.0002%	0.0124%	0.1723%	0.9149%	2.1431%	3.6146%	4.4677%	4.4383%	1.0793%
White <sup>c</sup>	Both Sexes	0.6688%	-	-	0.0057%	0.0855%	0.4655%	1.1356%	2.0230%	2.6788%	3.0054%	0.6158%
	Males	0.0075%	-	-	-	0.0004%	0.0025%	0.0080%	0.0250%	0.0436%	0.0624%	0.0078%
	Females	1.3265%	-	-	0.0117%	0.1748%	0.9420%	2.2471%	3.8849%	4.8349%	4.7217%	1.1455%
Black <sup>c</sup>	Both Sexes	0.4382%	-	-	0.0082%	0.0983%	0.4615%	1.0042%	1.6418%	2.1296%	2.4425%	0.5256%
	Males	0.0063%	-	-	-	-	0.0039%	0.0150%	0.0298%	0.0413%	0.0580%	0.0089%
	Females	0.8304%	-	-	0.0158%	0.1833%	0.8665%	1.8577%	2.9161%	3.5651%	3.5705%	0.9111%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.4567%	-	-	0.0061%	0.0818%	0.4423%	0.9828%	1.4105%	1.7091%	1.7846%	0.4530%
	Males	0.0036%	-	-	-	-	0.0021%	0.0049%	0.0112%	0.0239%	0.0292%	0.0042%
	Females	0.8717%	-	-	0.0119%	0.1548%	0.8333%	1.8207%	2.5765%	2.9922%	2.8696%	0.8140%
Hispanic <sup>d</sup>	Both Sexes	0.2501%	-	-	0.0040%	0.0620%	0.3194%	0.8044%	1.3380%	1.7204%	1.8922%	0.4133%
	Males	0.0018%	-	-	-	-	0.0013%	0.0040%	0.0138%	0.0220%	0.0239%	0.0038%
	Females	0.5045%	-	-	0.0085%	0.1276%	0.6499%	1.5621%	2.4598%	2.9676%	2.9970%	0.7515%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 5.15  
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	38,963	33,983	36,474	32,451	26,672	20,674	141,872	208,065	41,431	249,496
White <sup>b</sup>	30,795	27,060	29,144	26,982	22,554	17,087	113,981	168,980	35,075	204,055
Black <sup>b</sup>	5,250	4,714	4,710	3,551	2,736	2,710	18,225	26,334	+	+
Asian/Pacific Islander <sup>c</sup>	2,098	1,689	1,711	+	+	+	6,356	+	+	+
Hispanic <sup>d</sup>	8,092	7,461	6,298	+	+	+	21,851	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0766%	-	-	0.0096%	0.0773%	0.1554%	0.1552%	0.1319%	0.1229%	0.0747%	0.0737%	
White <sup>c</sup>	0.0792%	-	-	0.0114%	0.0884%	0.1702%	0.1555%	0.1234%	0.1109%	0.0616%	0.0763%	
Black <sup>c</sup>	0.0665%	-	-	0.0069%	0.0604%	0.1221%	0.1581%	0.1445%	0.1415%	0.1281%	0.0698%	
Asian/Pacific Islander <sup>c</sup>	0.0687%	-	-	0.0032%	0.0346%	0.1020%	0.1467%	0.1632%	0.1742%	0.1319%	0.0641%	
Hispanic <sup>d</sup>	0.0912%	-	-	0.0106%	0.0834%	0.2113%	0.2694%	0.2630%	0.2654%	0.2210%	0.1190%	

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 6.28  
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	407,833	280,179	185,283	117,126	75,687	40,740	990,422	1,129,666	24,815	1,154,481
	Males	207,302	141,885	91,940	57,815	36,542	17,157	498,942	561,245	8,685	569,930
	Females	200,531	138,294	93,343	59,311	39,145	23,583	491,480	568,421	16,130	584,551
White <sup>b</sup>	Both Sexes	336,917	239,200	159,963	102,083	66,617	36,354	838,164	961,429	21,884	983,313
	Males	173,422	122,564	80,173	50,857	32,595	15,447	427,017	482,801	7,763	490,564
	Females	163,495	116,636	79,790	51,226	34,022	20,907	411,147	478,628	14,121	492,749
Black <sup>b</sup>	Both Sexes	46,655	26,895	16,978	10,215	6,465	3,126	100,742	111,919	1,832	113,751
	Males	20,957	12,002	7,311	4,525	2,581	1,082	44,795	48,922	581	49,503
	Females	25,698	14,893	9,667	5,690	3,884	2,044	55,947	62,997	1,251	64,248
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	16,814	9,983	6,174	+	+	+	35,254	+	+	+
	Males	8,770	5,066	3,175	+	+	+	18,147	+	+	+
	Females	8,044	4,917	2,999	+	+	+	17,107	+	+	+
Hispanic <sup>d</sup>	Both Sexes	29,358	15,120	8,943	+	+	+	53,421	+	+	+
	Males	16,031	7,903	4,688	+	+	+	28,622	+	+	+
	Females	13,327	7,217	4,255	+	+	+	24,799	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.2614%	-	0.0003%	0.0053%	0.0251%	0.1005%	0.3250%	0.7145%	1.4138%	2.0252%	0.2591%
	Males	0.2685%	-	0.0004%	0.0052%	0.0257%	0.1068%	0.3626%	0.8487%	1.6776%	2.2478%	0.2983%
	Females	0.2546%	-	0.0003%	0.0054%	0.0244%	0.0943%	0.2892%	0.5931%	1.2032%	1.8962%	0.2276%
White <sup>c</sup>	Both Sexes	0.2736%	-	0.0003%	0.0056%	0.0256%	0.0983%	0.3114%	0.6933%	1.4150%	2.0435%	0.2563%
	Males	0.2810%	-	0.0003%	0.0053%	0.0264%	0.1026%	0.3510%	0.8307%	1.6819%	2.2624%	0.2957%
	Females	0.2663%	-	0.0003%	0.0059%	0.0248%	0.0938%	0.2724%	0.5652%	1.1966%	1.9159%	0.2238%
Black <sup>c</sup>	Both Sexes	0.2184%	-	-	0.0040%	0.0213%	0.1130%	0.3874%	0.8754%	1.5139%	1.9608%	0.2836%
	Males	0.2073%	-	-	0.0043%	0.0183%	0.1159%	0.4014%	0.9896%	1.6714%	2.0416%	0.3066%
	Females	0.2285%	-	-	0.0038%	0.0239%	0.1104%	0.3754%	0.7851%	1.4056%	1.9225%	0.2674%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.2362%	-	-	0.0041%	0.0239%	0.0965%	0.3314%	0.6903%	1.3093%	1.8890%	0.2470%
	Males	0.2519%	-	-	0.0051%	0.0260%	0.1154%	0.3695%	0.8182%	1.5943%	2.1933%	0.2919%
	Females	0.2218%	-	-	0.0032%	0.0220%	0.0798%	0.2988%	0.5838%	1.0923%	1.7009%	0.2129%
Hispanic <sup>d</sup>	Both Sexes	0.1094%	-	-	0.0035%	0.0205%	0.0736%	0.2555%	0.5982%	1.1051%	1.5088%	0.2042%
	Males	0.1148%	-	-	0.0036%	0.0223%	0.0754%	0.2869%	0.7358%	1.4419%	1.7804%	0.2482%
	Females	0.1038%	-	-	0.0035%	0.0186%	0.0718%	0.2257%	0.4817%	0.8578%	1.3482%	0.1713%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 7.15  
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	169,126	122,683	97,293	68,048	47,404	33,900	457,150	569,448	30,898	600,346
White <sup>b</sup>	146,097	109,780	88,760	63,323	44,222	32,089	407,960	513,677	26,391	540,068
Black <sup>b</sup>	13,514	7,005	4,350	2,378	1,764	816	27,246	30,652	860	31,512
Asian/Pacific Islander <sup>c</sup>	6,030	3,872	2,588	+	+	+	13,519	+	+	+
Hispanic <sup>d</sup>	11,595	7,293	4,739	+	+	+	23,627	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.2233%	-	-	0.0030%	0.0286%	0.1082%	0.3591%	0.8043%	0.9438%	0.7723%	0.2011%
White <sup>c</sup>	0.2542%	-	-	0.0025%	0.0283%	0.1103%	0.3816%	0.8801%	1.0543%	0.8514%	0.2184%
Black <sup>c</sup>	0.1131%	-	-	0.0043%	0.0242%	0.0676%	0.1859%	0.5083%	0.6896%	0.4512%	0.1287%
Asian/Pacific Islander <sup>c</sup>	0.1601%	-	-	0.0039%	0.0296%	0.1233%	0.3713%	0.5615%	0.4863%	0.3783%	0.1466%
Hispanic <sup>d</sup>	0.0987%	-	-	0.0043%	0.0340%	0.1071%	0.2848%	0.5784%	0.5676%	0.4666%	0.1455%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 8.21  
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	21,268	7,218	3,178	1,313	414	249	32,977	33,702	137	33,839
	Males	16,808	5,476	2,509	1,000	279	197	25,792	26,317	71	26,388
	Females	4,460	1,742	669	313	135	52	7,185	7,385	66	7,451
White <sup>b</sup>	Both Sexes	18,992	6,572	2,761	1,139	368	208	29,465	30,102	110	30,212
	Males	15,359	5,131	2,235	873	240	168	23,598	24,054	65	24,119
	Females	3,633	1,441	526	266	128	40	5,867	6,048	45	6,093
Black <sup>b</sup>	Both Sexes	1,753	446	300	133	39	39	2,632	2,711	34	2,745
	Males	1,061	196	189	96	39	28	1,541	1,609	9	1,618
	Females	692	250	111	37	0	11	1,091	1,102	25	1,127
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	364	114	70	+	+	+	578	+	+	+
	Males	261	75	49	+	+	+	410	+	+	+
	Females	103	39	21	+	+	+	168	+	+	+
Hispanic <sup>d</sup>	Both Sexes	833	229	124	+	+	+	1,186	+	+	+
	Males	716	184	103	+	+	+	1,003	+	+	+
	Females	117	45	21	+	+	+	183	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0085%	-	-	-	0.0005%	0.0026%	0.0119%	0.0318%	0.0484%	0.0385%	0.0083%
	Males	0.0133%	-	-	-	0.0009%	0.0044%	0.0193%	0.0555%	0.0802%	0.0734%	0.0142%
	Females	0.0037%	-	-	-	0.0002%	0.0009%	0.0049%	0.0104%	0.0230%	0.0183%	0.0034%
White <sup>c</sup>	Both Sexes	0.0098%	-	-	-	0.0007%	0.0030%	0.0135%	0.0355%	0.0531%	0.0403%	0.0092%
	Males	0.0157%	-	-	-	0.0012%	0.0051%	0.0217%	0.0629%	0.0894%	0.0783%	0.0159%
	Females	0.0040%	-	-	-	-	0.0009%	0.0054%	0.0099%	0.0235%	0.0181%	0.0035%
Black <sup>c</sup>	Both Sexes	0.0055%	-	-	-	-	0.0023%	0.0095%	0.0283%	0.0429%	0.0243%	0.0068%
	Males	0.0070%	-	-	-	-	0.0031%	0.0132%	0.0387%	0.0625%	0.0461%	0.0100%
	Females	0.0041%	-	-	-	-	0.0016%	0.0064%	0.0202%	0.0295%	0.0140%	0.0046%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0040%	-	-	-	-	0.0011%	0.0052%	0.0124%	0.0233%	0.0331%	0.0042%
	Males	0.0058%	-	-	-	-	0.0018%	0.0100%	0.0199%	0.0334%	0.0532%	0.0066%
	Females	0.0023%	-	-	-	-	-	-	0.0061%	0.0157%	0.0208%	0.0024%
Hispanic <sup>d</sup>	Both Sexes	0.0024%	-	-	-	0.0007%	0.0016%	0.0054%	0.0144%	0.0265%	0.0248%	0.0044%
	Males	0.0040%	-	-	-	0.0012%	0.0030%	0.0090%	0.0283%	0.0494%	0.0524%	0.0084%
	Females	0.0008%	-	-	-	-	-	0.0019%	0.0027%	0.0097%	0.0085%	0.0013%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 9.21  
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	39,048	31,542	27,862	24,612	20,519	14,967	123,063	168,309	13,619	181,928
	Males	21,122	16,396	14,182	12,479	10,024	8,020	64,179	87,084	6,806	93,890
	Females	17,926	15,146	13,680	12,133	10,495	6,947	58,884	81,225	6,813	88,038
White <sup>b</sup>	Both Sexes	32,588	27,082	24,516	22,107	19,002	13,946	106,294	148,513	15,113	163,626
	Males	17,710	14,158	12,461	11,292	9,244	7,318	55,622	76,721	7,802	84,523
	Females	14,878	12,924	12,055	10,815	9,758	6,628	50,672	71,792	7,311	79,103
Black <sup>b</sup>	Both Sexes	4,937	3,304	2,702	2,117	1,171	821	13,061	15,416	1,054	16,470
	Males	2,605	1,579	1,371	1,027	609	602	6,582	8,038	723	8,761
	Females	2,332	1,725	1,331	1,090	562	219	6,479	7,378	331	7,709
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	891	724	366	+	+	+	2,147	+	+	+
	Males	451	373	198	+	+	+	1,081	+	+	+
	Females	440	351	168	+	+	+	1,066	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,923	2,821	2,211	+	+	+	8,955	+	+	+
	Males	2,125	1,667	1,191	+	+	+	4,983	+	+	+
	Females	1,798	1,154	1,020	+	+	+	3,972	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0292%	0.0007%	0.0100%	0.0384%	0.0528%	0.0449%	0.0311%	0.0270%	0.0264%	0.0202%	0.0292%
	Males	0.0313%	0.0009%	0.0111%	0.0372%	0.0519%	0.0492%	0.0380%	0.0322%	0.0303%	0.0240%	0.0312%
	Females	0.0272%	0.0006%	0.0088%	0.0396%	0.0537%	0.0406%	0.0245%	0.0223%	0.0234%	0.0181%	0.0273%
White <sup>c</sup>	Both Sexes	0.0330%	0.0007%	0.0111%	0.0435%	0.0611%	0.0507%	0.0347%	0.0298%	0.0300%	0.0230%	0.0331%
	Males	0.0352%	0.0008%	0.0124%	0.0407%	0.0598%	0.0547%	0.0421%	0.0356%	0.0342%	0.0273%	0.0350%
	Females	0.0309%	0.0007%	0.0097%	0.0465%	0.0624%	0.0466%	0.0273%	0.0245%	0.0266%	0.0205%	0.0313%
Black <sup>c</sup>	Both Sexes	0.0261%	-	0.0087%	0.0309%	0.0493%	0.0429%	0.0323%	0.0278%	0.0220%	0.0122%	0.0268%
	Males	0.0278%	-	0.0082%	0.0338%	0.0473%	0.0485%	0.0404%	0.0298%	0.0251%	0.0165%	0.0292%
	Females	0.0246%	-	0.0093%	0.0282%	0.0511%	0.0380%	0.0253%	0.0262%	0.0198%	0.0102%	0.0249%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0116%	0.0009%	0.0045%	0.0199%	0.0193%	0.0158%	0.0093%	0.0085%	0.0087%	0.0052%	0.0110%
	Males	0.0124%	-	0.0059%	0.0231%	0.0177%	0.0174%	0.0092%	0.0115%	0.0087%	-	0.0120%
	Females	0.0107%	-	0.0029%	0.0168%	0.0209%	0.0143%	0.0094%	0.0060%	0.0087%	0.0054%	0.0101%
Hispanic <sup>d</sup>	Both Sexes	0.0182%	0.0008%	0.0094%	0.0240%	0.0276%	0.0268%	0.0228%	0.0282%	0.0305%	0.0202%	0.0200%
	Males	0.0199%	0.0007%	0.0117%	0.0248%	0.0266%	0.0305%	0.0302%	0.0293%	0.0367%	0.0380%	0.0226%
	Females	0.0164%	0.0009%	0.0070%	0.0231%	0.0287%	0.0229%	0.0157%	0.0272%	0.0260%	0.0097%	0.0177%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 10.15  
Kaposi SarcomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	7,319	5,335	6,995	3,499	1,165	334	23,149	24,815	+	+
	Males	6,865	5,044	6,752	3,293	1,143	311	21,954	23,533	+	+
	Females	454	291	243	206	22	23	1,195	1,282	+	+
White <sup>b</sup>	Both Sexes	5,002	3,946	5,497	2,916	963	258	17,363	18,729	+	+
	Males	4,685	3,683	5,346	2,755	941	247	16,470	17,761	+	+
	Females	317	263	151	161	22	11	893	968	+	+
Black <sup>b</sup>	Both Sexes	1,915	1,174	1,068	374	93	40	4,531	4,676	+	+
	Males	1,814	1,151	1,022	352	93	40	4,339	4,484	+	+
	Females	101	23	46	22	0	0	192	192	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	101	83	99	+	+	+	302	+	+	+
	Males	101	77	99	+	+	+	296	+	+	+
	Females	0	6	0	+	+	+	6	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,279	974	1,300	+	+	+	3,553	+	+	+
	Males	1,166	929	1,259	+	+	+	3,354	+	+	+
	Females	113	45	41	+	+	+	199	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0071%	-	-	0.0010%	0.0062%	0.0191%	0.0138%	0.0081%	0.0080%	0.0127%	0.0070%
	Males	0.0138%	-	-	0.0019%	0.0122%	0.0377%	0.0281%	0.0164%	0.0152%	0.0249%	0.0139%
	Females	0.0006%	-	-	-	0.0002%	0.0007%	0.0003%	0.0007%	0.0022%	0.0057%	0.0006%
White <sup>c</sup>	Both Sexes	0.0072%	-	-	0.0007%	0.0055%	0.0187%	0.0144%	0.0085%	0.0086%	0.0133%	0.0070%
	Males	0.0138%	-	-	0.0014%	0.0105%	0.0361%	0.0288%	0.0169%	0.0163%	0.0260%	0.0136%
	Females	0.0006%	-	-	-	-	0.0007%	-	0.0007%	0.0024%	0.0060%	0.0006%
Black <sup>c</sup>	Both Sexes	0.0103%	-	-	0.0032%	0.0148%	0.0322%	0.0173%	0.0102%	0.0042%	0.0098%	0.0109%
	Males	0.0209%	-	-	0.0065%	0.0313%	0.0674%	0.0365%	0.0211%	0.0088%	-	0.0227%
	Females	0.0007%	-	-	-	-	-	-	-	-	0.0082%	0.0008%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0018%	-	-	-	0.0016%	0.0043%	0.0035%	0.0015%	0.0035%	0.0045%	0.0018%
	Males	0.0037%	-	-	-	0.0034%	0.0089%	0.0076%	0.0034%	0.0080%	0.0097%	0.0038%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic <sup>d</sup>	Both Sexes	0.0073%	-	-	0.0012%	0.0073%	0.0229%	0.0178%	0.0110%	0.0168%	0.0327%	0.0098%
	Males	0.0135%	-	-	0.0022%	0.0141%	0.0438%	0.0361%	0.0230%	0.0288%	0.0570%	0.0187%
	Females	0.0008%	-	-	-	-	0.0013%	-	-	0.0080%	0.0183%	0.0015%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 11.21  
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	145,766	81,351	44,667	28,708	15,725	9,247	300,492	330,450	11,055	341,505
	Males	88,557	48,431	26,131	16,745	8,694	5,210	179,864	196,335	6,181	202,516
	Females	57,209	32,920	18,536	11,963	7,031	4,037	120,628	134,115	4,874	138,989
White <sup>b</sup>	Both Sexes	122,061	69,978	38,105	25,287	14,060	8,135	255,431	282,105	12,488	294,593
	Males	74,457	41,916	22,801	14,953	7,928	4,731	154,127	169,131	5,584	174,715
	Females	47,604	28,062	15,304	10,334	6,132	3,404	101,304	112,974	6,904	119,878
Black <sup>b</sup>	Both Sexes	16,284	8,176	4,948	2,646	1,395	896	32,053	34,748	1,999	36,747
	Males	9,579	4,611	2,437	1,357	614	377	17,984	19,138	699	19,837
	Females	6,705	3,565	2,511	1,289	781	519	14,069	15,610	1,300	16,910
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,497	1,764	849	+	+	+	6,374	+	+	+
	Males	2,147	1,080	501	+	+	+	3,889	+	+	+
	Females	1,350	684	348	+	+	+	2,485	+	+	+
Hispanic <sup>d</sup>	Both Sexes	13,030	6,714	3,349	+	+	+	23,093	+	+	+
	Males	7,625	3,806	1,790	+	+	+	13,221	+	+	+
	Females	5,405	2,908	1,559	+	+	+	9,872	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0788%	0.0069%	0.0086%	0.0028%	0.0140%	0.0507%	0.1272%	0.2545%	0.3581%	0.3158%	0.0769%
	Males	0.0966%	0.0068%	0.0074%	0.0026%	0.0152%	0.0631%	0.1665%	0.3453%	0.4812%	0.4455%	0.1013%
	Females	0.0614%	0.0069%	0.0099%	0.0030%	0.0127%	0.0385%	0.0897%	0.1723%	0.2598%	0.2406%	0.0565%
White <sup>c</sup>	Both Sexes	0.0868%	0.0072%	0.0094%	0.0027%	0.0142%	0.0544%	0.1341%	0.2690%	0.3788%	0.3278%	0.0811%
	Males	0.1062%	0.0071%	0.0086%	0.0025%	0.0154%	0.0669%	0.1734%	0.3608%	0.5080%	0.4672%	0.1063%
	Females	0.0676%	0.0072%	0.0102%	0.0029%	0.0131%	0.0416%	0.0953%	0.1835%	0.2730%	0.2465%	0.0595%
Black <sup>c</sup>	Both Sexes	0.0674%	0.0080%	0.0098%	0.0039%	0.0172%	0.0495%	0.1395%	0.2746%	0.3624%	0.3039%	0.0805%
	Males	0.0803%	0.0088%	0.0057%	0.0034%	0.0177%	0.0652%	0.1847%	0.3795%	0.4894%	0.3768%	0.1048%
	Females	0.0557%	0.0072%	0.0139%	0.0043%	0.0167%	0.0356%	0.1004%	0.1917%	0.2751%	0.2694%	0.0622%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0421%	0.0039%	0.0024%	0.0017%	0.0070%	0.0263%	0.0673%	0.1347%	0.2093%	0.2122%	0.0430%
	Males	0.0534%	0.0031%	0.0016%	0.0020%	0.0079%	0.0317%	0.0949%	0.1989%	0.2786%	0.3068%	0.0588%
	Females	0.0316%	0.0047%	0.0032%	0.0015%	0.0062%	0.0215%	0.0436%	0.0812%	0.1566%	0.1538%	0.0306%
Hispanic <sup>d</sup>	Both Sexes	0.0471%	0.0060%	0.0071%	0.0019%	0.0136%	0.0466%	0.1204%	0.2525%	0.3771%	0.3818%	0.0786%
	Males	0.0530%	0.0054%	0.0055%	0.0017%	0.0131%	0.0553%	0.1435%	0.3326%	0.5292%	0.4923%	0.1007%
	Females	0.0411%	0.0066%	0.0087%	0.0022%	0.0141%	0.0376%	0.0986%	0.1847%	0.2653%	0.3164%	0.0614%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 12.21  
Cancer of the Larynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	30,947	19,769	14,225	9,955	6,320	4,078	74,897	87,135	1,894	89,029
	Males	25,107	16,026	11,257	7,874	5,166	3,147	60,264	70,117	1,542	71,659
	Females	5,840	3,743	2,968	2,081	1,154	931	14,633	17,018	352	17,370
White <sup>b</sup>	Both Sexes	25,952	16,847	12,347	8,585	5,403	3,703	63,731	74,466	1,501	75,967
	Males	21,132	13,670	9,829	6,781	4,520	2,916	51,412	60,197	1,218	61,415
	Females	4,820	3,177	2,518	1,804	883	787	12,319	14,269	283	14,552
Black <sup>b</sup>	Both Sexes	4,146	2,343	1,538	1,173	761	288	9,200	10,401	125	10,526
	Males	3,276	1,849	1,135	950	571	163	7,210	8,085	106	8,191
	Females	870	494	403	223	190	125	1,990	2,316	19	2,335
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	469	367	227	+	+	+	1,138	+	+	+
	Males	410	322	197	+	+	+	994	+	+	+
	Females	59	45	30	+	+	+	144	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,815	1,000	659	+	+	+	3,474	+	+	+
	Males	1,547	813	544	+	+	+	2,903	+	+	+
	Females	268	187	115	+	+	+	571	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0179%	-	-	0.0002%	0.0008%	0.0053%	0.0258%	0.0665%	0.1052%	0.0793%	0.0176%
	Males	0.0295%	-	-	0.0004%	0.0010%	0.0071%	0.0415%	0.1172%	0.1956%	0.1803%	0.0321%
	Females	0.0065%	-	-	-	0.0007%	0.0034%	0.0107%	0.0206%	0.0330%	0.0207%	0.0060%
White <sup>c</sup>	Both Sexes	0.0197%	-	-	0.0003%	0.0010%	0.0058%	0.0279%	0.0682%	0.1104%	0.0814%	0.0184%
	Males	0.0322%	-	-	0.0005%	0.0011%	0.0077%	0.0448%	0.1180%	0.2018%	0.1824%	0.0330%
	Females	0.0073%	-	-	-	0.0010%	0.0038%	0.0112%	0.0218%	0.0355%	0.0224%	0.0064%
Black <sup>c</sup>	Both Sexes	0.0189%	-	-	-	0.0007%	0.0058%	0.0351%	0.1054%	0.1350%	0.0874%	0.0236%
	Males	0.0314%	-	-	-	-	0.0071%	0.0543%	0.1974%	0.2710%	0.2293%	0.0459%
	Females	0.0076%	-	-	-	-	0.0047%	0.0185%	0.0327%	0.0416%	0.0203%	0.0083%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0077%	-	-	-	-	0.0021%	0.0070%	0.0271%	0.0510%	0.0595%	0.0081%
	Males	0.0140%	-	-	-	-	0.0039%	0.0124%	0.0537%	0.1007%	0.1403%	0.0167%
	Females	0.0019%	-	-	-	-	-	0.0025%	0.0049%	0.0132%	0.0095%	0.0018%
Hispanic <sup>d</sup>	Both Sexes	0.0071%	-	-	-	0.0007%	0.0032%	0.0148%	0.0442%	0.0872%	0.0812%	0.0134%
	Males	0.0116%	-	-	-	0.0008%	0.0043%	0.0234%	0.0862%	0.1758%	0.1853%	0.0264%
	Females	0.0024%	-	-	-	-	0.0021%	0.0068%	0.0086%	0.0221%	0.0196%	0.0038%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.29  
LeukemiaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	112,120	70,289	41,386	26,108	16,514	9,787	249,903	282,150	5,813	287,963
	Males	64,915	40,010	23,413	14,348	9,401	5,114	142,686	160,039	2,612	162,651
	Females	47,205	30,279	17,973	11,760	7,113	4,673	107,217	122,111	3,201	125,312
White <sup>b</sup>	Both Sexes	97,704	62,409	36,830	23,468	14,839	8,940	220,411	249,664	4,882	254,546
	Males	56,696	35,643	20,766	13,104	8,543	4,712	126,208	142,135	1,856	143,991
	Females	41,008	26,766	16,064	10,364	6,296	4,228	94,203	107,529	3,026	110,555
Black <sup>b</sup>	Both Sexes	8,026	4,627	2,484	1,422	799	475	16,557	18,134	732	18,866
	Males	4,606	2,407	1,330	645	387	245	8,987	9,717	346	10,063
	Females	3,420	2,220	1,154	777	412	230	7,570	8,417	386	8,803
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,864	1,629	1,171	+	+	+	6,168	+	+	+
	Males	1,611	894	661	+	+	+	3,414	+	+	+
	Females	1,253	735	510	+	+	+	2,754	+	+	+
Hispanic <sup>d</sup>	Both Sexes	10,781	6,304	4,323	+	+	+	21,409	+	+	+
	Males	6,222	3,445	2,454	+	+	+	12,121	+	+	+
	Females	4,559	2,859	1,869	+	+	+	9,288	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0663%	0.0316%	0.0534%	0.0244%	0.0227%	0.0368%	0.0690%	0.1453%	0.2381%	0.2775%	0.0663%
	Males	0.0772%	0.0341%	0.0583%	0.0268%	0.0255%	0.0443%	0.0853%	0.1915%	0.3145%	0.3639%	0.0819%
	Females	0.0557%	0.0290%	0.0481%	0.0219%	0.0199%	0.0293%	0.0535%	0.1036%	0.1771%	0.2275%	0.0531%
White <sup>c</sup>	Both Sexes	0.0756%	0.0355%	0.0617%	0.0274%	0.0250%	0.0381%	0.0749%	0.1610%	0.2648%	0.3047%	0.0733%
	Males	0.0875%	0.0381%	0.0671%	0.0299%	0.0280%	0.0456%	0.0921%	0.2104%	0.3501%	0.3962%	0.0900%
	Females	0.0637%	0.0327%	0.0560%	0.0248%	0.0219%	0.0304%	0.0580%	0.1149%	0.1950%	0.2513%	0.0588%
Black <sup>c</sup>	Both Sexes	0.0355%	0.0157%	0.0231%	0.0162%	0.0146%	0.0284%	0.0469%	0.0914%	0.1575%	0.1597%	0.0410%
	Males	0.0407%	0.0184%	0.0246%	0.0175%	0.0178%	0.0360%	0.0564%	0.1215%	0.1911%	0.2173%	0.0506%
	Females	0.0307%	0.0129%	0.0215%	0.0151%	0.0118%	0.0216%	0.0387%	0.0676%	0.1344%	0.1325%	0.0337%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0353%	0.0268%	0.0416%	0.0163%	0.0176%	0.0321%	0.0371%	0.0580%	0.0841%	0.1025%	0.0365%
	Males	0.0413%	0.0290%	0.0476%	0.0185%	0.0182%	0.0395%	0.0451%	0.0706%	0.1101%	0.1490%	0.0441%
	Females	0.0297%	0.0244%	0.0353%	0.0141%	0.0172%	0.0254%	0.0301%	0.0474%	0.0643%	0.0738%	0.0302%
Hispanic <sup>d</sup>	Both Sexes	0.0430%	0.0354%	0.0599%	0.0274%	0.0232%	0.0311%	0.0469%	0.0878%	0.1225%	0.1576%	0.0500%
	Males	0.0479%	0.0389%	0.0651%	0.0307%	0.0274%	0.0372%	0.0523%	0.1074%	0.1499%	0.2049%	0.0584%
	Females	0.0380%	0.0318%	0.0544%	0.0237%	0.0187%	0.0247%	0.0417%	0.0712%	0.1023%	0.1297%	0.0427%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 13.30  
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	16,205	12,645	11,446	8,831	7,489	5,660	49,128	66,030	3,537	69,567
	Males	9,215	7,094	6,351	4,802	4,192	2,878	27,462	36,418	1,821	38,239
	Females	6,990	5,551	5,095	4,029	3,297	2,782	21,666	29,612	1,716	31,328
White <sup>b</sup>	Both Sexes	13,564	11,005	9,948	7,750	6,558	5,125	42,265	57,462	+	+
	Males	7,616	6,265	5,471	4,321	3,715	2,608	23,672	31,795	+	+
	Females	5,948	4,740	4,477	3,429	2,843	2,517	18,593	25,667	+	+
Black <sup>b</sup>	Both Sexes	1,375	800	776	557	435	277	3,509	4,352	+	+
	Males	907	360	432	230	214	150	1,929	2,344	+	+
	Females	468	440	344	327	221	127	1,580	2,008	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	805	556	490	+	+	+	2,104	+	+	+
	Males	450	298	311	+	+	+	1,176	+	+	+
	Females	355	258	179	+	+	+	928	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,880	3,078	2,543	+	+	+	10,501	+	+	+
	Males	2,902	1,669	1,431	+	+	+	6,002	+	+	+
	Females	1,978	1,409	1,112	+	+	+	4,499	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0141%	0.0266%	0.0458%	0.0143%	0.0049%	0.0040%	0.0030%	0.0036%	0.0024%	0.0018%	0.0145%
	Males	0.0161%	0.0290%	0.0499%	0.0166%	0.0060%	0.0044%	0.0034%	0.0042%	0.0026%	0.0021%	0.0161%
	Females	0.0121%	0.0242%	0.0414%	0.0118%	0.0038%	0.0036%	0.0025%	0.0030%	0.0023%	0.0016%	0.0129%
White <sup>c</sup>	Both Sexes	0.0159%	0.0302%	0.0537%	0.0167%	0.0058%	0.0042%	0.0031%	0.0037%	0.0024%	0.0016%	0.0167%
	Males	0.0179%	0.0326%	0.0579%	0.0193%	0.0070%	0.0048%	0.0034%	0.0043%	0.0029%	0.0016%	0.0184%
	Females	0.0138%	0.0276%	0.0493%	0.0139%	0.0046%	0.0037%	0.0027%	0.0031%	0.0019%	0.0015%	0.0149%
Black <sup>c</sup>	Both Sexes	0.0068%	0.0119%	0.0172%	0.0083%	0.0013%	0.0019%	0.0015%	0.0021%	-	-	0.0063%
	Males	0.0082%	0.0152%	0.0186%	0.0094%	0.0016%	0.0019%	-	-	-	-	0.0071%
	Females	0.0057%	0.0085%	0.0158%	0.0073%	-	0.0019%	0.0016%	-	-	-	0.0054%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0104%	0.0226%	0.0339%	0.0075%	0.0037%	0.0045%	0.0032%	0.0031%	0.0031%	0.0031%	0.0114%
	Males	0.0126%	0.0237%	0.0398%	0.0094%	0.0047%	0.0048%	0.0049%	0.0034%	-	-	0.0130%
	Females	0.0085%	0.0215%	0.0278%	0.0055%	0.0028%	0.0041%	0.0018%	0.0028%	0.0048%	-	0.0097%
Hispanic <sup>d</sup>	Both Sexes	0.0208%	0.0301%	0.0533%	0.0174%	0.0069%	0.0055%	0.0041%	0.0061%	0.0044%	-	0.0175%
	Males	0.0234%	0.0337%	0.0569%	0.0206%	0.0081%	0.0074%	0.0045%	0.0055%	-	-	0.0194%
	Females	0.0181%	0.0265%	0.0496%	0.0138%	0.0055%	0.0036%	0.0037%	0.0066%	0.0038%	-	0.0155%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 14.21  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	30,577	6,193	2,076	1,020	640	290	39,864	40,940	464	41,404
	Males	22,266	4,259	1,270	638	260	125	28,432	28,891	76	28,967
	Females	8,311	1,934	806	382	380	165	11,432	12,049	388	12,437
White <sup>b</sup>	Both Sexes	22,207	4,560	1,471	757	496	255	28,995	29,837	462	30,299
	Males	16,307	3,183	900	467	171	106	20,857	21,190	196 <sup>i</sup>	21,386
	Females	5,900	1,377	571	290	325	149	8,138	8,647	266 <sup>i</sup>	8,913
Black <sup>b</sup>	Both Sexes	4,513	532	143	108	72	23	5,296	5,425	+	+
	Males	3,283	283	68	61	36	13	3,695	3,755	+	+
	Females	1,230	249	75	47	36	10	1,601	1,670	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,343	938	367	+	+	+	4,720	+	+	+
	Males	2,343	635	262	+	+	+	3,290	+	+	+
	Females	1,000	303	105	+	+	+	1,430	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,323	802	280	+	+	+	5,406	+	+	+
	Males	3,165	564	165	+	+	+	3,895	+	+	+
	Females	1,158	238	115	+	+	+	1,511	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0147%	0.0027%	0.0018%	0.0008%	0.0014%	0.0064%	0.0327%	0.0510%	0.0576%	0.0374%	0.0139%
	Males	0.0211%	0.0029%	0.0019%	0.0010%	0.0016%	0.0090%	0.0537%	0.0807%	0.0810%	0.0579%	0.0208%
	Females	0.0083%	0.0024%	0.0018%	0.0005%	0.0013%	0.0037%	0.0126%	0.0242%	0.0390%	0.0255%	0.0078%
White <sup>c</sup>	Both Sexes	0.0119%	0.0028%	0.0018%	0.0008%	0.0010%	0.0046%	0.0279%	0.0402%	0.0374%	0.0267%	0.0107%
	Males	0.0173%	0.0029%	0.0018%	0.0011%	0.0008%	0.0064%	0.0459%	0.0640%	0.0535%	0.0408%	0.0161%
	Females	0.0065%	0.0027%	0.0017%	0.0005%	0.0013%	0.0028%	0.0102%	0.0181%	0.0242%	0.0185%	0.0058%
Black <sup>c</sup>	Both Sexes	0.0120%	0.0007%	0.0011%	0.0010%	0.0021%	0.0046%	0.0355%	0.0617%	0.0446%	0.0256%	0.0132%
	Males	0.0175%	-	0.0017%	-	0.0036%	0.0057%	0.0559%	0.1024%	0.0638%	0.0431%	0.0205%
	Females	0.0070%	-	-	-	-	0.0037%	0.0179%	0.0296%	0.0314%	0.0173%	0.0075%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0318%	0.0032%	0.0026%	-	0.0028%	0.0167%	0.0553%	0.1079%	0.1808%	0.1183%	0.0325%
	Males	0.0460%	0.0039%	0.0020%	-	0.0039%	0.0260%	0.0960%	0.1718%	0.2563%	0.1811%	0.0499%
	Females	0.0188%	0.0024%	0.0032%	-	0.0018%	0.0085%	0.0204%	0.0547%	0.1233%	0.0795%	0.0187%
Hispanic <sup>d</sup>	Both Sexes	0.0110%	0.0029%	0.0017%	0.0005%	0.0008%	0.0065%	0.0393%	0.0709%	0.0679%	0.0575%	0.0174%
	Males	0.0156%	0.0032%	0.0019%	-	0.0008%	0.0100%	0.0680%	0.1161%	0.0889%	0.0824%	0.0266%
	Females	0.0062%	0.0025%	0.0015%	-	0.0008%	0.0029%	0.0120%	0.0327%	0.0524%	0.0428%	0.0095%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 15.27  
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	231,996	75,108	41,458	22,038	11,527	5,962	370,600	391,564	7,867	399,431
	Males	108,211	33,462	17,795	9,916	5,160	2,716	169,384	178,912	4,498	183,410
	Females	123,785	41,646	23,663	12,122	6,367	3,246	201,216	212,652	3,369	216,021
White <sup>b</sup>	Both Sexes	197,163	65,648	36,169	19,441	10,233	5,274	318,423	336,910	3,296	340,206
	Males	91,187	28,787	15,084	8,691	4,482	2,396	143,750	152,043	1,552	153,595
	Females	105,976	36,861	21,085	10,750	5,751	2,878	174,673	184,867	1,744	186,611
Black <sup>b</sup>	Both Sexes	25,150	6,834	3,960	2,047	971	535	37,990	39,861	310	40,171
	Males	12,105	3,399	1,952	942	518	247	18,397	19,346	162	19,508
	Females	13,045	3,435	2,008	1,105	453	288	19,593	20,515	148	20,663
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	7,002	1,951	979	+	+	+	10,187	+	+	+
	Males	3,589	942	536	+	+	+	5,196	+	+	+
	Females	3,413	1,009	443	+	+	+	4,991	+	+	+
Hispanic <sup>d</sup>	Both Sexes	8,530	2,434	1,169	+	+	+	12,133	+	+	+
	Males	4,159	1,083	578	+	+	+	5,819	+	+	+
	Females	4,371	1,351	591	+	+	+	6,314	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0960%	0.0001%	0.0003%	0.0017%	0.0056%	0.0243%	0.1067%	0.3218%	0.6243%	0.5644%	0.0965%
	Males	0.0898%	-	0.0003%	0.0016%	0.0050%	0.0210%	0.1030%	0.3304%	0.6495%	0.6464%	0.1002%
	Females	0.1021%	-	0.0003%	0.0018%	0.0062%	0.0275%	0.1103%	0.3141%	0.6042%	0.5169%	0.0943%
White <sup>c</sup>	Both Sexes	0.1038%	0.0002%	0.0003%	0.0019%	0.0057%	0.0245%	0.1052%	0.3289%	0.6485%	0.5773%	0.0989%
	Males	0.0940%	-	0.0003%	0.0018%	0.0051%	0.0212%	0.0983%	0.3253%	0.6528%	0.6405%	0.0994%
	Females	0.1135%	-	0.0004%	0.0020%	0.0062%	0.0279%	0.1119%	0.3323%	0.6450%	0.5404%	0.0992%
Black <sup>c</sup>	Both Sexes	0.0830%	-	-	0.0015%	0.0060%	0.0299%	0.1536%	0.3979%	0.6314%	0.4834%	0.1060%
	Males	0.0851%	-	-	0.0017%	0.0066%	0.0236%	0.1717%	0.4555%	0.7500%	0.6533%	0.1238%
	Females	0.0811%	-	-	0.0013%	0.0055%	0.0354%	0.1380%	0.3524%	0.5498%	0.4031%	0.0942%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0713%	-	-	0.0009%	0.0052%	0.0197%	0.0792%	0.2294%	0.4858%	0.5170%	0.0758%
	Males	0.0758%	-	-	-	0.0036%	0.0194%	0.0798%	0.2816%	0.5681%	0.6767%	0.0896%
	Females	0.0672%	-	-	-	0.0067%	0.0200%	0.0788%	0.1860%	0.4232%	0.4183%	0.0658%
Hispanic <sup>d</sup>	Both Sexes	0.0248%	-	-	0.0011%	0.0037%	0.0121%	0.0448%	0.1413%	0.3111%	0.3464%	0.0482%
	Males	0.0232%	-	-	0.0007%	0.0042%	0.0111%	0.0432%	0.1412%	0.3767%	0.4093%	0.0539%
	Females	0.0263%	-	-	0.0017%	0.0032%	0.0132%	0.0463%	0.1414%	0.2628%	0.3092%	0.0444%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 16.21  
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	276,312	198,547	148,808	102,930	77,036	50,041	726,597	884,259	37,521	921,780
	Males	148,749	101,316	73,983	50,016	33,809	20,557	374,064	440,298	12,242	452,540
	Females	127,563	97,231	74,825	52,914	43,227	29,484	352,533	443,961	25,279	469,240
White <sup>b</sup>	Both Sexes	266,098	192,097	143,074	98,849	73,365	48,596	700,118	851,305	35,396	886,701
	Males	143,731	98,225	71,149	48,246	32,417	19,982	361,352	425,076	11,666	436,742
	Females	122,367	93,872	71,925	50,603	40,948	28,614	338,766	426,229	23,730	449,959
Black <sup>b</sup>	Both Sexes	1,095	636	639	324	257	98	2,694	3,148	107	3,255
	Males	462	320	275	146	56	25	1,203	1,323	44	1,367
	Females	633	316	364	178	201	73	1,491	1,825	63	1,888
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	709	553	310	+	+	+	1,700	+	+	+
	Males	338	238	128	+	+	+	764	+	+	+
	Females	371	315	182	+	+	+	936	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,721	3,527	2,598	+	+	+	10,847	+	+	+
	Males	1,951	1,179	885	+	+	+	4,016	+	+	+
	Females	2,770	2,348	1,713	+	+	+	6,831	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1669%	0.0004%	0.0033%	0.0289%	0.0934%	0.1823%	0.2918%	0.4392%	0.5380%	0.5894%	0.1623%
	Males	0.1777%	0.0003%	0.0028%	0.0184%	0.0672%	0.1530%	0.3037%	0.5527%	0.7612%	0.9293%	0.1868%
	Females	0.1565%	0.0006%	0.0038%	0.0398%	0.1197%	0.2112%	0.2804%	0.3366%	0.3599%	0.3926%	0.1463%
White <sup>c</sup>	Both Sexes	0.2165%	0.0005%	0.0041%	0.0380%	0.1230%	0.2344%	0.3699%	0.5437%	0.6664%	0.6972%	0.2035%
	Males	0.2287%	0.0004%	0.0034%	0.0239%	0.0863%	0.1931%	0.3792%	0.6753%	0.9327%	1.0960%	0.2295%
	Females	0.2044%	0.0006%	0.0048%	0.0530%	0.1615%	0.2768%	0.3608%	0.4211%	0.4486%	0.4647%	0.1874%
Black <sup>c</sup>	Both Sexes	0.0054%	-	-	0.0014%	0.0031%	0.0062%	0.0069%	0.0194%	0.0264%	0.0350%	0.0066%
	Males	0.0052%	-	-	-	0.0018%	0.0049%	0.0071%	0.0210%	0.0360%	0.0626%	0.0077%
	Females	0.0055%	-	-	0.0018%	0.0041%	0.0074%	0.0067%	0.0182%	0.0198%	0.0219%	0.0061%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0104%	-	0.0016%	0.0020%	0.0062%	0.0130%	0.0143%	0.0278%	0.0286%	0.0468%	0.0104%
	Males	0.0099%	-	0.0022%	0.0017%	0.0052%	0.0088%	0.0142%	0.0305%	0.0354%	0.0616%	0.0106%
	Females	0.0109%	-	-	0.0024%	0.0071%	0.0168%	0.0143%	0.0255%	0.0234%	0.0377%	0.0104%
Hispanic <sup>d</sup>	Both Sexes	0.0223%	-	0.0011%	0.0047%	0.0165%	0.0365%	0.0615%	0.0828%	0.1107%	0.1537%	0.0336%
	Males	0.0162%	-	0.0012%	0.0024%	0.0089%	0.0220%	0.0452%	0.0682%	0.1284%	0.2032%	0.0300%
	Females	0.0286%	-	0.0011%	0.0072%	0.0246%	0.0515%	0.0770%	0.0951%	0.0977%	0.1244%	0.0382%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 17.13  
MesotheliomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	2,968	471	387	292	170	130	4,118	4,538	+	+
	Males	1,998	246	129	83	51	70	2,456	2,601	+	+
	Females	970	225	258	209	119	60	1,662	1,937	+	+
White <sup>b</sup>	Both Sexes	2,815	382	326	256	167	118	3,780	4,159	+	+
	Males	1,882	202	105	71	48	58	2,261	2,377	+	+
	Females	933	180	221	185	119	60	1,519	1,782	+	+
Black <sup>b</sup>	Both Sexes	111	58	48	25	0	12	241	278	+	+
	Males	86	35	24	13	0	12	157	182	+	+
	Females	25	23	24	12	0	0	84	96	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	16	20	7	+	+	+	42	+	+	+
	Males	8	10	4	+	+	+	21	+	+	+
	Females	8	10	3	+	+	+	21	+	+	+
Hispanic <sup>d</sup>	Both Sexes	260	36	41	+	+	+	338	+	+	+
	Males	150	29	36	+	+	+	216	+	+	+
	Females	110	7	5	+	+	+	122	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0011%	-	-	-	0.0003%	0.0005%	0.0013%	0.0036%	0.0058%	0.0055%	0.0011%
	Males	0.0014%	-	-	-	-	0.0004%	0.0017%	0.0051%	0.0089%	0.0100%	0.0015%
	Females	0.0008%	-	-	-	0.0005%	0.0006%	0.0009%	0.0023%	0.0034%	0.0029%	0.0007%
White <sup>c</sup>	Both Sexes	0.0013%	-	-	-	0.0003%	0.0006%	0.0015%	0.0043%	0.0068%	0.0059%	0.0012%
	Males	0.0017%	-	-	-	-	0.0005%	0.0020%	0.0060%	0.0103%	0.0106%	0.0017%
	Females	0.0010%	-	-	-	0.0005%	0.0007%	0.0011%	0.0027%	0.0039%	0.0032%	0.0009%
Black <sup>c</sup>	Both Sexes	0.0004%	-	-	-	-	-	0.0010%	-	0.0042%	-	0.0006%
	Males	0.0006%	-	-	-	-	-	0.0019%	-	-	-	0.0009%
	Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0004%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0003%	-	-	-	-	-	-	0.0013%	-	0.0031%	0.0003%
	Males	0.0003%	-	-	-	-	-	-	-	-	-	0.0004%
	Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0002%
Hispanic <sup>d</sup>	Both Sexes	0.0007%	-	-	-	-	0.0005%	0.0023%	0.0040%	0.0051%	0.0066%	0.0012%
	Males	0.0009%	-	-	-	-	-	0.0040%	0.0057%	0.0073%	0.0133%	0.0018%
	Females	0.0005%	-	-	-	-	-	-	0.0025%	-	-	0.0008%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 18.21  
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	49,451	17,820	6,075	2,600	975	316	75,945	77,360	257	77,617	
	Males	26,622	9,775	3,552	1,373	568	177	41,322	42,092	93	42,185	
	Females	22,829	8,045	2,523	1,227	407	139	34,623	35,268	164	35,432	
White <sup>b</sup>	Both Sexes	37,208	14,041	4,702	1,975	715	195	57,927	58,934	103	59,037	
	Males	21,060	7,936	2,881	1,028	421	116	32,906	33,468	0 <sup>i</sup>	33,468	
	Females	16,148	6,105	1,821	947	294	79	25,021	25,466	103 <sup>i</sup>	25,569	
Black <sup>b</sup>	Both Sexes	10,159	3,009	1,157	525	205	106	14,849	15,184	+	+	
	Males	4,573	1,459	534	304	131	52	6,869	7,053	+	+	
	Females	5,586	1,550	623	221	74	54	7,980	8,131	+	+	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1,164	399	143	+	+	+	1,757	+	+	+	
	Males	602	208	81	+	+	+	911	+	+	+	
	Females	562	191	62	+	+	+	846	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	3,511	1,308	445	+	+	+	5,264	+	+	+	
	Males	1,915	724	240	+	+	+	2,879	+	+	+	
	Females	1,596	584	205	+	+	+	2,385	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0214%	-	-	0.0003%	0.0020%	0.0106%	0.0317%	0.0767%	0.1108%	0.0982%	0.0209%
	Males	0.0235%	-	-	0.0005%	0.0026%	0.0121%	0.0365%	0.0902%	0.1347%	0.1233%	0.0250%
	Females	0.0194%	-	-	-	0.0015%	0.0091%	0.0270%	0.0644%	0.0917%	0.0837%	0.0176%
White <sup>c</sup>	Both Sexes	0.0210%	-	-	0.0003%	0.0018%	0.0097%	0.0293%	0.0709%	0.1059%	0.0910%	0.0195%
	Males	0.0239%	-	-	0.0006%	0.0023%	0.0118%	0.0343%	0.0858%	0.1339%	0.1227%	0.0243%
	Females	0.0180%	-	-	-	0.0013%	0.0074%	0.0244%	0.0570%	0.0830%	0.0725%	0.0155%
Black <sup>c</sup>	Both Sexes	0.0339%	-	-	-	0.0041%	0.0232%	0.0648%	0.1578%	0.2045%	0.2081%	0.0422%
	Males	0.0324%	-	-	-	0.0051%	0.0200%	0.0720%	0.1739%	0.2153%	0.2104%	0.0444%
	Females	0.0353%	-	-	-	0.0033%	0.0260%	0.0586%	0.1451%	0.1971%	0.2071%	0.0404%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0120%	-	-	-	0.0012%	0.0045%	0.0163%	0.0461%	0.0660%	0.0694%	0.0123%
	Males	0.0129%	-	-	-	0.0017%	0.0057%	0.0189%	0.0550%	0.0794%	0.0695%	0.0144%
	Females	0.0112%	-	-	-	-	0.0035%	0.0141%	0.0386%	0.0558%	0.0693%	0.0107%
Hispanic <sup>d</sup>	Both Sexes	0.0107%	-	-	0.0006%	0.0021%	0.0085%	0.0274%	0.0657%	0.1092%	0.0901%	0.0190%
	Males	0.0115%	-	-	0.0010%	0.0028%	0.0102%	0.0327%	0.0797%	0.1226%	0.1065%	0.0224%
	Females	0.0099%	-	-	-	0.0013%	0.0067%	0.0224%	0.0539%	0.0994%	0.0803%	0.0163%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 19.25  
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	198,946	132,700	78,129	43,600	25,113	13,936	453,373	499,712	9,353	509,065	
	Males	105,553	68,564	40,496	22,771	13,154	7,378	237,383	261,560	4,918	266,478	
	Females	93,393	64,136	37,633	20,829	11,959	6,558	215,990	238,152	4,435	242,587	
White <sup>b</sup>	Both Sexes	173,392	117,218	69,017	38,851	22,746	12,707	398,479	440,727	9,713	450,440	
	Males	92,730	60,876	35,713	20,357	11,942	6,714	209,677	231,746	5,482	237,228	
	Females	80,662	56,342	33,304	18,494	10,804	5,993	188,802	208,981	4,231	213,212	
Black <sup>b</sup>	Both Sexes	15,619	10,026	5,921	2,948	1,511	749	34,515	37,133	747	37,880	
	Males	7,588	4,978	3,116	1,449	795	419	17,132	18,495	311	18,806	
	Females	8,031	5,048	2,805	1,499	716	330	17,383	18,638	436	19,074	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	5,408	3,518	1,911	+	+	+	11,562	+	+	+	
	Males	2,874	1,779	984	+	+	+	5,983	+	+	+	
	Females	2,534	1,739	927	+	+	+	5,579	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	14,335	8,665	4,824	+	+	+	27,824	+	+	+	
	Males	7,183	4,586	2,545	+	+	+	14,314	+	+	+	
	Females	7,152	4,079	2,279	+	+	+	13,510	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.1199%	0.0028%	0.0111%	0.0215%	0.0416%	0.0886%	0.1741%	0.3276%	0.5227%	0.5731%	0.1181%
	Males	0.1277%	0.0036%	0.0146%	0.0260%	0.0486%	0.1042%	0.1999%	0.3731%	0.5834%	0.6487%	0.1352%
	Females	0.1123%	0.0019%	0.0074%	0.0169%	0.0347%	0.0732%	0.1496%	0.2865%	0.4742%	0.5292%	0.1032%
White <sup>c</sup>	Both Sexes	0.1346%	0.0030%	0.0114%	0.0225%	0.0436%	0.0940%	0.1875%	0.3550%	0.5717%	0.6097%	0.1269%
	Males	0.1427%	0.0041%	0.0151%	0.0277%	0.0509%	0.1104%	0.2143%	0.4023%	0.6341%	0.6837%	0.1448%
	Females	0.1266%	0.0019%	0.0074%	0.0170%	0.0359%	0.0771%	0.1611%	0.3110%	0.5205%	0.5665%	0.1110%
Black <sup>c</sup>	Both Sexes	0.0744%	0.0022%	0.0114%	0.0217%	0.0434%	0.0860%	0.1466%	0.2401%	0.3073%	0.2761%	0.0859%
	Males	0.0790%	0.0019%	0.0148%	0.0259%	0.0531%	0.1016%	0.1686%	0.2757%	0.2863%	0.3231%	0.0958%
	Females	0.0703%	0.0025%	0.0080%	0.0177%	0.0350%	0.0721%	0.1276%	0.2120%	0.3217%	0.2538%	0.0776%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0739%	0.0023%	0.0085%	0.0160%	0.0270%	0.0550%	0.1050%	0.1908%	0.3318%	0.4270%	0.0756%
	Males	0.0796%	0.0031%	0.0112%	0.0172%	0.0293%	0.0631%	0.1199%	0.2156%	0.4064%	0.5053%	0.0883%
	Females	0.0686%	0.0015%	0.0056%	0.0148%	0.0250%	0.0478%	0.0922%	0.1701%	0.2751%	0.3785%	0.0656%
Hispanic <sup>d</sup>	Both Sexes	0.0569%	0.0025%	0.0081%	0.0157%	0.0305%	0.0660%	0.1351%	0.2491%	0.3956%	0.4649%	0.0910%
	Males	0.0574%	0.0034%	0.0107%	0.0185%	0.0364%	0.0752%	0.1491%	0.2529%	0.3999%	0.4829%	0.0970%
	Females	0.0564%	0.0015%	0.0054%	0.0126%	0.0243%	0.0565%	0.1219%	0.2459%	0.3925%	0.4543%	0.0853%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 20.27  
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	102,019	62,468	40,413	25,710	16,622	10,222	230,609	264,620	10,573	275,193	
	Males	71,123	42,078	25,672	16,391	9,818	5,832	155,263	175,045	6,039	181,084	
	Females	30,896	20,390	14,741	9,319	6,804	4,390	75,346	89,575	4,534	94,109	
White <sup>b</sup>	Both Sexes	87,884	54,203	35,113	22,566	14,701	9,004	199,766	229,701	9,027	238,728	
	Males	61,858	37,081	22,730	14,681	8,836	5,330	136,350	154,201	5,183	159,384	
	Females	26,026	17,122	12,383	7,885	5,865	3,674	63,416	75,500	3,844	79,344	
Black <sup>b</sup>	Both Sexes	7,824	4,802	2,811	1,702	1,091	703	17,139	19,487	523	20,010	
	Males	5,196	2,880	1,475	841	555	226	10,392	11,378	202	11,580	
	Females	2,628	1,922	1,336	861	536	477	6,747	8,109	321	8,430	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,791	2,306	1,579	+	+	+	8,305	+	+	+	
	Males	2,415	1,406	925	+	+	+	5,113	+	+	+	
	Females	1,376	900	654	+	+	+	3,192	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	5,447	2,800	1,887	+	+	+	10,134	+	+	+	
	Males	3,466	1,726	1,119	+	+	+	6,312	+	+	+	
	Females	1,981	1,074	768	+	+	+	3,822	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0603%	0.0005%	0.0021%	0.0057%	0.0151%	0.0436%	0.1185%	0.1993%	0.2220%	0.2053%	0.0576%
	Males	0.0823%	0.0004%	0.0019%	0.0056%	0.0150%	0.0565%	0.1768%	0.3061%	0.3309%	0.2994%	0.0832%
	Females	0.0389%	0.0005%	0.0023%	0.0059%	0.0153%	0.0308%	0.0628%	0.1027%	0.1351%	0.1508%	0.0356%
White <sup>c</sup>	Both Sexes	0.0655%	0.0005%	0.0019%	0.0056%	0.0139%	0.0418%	0.1239%	0.2120%	0.2375%	0.2170%	0.0599%
	Males	0.0903%	0.0005%	0.0015%	0.0051%	0.0139%	0.0553%	0.1874%	0.3248%	0.3549%	0.3164%	0.0872%
	Females	0.0409%	0.0005%	0.0024%	0.0061%	0.0138%	0.0278%	0.0614%	0.1070%	0.1415%	0.1590%	0.0358%
Black <sup>c</sup>	Both Sexes	0.0374%	-	0.0025%	0.0060%	0.0119%	0.0362%	0.0935%	0.1502%	0.1535%	0.1064%	0.0425%
	Males	0.0490%	-	0.0031%	0.0081%	0.0091%	0.0412%	0.1326%	0.2373%	0.2341%	0.1695%	0.0603%
	Females	0.0269%	-	0.0019%	0.0040%	0.0143%	0.0318%	0.0598%	0.0813%	0.0981%	0.0765%	0.0290%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0507%	-	0.0021%	0.0060%	0.0227%	0.0560%	0.1025%	0.1449%	0.1627%	0.1578%	0.0499%
	Males	0.0655%	-	0.0028%	0.0057%	0.0247%	0.0719%	0.1422%	0.2194%	0.2244%	0.2218%	0.0684%
	Females	0.0371%	-	-	0.0063%	0.0209%	0.0418%	0.0684%	0.0828%	0.1158%	0.1183%	0.0350%
Hispanic <sup>d</sup>	Both Sexes	0.0207%	0.0004%	0.0013%	0.0041%	0.0088%	0.0199%	0.0592%	0.1128%	0.1387%	0.1521%	0.0334%
	Males	0.0253%	0.0006%	0.0008%	0.0041%	0.0091%	0.0234%	0.0825%	0.1603%	0.2138%	0.1969%	0.0459%
	Females	0.0160%	-	0.0018%	0.0040%	0.0086%	0.0164%	0.0371%	0.0726%	0.0836%	0.1257%	0.0233%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 21.15  
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	57,207	31,228	23,911	19,604	16,095	12,285	131,950	169,398	16,740	186,138
White <sup>b</sup>	49,594	27,354	20,967	17,364	14,538	10,962	115,279	148,810	14,272	163,082
Black <sup>b</sup>	4,179	1,949	1,611	1,317	997	835	9,056	11,639	1,633	13,272
Asian/Pacific Islander <sup>c</sup>	2,514	1,445	957	+	+	+	5,372	+	+	+
Hispanic <sup>d</sup>	4,917	2,602	1,711	+	+	+	9,230	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0687%	0.0003%	0.0063%	0.0164%	0.0279%	0.0615%	0.1313%	0.1941%	0.2068%	0.1621%	0.0630%	
White <sup>c</sup>	0.0757%	0.0004%	0.0067%	0.0175%	0.0277%	0.0639%	0.1384%	0.2122%	0.2267%	0.1711%	0.0671%	
Black <sup>c</sup>	0.0368%	-	0.0044%	0.0105%	0.0211%	0.0415%	0.0729%	0.1056%	0.1287%	0.1401%	0.0397%	
Asian/Pacific Islander <sup>c</sup>	0.0609%	-	0.0064%	0.0160%	0.0330%	0.0635%	0.1382%	0.1561%	0.1419%	0.1046%	0.0562%	
Hispanic <sup>d</sup>	0.0384%	-	0.0087%	0.0182%	0.0251%	0.0495%	0.0998%	0.1467%	0.1602%	0.1538%	0.0516%	

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.  
<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census.  
<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.  
<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.  
<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.  
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
 + Not available.

Table 22.21  
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	31,442	4,797	2,320	1,397	700	406	39,955	41,284	325	41,609	
	Males	15,733	2,176	1,119	606	287	119	19,633	20,155	138	20,293	
	Females	15,709	2,621	1,201	791	413	287	20,322	21,129	187	21,316	
White <sup>b</sup>	Both Sexes	26,398	4,019	1,786	1,142	469	329	33,346	34,332	266	34,598	
	Males	13,417	1,938	925	524	209	108	16,804	17,203	123	17,326	
	Females	12,981	2,081	861	618	260	221	16,542	17,129	143	17,272	
Black <sup>b</sup>	Both Sexes	3,651	557	287	141	192	46	4,636	4,902	147	5,049	
	Males	1,680	183	93	24	78	0	1,980	2,087	62	2,149	
	Females	1,971	374	194	117	114	46	2,656	2,815	85	2,900	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	988	175	117	+	+	+	1,310	+	+	+	
	Males	451	60	41	+	+	+	565	+	+	+	
	Females	537	115	76	+	+	+	745	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	2,003	295	197	+	+	+	2,494	+	+	+	
	Males	932	131	62	+	+	+	1,125	+	+	+	
	Females	1,071	164	135	+	+	+	1,369	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0113%	-	0.0001%	0.0006%	0.0016%	0.0061%	0.0171%	0.0366%	0.0570%	0.0549%	0.0111%
	Males	0.0112%	-	-	0.0005%	0.0015%	0.0065%	0.0189%	0.0417%	0.0592%	0.0576%	0.0119%
	Females	0.0114%	-	0.0002%	0.0008%	0.0018%	0.0057%	0.0154%	0.0321%	0.0554%	0.0533%	0.0104%
White <sup>c</sup>	Both Sexes	0.0119%	-	0.0001%	0.0006%	0.0017%	0.0060%	0.0175%	0.0378%	0.0568%	0.0530%	0.0111%
	Males	0.0122%	-	-	0.0004%	0.0014%	0.0068%	0.0196%	0.0435%	0.0613%	0.0572%	0.0122%
	Females	0.0116%	-	-	0.0007%	0.0019%	0.0051%	0.0154%	0.0324%	0.0532%	0.0505%	0.0101%
Black <sup>c</sup>	Both Sexes	0.0105%	-	-	0.0008%	0.0021%	0.0081%	0.0201%	0.0406%	0.0644%	0.0714%	0.0131%
	Males	0.0096%	-	-	-	0.0025%	0.0085%	0.0224%	0.0392%	0.0562%	0.0791%	0.0132%
	Females	0.0113%	-	-	-	0.0017%	0.0077%	0.0180%	0.0418%	0.0700%	0.0678%	0.0130%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0090%	-	-	0.0009%	0.0012%	0.0046%	0.0126%	0.0263%	0.0521%	0.0584%	0.0094%
	Males	0.0080%	-	-	-	0.0013%	0.0031%	0.0123%	0.0319%	0.0476%	0.0493%	0.0089%
	Females	0.0099%	-	-	0.0012%	0.0011%	0.0060%	0.0128%	0.0217%	0.0554%	0.0640%	0.0097%
Hispanic <sup>d</sup>	Both Sexes	0.0051%	-	-	0.0006%	0.0008%	0.0050%	0.0126%	0.0289%	0.0425%	0.0576%	0.0090%
	Males	0.0045%	-	-	-	-	0.0054%	0.0128%	0.0298%	0.0424%	0.0462%	0.0087%
	Females	0.0057%	-	-	0.0009%	0.0011%	0.0045%	0.0124%	0.0280%	0.0425%	0.0644%	0.0092%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 23.15  
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	1,026,391	792,069	487,305	251,456	45,972	10,179	2,557,220	2,616,190	1,492	2,617,682
White <sup>b</sup>	829,641	664,092	414,049	221,725	41,824	9,368	2,129,506	2,183,280	1,708	2,184,988
Black <sup>b</sup>	143,372	101,453	58,873	24,343	3,334	704	328,040	332,272	143	332,415
Asian/Pacific Islander <sup>c</sup>	20,238	14,164	6,970	+	+	+	43,551	+	+	+
Hispanic <sup>d</sup>	56,919	40,651	21,777	+	+	+	119,346	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	1.3222%	-	0.0002%	0.0002%	0.0013%	0.0852%	1.0145%	4.8944%	10.9028%	11.6130%	1.5246%
White <sup>c</sup>	1.3915%	-	-	-	0.0008%	0.0746%	0.9740%	4.8637%	10.9694%	11.2939%	1.5083%
Black <sup>c</sup>	1.4790%	-	-	-	0.0038%	0.2140%	1.9558%	8.2407%	16.1070%	15.8241%	2.3520%
Asian/Pacific Islander <sup>c</sup>	0.6478%	-	-	-	-	0.0213%	0.3416%	1.9750%	5.6894%	8.6466%	0.8183%
Hispanic <sup>d</sup>	0.4767%	-	-	-	0.0008%	0.0396%	0.6030%	3.3919%	8.9218%	10.5882%	1.2170%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.21  
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	35,420	15,772	8,440	4,999	3,645	1,726	64,632	70,821	1,448	72,269
	Males	21,286	8,829	4,744	2,527	1,825	857	37,387	40,560	826	41,386
	Females	14,134	6,943	3,696	2,472	1,820	869	27,245	30,261	622	30,883
White <sup>b</sup>	Both Sexes	26,260	11,637	6,058	3,688	2,809	1,276	47,643	52,416	1,156	53,572
	Males	16,450	6,700	3,393	1,779	1,426	695	28,322	30,856	635	31,491
	Females	9,810	4,937	2,665	1,909	1,383	581	19,321	21,560	521	22,081
Black <sup>b</sup>	Both Sexes	5,652	2,502	1,341	626	459	271	10,121	10,904	206	11,110
	Males	2,842	1,238	671	401	213	113	5,152	5,519	120	5,639
	Females	2,810	1,264	670	225	246	158	4,969	5,385	86	5,471
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,206	1,595	943	+	+	+	6,146	+	+	+
	Males	1,806	853	567	+	+	+	3,449	+	+	+
	Females	1,400	742	376	+	+	+	2,697	+	+	+
Hispanic <sup>d</sup>	Both Sexes	5,061	2,000	1,112	+	+	+	8,175	+	+	+
	Males	2,908	1,086	517	+	+	+	4,512	+	+	+
	Females	2,153	914	595	+	+	+	3,663	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0219%	-	-	0.0005%	0.0031%	0.0120%	0.0279%	0.0592%	0.1164%	0.1535%	0.0218%
	Males	0.0255%	-	-	0.0003%	0.0032%	0.0132%	0.0350%	0.0764%	0.1587%	0.2050%	0.0283%
	Females	0.0185%	-	-	0.0008%	0.0029%	0.0108%	0.0212%	0.0436%	0.0826%	0.1236%	0.0167%
White <sup>c</sup>	Both Sexes	0.0185%	-	-	0.0006%	0.0028%	0.0105%	0.0232%	0.0485%	0.0907%	0.1133%	0.0174%
	Males	0.0220%	-	-	0.0004%	0.0030%	0.0118%	0.0298%	0.0647%	0.1267%	0.1508%	0.0229%
	Females	0.0150%	-	-	0.0008%	0.0026%	0.0092%	0.0168%	0.0334%	0.0612%	0.0914%	0.0128%
Black <sup>c</sup>	Both Sexes	0.0216%	-	-	-	0.0037%	0.0142%	0.0396%	0.0778%	0.1425%	0.1982%	0.0279%
	Males	0.0226%	-	-	-	0.0038%	0.0149%	0.0437%	0.0959%	0.1775%	0.2519%	0.0339%
	Females	0.0208%	-	-	-	0.0036%	0.0135%	0.0360%	0.0635%	0.1184%	0.1728%	0.0240%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0419%	-	-	-	0.0039%	0.0184%	0.0446%	0.1076%	0.2484%	0.4104%	0.0447%
	Males	0.0489%	-	-	-	0.0038%	0.0198%	0.0579%	0.1349%	0.3373%	0.5580%	0.0588%
	Females	0.0355%	-	-	-	0.0040%	0.0172%	0.0332%	0.0848%	0.1807%	0.3191%	0.0345%
Hispanic <sup>d</sup>	Both Sexes	0.0167%	-	-	0.0008%	0.0045%	0.0152%	0.0362%	0.0854%	0.1652%	0.2185%	0.0307%
	Males	0.0181%	-	-	-	0.0049%	0.0153%	0.0449%	0.1108%	0.2139%	0.2667%	0.0382%
	Females	0.0153%	-	-	0.0011%	0.0042%	0.0152%	0.0279%	0.0639%	0.1294%	0.1900%	0.0250%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 25.15  
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	42,302	38,951	34,404	31,472	26,664	19,419	147,128	205,130	15,890	221,020
White <sup>b</sup>	39,059	36,727	32,442	30,099	25,742	18,757	138,326	194,284	15,209	209,493
Black <sup>b</sup>	1,240	964	874	477	389	311	3,555	4,490	172	4,662
Asian/Pacific Islander <sup>c</sup>	855	621	506	+	+	+	2,264	+	+	+
Hispanic <sup>d</sup>	6,181	4,244	3,138	+	+	+	13,563	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0724%	0.0023%	0.0069%	0.0688%	0.1539%	0.1572%	0.0948%	0.0361%	0.0153%	0.0084%	0.0724%
White <sup>c</sup>	0.0889%	0.0024%	0.0084%	0.0851%	0.1925%	0.1939%	0.1127%	0.0426%	0.0175%	0.0101%	0.0890%
Black <sup>c</sup>	0.0158%	0.0016%	0.0027%	0.0109%	0.0310%	0.0338%	0.0300%	0.0102%	-	-	0.0164%
Asian/Pacific Islander <sup>c</sup>	0.0243%	0.0034%	0.0033%	0.0273%	0.0450%	0.0449%	0.0339%	0.0121%	0.0062%	-	0.0233%
Hispanic <sup>d</sup>	0.0542%	0.0024%	0.0095%	0.0704%	0.1296%	0.0987%	0.0513%	0.0195%	-	-	0.0532%

- <sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
- <sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
- <sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table 26.21  
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	165,253	108,492	72,789	52,405	39,963	29,662	398,939	494,969	40,004	534,973
	Males	37,076	23,208	15,370	11,746	8,524	6,003	87,400	107,747	9,115	116,862
	Females	128,177	85,284	57,419	40,659	31,439	23,659	311,539	387,222	30,889	418,111
White <sup>b</sup>	Both Sexes	143,323	95,066	63,951	46,856	35,950	26,905	349,196	435,879	36,272	472,151
	Males	33,136	20,812	13,946	10,718	7,869	5,356	78,612	97,244	7,682	104,926
	Females	110,187	74,254	50,005	36,138	28,081	21,549	270,584	338,635	28,590	367,225
Black <sup>b</sup>	Both Sexes	10,318	6,600	4,050	2,688	1,754	1,345	23,655	28,019	1,674	29,693
	Males	1,687	1,227	596	551	307	308	4,060	4,796	185	4,981
	Females	8,631	5,373	3,454	2,137	1,447	1,037	19,595	23,223	1,489	24,712
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	7,791	4,908	3,245	+	+	+	17,378	+	+	+
	Males	1,462	877	533	+	+	+	3,120	+	+	+
	Females	6,329	4,031	2,712	+	+	+	14,258	+	+	+
Hispanic <sup>d</sup>	Both Sexes	16,337	10,194	6,279	+	+	+	32,811	+	+	+
	Males	2,747	1,743	1,014	+	+	+	5,505	+	+	+
	Females	13,590	8,451	5,265	+	+	+	27,306	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1065%	0.0001%	0.0049%	0.0377%	0.1163%	0.1798%	0.2041%	0.2107%	0.1830%	0.1185%	0.1038%
	Males	0.0463%	-	0.0018%	0.0112%	0.0367%	0.0700%	0.0926%	0.1174%	0.1145%	0.0795%	0.0463%
	Females	0.1651%	-	0.0081%	0.0652%	0.1961%	0.2886%	0.3105%	0.2951%	0.2377%	0.1412%	0.1587%
White <sup>c</sup>	Both Sexes	0.1164%	0.0002%	0.0056%	0.0427%	0.1302%	0.1983%	0.2182%	0.2188%	0.1864%	0.1165%	0.1118%
	Males	0.0521%	-	0.0021%	0.0129%	0.0406%	0.0781%	0.1022%	0.1271%	0.1228%	0.0767%	0.0506%
	Females	0.1803%	-	0.0092%	0.0746%	0.2242%	0.3219%	0.3325%	0.3043%	0.2384%	0.1397%	0.1727%
Black <sup>c</sup>	Both Sexes	0.0503%	-	0.0010%	0.0125%	0.0492%	0.0902%	0.1057%	0.1384%	0.1226%	0.0923%	0.0552%
	Males	0.0174%	-	-	0.0032%	0.0114%	0.0336%	0.0348%	0.0562%	0.0581%	0.0805%	0.0213%
	Females	0.0803%	-	0.0015%	0.0212%	0.0820%	0.1403%	0.1669%	0.2034%	0.1669%	0.0979%	0.0835%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.1004%	-	0.0045%	0.0309%	0.1018%	0.1529%	0.1972%	0.2069%	0.1978%	0.1412%	0.0970%
	Males	0.0383%	-	0.0014%	0.0080%	0.0361%	0.0538%	0.0761%	0.0953%	0.0980%	0.0891%	0.0393%
	Females	0.1572%	-	0.0077%	0.0533%	0.1608%	0.2409%	0.3010%	0.2999%	0.2738%	0.1733%	0.1466%
Hispanic <sup>d</sup>	Both Sexes	0.0673%	-	0.0049%	0.0302%	0.0841%	0.1343%	0.1776%	0.1897%	0.1720%	0.1374%	0.0865%
	Males	0.0221%	-	0.0016%	0.0065%	0.0221%	0.0391%	0.0665%	0.0892%	0.0942%	0.0756%	0.0324%
	Females	0.1136%	-	0.0083%	0.0568%	0.1500%	0.2331%	0.2827%	0.2749%	0.2292%	0.1739%	0.1382%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 27.21  
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	198,998	134,705	90,043	58,795	36,321	21,372	482,542	551,278	12,362	563,640
	Males	151,489	100,571	66,397	43,603	26,844	15,440	362,060	411,638	8,010	419,648
	Females	47,509	34,134	23,646	15,192	9,477	5,932	120,482	139,640	4,352	143,992
White <sup>b</sup>	Both Sexes	181,710	125,929	84,624	55,622	34,763	20,458	447,885	513,591	9,653	523,244
	Males	139,169	94,374	62,500	41,198	25,612	14,794	337,241	384,585	6,140	390,725
	Females	42,541	31,555	22,124	14,424	9,151	5,664	110,644	129,006	3,513	132,519
Black <sup>b</sup>	Both Sexes	8,909	5,139	3,063	1,986	976	569	19,098	20,897	204	21,101
	Males	6,065	3,338	2,187	1,474	826	379	13,065	14,422	166	14,588
	Females	2,844	1,801	876	512	150	190	6,033	6,475	38	6,513
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,498	1,959	1,173	+	+	+	7,124	+	+	+
	Males	2,706	1,515	830	+	+	+	5,447	+	+	+
	Females	792	444	343	+	+	+	1,677	+	+	+
Hispanic <sup>d</sup>	Both Sexes	7,356	4,187	2,631	+	+	+	14,174	+	+	+
	Males	5,467	3,088	2,007	+	+	+	10,562	+	+	+
	Females	1,889	1,099	624	+	+	+	3,612	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1170%	0.0001%	0.0003%	0.0015%	0.0066%	0.0274%	0.1031%	0.3322%	0.7219%	1.0087%	0.1178%
	Males	0.1791%	-	0.0005%	0.0021%	0.0097%	0.0416%	0.1620%	0.5415%	1.2502%	1.9863%	0.2070%
	Females	0.0565%	-	0.0002%	0.0008%	0.0036%	0.0134%	0.0469%	0.1429%	0.3002%	0.4424%	0.0508%
White <sup>c</sup>	Both Sexes	0.1407%	0.0002%	0.0003%	0.0017%	0.0079%	0.0315%	0.1175%	0.3818%	0.8292%	1.1024%	0.1333%
	Males	0.2140%	-	0.0006%	0.0025%	0.0112%	0.0462%	0.1817%	0.6141%	1.4222%	2.1802%	0.2321%
	Females	0.0679%	-	-	0.0010%	0.0045%	0.0164%	0.0542%	0.1653%	0.3441%	0.4738%	0.0575%
Black <sup>c</sup>	Both Sexes	0.0422%	-	-	-	0.0038%	0.0150%	0.0585%	0.1671%	0.3235%	0.4866%	0.0573%
	Males	0.0612%	-	-	-	0.0070%	0.0255%	0.0931%	0.2797%	0.5350%	0.8867%	0.0976%
	Females	0.0250%	-	-	-	-	0.0057%	0.0285%	0.0781%	0.1781%	0.2973%	0.0307%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0496%	-	-	0.0006%	0.0026%	0.0118%	0.0422%	0.1267%	0.3004%	0.5825%	0.0536%
	Males	0.0783%	-	-	-	0.0038%	0.0201%	0.0719%	0.2165%	0.5434%	1.0784%	0.0965%
	Females	0.0232%	-	-	-	0.0015%	0.0044%	0.0167%	0.0519%	0.1154%	0.2760%	0.0226%
Hispanic <sup>d</sup>	Both Sexes	0.0289%	-	-	0.0012%	0.0033%	0.0130%	0.0466%	0.1520%	0.3457%	0.5502%	0.0584%
	Males	0.0424%	-	-	0.0015%	0.0044%	0.0172%	0.0728%	0.2493%	0.6291%	1.0499%	0.1036%
	Females	0.0151%	-	-	0.0008%	0.0022%	0.0087%	0.0219%	0.0696%	0.1375%	0.2548%	0.0262%

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 28.12  
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>f</sup>	Complete <sup>g</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	63,904	55,339	48,098	43,098	36,804	30,537	210,439	304,247	+	+
	Males	33,948	29,297	25,078	22,583	18,590	15,258	110,906	156,819	+	+
	Females	29,956	26,042	23,020	20,515	18,214	15,279	99,533	147,428	+	+
White <sup>b</sup>	Both Sexes	51,809	46,650	40,429	36,395	31,853	26,845	175,284	257,416	+	+
	Males	27,680	24,866	20,948	19,277	16,133	13,540	92,772	133,285	+	+
	Females	24,129	21,784	19,481	17,118	15,720	13,305	82,512	124,131	+	+
Black <sup>b</sup>	Both Sexes	7,556	5,571	4,739	4,240	3,173	2,580	22,104	30,016	+	+
	Males	3,977	2,757	2,516	2,069	1,522	1,189	11,317	14,861	+	+
	Females	3,579	2,814	2,223	2,171	1,651	1,391	10,787	15,155	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,583	1,967	1,774	+	+	+	7,257	+	+	+
	Males	1,329	1,047	968	+	+	+	3,784	+	+	+
	Females	1,254	920	806	+	+	+	3,473	+	+	+
Hispanic <sup>d</sup>	Both Sexes	12,879	9,562	7,492	+	+	+	29,934	+	+	+
	Males	7,136	5,215	3,895	+	+	+	16,246	+	+	+
	Females	5,743	4,347	3,597	+	+	+	13,688	+	+	+

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census.

<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f g</sup> (f) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <35 and >=35.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 29.7  
U.S. Childhood Cancer Survivors at January 1, 2010  
Number of People Previously Diagnosed with Cancer as Children (Ages 0-19 Years) in the United States and Alive January 1, 2010  
By International Classification of Childhood Cancer (ICCC)<sup>a</sup> Selected Group and Subgroup, Age Group, and Sex

Site	Sex	Complete Prevalence Counts by Age at Prevalence							All Ages		
		0-19	20-29	30-39	40-49	50-59	60+	All Ages	Diagnosed between 1975-2009	Diagnosed before 1975	% <sup>b</sup>
									Count	Count	% <sup>b</sup>
All Sites	Both	113,782	93,658	62,583	55,044	36,545	17,500	379,112	304,180	74,932	20
	Male	60,936	49,379	30,706	24,513	15,657	9,268	190,459	156,993	33,466	18
	Female <sup>c</sup>	52,846	44,279	31,877	30,531	20,888	8,232	188,653	147,187	41,466	22
I Leukemia	Both	35,749	20,000	11,075	4,605	1,422	160	73,011	68,730	4,281	6
	Male	19,543	10,789	5,622	2,377	667	72	39,070	37,116	1,954	5
	Female	16,206	9,211	5,453	2,228	755	88	33,941	31,614	2,327	7
Acute lymphocytic leukemia <sup>d</sup>	Both	30,171	16,529	9,420	3,992	375	2	60,489	57,232	3,257	5
	Male	16,573	9,028	4,935	2,252	215	1	33,004	31,293	1,711	5
	Female	13,598	7,501	4,485	1,740	160	1	27,485	25,939	1,546	6
I(b) Acute myeloid leukemia	Both	4,045	2,249	1,145	619	207	2	8,267	7,771	496	6
	Male	2,068	1,135	450	296	113	2	4,064	3,799	265	7
	Female	1,977	1,114	695	323	94	0	4,203	3,972	231	5
II(a) Hodgkin lymphoma	Both	4,514	8,955	8,202	7,786	4,629	1,167	35,253	29,702	5,551	16
	Male	2,501	4,496	4,124	3,816	2,328	594	17,859	14,888	2,971	17
	Female	2,013	4,459	4,078	3,970	2,301	573	17,394	14,814	2,580	15
II(b,c,e) Non-Hodgkin lymphoma	Both	6,442	6,675	4,192	2,780	1,601	1,053	22,743	19,263	3,480	15
	Male	4,223	4,450	2,886	1,743	817	622	14,741	12,861	1,880	13
	Female	2,219	2,225	1,306	1,037	784	431	8,002	6,402	1,600	20
III CNS and misc intracranial & intraspinal neoplasms	Both	20,430	15,174	8,801	8,415	4,806	1,457	59,083	47,793	11,290	19
	Male	11,018	8,350	4,754	4,074	2,243	650	31,089	25,904	5,185	17
	Female	9,412	6,824	4,047	4,341	2,563	807	27,994	21,889	6,105	22
IV Neuroblastoma & other peripheral nervous cell tumor	Both	9,704	3,628	2,190	1,617	1,573	740	19,452	14,922	4,530	23
	Male	4,959	1,854	1,013	517	499	193	9,035	7,717	1,318	15
	Female	4,745	1,774	1,177	1,100	1,074	547	10,417	7,205	3,212	31
VI Renal tumors	Both	7,831	4,651	3,801	4,192	2,323	740	23,538	15,638	7,900	34
	Male	3,894	2,167	1,718	2,356	1,288	431	11,854	7,392	4,462	38
	Female	3,937	2,484	2,083	1,836	1,035	309	11,684	8,246	3,438	29
VIII Malignant bone tumors	Both	3,766	4,194	3,184	2,489	2,320	815	16,768	13,132	3,636	22
	Male	2,252	2,433	1,755	1,237	1,009	434	9,120	7,438	1,682	18
	Female	1,514	1,761	1,429	1,252	1,311	381	7,648	5,694	1,954	26
VIII(a) Osteosarcoma	Both	1,895	2,216	1,804	1,246	1,172	770	9,103	6,951	2,152	24
	Male	1,062	1,285	1,017	601	446	377	4,788	3,831	957	20
	Female	833	931	787	645	726	393	4,315	3,120	1,195	28
VIII(c) Ewing tumor & related sarcomas of bone	Both	1,394	1,377	946	670	405	94	4,886	4,265	621	13
	Male	860	796	462	323	211	34	2,686	2,417	269	10
	Female	534	581	484	347	194	60	2,200	1,848	352	16
IX Soft tissue & other extraosseous sarcomas	Both	6,849	6,964	4,239	4,829	4,161	4,406	31,448	20,922	10,526	33
	Male	3,860	3,714	2,211	2,289	1,962	2,442	16,478	11,169	5,309	32
	Female	2,989	3,250	2,028	2,540	2,199	1,964	14,970	9,753	5,217	35
X Germ cell & trophoblastic tumors & neoplasms of gonads	Both	5,219	7,632	5,531	6,273	6,614	6,468	37,737	23,603	14,134	37
	Male	2,755	5,031	3,302	3,296	3,326	2,935	20,645	13,959	6,686	32
	Female	2,464	2,601	2,229	2,977	3,288	3,533	17,092	9,644	7,448	44

<sup>a</sup> International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.

<sup>b</sup> Percent diagnosed before 1975 of all ages complete prevalence counts.

<sup>c</sup> All sites females excludes breast cancer.

<sup>d</sup> SEER site recode Acute Lymphocytic Leukemia was used instead of the ICCC classification lymphocytic leukemia (Ia) because of statistical modeling requirements.

For more information regarding the modeling of the prevalence estimates, see Mariotto et al. Long-Term Survivors of Childhood Cancers in the United States. Cancer Epidemiol Biomarkers Prev 2009, page 1034.