

Table 19.15

Non-Hodgkin Lymphoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2016

All Races, Males, Ages 55+

Year of Diagnosis

SEER Location	1975- 1979	1980- 1981	1982- 1983	1984- 1985	1986- 1987	1988- 1989	1990- 1991	1992- 1993	1994- 1995	1996- 1997	1998- 1999	2000- 2001	2002- 2003	2004- 2005	2006- 2007	2008- 2009	2010- 2011	2012- 2013	2014- 2016
SEER 9 ^b	3,536	1,732	1,873	2,127	2,322	2,497	2,704	2,810	3,116	3,385	3,551	3,747	4,025	4,397	4,640	4,868	5,190	5,321	8,672
SEER 9 except SF-Oakland	2,886	1,447	1,584	1,791	1,912	2,092	2,226	2,326	2,569	2,840	3,002	3,088	3,395	3,709	3,928	4,113	4,320	4,558	7,317
San Francisco-Oakland SMSA ^c	650	285	289	336	410	405	478	484	547	545	549	659	630	688	712	755	870	763	1,355
San Francisco City/County	178	65	88	104	109	105	132	106	130	126	106	141	108	169	131	136	164	148	261

Age-Adjusted Cancer Incidence Rates^a

By SEER Location and Year of Diagnosis, 1975-2016

All Races, Males, Ages 55+

Year of Diagnosis

SEER Location	1975- 1979	1980- 1981	1982- 1983	1984- 1985	1986- 1987	1988- 1989	1990- 1991	1992- 1993	1994- 1995	1996- 1997	1998- 1999	2000- 2001	2002- 2003	2004- 2005	2006- 2007	2008- 2009	2010- 2011	2012- 2013	2014- 2016
SEER 9 ^b	45.1	52.2	54.0	59.6	63.3	67.3	70.1	70.6	76.1	80.0	81.1	83.1	85.8	90.0	90.1	89.1	89.9	86.2	87.1
SEER 9 except SF-Oakland	44.0	52.1	54.4	60.1	62.0	66.7	68.7	69.2	74.4	79.7	81.3	81.3	85.8	90.1	90.4	89.1	88.8	87.4	87.4
San Francisco-Oakland SMSA ^c	50.8	52.8	52.0	57.1	70.2	70.3	77.3	78.3	85.4	82.1	80.0	93.2	85.5	89.4	88.8	89.0	95.9	80.0	86.0
San Francisco City/County	48.2	44.6	59.8	72.7	73.9	75.7	93.4	74.7	91.4	87.2	72.7	93.4	69.0	104.7	76.4	76.9	88.9	77.6	82.6

^a Rates are per 100,000 males aged 55+ and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).^b The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.^c The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Non-Hodgkin Lymphoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.