

Figure 2a - Incidence

Males, All Cancers, SEER[^] Incidence Rates,
1975-2000 for Whites and Blacks
1992-2000 for American Indian/Alaska Natives, Asian/Pacific Islanders, and Hispanics

Year of Death	White Joinpoint Regression Line Value	Delay - Adjusted White Joinpoint Regression Line Value	Black Joinpoint Regression Line Value	Delay - Adjusted Black Joinpoint Regression Line Value	American Indian/Alaska Native [†] 2 - Year Average Rate	Asian/Pacific Islander [†] 2 - Year Average Rate	Hispanic ^{†#} 2 - Year Average Rate
1975	469.245	468.735	532.341	532.471			
1976	475.891	475.524	547.351	547.411			
1977	482.632	482.411	562.785	562.770			
1978	489.468	489.397	578.654	578.560			
1979	496.401	496.485	594.970	594.793			
1980	503.432	503.675	611.746	611.481			
1981	510.563	510.970	628.995	628.638			
1982	517.795	518.370	632.895	632.967			
1983	525.129	525.877	636.820	637.325			
1984	532.567	533.494	640.769	641.714			
1985	540.111	541.220	644.742	646.132			
1986	547.761	549.058	648.740	650.582			
1987	555.520	557.010	652.762	655.061			
1988	563.389	565.077	656.810	659.572			
1989	571.369	573.261	660.883	664.114			
1990	599.428	602.579	706.340	709.329			
1991	628.865	633.396	754.925	757.623			
1992	659.748	665.790	806.851	809.205			
1993	626.376	632.143	789.492	795.504	336.380	428.701	451.693
1994	594.693	600.196	772.506	782.034	334.402	419.026	443.195
1995	564.612	569.864	755.886	768.793	314.096	404.647	438.859
1996	564.641	574.016	739.623	755.776	295.735	392.459	432.170
1997	564.670	578.199	723.710	742.979	291.202	393.864	421.916
1998	564.699	582.412	708.139	730.399	281.436	391.422	420.503
1999	564.728	586.656	692.904	718.032	265.189	389.014	423.271
2000	564.757	590.930	677.996	705.874	233.993	394.268	414.414

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard.

[†] Rates listed for each year are the average rate from the year listed and the prior year.

[#] Hispanic is not mutually exclusive from whites, blacks, Asian/ Pacific Islanders, and American Indians/Alaska Natives.

[^] Data Source

White and Black Data: SEER 9 (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Asian/Pacific Islanders: SEER 11 (SEER 9 areas, San Jose-Monterey and Los Angeles).

American Indians/Alaskan Natives: SEER 11 and Alaska.

Hispanic: SEER 11 areas excluding cases diagnosed in Detroit and Hawaii.

Figure 2a - Mortality

Males, All Cancers, U.S. Death Rates,
1975-2000 for Whites and Blacks
1992-2000 for American Indian/Alaska Natives, Asian/Pacific Islanders, and Hispanics

Year of Death	White Joinpoint Regression Line Value	Black Joinpoint Regression Line Value	American Indian/Alaska Native [†] 2 - Year Average Rate	Asian/Pacific Islander [†] 2 - Year Average Rate	Hispanic ^{†#} 2 - Year Average Rate
1975	254.786	322.551			
1976	256.835	328.428			
1977	258.900	334.412			
1978	260.981	340.505			
1979	263.079	346.709			
1980	265.195	353.026			
1981	265.703	359.458			
1982	266.213	366.007			
1983	266.724	369.677			
1984	267.236	373.384			
1985	267.749	377.128			
1986	268.262	380.910			
1987	268.777	384.730			
1988	269.293	388.588			
1989	269.809	392.485			
1990	270.327	396.420			
1991	270.846	395.308			
1992	271.366	394.199			
1993	267.656	393.093	169.789	171.433	179.429
1994	263.998	385.435	164.971	170.201	181.241
1995	260.389	377.926	166.059	166.731	182.076
1996	256.830	370.564	173.509	162.817	181.235
1997	253.320	363.345	179.578	157.794	178.683
1998	249.857	356.266	179.083	154.735	178.962
1999	246.442	349.326	173.717	153.705	176.115
2000	243.073	342.521	164.475	152.236	173.395

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard.

[†] Rates listed for each year are the average rate from the year listed and the prior year.

[#] Hispanic is not mutually exclusive from whites, blacks, Asian/ Pacific Islanders, and American Indians/Alaska Natives. Mortality data for Hispanics is included for all states except Connecticut, Oklahoma, New York and New Hampshire

Figure 2b - Incidence

Females, All Cancers, SEER[^] Incidence Rates,
1975-2000 for Whites and Blacks
1992-2000 for American Indian/Alaska Natives, Asian/Pacific Islanders, and Hispanics

Year of Death	White Joinpoint Regression Line Value	Delay - Adjusted White Joinpoint Regression Line Value	Black Joinpoint Regression Line Value	Delay - Adjusted Black Joinpoint Regression Line Value	American Indian/Alaska Native [†] 2 - Year Average Rate	Asian/Pacific Islander [†] 2 - Year Average Rate	Hispanic ^{†#} 2 - Year Average Rate
1975	370.885	370.720	342.789	342.773			
1976	370.000	369.998	346.720	346.743			
1977	369.116	369.277	350.695	350.758			
1978	368.234	368.557	354.716	354.820			
1979	367.355	367.839	358.783	358.929			
1980	373.478	373.792	362.896	363.086			
1981	379.703	379.841	367.057	367.291			
1982	386.032	385.988	371.265	371.545			
1983	392.466	392.234	375.522	375.847			
1984	399.007	398.582	379.827	380.200			
1985	405.658	405.032	384.182	384.603			
1986	412.419	411.587	388.587	389.057			
1987	419.293	418.247	393.042	393.563			
1988	420.610	420.637	397.548	398.120			
1989	421.931	423.041	402.106	402.731			
1990	423.256	425.459	406.717	407.395			
1991	424.586	427.890	411.380	412.113			
1992	425.919	430.335	410.501	412.755			
1993	427.257	432.794	409.623	413.397	238.852	300.616	305.572
1994	428.599	435.267	408.748	414.041	229.051	298.351	305.095
1995	429.945	437.755	407.874	414.686	233.773	298.601	310.310
1996	431.296	440.256	407.003	415.332	238.997	299.646	310.651
1997	432.650	442.772	406.133	415.978	240.660	306.354	308.260
1998	434.009	445.302	405.265	416.626	241.253	312.167	314.469
1999	435.373	447.847	404.399	417.275	231.630	310.556	319.768
2000	436.740	450.406	403.535	417.925	213.812	304.853	311.126

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard.

[†] Rates listed for each year are the average rate from the year listed and the prior year.

[#] Hispanic is not mutually exclusive from whites, blacks, Asian/ Pacific Islanders, and American Indians/Alaska Natives.

[^] Data Source

White and Black Data: SEER 9 (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Asian/Pacific Islanders: SEER 11 (SEER 9 areas, San Jose-Monterey and Los Angeles).

American Indians/Alaskan Natives: SEER 11 and Alaska.

Hispanic: SEER 11 areas excluding cases diagnosed in Detroit and Hawaii.

Figure 2b - Mortality

Females, All Cancers, U.S. Death Rates,
1975-2000 for Whites and Blacks
1992-2000 for American Indian/Alaska Natives, Asian/Pacific Islanders, and Hispanics

Year of Death	White Joinpoint Regression Line Value	Black Joinpoint Regression Line Value	American Indian/Alaska Native [†] 2 - Year Average Rate	Asian/Pacific Islander [†] 2 - Year Average Rate	Hispanic ^{†#} 2 - Year Average Rate
1975	159.670	177.029			
1976	160.553	178.822			
1977	161.440	180.634			
1978	162.333	182.464			
1979	163.230	184.313			
1980	164.132	186.180			
1981	165.040	188.066			
1982	165.952	189.971			
1983	166.869	191.896			
1984	167.792	193.840			
1985	168.719	195.804			
1986	169.652	197.787			
1987	170.590	199.791			
1988	171.533	201.815			
1989	172.481	203.860			
1990	173.435	205.925			
1991	173.154	208.011			
1992	172.874	206.667			
1993	172.594	205.331	115.011	106.375	113.718
1994	172.314	204.004	113.272	108.438	111.798
1995	172.035	202.685	115.439	108.516	112.003
1996	169.764	201.375	122.159	105.696	113.009
1997	167.523	200.074	119.565	103.150	112.962
1998	165.311	198.780	115.891	101.564	111.406
1999	165.888	197.496	115.946	101.570	111.696
2000	166.467	196.219	111.680	101.781	112.712

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard.

[†] Rates listed for each year are the average rate from the year listed and the prior year.

[#] Hispanic is not mutually exclusive from whites, blacks, Asian/ Pacific Islanders, and American Indians/Alaska Natives. Mortality data for Hispanics is included for all states except Connecticut, Oklahoma, New York and New Hampshire