

Table XVI-1
MELANOMAS OF SKIN (Invasive)

TRENDS IN SEER INCIDENCE AND U.S. MORTALITY, BY RACE AND SEX

| | All Races | | | Whites | | | Blacks | | |
|---|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>INCIDENCE TRENDS (expressed in percents)§:</u> | | | | | | | | | |
| <u>1973-94</u> | | | | | | | | | |
| All Ages | | | | | | | | | |
| Percent Change (PC) | 113.0 | 149.0 | 81.9 | 119.7 | 154.2 | 89.5 | 25.5 | 95.7 | -1.8 |
| Est. Annual PC (EAPC) | 3.7★ | 4.6★ | 2.9★ | 3.9★ | 4.7★ | 3.2★ | -0.5 | 0.7 | -1.3 |
| Under 65 | | | | | | | | | |
| Percent Change (PC) | 91.6 | 112.3 | 72.4 | 97.1 | 114.5 | 80.3 | 57.2 | 285.3 | -30.0 |
| Est. Annual PC (EAPC) | 3.2★ | 3.6★ | 2.7★ | 3.3★ | 3.8★ | 2.9★ | -0.4 | - | -2.9 |
| 65 and over | | | | | | | | | |
| Percent Change (PC) | 177.1 | 240.0 | 115.4 | 190.2 | 256.5 | 123.5 | -0.6 | -45.1 | 27.8 |
| Est. Annual PC (EAPC) | 5.3★ | 6.5★ | 3.7★ | 5.5★ | 6.8★ | 4.0★ | -0.6 | - | 0.0 |
| <u>1975-79 - ALL AGES</u> | | | | | | | | | |
| Percent Change (PC) | 15.2 | 15.9 | 16.5 | 18.0 | 18.5 | 19.8 | -36.8 | -65.1 | -13.0 |
| Est. Annual PC (EAPC) | 4.9★ | 5.3★ | 4.9★ | 5.6★ | 5.9★ | 5.8★ | -9.4 | -23.8 | -0.1 |
| <u>1990-94 - ALL AGES</u> | | | | | | | | | |
| Percent Change (PC) | 4.5 | 8.5 | -0.3 | 2.9 | 6.7 | -1.6 | -19.4 | -42.9 | 6.7 |
| Est. Annual PC (EAPC) | 1.7♦ | 3.0★ | 0.2♦ | 1.4♦ | 2.6♦ | -0.1♦ | -4.4 | -14.2 | 3.1 |
| <u>MORTALITY TRENDS (expressed in percents)*:</u> | | | | | | | | | |
| <u>1973-94</u> | | | | | | | | | |
| All Ages | | | | | | | | | |
| Percent Change (PC) | 33.2 | 46.1 | 17.7 | 37.0 | 50.0 | 20.9 | -2.9 | -25.6 | 18.1 |
| Est. Annual PC (EAPC) | 1.4★ | 2.0★ | 0.7★ | 1.6★ | 2.1★ | 0.8★ | -0.2 | -0.9 | 0.4 |
| Under 65 | | | | | | | | | |
| Percent Change (PC) | 9.9 | 16.5 | -0.5 | 14.6 | 20.9 | 4.0 | -27.1 | -25.6 | -27.5 |
| Est. Annual PC (EAPC) | 0.5★ | 0.8★ | -0.1 | 0.7★ | 1.0★ | 0.2 | -1.7★ | -1.8★ | -1.4 |
| 65 and over | | | | | | | | | |
| Percent Change (PC) | 76.7 | 97.0 | 51.9 | 80.5 | 101.5 | 53.7 | 20.3 | -25.7 | 71.9 |
| Est. Annual PC (EAPC) | 2.8★ | 3.6★ | 1.8★ | 2.9★ | 3.7★ | 1.8★ | 0.9 | -0.1 | 1.8★ |
| <u>1975-79 - ALL AGES</u> | | | | | | | | | |
| Percent Change (PC) | 9.7 | 12.0 | 7.0 | 10.3 | 12.8 | 7.3 | -0.7 | -10.1 | 16.0 |
| Est. Annual PC (EAPC) | 3.5★ | 4.1★ | 2.9 | 3.7★ | 4.3★ | 3.0 | 0.0 | -1.8 | 3.8 |
| <u>1990-94 - ALL AGES</u> | | | | | | | | | |
| Percent Change (PC) | -1.3 | -0.5 | -2.4 | -0.7 | 0.3 | -2.1 | -14.1 | -31.5 | -1.5 |
| Est. Annual PC (EAPC) | -0.6♦ | -0.1♦ | -1.3♦ | -0.4♦ | 0.1♦ | -1.2♦ | -4.8 | -10.1 | -1.8 |

The Est. Annual PC is the Estimated Annual Percent Change (EAPC) over the time interval.
SEER Program.

* NCHS public use tape.

★ The EAPC is significantly different from zero ($p < .05$).

◆ The EAPC for 1990-94 is significantly different from the EAPC for 1975-79 ($p < .05$).

◊ The EAPC for 1990-94 is significantly different from the EAPC for 1975-79 ($p < .10$).

- Statistic could not be calculated.

Table XVI-2
MELANOMAS OF SKIN (Invasive)

SEER INCIDENCE RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE AND SEX

| | All Races | | | Whites | | | Blacks | | |
|------------------------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>AGE-ADJUSTED RATES</u> | | | | | | | | | |
| YEAR OF DIAGNOSIS: | | | | | | | | | |
| 1973 | 5.7 | 6.1 | 5.4 | 6.2 | 6.7 | 5.9 | 0.7 | 0.6 | 0.7 |
| 1974 | 5.9 | 6.2 | 5.7 | 6.5 | 6.9 | 6.3 | 0.5 | 0.2 | 0.7 |
| 1975 | 6.7 | 7.1 | 6.3 | 7.3 | 7.8 | 6.9 | 0.9 | 1.1 | 0.8 |
| 1976 | 6.9 | 7.4 | 6.5 | 7.6 | 8.1 | 7.1 | 1.4 | 1.7 | 1.2 |
| 1977 | 7.5 | 8.2 | 7.0 | 8.3 | 9.1 | 7.8 | 0.7 | 0.7 | 0.7 |
| 1978 | 7.5 | 8.0 | 7.3 | 8.5 | 8.9 | 8.3 | 0.6 | 0.3 | 0.8 |
| 1979 | 8.1 | 8.9 | 7.6 | 9.0 | 9.9 | 8.5 | 0.8 | 0.7 | 1.0 |
| 1980 | 8.9 | 9.9 | 8.2 | 10.1 | 11.2 | 9.3 | 0.7 | 0.7 | 0.7 |
| 1981 | 9.4 | 10.2 | 8.8 | 10.4 | 11.2 | 10.0 | 1.1 | 1.7 | 0.6 |
| 1982 | 9.4 | 10.5 | 8.6 | 10.6 | 11.7 | 9.8 | 0.8 | 1.0 | 1.0 |
| 1983 | 9.3 | 10.5 | 8.4 | 10.4 | 11.8 | 9.4 | 0.8 | 0.4 | 0.4 |
| 1984 | 9.6 | 10.9 | 8.7 | 10.7 | 12.2 | 9.7 | 0.6 | 0.1 | 1.0 |
| 1985 | 10.7 | 12.6 | 9.4 | 11.9 | 14.0 | 10.4 | 0.8 | 0.2 | 1.0 |
| 1986 | 11.2 | 13.1 | 9.8 | 12.3 | 14.3 | 10.8 | 0.8 | 0.2 | 0.9 |
| 1987 | 11.4 | 13.2 | 10.1 | 12.6 | 14.5 | 11.4 | 1.1 | 1.1 | 0.9 |
| 1988 | 10.7 | 12.5 | 9.5 | 12.3 | 14.2 | 10.9 | 0.9 | 0.9 | 0.9 |
| 1989 | 11.4 | 13.6 | 9.8 | 13.0 | 15.5 | 11.2 | 1.0 | 1.2 | 0.8 |
| 1990 | 11.5 | 13.8 | 9.8 | 13.1 | 15.6 | 11.3 | 1.0 | 1.2 | 0.6 |
| 1991 | 12.2 | 14.6 | 10.5 | 14.1 | 16.7 | 12.1 | 1.0 | 1.3 | 0.7 |
| 1992 | 12.3 | 15.1 | 10.2 | 14.0 | 17.1 | 11.7 | 0.9 | 0.3 | 0.3 |
| 1993 | 12.2 | 15.0 | 10.1 | 13.8 | 16.8 | 11.5 | 0.6 | 0.3 | 0.7 |
| 1994 | 12.5 | 15.7 | 10.1 | 14.2 | 17.7 | 11.6 | 0.9 | 1.1 | 0.7 |
| <u>AGE-ADJUSTED RATES, 1990-94</u> | | | | | | | | | |
| AGE AT DIAGNOSIS: | | | | | | | | | |
| All ages | 12.2 | 14.9 | 10.1 | 13.8 | 16.8 | 11.6 | 0.7 | 0.9 | 0.6 |
| Under 65 | 9.3 | 10.3 | 8.5 | 10.7 | 11.6 | 9.8 | 0.4 | 0.5 | 0.3 |
| 65 and over | 38.4 | 57.0 | 25.5 | 42.7 | 63.9 | 28.2 | 3.7 | 3.6 | 3.6 |
| All ages (world std.) ⁸ | 10.8 | 12.8 | 9.3 | 12.3 | 14.5 | 10.7 | 0.6 | 0.7 | 0.4 |
| <u>AGE-SPECIFIC RATES, 1990-94</u> | | | | | | | | | |
| AGE AT DIAGNOSIS: | | | | | | | | | |
| 0-4 | 0.1 | 0.0 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 |
| 5-9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 |
| 10-14 | 0.3 | 0.3 | 0.4 | 0.4 | 0.3 | 0.4 | 0.0 | 0.0 | 0.0 |
| 15-19 | 1.6 | 1.2 | 2.0 | 1.8 | 1.4 | 2.3 | 0.0 | 0.0 | 0.0 |
| 20-24 | 4.0 | 2.7 | 5.4 | 4.6 | 3.2 | 6.0 | 0.0 | 0.0 | 0.0 |
| 25-29 | 6.6 | 4.9 | 8.1 | 7.7 | 9.5 | 9.6 | 0.0 | 0.0 | 0.1 |
| 30-34 | 10.2 | 8.3 | 12.2 | 11.9 | 10.5 | 14.4 | 0.3 | 0.5 | 0.1 |
| 35-39 | 13.3 | 12.9 | 13.8 | 15.6 | 15.0 | 16.2 | 0.4 | 0.9 | 0.0 |
| 40-44 | 16.8 | 17.3 | 16.4 | 19.5 | 19.8 | 19.3 | 0.2 | 1.0 | 0.0 |
| 45-49 | 20.8 | 22.7 | 18.9 | 23.6 | 25.2 | 21.9 | 0.4 | 0.9 | 0.4 |
| 50-54 | 22.4 | 27.3 | 17.7 | 25.7 | 31.0 | 20.5 | 1.2 | 2.1 | 2.2 |
| 55-59 | 27.6 | 35.3 | 20.4 | 31.5 | 40.1 | 23.3 | 2.4 | 2.2 | 2.6 |
| 60-64 | 31.5 | 42.2 | 21.9 | 35.5 | 46.9 | 24.9 | 1.8 | 2.9 | 0.9 |
| 65-69 | 34.0 | 48.5 | 22.0 | 38.6 | 54.9 | 24.9 | 1.7 | 2.6 | 1.0 |
| 70-74 | 40.2 | 56.3 | 27.9 | 44.9 | 63.1 | 30.9 | 2.6 | 4.2 | 2.5 |
| 75-79 | 40.3 | 63.5 | 24.6 | 44.6 | 71.0 | 26.9 | 4.8 | 5.1 | 4.4 |
| 80-84 | 41.1 | 64.6 | 28.1 | 44.8 | 71.9 | 30.2 | 7.4 | 8.4 | 7.5 |
| 85+ | 42.7 | 70.9 | 31.8 | 46.3 | 79.0 | 34.1 | 7.9 | 8.1 | 7.5 |

SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, except where noted.
 SEER Program. Rates are per 100,000 and are age-adjusted to the world standard population.

Table XVI-3
MELANOMAS OF SKIN (Invasive)

U.S. MORTALITY RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES*, BY RACE AND SEX

| | All Races | | | Whites | | | Blacks | | |
|------------------------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>AGE-ADJUSTED RATES</u> | | | | | | | | | |
| YEAR OF DEATH: | | | | | | | | | |
| 1973 | 1.6 | 2.0 | 1.3 | 1.7 | 2.2 | 1.4 | 0.4 | 0.6 | 0.4 |
| 1974 | 1.7 | 2.2 | 1.3 | 1.9 | 2.5 | 1.4 | 0.3 | 0.4 | 0.3 |
| 1975 | 1.7 | 2.2 | 1.3 | 1.9 | 2.4 | 1.4 | 0.4 | 0.5 | 0.4 |
| 1976 | 1.9 | 2.4 | 1.4 | 2.0 | 2.6 | 1.6 | 0.4 | 0.6 | 0.3 |
| 1977 | 1.9 | 2.4 | 1.4 | 2.1 | 2.6 | 1.6 | 0.4 | 0.5 | 0.3 |
| 1978 | 1.9 | 2.5 | 1.5 | 2.1 | 2.7 | 1.7 | 0.4 | 0.5 | 0.4 |
| 1979 | 2.0 | 2.6 | 1.5 | 2.2 | 2.9 | 1.7 | 0.4 | 0.5 | 0.4 |
| 1980 | 1.9 | 2.5 | 1.4 | 2.1 | 2.8 | 1.5 | 0.5 | 0.4 | 0.5 |
| 1981 | 2.0 | 2.6 | 1.5 | 2.2 | 2.8 | 1.7 | 0.4 | 0.5 | 0.4 |
| 1982 | 2.0 | 2.7 | 1.5 | 2.2 | 3.0 | 1.6 | 0.4 | 0.5 | 0.4 |
| 1983 | 2.0 | 2.7 | 1.5 | 2.3 | 3.0 | 1.7 | 0.4 | 0.5 | 0.3 |
| 1984 | 2.1 | 2.8 | 1.5 | 2.3 | 3.1 | 1.6 | 0.4 | 0.4 | 0.4 |
| 1985 | 2.1 | 2.8 | 1.5 | 2.3 | 3.1 | 1.7 | 0.4 | 0.5 | 0.4 |
| 1986 | 2.1 | 2.9 | 1.5 | 2.4 | 3.2 | 1.7 | 0.4 | 0.6 | 0.4 |
| 1987 | 2.2 | 3.1 | 1.5 | 2.4 | 3.4 | 1.7 | 0.4 | 0.5 | 0.4 |
| 1988 | 2.2 | 3.0 | 1.5 | 2.4 | 3.3 | 1.7 | 0.4 | 0.4 | 0.3 |
| 1989 | 2.2 | 3.1 | 1.5 | 2.5 | 3.4 | 1.7 | 0.4 | 0.5 | 0.4 |
| 1990 | 2.3 | 3.1 | 1.6 | 2.5 | 3.4 | 1.8 | 0.4 | 0.5 | 0.4 |
| 1991 | 2.2 | 3.2 | 1.5 | 2.5 | 3.5 | 1.6 | 0.5 | 0.5 | 0.4 |
| 1992 | 2.2 | 3.1 | 1.5 | 2.5 | 3.5 | 1.7 | 0.4 | 0.5 | 0.3 |
| 1993 | 2.2 | 3.2 | 1.5 | 2.5 | 3.5 | 1.7 | 0.4 | 0.5 | 0.3 |
| 1994 | 2.2 | 3.1 | 1.5 | 2.5 | 3.5 | 1.7 | 0.3 | 0.4 | 0.3 |
| <u>AGE-ADJUSTED RATES, 1990-94</u> | | | | | | | | | |
| AGE AT DEATH: | | | | | | | | | |
| All ages | 2.2 | 3.1 | 1.5 | 2.5 | 3.5 | 1.7 | 0.4 | 0.4 | 0.4 |
| Under 65 | 1.4 | 1.8 | 0.9 | 1.6 | 2.1 | 1.1 | 0.2 | 0.1 | 0.1 |
| 65 and over | 10.1 | 15.2 | 6.8 | 11.0 | 16.6 | 7.3 | 2.6 | 2.6 | 2.7 |
| All ages (world std.)N | 1.8 | 2.6 | 1.3 | 2.1 | 2.9 | 1.4 | 0.3 | 0.3 | 0.3 |
| <u>AGE-SPECIFIC RATES, 1990-94</u> | | | | | | | | | |
| AGE AT DEATH: | | | | | | | | | |
| 0-4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 5-9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 10-14 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 15-19 | 0.1 | 0.1 | 0.0 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 |
| 20-24 | 0.2 | 0.2 | 0.1 | 0.3 | 0.3 | 0.2 | 0.0 | 0.0 | 0.0 |
| 25-29 | 0.5 | 0.5 | 0.5 | 0.6 | 0.6 | 0.5 | 0.1 | 0.1 | 0.1 |
| 30-34 | 0.9 | 1.1 | 0.7 | 1.1 | 1.3 | 0.8 | 0.1 | 0.1 | 0.1 |
| 35-39 | 1.5 | 1.9 | 1.1 | 1.8 | 2.3 | 1.3 | 0.2 | 0.2 | 0.1 |
| 40-44 | 2.2 | 2.8 | 1.7 | 2.3 | 3.2 | 2.0 | 0.2 | 0.2 | 0.2 |
| 45-49 | 3.0 | 3.9 | 2.1 | 3.5 | 4.5 | 2.5 | 0.3 | 0.3 | 0.3 |
| 50-54 | 3.8 | 5.0 | 2.6 | 4.4 | 5.7 | 3.0 | 0.4 | 0.4 | 0.3 |
| 55-59 | 4.9 | 6.9 | 3.2 | 5.6 | 7.8 | 3.6 | 0.7 | 0.7 | 0.7 |
| 60-64 | 6.5 | 9.3 | 4.0 | 7.2 | 10.3 | 4.5 | 1.2 | 1.8 | 0.7 |
| 65-69 | 7.7 | 11.2 | 4.9 | 8.5 | 12.3 | 5.4 | 1.6 | 1.8 | 1.8 |
| 70-74 | 9.6 | 14.2 | 6.2 | 10.5 | 15.4 | 6.7 | 2.2 | 2.2 | 2.2 |
| 75-79 | 11.4 | 17.5 | 7.3 | 12.2 | 18.9 | 7.8 | 3.5 | 4.1 | 3.1 |
| 80-84 | 13.1 | 19.8 | 9.5 | 14.0 | 21.4 | 10.0 | 3.6 | 4.1 | 4.1 |
| 85+ | 15.6 | 25.2 | 12.0 | 16.7 | 27.4 | 12.6 | 5.0 | 4.5 | 5.2 |

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, except where noted.

N NCHS public use tape. Rates are per 100,000 and are age-adjusted to the world standard population.

Table XVI-4
MELANOMAS OF SKIN (Invasive)

SURVIVAL RATES, BY RACE, SEX, DIAGNOSIS YEAR, STAGE AND AGE

| | All Races | | | Whites | | | Blacks | | |
|--|-----------|--------|---------|--------|--------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>5-YR RELATIVE SURVIVAL RATES</u> | | | | | | | | | |
| YEAR OF DIAGNOSIS: | | | | | | | | | |
| 1960-63 ^a | - | - | - | 60 | 51 | 68 | - | - | - |
| 1970-73 ^a | - | - | - | 68 | 62 | 75 | - | - | - |
| 1974-76 ^b | 79.9 | 74.8 | 84.9 | 80.2 | 75.2 | 85.1 | 66.4↑ | - | - |
| 1977-79 ^b | 81.8 | 77.4 | 86.0 | 82.1 | 77.7 | 86.3 | 50.8↑ | - | - |
| 1980-82 ^b | 82.7 | 77.2 | 88.1 | 82.9 | 77.6 | 88.1 | 59.5↑ | - | - |
| 1983-85 ^b | 84.8 | 80.6 | 89.1 | 84.9 | 80.7 | 89.3 | 75.6↑ | - | 71.6↑ |
| 1986-93 ^b | 87.9S | 85.1S | 90.9S | 88.1S | 85.4S | 91.1S | 66.8↑ | 50.0↑ | 78.7↑ |
| 1986-93 ^b | | | | | | | | | |
| ALL STAGES | 87.9 | 85.1 | 90.9 | 88.1 | 85.4 | 91.1 | 66.8↑ | 50.0↑ | 78.7↑ |
| LOCALIZED | 95.0 | 93.9 | 96.2 | 95.1 | 94.0 | 96.2 | 86.5↑ | - | 92.5↑ |
| REGIONAL | 60.8 | 57.4 | 65.7 | 61.2 | 57.7 | 66.1 | - | - | - |
| DISTANT | 15.9 | 15.8 | 16.0 | 16.1 | 15.8 | 16.7 | - | - | - |
| UNSTAGED | 70.5 | 66.5 | 76.0 | 70.9 | 66.8 | 76.4 | - | - | - |
| <u>STAGE DISTRIBUTION (%) 1986-93^b</u> | | | | | | | | | |
| ALL STAGES | | | | | | | | | |
| Number of cases | 19,910 | 10,696 | 9,214 | 19,709 | 10,597 | 9,112 | 102 | 49 | 53 |
| Percent | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% |
| LOCALIZED | 82 | 79 | 84 | 82 | 79 | 84 | 60 | 39 | 79 |
| REGIONAL | 8 | 9 | 7 | 8 | 9 | 7 | 20 | 33 | 8 |
| DISTANT | 4 | 5 | 3 | 4 | 5 | 3 | 14 | 24 | 4 |
| UNSTAGED | 6 | 7 | 6 | 6 | 7 | 6 | 7 | 4 | 9 |
| <u>5-YR RELATIVE SURVIVAL RATES, 1986-93^b</u> | | | | | | | | | |
| AGE AT DIAGNOSIS: | | | | | | | | | |
| <45 | 90.1 | 85.4 | 93.9 | 90.2 | 85.5 | 94.0 | - | - | - |
| 45-54 | 87.0 | 83.9 | 90.8 | 87.2 | 84.0 | 91.0 | - | - | - |
| 55-64 | 87.6 | 85.3 | 90.9 | 87.7 | 85.5 | 90.9 | - | - | - |
| 65-74 | 87.5 | 86.8 | 88.8 | 87.9 | 87.2 | 89.0 | - | - | - |
| 75+ | 82.2 | 81.5 | 81.1 | 82.9 | 82.2 | 81.8 | 63.1↑ | - | - |
| Under 65 | 88.7 | 84.9 | 92.6 | 88.8 | 85.1 | 92.6 | 71.3↑ | 53.3↑ | 87.6↑ |
| 65 and over | 85.8 | 86.0 | 85.6 | 86.3 | 86.5 | 86.0 | 65.3↑ | - | 71.0↑ |

^a Rates are based on End Results data from a series of hospital registries and one population-based registry.

^b Rates are from the SEER Program. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients through 1994.

S The difference in rates between 1974-76 and 1986-93 is statistically significant ($p<.05$).

\uparrow The standard error of the survival rate is between 5 and 10 percentage points.

\downarrow The standard error of the survival rate is greater than 10 percentage points.

- Statistic could not be calculated.

Table XVI-5
MELANOMAS OF SKIN (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Males and Females

Year of Diagnosis/Death

| | 1973 | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 |
|---------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Incidence | | | | | | | | | | | | | | | | | | | | | | |
| SEER | 5.7 | 5.9 | 6.7 | 6.9 | 7.5 | 7.5 | 8.1 | 8.9 | 9.4 | 9.4 | 9.3 | 9.6 | 10.7 | 11.2 | 11.4 | 10.7 | 11.4 | 11.5 | 12.2 | 12.3 | 12.2 | 12.5 |
| Mortality | | | | | | | | | | | | | | | | | | | | | | |
| U.S. | 1.6 | 1.7 | 1.7 | 1.9 | 1.9 | 1.9 | 2.0 | 1.9 | 2.0 | 2.0 | 2.0 | 2.1 | 2.1 | 2.1 | 2.1 | 2.2 | 2.2 | 2.3 | 2.2 | 2.2 | 2.2 | 2.2 |
| SEER | 1.5 | 1.6 | 1.6 | 1.7 | 1.7 | 1.8 | 1.8 | 1.9 | 2.0 | 1.9 | 1.9 | 2.1 | 1.9 | 1.9 | 2.1 | 2.1 | 2.0 | 2.0 | 2.2 | 2.0 | 2.1 | 1.9 |
| Relative Survival Rates (SEER) | | | | | | | | | | | | | | | | | | | | | | |
| 1-year | 90.9 | 92.2 | 94.9 | 93.5 | 95.1 | 93.8 | 94.6 | 96.7 | 94.4 | 95.2 | 96.5 | 96.3 | 95.7 | 96.8 | 96.2 | 97.0 | 95.9 | 96.5 | 97.2 | 96.2 | 96.6 | |
| 2-year | 84.5 | 87.6 | 89.7 | 89.1 | 89.4 | 89.5 | 90.9 | 92.4 | 90.2 | 91.0 | 92.1 | 92.7 | 92.7 | 93.1 | 92.8 | 93.2 | 93.5 | 93.4 | 94.1 | 93.2 | | |
| 3-year | 78.6 | 82.8 | 85.6 | 85.4 | 85.9 | 85.7 | 87.0 | 88.3 | 84.6 | 87.8 | 88.6 | 89.9 | 89.4 | 90.8 | 90.2 | 91.0 | 90.7 | 91.3 | 91.6 | | | |
| 4-year | 74.7 | 79.6 | 82.1 | 83.1 | 83.1 | 83.3 | 83.6 | 85.7 | 82.0 | 85.4 | 86.4 | 86.9 | 87.5 | 89.4 | 88.5 | 89.7 | 88.8 | 89.0 | | | | |
| 5-year | 72.0 | 76.0 | 81.1 | 81.9 | 81.5 | 81.2 | 82.5 | 83.9 | 79.7 | 84.4 | 83.7 | 84.7 | 85.7 | 87.9 | 86.8 | 89.0 | 86.9 | | | | | |
| 6-year | 70.1 | 73.4 | 79.1 | 80.4 | 80.6 | 79.3 | 80.2 | 82.9 | 78.7 | 83.7 | 82.2 | 83.6 | 84.5 | 86.9 | 86.0 | 87.0 | | | | | | |
| 7-year | 69.4 | 72.1 | 77.9 | 78.7 | 79.4 | 78.2 | 79.5 | 82.0 | 78.2 | 82.4 | 81.9 | 83.4 | 83.4 | 86.4 | 85.0 | | | | | | | |
| 8-year | 68.4 | 71.3 | 76.4 | 77.4 | 78.5 | 77.4 | 78.9 | 80.9 | 78.2 | 81.8 | 80.6 | 83.0 | 83.2 | 85.5 | | | | | | | | |
| 9-year | 68.0 | 69.9 | 74.5 | 76.6 | 77.7 | 77.4 | 78.2 | 80.4 | 77.3 | 81.3 | 80.6 | 81.8 | 82.7 | | | | | | | | | |
| 10-year | 67.6 | 69.3 | 72.8 | 76.6 | 76.9 | 77.4 | 78.2 | 80.2 | 77.3 | 81.2 | 80.1 | 80.8 | | | | | | | | | | |
| 11-year | 67.4 | 69.2 | 72.3 | 75.0 | 76.5 | 76.9 | 78.2 | 79.8 | 77.1 | 80.8 | 80.1 | | | | | | | | | | | |
| 12-year | 66.6 | 69.2 | 71.0 | 75.0 | 76.4 | 76.6 | 78.2 | 78.7 | 77.1 | 80.7 | | | | | | | | | | | | |
| 13-year | 66.1 | 69.2 | 71.0 | 74.7 | 75.9 | 76.6 | 78.1 | 78.2 | 77.1 | | | | | | | | | | | | | |
| 14-year | 66.1 | 69.1 | 69.4 | 74.3 | 75.8 | 76.4 | 78.1 | 77.8 | | | | | | | | | | | | | | |
| 15-year | 66.1 | 69.0 | 69.1 | 74.3 | 75.1 | 75.1 | 78.1 | | | | | | | | | | | | | | | |
| 16-year | 66.0 | 68.0 | 69.1 | 74.1 | 75.1 | 74.2 | | | | | | | | | | | | | | | | |
| 17-year | 66.0 | 67.1 | 69.1 | 73.7 | 75.1 | | | | | | | | | | | | | | | | | |
| 18-year | 66.0 | 67.1 | 69.1 | 72.8 | | | | | | | | | | | | | | | | | | |
| 19-year | 66.0 | 67.1 | 69.1 | | | | | | | | | | | | | | | | | | | |
| 20-year | 66.0 | 67.1 | | | | | | | | | | | | | | | | | | | | |
| 21-year | 66.0 | | | | | | | | | | | | | | | | | | | | | |

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table XVI-6
MELANOMAS OF SKIN (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Males

Year of Diagnosis/Death

| | 1973 | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 |
|---------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Incidence | | | | | | | | | | | | | | | | | | | | | | |
| SEER | 6.1 | 6.2 | 7.1 | 7.4 | 8.2 | 8.0 | 8.9 | 9.9 | 10.2 | 10.5 | 10.5 | 10.9 | 12.6 | 13.1 | 13.2 | 12.5 | 13.6 | 13.8 | 14.6 | 15.1 | 15.0 | 15.7 |
| Mortality | | | | | | | | | | | | | | | | | | | | | | |
| U.S. | 2.0 | 2.2 | 2.2 | 2.4 | 2.4 | 2.5 | 2.6 | 2.5 | 2.6 | 2.7 | 2.7 | 2.8 | 2.8 | 2.9 | 3.1 | 3.0 | 3.1 | 3.1 | 3.2 | 3.1 | 3.2 | 3.1 |
| SEER | 1.8 | 2.0 | 1.9 | 2.1 | 2.1 | 2.3 | 2.3 | 2.5 | 2.6 | 2.7 | 2.7 | 2.9 | 2.5 | 2.5 | 3.0 | 2.8 | 2.5 | 2.6 | 3.2 | 2.8 | 2.9 | 2.7 |
| Relative Survival Rates (SEER) | | | | | | | | | | | | | | | | | | | | | | |
| 1-year | 90.5 | 89.6 | 92.6 | 91.6 | 93.7 | 92.3 | 92.6 | 95.8 | 93.2 | 92.7 | 95.2 | 94.2 | 94.6 | 95.8 | 95.3 | 95.8 | 94.6 | 95.7 | 96.3 | 95.9 | 95.8 | |
| 2-year | 80.7 | 83.3 | 84.9 | 85.4 | 87.2 | 86.5 | 87.8 | 89.6 | 88.1 | 87.2 | 88.8 | 89.5 | 90.6 | 91.1 | 90.6 | 91.8 | 92.1 | 91.6 | 93.3 | 92.1 | | |
| 3-year | 73.0 | 78.2 | 80.4 | 81.5 | 83.3 | 81.7 | 83.1 | 84.6 | 80.3 | 84.0 | 85.5 | 85.7 | 86.5 | 88.2 | 87.6 | 88.9 | 88.5 | 89.2 | 90.7 | | | |
| 4-year | 69.7 | 75.0 | 76.6 | 78.8 | 80.0 | 78.4 | 78.3 | 81.4 | 77.2 | 81.4 | 83.3 | 81.7 | 84.1 | 86.3 | 85.7 | 87.5 | 86.1 | 86.3 | | | | |
| 5-year | 66.5 | 70.3 | 75.5 | 77.4 | 77.9 | 76.5 | 77.7 | 77.6 | 74.5 | 79.4 | 80.5 | 79.1 | 82.1 | 84.4 | 83.2 | 86.9 | 83.9 | | | | | |
| 6-year | 63.9 | 66.7 | 73.0 | 75.4 | 76.7 | 74.1 | 75.2 | 76.8 | 72.9 | 78.4 | 79.4 | 78.2 | 81.0 | 82.9 | 82.6 | 84.7 | | | | | | |
| 7-year | 62.9 | 64.6 | 71.7 | 73.7 | 75.2 | 72.5 | 73.2 | 75.7 | 72.9 | 77.3 | 79.4 | 78.2 | 80.1 | 82.3 | 81.0 | | | | | | | |
| 8-year | 61.2 | 63.1 | 69.7 | 72.6 | 74.7 | 71.6 | 71.7 | 74.3 | 72.7 | 76.8 | 78.0 | 78.2 | 79.8 | 81.5 | | | | | | | | |
| 9-year | 60.4 | 62.2 | 67.8 | 71.9 | 74.1 | 71.6 | 70.8 | 73.5 | 71.1 | 76.4 | 78.0 | 77.1 | 79.4 | | | | | | | | | |
| 10-year | 59.8 | 61.8 | 65.8 | 71.9 | 73.0 | 71.6 | 70.8 | 73.4 | 71.1 | 76.4 | 78.0 | 76.1 | | | | | | | | | | |
| 11-year | 59.8 | 61.4 | 65.0 | 71.4 | 71.8 | 70.0 | 70.8 | 73.0 | 71.0 | 76.4 | 78.0 | | | | | | | | | | | |
| 12-year | 59.8 | 61.4 | 62.8 | 71.4 | 71.8 | 69.8 | 70.8 | 72.0 | 71.0 | 76.4 | | | | | | | | | | | | |
| 13-year | 59.8 | 61.0 | 62.8 | 70.9 | 71.5 | 69.5 | 70.8 | 72.0 | 71.0 | | | | | | | | | | | | | |
| 14-year | 59.8 | 61.0 | 61.8 | 70.3 | 71.5 | 69.5 | 70.8 | 71.0 | | | | | | | | | | | | | | |
| 15-year | 59.8 | 61.0 | 61.8 | 70.3 | 71.1 | 68.5 | 70.8 | | | | | | | | | | | | | | | |
| 16-year | 59.8 | 60.3 | 61.8 | 70.2 | 71.1 | 67.9 | | | | | | | | | | | | | | | | |
| 17-year | 59.8 | 60.0 | 61.8 | 69.4 | 71.1 | | | | | | | | | | | | | | | | | |
| 18-year | 59.8 | 60.0 | 61.8 | 68.9 | | | | | | | | | | | | | | | | | | |
| 19-year | 59.8 | 60.0 | 61.8 | | | | | | | | | | | | | | | | | | | |
| 20-year | 59.8 | 60.0 | | | | | | | | | | | | | | | | | | | | |
| 21-year | 59.8 | | | | | | | | | | | | | | | | | | | | | |

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table XVI-7
MELANOMAS OF SKIN (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

| | 1973 | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 |
|---------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Incidence | | | | | | | | | | | | | | | | | | | | | | |
| SEER | 5.4 | 5.7 | 6.3 | 6.5 | 7.0 | 7.3 | 7.6 | 8.2 | 8.8 | 8.6 | 8.4 | 8.7 | 9.4 | 9.8 | 10.1 | 9.5 | 9.8 | 9.8 | 10.5 | 10.2 | 10.1 | 10.1 |
| Mortality | | | | | | | | | | | | | | | | | | | | | | |
| U.S. | 1.3 | 1.3 | 1.3 | 1.4 | 1.4 | 1.4 | 1.5 | 1.4 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.6 | 1.5 | 1.5 | 1.5 | 1.5 |
| SEER | 1.3 | 1.2 | 1.3 | 1.3 | 1.4 | 1.3 | 1.4 | 1.3 | 1.4 | 1.3 | 1.3 | 1.4 | 1.5 | 1.5 | 1.4 | 1.5 | 1.5 | 1.6 | 1.5 | 1.5 | 1.4 | 1.2 |
| Relative Survival Rates (SEER) | | | | | | | | | | | | | | | | | | | | | | |
| 1-year | 91.2 | 94.6 | 97.2 | 95.4 | 96.6 | 95.2 | 96.7 | 97.5 | 95.7 | 97.8 | 98.0 | 98.4 | 97.0 | 97.9 | 97.2 | 98.2 | 97.5 | 97.5 | 98.3 | 96.5 | 97.5 | |
| 2-year | 88.2 | 91.5 | 94.6 | 92.8 | 91.6 | 92.2 | 94.0 | 95.3 | 92.2 | 94.9 | 95.6 | 96.1 | 95.1 | 95.3 | 95.2 | 94.7 | 95.2 | 95.5 | 95.0 | 94.6 | | |
| 3-year | 84.0 | 87.0 | 90.8 | 89.3 | 88.4 | 89.4 | 90.9 | 92.3 | 88.8 | 91.7 | 91.9 | 94.3 | 92.7 | 93.8 | 92.9 | 93.2 | 93.2 | 93.8 | 92.7 | | | |
| 4-year | 79.5 | 83.8 | 87.6 | 87.4 | 86.3 | 87.8 | 88.8 | 90.2 | 86.5 | 89.5 | 89.6 | 92.3 | 91.2 | 92.7 | 91.4 | 91.9 | 92.0 | 92.1 | | | | |
| 5-year | 77.4 | 81.2 | 86.6 | 86.4 | 85.2 | 85.4 | 87.3 | 90.2 | 84.7 | 89.5 | 87.2 | 90.5 | 89.5 | 91.7 | 90.7 | 91.2 | 90.5 | | | | | |
| 6-year | 76.1 | 79.5 | 85.0 | 85.3 | 84.5 | 83.9 | 85.1 | 89.3 | 84.1 | 89.0 | 85.2 | 89.2 | 88.3 | 91.2 | 89.5 | 89.4 | | | | | | |
| 7-year | 75.7 | 79.0 | 83.8 | 83.6 | 83.5 | 83.1 | 85.1 | 88.5 | 83.1 | 87.6 | 84.2 | 88.5 | 87.0 | 90.9 | 89.1 | | | | | | | |
| 8-year | 75.4 | 78.7 | 82.7 | 82.1 | 82.3 | 82.4 | 85.1 | 87.7 | 83.1 | 86.7 | 83.2 | 87.3 | 86.9 | 89.8 | | | | | | | | |
| 9-year | 75.3 | 76.9 | 80.8 | 81.2 | 81.2 | 82.4 | 85.1 | 87.2 | 82.9 | 86.2 | 83.1 | 86.4 | 86.2 | | | | | | | | | |
| 10-year | 74.9 | 76.1 | 79.4 | 81.0 | 80.7 | 82.4 | 85.0 | 87.1 | 82.6 | 85.9 | 82.0 | 85.5 | | | | | | | | | | |
| 11-year | 73.6 | 76.1 | 79.2 | 78.5 | 80.7 | 82.4 | 85.0 | 86.6 | 82.5 | 84.6 | 82.0 | | | | | | | | | | | |
| 12-year | 72.3 | 76.1 | 78.5 | 78.2 | 80.2 | 82.3 | 84.7 | 85.3 | 82.4 | 84.6 | | | | | | | | | | | | |
| 13-year | 71.5 | 76.1 | 78.5 | 78.2 | 80.0 | 82.3 | 84.4 | 84.1 | 82.4 | | | | | | | | | | | | | |
| 14-year | 71.5 | 76.1 | 76.3 | 78.1 | 79.8 | 81.6 | 84.4 | 84.1 | | | | | | | | | | | | | | |
| 15-year | 71.1 | 75.4 | 75.5 | 78.1 | 78.9 | 80.4 | 84.4 | | | | | | | | | | | | | | | |
| 16-year | 71.1 | 74.5 | 75.5 | 77.8 | 78.9 | 79.2 | | | | | | | | | | | | | | | | |
| 17-year | 71.1 | 73.1 | 75.5 | 77.8 | 77.5 | | | | | | | | | | | | | | | | | |
| 18-year | 71.1 | 73.1 | 75.5 | 76.5 | | | | | | | | | | | | | | | | | | |
| 19-year | 71.1 | 73.1 | 75.5 | | | | | | | | | | | | | | | | | | | |
| 20-year | 71.1 | 73.1 | | | | | | | | | | | | | | | | | | | | |
| 21-year | 71.1 | | | | | | | | | | | | | | | | | | | | | |

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table XVI-8
MELANOMAS OF SKIN (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed With Cancer and
Lifetime Risk (Percent) of Dying From Cancer

SEER Areas, 1992-94

All Races

| Current Age | Males | | | | Females | | | |
|----------------|---------|---------|---------|------------|---------|---------|---------|------------|
| | +10 yrs | +20 yrs | +30 yrs | Eventually | +10 yrs | +20 yrs | +30 yrs | Eventually |
| 0 | 0.00 | 0.01 | 0.04 | 1.46 | 0.00 | 0.01 | 0.08 | 1.07 |
| 10 | 0.01 | 0.04 | 0.13 | 1.48 | 0.01 | 0.08 | 0.20 | 1.08 |
| 20 | 0.03 | 0.13 | 0.31 | 1.48 | 0.07 | 0.19 | 0.36 | 1.07 |
| 30 | 0.10 | 0.29 | 0.56 | 1.48 | 0.12 | 0.29 | 0.46 | 1.01 |
| 40 | 0.19 | 0.48 | 0.85 | 1.42 | 0.17 | 0.34 | 0.53 | 0.89 |
| 50 | 0.30 | 0.69 | 1.05 | 1.28 | 0.18 | 0.37 | 0.57 | 0.74 |
| 60 | 0.42 | 0.82 | 1.02 | 1.07 | 0.21 | 0.41 | 0.55 | 0.59 |

Lifetime Risk of Being Diagnosed = 1.46% Lifetime Risk of Being Diagnosed = 1.07%
Lifetime Risk of Dying = 0.30% Lifetime Risk of Dying = 0.19%

Whites

| Current Age | Males | | | | Females | | | |
|----------------|---------|---------|---------|------------|---------|---------|---------|------------|
| | +10 yrs | +20 yrs | +30 yrs | Eventually | +10 yrs | +20 yrs | +30 yrs | Eventually |
| 0 | 0.00 | 0.01 | 0.04 | 1.67 | 0.00 | 0.01 | 0.09 | 1.22 |
| 10 | 0.01 | 0.05 | 0.16 | 1.69 | 0.01 | 0.09 | 0.23 | 1.22 |
| 20 | 0.04 | 0.15 | 0.36 | 1.69 | 0.08 | 0.22 | 0.41 | 1.21 |
| 30 | 0.11 | 0.32 | 0.64 | 1.68 | 0.15 | 0.34 | 0.54 | 1.15 |
| 40 | 0.22 | 0.54 | 0.96 | 1.60 | 0.20 | 0.40 | 0.62 | 1.01 |
| 50 | 0.34 | 0.77 | 1.18 | 1.44 | 0.20 | 0.43 | 0.64 | 0.83 |
| 60 | 0.47 | 0.92 | 1.15 | 1.20 | 0.23 | 0.46 | 0.61 | 0.65 |

Lifetime Risk of Being Diagnosed = 1.67% Lifetime Risk of Being Diagnosed = 1.22%
Lifetime Risk of Dying = 0.35% Lifetime Risk of Dying = 0.22%

Blacks

| Current Age | Males | | | | Females | | | |
|----------------|---------|---------|---------|------------|---------|---------|---------|------------|
| | +10 yrs | +20 yrs | +30 yrs | Eventually | +10 yrs | +20 yrs | +30 yrs | Eventually |
| 0 | 0.00 | 0.00 | 0.00 | 0.04 | 0.00 | 0.00 | 0.00 | 0.07 |
| 10 | 0.00 | 0.00 | 0.00 | 0.04 | 0.00 | 0.00 | 0.00 | 0.07 |
| 20 | 0.00 | 0.00 | 0.01 | 0.04 | 0.00 | 0.00 | 0.01 | 0.07 |
| 30 | 0.00 | 0.01 | 0.02 | 0.05 | 0.00 | 0.01 | 0.02 | 0.07 |
| 40 | 0.01 | 0.02 | 0.04 | 0.04 | 0.01 | 0.02 | 0.02 | 0.07 |
| 50 | 0.01 | 0.04 | 0.04 | 0.04 | 0.01 | 0.02 | 0.04 | 0.07 |
| 60 | 0.03 | 0.03 | 0.03 | 0.03 | 0.01 | 0.03 | 0.05 | 0.06 |

Lifetime Risk of Being Diagnosed = 0.04% Lifetime Risk of Being Diagnosed = 0.07%
Lifetime Risk of Dying = 0.03% Lifetime Risk of Dying = 0.04%

Table XVI-9
MELANOMAS OF SKIN (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER MORTALITY* RATES BY STATE, 1990-94

All Races, Males and Females

| <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> | <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> |
|----------------------|-------------|-----------|-------------|-----------|----------------|-------------|-----------|-------------|-----------|
| TOTAL U.S. | 2.2 | 0.01 | | | Montana | 2.2 | 0.22 | (31) | 0.0 |
| High Five States | | | | | Nebraska | 2.0 | 0.15 | (38) | -9.1 |
| Vermont | 3.0* | 0.31 | (01) | 36.4 | Nevada | 2.8* | 0.20 | (03) | 27.3 |
| Delaware | 3.0* | 0.28 | (02) | 36.4 | New Hampshire | 2.5 | 0.21 | (12) | 13.6 |
| Nevada | 2.8* | 0.20 | (03) | 27.3 | New Jersey | 2.4 | 0.07 | (16) | 9.1 |
| Utah | 2.8* | 0.19 | (04) | 27.3 | New Mexico | 2.4 | 0.17 | (19) | 9.1 |
| Massachusetts | 2.7⊕* | 0.09 | (05) | 22.7 | New York | 2.0⊕ | 0.04 | (41) | -9.1 |
| Low Five States | | | | | North Carolina | 2.5 | 0.08 | (11) | 13.6 |
| Michigan | 1.8⊕* | 0.06 | (47) | -18.2 | North Dakota | 1.7* | 0.21 | (48) | -22.7 |
| North Dakota | 1.7* | 0.21 | (48) | -22.7 | Ohio | 2.1 | 0.06 | (36) | -4.5 |
| Mississippi | 1.6⊕* | 0.10 | (49) | -27.3 | Oklahoma | 2.5 | 0.12 | (06) | 13.6 |
| Hawaii | 1.2⊕* | 0.14 | (50) | -45.5 | Oregon | 2.5 | 0.12 | (09) | 13.6 |
| District of Columbia | 0.9⊕* | 0.16 | (51) | -59.1 | Pennsylvania | 2.3 | 0.06 | (29) | 4.5 |
| Alabama | 2.3 | 0.10 | (27) | 4.5 | Rhode Island | 2.0 | 0.18 | (40) | -9.1 |
| Alaska | 1.8* | 0.34 | (45) | -18.2 | South Carolina | 2.1 | 0.11 | (35) | -4.5 |
| Arizona | 2.4 | 0.11 | (17) | 9.1 | South Dakota | 2.3 | 0.24 | (28) | 4.5 |
| Arkansas | 2.2 | 0.12 | (34) | 0.0 | Tennessee | 2.5 | 0.09 | (13) | 13.6 |
| California | 2.4 | 0.04 | (14) | 9.1 | Texas | 2.2 | 0.05 | (32) | 0.0 |
| Colorado | 2.5 | 0.12 | (08) | 13.6 | Utah | 2.8* | 0.19 | (04) | 27.3 |
| Connecticut | 2.3 | 0.11 | (24) | 4.5 | Vermont | 3.0* | 0.31 | (01) | 36.4 |
| Delaware | 3.0* | 0.28 | (02) | 36.4 | Virginia | 2.3 | 0.08 | (22) | 4.5 |
| District of Columbia | 0.9⊕* | 0.16 | (51) | -59.1 | Washington | 2.3 | 0.09 | (25) | 4.5 |
| Florida | 2.4 | 0.05 | (18) | 9.1 | West Virginia | 2.4 | 0.15 | (15) | 9.1 |
| Georgia | 2.3 | 0.08 | (23) | 4.5 | Wisconsin | 1.9 | 0.08 | (42) | -13.6 |
| Hawaii | 1.2⊕* | 0.14 | (50) | -45.5 | Wyoming | 2.5 | 0.33 | (07) | 13.6 |
| Idaho | 2.4 | 0.20 | (20) | 9.1 | | | | | |
| Illinois | 1.9⊕ | 0.05 | (43) | -13.6 | | | | | |
| Indiana | 2.2 | 0.08 | (33) | 0.0 | | | | | |
| Iowa | 2.0 | 0.11 | (39) | -9.1 | | | | | |
| Kansas | 2.5 | 0.13 | (10) | 13.6 | | | | | |
| Kentucky | 2.3 | 0.10 | (26) | 4.5 | | | | | |
| Louisiana | 1.8* | 0.09 | (46) | -18.2 | | | | | |
| Maine | 2.2 | 0.18 | (30) | 0.0 | | | | | |
| Maryland | 2.1 | 0.09 | (37) | -4.5 | | | | | |
| Massachusetts | 2.7⊕* | 0.09 | (05) | 22.7 | | | | | |
| Michigan | 1.8⊕* | 0.06 | (47) | -18.2 | | | | | |
| Minnesota | 1.9 | 0.09 | (44) | -13.6 | | | | | |
| Mississippi | 1.6⊕* | 0.10 | (49) | -27.3 | | | | | |
| Missouri | 2.3 | 0.09 | (21) | 4.5 | | | | | |

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

* Absolute difference between state rate and total U.S. rate is 15% or more.

⊕ Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

Table XVI-10
MELANOMAS OF SKIN (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER MORTALITY* RATES BY STATE, 1990-94

All Races, Males

| <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> | <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> |
|----------------------|-------------|-----------|-------------|-----------|----------------|-------------|-----------|-------------|-----------|
| TOTAL U.S. | 3.1 | 0.02 | | | | | | | |
| High Five States | | | | | | | | | |
| Vermont | 4.0* | 0.54 | (01) | 29.0 | | | | | |
| Delaware | 3.9* | 0.47 | (02) | 25.8 | | | | | |
| Wyoming | 3.9* | 0.59 | (03) | 25.8 | | | | | |
| Utah | 3.9* | 0.33 | (04) | 25.8 | | | | | |
| Massachusetts | 3.7* | 0.16 | (05) | 19.4 | | | | | |
| Low Five States | | | | | | | | | |
| Mississippi | 2.3* | 0.19 | (47) | -25.8 | | | | | |
| North Dakota | 2.1* | 0.34 | (48) | -32.3 | | | | | |
| Alaska | 1.6* | 0.40 | (49) | -48.4 | | | | | |
| Hawaii | 1.6⊕* | 0.24 | (50) | -48.4 | | | | | |
| District of Columbia | 1.20* | 0.29 | (51) | -61.3 | | | | | |
| State | Rate | SE | Rank | PD | State | Rate | SE | Rank | PD |
| Alabama | 3.2 | 0.18 | (25) | 3.2 | Montana | 2.6* | 0.35 | (43) | -16.1 |
| Alaska | 1.6* | 0.40 | (49) | -48.4 | Nebraska | 2.7 | 0.26 | (40) | -12.9 |
| Arizona | 3.4 | 0.18 | (16) | 9.7 | Nevada | 3.6* | 0.33 | (08) | 16.1 |
| Arkansas | 2.8 | 0.21 | (37) | -9.7 | New Hampshire | 3.5 | 0.36 | (10) | 12.9 |
| California | 3.4 | 0.07 | (14) | 9.7 | New Jersey | 3.5 | 0.13 | (11) | 12.9 |
| Colorado | 3.4 | 0.21 | (12) | 9.7 | New Mexico | 3.3 | 0.30 | (18) | 6.5 |
| Connecticut | 3.0 | 0.19 | (34) | -3.2 | New York | 2.8 | 0.08 | (38) | -9.7 |
| Delaware | 3.9* | 0.47 | (02) | 25.8 | North Carolina | 3.4 | 0.14 | (15) | 9.7 |
| District of Columbia | 1.20* | 0.29 | (51) | -61.3 | North Dakota | 2.1* | 0.34 | (48) | -32.3 |
| Florida | 3.6⊕* | 0.09 | (07) | 16.1 | Ohio | 2.9 | 0.10 | (35) | -6.5 |
| Georgia | 3.3 | 0.15 | (19) | 6.5 | Oklahoma | 3.6* | 0.21 | (09) | 16.1 |
| Hawaii | 1.60* | 0.24 | (50) | -48.4 | Oregon | 3.3 | 0.20 | (21) | 6.5 |
| Idaho | 3.6* | 0.37 | (06) | 16.1 | Pennsylvania | 3.2 | 0.10 | (26) | 3.2 |
| Illinois | 2.7 | 0.10 | (41) | -12.9 | Rhode Island | 2.9 | 0.33 | (36) | -6.5 |
| Indiana | 3.1 | 0.15 | (28) | 0.0 | South Carolina | 3.4 | 0.20 | (17) | 9.7 |
| Iowa | 2.8 | 0.19 | (39) | -9.7 | South Dakota | 3.1 | 0.41 | (31) | 0.0 |
| Kansas | 3.3 | 0.23 | (20) | 6.5 | Tennessee | 3.4 | 0.17 | (13) | 9.7 |
| Kentucky | 3.1 | 0.18 | (27) | 0.0 | Texas | 3.1 | 0.09 | (30) | 0.0 |
| Louisiana | 2.5* | 0.16 | (44) | -19.4 | Utah | 3.9* | 0.33 | (04) | 25.8 |
| Maine | 3.2 | 0.31 | (23) | 3.2 | Vermont | 4.0* | 0.54 | (01) | 29.0 |
| Maryland | 3.1 | 0.17 | (29) | 0.0 | Virginia | 3.0 | 0.14 | (33) | -3.2 |
| Massachusetts | 3.7* | 0.16 | (05) | 19.4 | Washington | 3.1 | 0.16 | (32) | 0.0 |
| Michigan | 2.50* | 0.10 | (46) | -19.4 | West Virginia | 3.2 | 0.25 | (24) | 3.2 |
| Minnesota | 2.5* | 0.15 | (45) | -19.4 | Wisconsin | 2.6* | 0.14 | (42) | -16.1 |
| Mississippi | 2.3* | 0.19 | (47) | -25.8 | Wyoming | 3.9* | 0.59 | (03) | 25.8 |
| Missouri | 3.2 | 0.16 | (22) | 3.2 | | | | | |

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

* Absolute difference between state rate and total U.S. rate is 15% or more.

⊕ Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

Table XVI-11
MELANOMAS OF SKIN (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER MORTALITY* RATES BY STATE, 1990-94

All Races, Females

| <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> | <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> |
|----------------------|-------------|-----------|-------------|-----------|----------------|-------------|-----------|-------------|-----------|
| TOTAL U.S. | 1.5 | 0.01 | | | Montana | 1.9* | 0.28 | (07) | 26.7 |
| High Five States | | | | | Nebraska | 1.5 | 0.17 | (31) | 0.0 |
| Delaware | 2.2* | 0.33 | (01) | 46.7 | Nevada | 2.1* | 0.24 | (03) | 40.0 |
| Vermont | 2.1* | 0.36 | (02) | 40.0 | New Hampshire | 1.7 | 0.24 | (15) | 13.3 |
| Nevada | 2.1* | 0.24 | (03) | 40.0 | New Jersey | 1.6 | 0.08 | (19) | 6.7 |
| Massachusetts | 2.0* | 0.10 | (04) | 33.3 | New Mexico | 1.6 | 0.19 | (23) | 6.7 |
| Oregon | 1.9* | 0.14 | (05) | 26.7 | New York | 1.4 | 0.05 | (39) | -6.7 |
| Low Five States | | | | | North Carolina | 1.8* | 0.09 | (11) | 20.0 |
| Michigan | 1.2* | 0.07 | (47) | -20.0 | North Dakota | 1.5 | 0.27 | (32) | 0.0 |
| South Carolina | 1.2* | 0.10 | (48) | -20.0 | Ohio | 1.4 | 0.06 | (35) | -6.7 |
| Mississippi | 1.1* | 0.11 | (49) | -26.7 | Oklahoma | 1.7 | 0.13 | (16) | 13.3 |
| Hawaii | 0.8* | 0.16 | (50) | -46.7 | Oregon | 1.9* | 0.14 | (05) | 26.7 |
| District of Columbia | 0.6⊕* | 0.18 | (51) | -60.0 | Pennsylvania | 1.6 | 0.06 | (22) | 6.7 |
| Alabama | 1.5 | 0.11 | (27) | 0.0 | Rhode Island | 1.3 | 0.19 | (44) | -13.3 |
| Alaska | 1.9* | 0.50 | (08) | 26.7 | South Carolina | 1.2* | 0.10 | (48) | -20.0 |
| Arizona | 1.6 | 0.12 | (25) | 6.7 | South Dakota | 1.5 | 0.26 | (28) | 0.0 |
| Arkansas | 1.7 | 0.15 | (17) | 13.3 | Tennessee | 1.7 | 0.10 | (18) | 13.3 |
| California | 1.6 | 0.04 | (20) | 6.7 | Texas | 1.5 | 0.05 | (34) | 0.0 |
| Colorado | 1.8* | 0.14 | (10) | 20.0 | Utah | 1.9* | 0.22 | (06) | 26.7 |
| Connecticut | 1.7 | 0.13 | (13) | 13.3 | Vermont | 2.1* | 0.36 | (02) | 40.0 |
| Delaware | 2.2* | 0.33 | (01) | 46.7 | Virginia | 1.7 | 0.10 | (14) | 13.3 |
| District of Columbia | 0.6⊕* | 0.18 | (51) | -60.0 | Washington | 1.6 | 0.10 | (26) | 6.7 |
| Florida | 1.3 | 0.05 | (41) | -13.3 | West Virginia | 1.7 | 0.16 | (12) | 13.3 |
| Georgia | 1.5 | 0.09 | (30) | 0.0 | Wisconsin | 1.3 | 0.09 | (42) | -13.3 |
| Hawaii | 0.8* | 0.16 | (50) | -46.7 | Wyoming | 1.4 | 0.33 | (40) | -6.7 |
| Idaho | 1.3 | 0.20 | (46) | -13.3 | | | | | |
| Illinois | 1.4 | 0.06 | (38) | -6.7 | | | | | |
| Indiana | 1.5 | 0.09 | (33) | 0.0 | | | | | |
| Iowa | 1.4 | 0.12 | (36) | -6.7 | | | | | |
| Kansas | 1.8* | 0.16 | (09) | 20.0 | | | | | |
| Kentucky | 1.6 | 0.12 | (24) | 6.7 | | | | | |
| Louisiana | 1.3 | 0.10 | (45) | -13.3 | | | | | |
| Maine | 1.5 | 0.20 | (29) | 0.0 | | | | | |
| Maryland | 1.3 | 0.10 | (43) | -13.3 | | | | | |
| Massachusetts | 2.0* | 0.10 | (04) | 33.3 | | | | | |
| Michigan | 1.2* | 0.07 | (47) | -20.0 | | | | | |
| Minnesota | 1.4 | 0.11 | (37) | -6.7 | | | | | |
| Mississippi | 1.1* | 0.11 | (49) | -26.7 | | | | | |
| Missouri | 1.6 | 0.10 | (21) | 6.7 | | | | | |

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

* Absolute difference between state rate and total U.S. rate is 15% or more.

⊕ Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

Table XVI-12
MELANOMAS OF SKIN (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES[§], SEER MORTALITY RATES*, AND 5-YEAR RELATIVE SURVIVAL RATES[†]
By Registry, Race and Sex

| | All Races | | | Whites | | | Blacks | | |
|--|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>SEER INCIDENCE RATES[§], 1990-94</u> | | | | | | | | | |
| REGISTRY | | | | | | | | | |
| Standard 9 registries | 12.2 | 14.9 | 10.1 | 13.8 | 16.8 | 11.6 | 0.7 | 0.9 | 0.6 |
| San Francisco-Oakland | 10.9 | 14.1 | 8.4 | 13.7 | 17.4 | 10.7 | 0.7 | 0.9 | 0.5 |
| Connecticut | 14.9 | 18.2 | 12.5 | 15.7 | 19.0 | 13.4 | 1.6 | 2.6 | 1.0 |
| Metropolitan Detroit | 9.7 | 12.6 | 7.4 | 11.2 | 14.5 | 8.7 | 0.5 | 0.6 | 0.4 |
| Hawaii | 8.8 | 10.9 | 6.9 | 24.3 | 30.1 | 18.7 | - | - | - |
| Iowa | 10.6 | 12.3 | 9.4 | 10.6 | 12.3 | 9.4 | - | - | - |
| New Mexico | 11.0 | 13.7 | 8.9 | 12.0 | 14.8 | 9.7 | - | - | - |
| Seattle-Puget Sound | 13.9 | 15.7 | 12.6 | 15.2 | 17.1 | 13.8 | 0.7 | 0.6 | 0.7 |
| Utah | 14.9 | 17.4 | 12.9 | 15.4 | 17.9 | 13.3 | - | - | - |
| Metropolitan Atlanta | 14.9 | 19.2 | 12.1 | 16.9 | 22.0 | 13.4 | 0.7 | 0.9 | 0.5 |
| San Jose-Monterey | 10.1 | 12.1 | 8.6 | 10.7 | 13.1 | 9.0 | 0.0 | 0.0 | 0.0 |
| Los Angeles | 9.5 | 12.2 | 7.5 | 11.2 | 14.3 | 9.0 | 0.9 | 1.0 | 0.9 |
| <u>SEER MORTALITY RATES*, 1990-94</u> | | | | | | | | | |
| REGISTRY | | | | | | | | | |
| Standard 9 registries | 2.1 | 2.8 | 1.4 | 2.4 | 3.3 | 1.7 | 0.4 | 0.5 | 0.4 |
| San Francisco-Oakland | 2.1 | 2.8 | 1.5 | 2.7 | 3.7 | 1.9 | 0.5 | 0.4 | 0.6 |
| Connecticut | 2.3 | 3.0 | 1.7 | 2.4 | 3.2 | 1.9 | 0.9 | 0.9 | 0.2 |
| Metropolitan Detroit | 1.5 | 2.1 | 1.1 | 1.9 | 2.6 | 1.3 | 0.3 | 0.4 | 0.3 |
| Hawaii | 1.2 | 1.6 | 0.8 | 3.3 | 4.5 | 2.0 | - | - | - |
| Iowa | 2.0 | 2.8 | 1.4 | 2.1 | 2.8 | 1.4 | - | - | - |
| New Mexico | 2.4 | 3.3 | 1.6 | 2.5 | 3.5 | 1.7 | - | - | - |
| Seattle-Puget Sound | 2.3 | 3.1 | 1.6 | 2.5 | 3.3 | 1.7 | 0.3 | 0.7 | 0.0 |
| Utah | 2.8 | 3.9 | 1.9 | 2.9 | 3.9 | 2.0 | - | - | - |
| Metropolitan Atlanta | 2.1 | 3.4 | 1.2 | 2.6 | 4.2 | 1.4 | 0.7 | 0.9 | 0.6 |
| San Jose-Monterey | 2.4 | 3.4 | 1.6 | 2.8 | 4.0 | 1.8 | 0.0 | 0.0 | 0.0 |
| Los Angeles | 2.2 | 3.0 | 1.5 | 2.7 | 3.7 | 1.9 | 0.5 | 0.7 | 0.3 |
| <u>SURVIVAL RATES†, 1986-93</u> | | | | | | | | | |
| REGISTRY | | | | | | | | | |
| Standard 9 registries | 87.9 | 85.1 | 90.9 | 88.1 | 85.4 | 91.1 | 66.8↑ | 50.0↑ | 78.7↑ |
| San Francisco-Oakland | 88.4 | 85.5 | 91.8 | 88.7 | 85.7 | 92.1 | - | - | - |
| Connecticut | 86.0 | 84.0 | 88.1 | 86.1 | 84.3 | 88.1 | - | - | - |
| Metropolitan Detroit | 88.0 | 86.0 | 90.3 | 88.1 | 86.3 | 90.3 | 70.4↑ | - | - |
| Hawaii | 91.1 | 89.1 | 92.7 | 93.0 | 90.8 | 94.9 | - | - | - |
| Iowa | 86.5 | 83.2 | 89.7 | 86.5 | 83.2 | 89.7 | - | - | - |
| New Mexico | 87.1 | 85.8 | 88.6 | 87.3 | 86.0 | 88.7 | - | - | - |
| Seattle-Puget Sound | 89.6 | 86.3 | 93.1 | 89.7 | 86.7 | 93.1 | - | - | - |
| Utah | 84.6 | 81.4 | 87.9 | 84.6 | 81.4 | 88.0 | - | - | - |
| Metropolitan Atlanta | 90.3 | 85.4 | 95.8 | 90.8 | 86.0 | 96.2 | - | - | - |

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in this table and not other tables in this section.

SEER Program. Incidence rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

NCHS public use tape. Mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents and are based on follow-up of patients through 1994.

The standard error of the survival rate is between 5 and 10 percentage points.

The standard error of the survival rate is greater than 10 percentage points.

Statistic could not be calculated or populations are inadequate.

§

*

†


↑

↔


-

SEER Incidence and U.S. Mortality Melanoma of the Skin, White By Sex

Male



Female



Melanoma of the Skin SEER Incidence by Age and Sex Whites 1973-75 vs. 1992-94

