

Table VII-1  
CORPUS AND UTERUS, NOS CANCER (Invasive)

## TRENDS IN SEER INCIDENCE AND U.S. MORTALITY, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>INCIDENCE TRENDS (expressed in percents)§:</u>									
<u>1973-96</u>									
All Ages									
Percent Change (PC)	-27.4	-33.3	-26.5	-27.9	-37.2	-26.5	12.2	-9.4	15.3
Est. Annual PC (EAPC)	-1.7★	-1.6★	-1.7★	-1.7★	-1.8★	-1.7★	0.0	-1.1	0.2
Under 65									
Percent Change (PC)	-39.9	-	-	-41.2	-	-	24.3	-	-
Est. Annual PC (EAPC)	-2.8★	-	-	-2.9★	-	-	0.2	-	-
65 and over									
Percent Change (PC)	-1.0	-	-	1.0	-	-	2.4	-	-
Est. Annual PC (EAPC)	0.0	-	-	0.1	-	-	0.0	-	-
<u>1975-79 - ALL AGES</u>									
Percent Change (PC)	-18.5	-10.5	-19.5	-18.5	-10.9	-19.4	-3.5	12.2	-5.7
Est. Annual PC (EAPC)	-6.4★	-4.2	-6.7★	-6.4★	-4.6	-6.7★	-2.3	3.2	-3.1
<u>1992-96 - ALL AGES</u>									
Percent Change (PC)	1.0	-9.2	2.6	-0.6	-11.5	1.0	7.4	-12.8	10.2
Est. Annual PC (EAPC)	0.0♦	-2.8	0.4♦	-0.5♦	-3.9	-0.1♦	2.2	-1.0	2.6
<u>MORTALITY TRENDS (expressed in percents)*:</u>									
<u>1973-96</u>									
All Ages									
Percent Change (PC)	-26.7	-47.3	-24.9	-26.8	-42.3	-25.5	-25.5	-69.1	-21.3
Est. Annual PC (EAPC)	-1.5★	-2.9★	-1.4★	-1.5★	-2.6★	-1.5★	-1.1★	-4.7★	-0.9★
Under 65									
Percent Change (PC)	-39.0	-	-	-37.6	-	-	-48.0	-	-
Est. Annual PC (EAPC)	-2.5★	-	-	-2.5★	-	-	-2.6★	-	-
65 and over									
Percent Change (PC)	-18.4	-	-	-19.8	-	-	-7.5	-	-
Est. Annual PC (EAPC)	-0.9★	-	-	-1.0★	-	-	-0.2	-	-
<u>1975-79 - ALL AGES</u>									
Percent Change (PC)	-4.7	-11.7	-4.2	-5.6	-11.6	-5.2	0.6	-12.3	1.7
Est. Annual PC (EAPC)	-1.5★	-5.1	-1.3★	-1.8★	-5.1	-1.6★	0.2	-4.9★	0.6
<u>1992-96 - ALL AGES</u>									
Percent Change (PC)	-1.0	3.9	-1.3	-0.4	9.3	-0.9	-4.3	-18.4	-3.7
Est. Annual PC (EAPC)	-0.3★♦	0.8	-0.4◇	-0.3♦	2.1	-0.4♦	-0.6	-5.6	-0.4

The Est. Annual PC is the Estimated Annual Percent Change (EAPC) over the time interval.  
SEER Program.  
NCHS public use tape.  
The EAPC is significantly different from zero (p<.05).  
The EAPC for 1992-96 is significantly different from the EAPC for 1975-79 (p<.05).  
The EAPC for 1992-96 is significantly different from the EAPC for 1975-79 (p<.10).  
Statistic could not be calculated.

Table VII-2  
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCES AND U.S. MORTALITY\* RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidences			US Mortality*		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-SPECIFIC RATES, 1992-96</u>						
AGE AT DIAGNOSIS/DEATH:						
0-4	0.0	0.0	0.0	0.0	0.0	0.0
5-9	0.0	0.0	0.0	0.0	0.0	0.0
10-14	0.0	0.0	0.0	0.0	0.0	0.0
15-19	0.1	0.1	0.2	0.0	0.0	0.0
20-24	0.2	0.2	0.3	0.0	0.0	0.0
25-29	0.9	0.7	1.3	0.1	0.0	0.1
30-34	2.3	1.9	1.2	0.1	0.1	0.2
35-39	5.9	5.5	3.9	0.4	0.3	0.7
40-44	12.0	11.8	8.8	0.8	0.8	0.9
45-49	24.7	25.0	12.1	1.6	1.6	1.8
50-54	44.1	46.1	24.6	3.4	3.3	4.8
55-59	63.7	67.2	39.5	5.8	5.3	9.5
60-64	83.9	89.3	63.2	10.9	9.8	21.1
65-69	92.4	97.6	78.6	16.0	14.6	31.0
70-74	107.0	113.1	83.9	21.8	20.3	41.4
75-79	109.3	115.9	70.8	25.9	24.6	43.9
80-84	91.9	96.3	76.1	31.0	29.8	48.7
85+	67.6	68.9	73.9	36.2	35.5	48.4
<u>AGE-ADJUSTED RATES, 1992-96</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	21.5	22.4	15.3	3.3	3.1	5.8
Under 65	13.1	13.6	8.4	1.2	1.1	2.0
65 and over	97.7	103.0	77.9	22.7	21.4	39.6
All ages (world std.) <sup>⊕</sup>	17.7	18.5	12.5	2.5	2.4	4.4

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, except where noted.  
\* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, except where noted.  
⊕ Rates are per 100,000 and are age-adjusted to the world standard population.

Table VII-3  
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCES AND U.S. MORTALITY\* RATES, AGE-ADJUSTED RATES, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1973	28.4	5.3	99.5	29.5	5.4	103.8	15.0	2.2	54.3
1974	30.8	4.9	110.6	32.4	5.0	116.6	13.0	2.3	46.1
1975	32.1	4.9	116.1	33.7	5.1	122.1	17.0	2.7	61.3
1976	31.0	4.7	112.0	32.6	4.7	118.8	14.4	2.4	51.2
1977	28.5	4.3	103.2	29.8	4.4	108.3	17.1	2.6	61.9
1978	26.5	4.5	94.3	28.0	4.7	99.8	15.9	2.8	56.1
1979	24.9	4.0	89.4	26.1	4.0	94.2	14.4	2.9	49.9
1980	24.2	3.7	87.5	25.3	3.7	92.0	14.1	1.8	52.0
1981	24.1	4.1	85.6	25.1	4.0	90.1	14.3	2.0	52.4
1982	23.6	3.8	85.0	24.8	3.8	89.6	14.7	1.9	54.0
1983	23.4	3.6	84.4	24.6	3.7	89.1	16.0	2.5	57.6
1984	22.6	3.4	81.9	23.9	3.6	86.6	14.7	1.2	56.3
1985	22.1	3.4	79.5	23.2	3.6	83.6	15.4	1.6	57.7
1986	21.3	3.6	75.8	22.4	3.7	80.1	14.2	1.9	52.0
1987	21.5	3.7	76.6	22.8	3.9	80.9	13.4	1.3	50.7
1988	20.5	3.4	73.3	21.4	3.3	77.4	14.1	2.1	50.9
1989	21.3	3.4	76.4	22.2	3.6	79.7	16.3	1.4	62.5
1990	21.8	3.3	78.6	23.1	3.5	83.6	14.4	1.6	54.0
1991	21.3	3.2	77.0	22.5	3.3	81.8	14.7	2.3	53.2
1992	21.5	3.6	76.8	22.7	3.6	81.6	14.6	2.0	53.5
1993	21.0	3.9	73.7	22.1	3.8	78.8	14.7	2.8	51.5
1994	21.8	3.8	77.5	22.7	3.7	81.3	15.7	2.1	57.7
1995	21.9	3.5	78.5	22.8	3.4	82.5	15.8	1.8	59.0
1996	21.1	3.3	75.9	21.8	3.1	79.5	15.7	2.3	56.7
1992-96	21.5	3.6	76.5	22.4	3.5	80.8	15.3	2.2	55.8
<u>U.S. MORTALITY RATES</u>									
<u>YEAR OF DEATH:</u>									
1973	4.6	0.5	17.4	4.3	0.4	16.4	8.0	1.1	29.5
1974	4.5	0.5	16.8	4.2	0.4	15.9	7.0	0.7	26.6
1975	4.3	0.4	16.4	4.1	0.4	15.7	6.6	0.7	24.8
1976	4.3	0.4	16.6	4.1	0.4	15.8	6.8	0.6	25.6
1977	4.3	0.4	16.2	4.0	0.4	15.4	7.0	0.6	26.5
1978	4.2	0.4	15.8	3.9	0.4	15.0	6.8	0.6	25.8
1979	4.1	0.3	15.7	3.8	0.3	14.9	6.6	0.6	25.4
1980	4.2	0.3	16.0	3.9	0.3	15.1	6.9	0.6	26.3
1981	4.0	0.3	15.3	3.8	0.3	14.6	6.1	0.4	23.8
1982	3.9	0.3	15.2	3.7	0.3	14.3	6.3	0.3	24.7
1983	3.9	0.3	15.1	3.7	0.2	14.4	6.0	0.5	23.1
1984	3.8	0.3	14.7	3.6	0.2	13.9	6.3	0.5	23.9
1985	3.7	0.3	14.5	3.6	0.3	13.7	6.1	0.4	23.6
1986	3.6	0.3	13.9	3.4	0.2	13.1	6.0	0.4	23.3
1987	3.6	0.3	13.7	3.4	0.2	13.0	5.9	0.4	22.9
1988	3.5	0.2	13.6	3.3	0.2	12.7	6.1	0.3	23.8
1989	3.4	0.3	13.2	3.2	0.3	12.3	5.9	0.4	23.0
1990	3.5	0.2	13.5	3.3	0.2	12.6	6.0	0.3	23.4
1991	3.4	0.2	12.9	3.1	0.2	12.0	6.0	0.4	23.3
1992	3.4	0.2	13.1	3.2	0.2	12.3	5.7	0.3	22.4
1993	3.4	0.2	12.9	3.1	0.2	12.0	6.0	0.3	23.4
1994	3.3	0.2	13.0	3.1	0.2	12.1	5.9	0.2	23.2
1995	3.3	0.3	12.8	3.2	0.3	12.1	5.3	0.3	20.9
1996	3.3	0.2	12.9	3.1	0.2	11.9	5.9	0.3	23.2
1992-96	3.3	0.2	12.9	3.1	0.2	12.1	5.8	0.3	22.6

\*§

SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Table VII-4  
CORPUS AND UTERUS, NOS CANCER (Invasive)

## SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<b>5-YR RELATIVE SURVIVAL RATES</b>									
YEAR OF DIAGNOSIS:									
1960-63 <sup>a</sup>	-	-	-	73	-	-	31	-	-
1970-73 <sup>a</sup>	-	-	-	81	-	-	44	-	-
1974-76 <sup>b</sup>	87.9	92.1	87.3	88.8	92.9	88.3	60.9	84.2↑	56.1
1977-79 <sup>b</sup>	84.9	92.0	84.0	86.3	92.9	85.4	57.6	80.8↑	52.3
1980-82 <sup>b</sup>	81.6	91.0	80.4	82.8	92.3	81.7	54.1	85.1↑	49.3
1983-85 <sup>b</sup>	82.8	92.6	81.6	84.5	93.4	83.4	54.0	81.0↑	50.1
1986-88 <sup>b</sup>	82.7	92.2	81.3	84.3	92.4	83.2	56.5	86.4	51.1
1989-95 <sup>b</sup>	83.5S	88.9S	82.7S	85.5S	90.9	84.7S	56.1	74.2	52.6
1989-95 <sup>b</sup>									
ALL STAGES	83.5	88.9	82.7	85.5	90.9	84.7	56.1	74.2	52.6
LOCALIZED	95.4	95.5	95.5	96.4	97.0	96.4	80.0	87.4	78.2
REGIONAL	64.0	77.3	62.0	66.6	79.8	64.8	41.2	54.4↑	39.0
DISTANT	25.4	42.3	23.3	27.5	48.7	25.1	11.8	-	9.5
UNSTAGED	54.1	75.9↑	50.7	54.4	72.0↑	51.8	48.4↑	-	42.8↑
<b>STAGE DISTRIBUTION (%) 1989-95<sup>b</sup></b>									
ALL STAGES									
Number of cases	18,542	2,504	16,038	16,380	2,028	14,352	1,128	167	961
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	73	77	73	75	78	74	51	65	48
REGIONAL	13	12	13	13	12	13	22	17	23
DISTANT	9	7	9	8	6	9	19	13	20
UNSTAGED	4	4	5	4	4	4	8	5	9
<b>5-YR RELATIVE SURVIVAL RATES, 1989-95<sup>b</sup></b>									
AGE AT DIAGNOSIS:									
<45	88.7	-	-	89.7	-	-	83.6	-	-
45-54	89.5	-	-	91.5	-	-	60.2↑	-	-
55-64	86.1	-	-	88.8	-	-	53.5	-	-
65-74	82.9	-	-	85.1	-	-	52.9	-	-
75+	73.5	-	-	74.9	-	-	46.6↑	-	-
Under 65	87.5	-	-	89.7	-	-	61.1	-	-
65 and over	79.7	-	-	81.7	-	-	50.6	-	-

<sup>a</sup> Rates are based on End Results data from a series of hospital registries and one population-based registry.  
<sup>b</sup> Rates are from the SEER Program. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients through 1996.  
**S** The difference in rates between 1974-76 and 1989-95 is statistically significant (p<.05).  
**↑** The standard error of the survival rate is between 5 and 10 percentage points.  
**↑↑** The standard error of the survival rate is greater than 10 percentage points.  
**-** Statistic could not be calculated.

Table VII-5  
 CORPUS AND UTERUS, NOS CANCER (Invasive)  
 INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Incidence																						
SEER	32.1	31.0	28.5	26.5	24.9	24.2	24.1	23.6	23.4	22.6	22.1	21.3	21.5	20.5	21.3	21.8	21.3	21.5	21.0	21.8	21.9	21.1
Mortality																						
U.S.	4.3	4.3	4.3	4.2	4.1	4.2	4.0	3.9	3.9	3.8	3.7	3.6	3.6	3.5	3.4	3.5	3.4	3.4	3.4	3.3	3.3	3.3
SEER	3.9	4.2	4.2	4.1	4.0	4.2	4.1	4.3	4.2	4.0	4.1	3.4	3.3	3.3	3.5	3.4	3.4	3.5	3.3	3.1	3.3	3.3
Relative Survival Rates (SEER)																						
1-year	94.5	94.2	93.9	94.0	92.7	92.3	91.0	91.5	91.9	91.5	93.2	93.0	92.1	92.3	92.4	92.7	92.2	93.0	92.4	91.9	92.9	
2-year	91.9	90.9	89.7	90.1	88.5	86.3	86.4	86.4	87.0	87.5	88.2	87.1	87.3	88.3	88.0	88.2	88.2	89.7	88.5	87.1		
3-year	90.0	88.9	87.5	87.2	85.9	83.3	84.0	83.9	83.9	85.2	85.5	85.4	84.2	85.3	85.9	85.3	85.8	87.3	85.7			
4-year	88.5	88.2	85.9	85.4	84.4	81.8	82.7	83.2	82.6	84.2	83.8	84.0	83.4	83.7	84.9	84.1	84.2	85.8				
5-year	88.3	87.7	85.7	85.0	84.0	80.9	81.9	81.9	81.6	83.6	83.4	82.9	82.3	83.0	84.1	83.3	83.6					
6-year	88.1	87.0	85.2	84.8	83.7	80.6	81.4	81.5	81.0	83.3	82.8	82.7	81.5	82.6	83.6	81.9						
7-year	87.9	87.0	84.7	84.8	83.7	80.6	80.9	81.4	80.2	82.5	82.2	82.6	81.5	82.3	83.6							
8-year	87.8	87.0	84.7	84.8	83.7	79.9	80.4	81.0	79.8	82.5	82.0	82.6	81.2	82.1								
9-year	87.8	87.0	84.7	84.8	83.7	79.9	80.0	81.0	79.4	82.0	81.5	82.3	81.2									
10-year	87.8	87.0	84.3	84.6	83.7	79.9	79.7	81.0	79.4	81.5	81.4	82.2										
11-year	87.8	87.0	83.9	84.3	83.6	79.7	79.7	81.0	79.1	81.0	81.4											
12-year	87.8	87.0	83.9	84.1	83.4	79.3	79.7	81.0	79.0	80.7												
13-year	87.8	87.0	83.9	84.1	83.2	78.9	79.6	81.0	78.6													
14-year	87.8	87.0	83.9	84.1	82.9	78.9	79.6	81.0														
15-year	87.8	87.0	83.9	84.1	82.0	78.9	79.6															
16-year	87.8	87.0	83.9	83.8	82.0	78.5																
17-year	87.8	87.0	83.9	83.8	82.0																	
18-year	87.8	87.0	83.9	83.8																		
19-year	87.8	87.0	83.9																			
20-year	87.8	87.0																				
21-year	87.8																					

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
 Survival rates are relative rates expressed as percents.

Table VII-6  
CORPUS AND UTERUS, NOS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,  
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed With Cancer and  
 Lifetime Risk (Percent) of Dying From Cancer

SEER Areas, 1994-96

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.69
10	0.00	0.01	0.05	2.71
20	0.01	0.04	0.22	2.72
30	0.04	0.22	0.75	2.73
40	0.18	0.71	1.49	2.71
50	0.55	1.34	2.13	2.59
60	0.84	1.67	2.10	2.16

Lifetime Risk of Being Diagnosed = 2.69%  
 Lifetime Risk of Dying = 0.51%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	2.83
10	0.00	0.00	0.04	2.85
20	0.00	0.04	0.21	2.86
30	0.04	0.21	0.77	2.87
40	0.18	0.74	1.56	2.86
50	0.58	1.41	2.25	2.73
60	0.88	1.76	2.20	2.27

Lifetime Risk of Being Diagnosed = 2.83%  
 Lifetime Risk of Dying = 0.52%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.71
10	0.00	0.01	0.03	1.75
20	0.01	0.03	0.12	1.75
30	0.02	0.12	0.42	1.76
40	0.10	0.40	1.00	1.78
50	0.32	0.94	1.43	1.76
60	0.69	1.23	1.53	1.59

Lifetime Risk of Being Diagnosed = 1.71%  
 Lifetime Risk of Dying = 0.56%

Table VII-7  
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. MORTALITY RATES

By Race/Ethnicity and Sex

Incidence<sup>i</sup>

RACE/ETHNICITY	Rate 1990-1996	Trend 1990-1996
	Rate per 100,000 persons	EAPC (%)
	<u>Females</u>	<u>Females</u>
All Races	21.1	-0.2
White	22.5	-0.6
White Hispanic	14.1	-1.7
White Non-Hispanic	23.3	-0.4
Black	14.9	1.8
Asian/Pacific Islander <sup>‡</sup>	13.6	4.3*
American Indian	7.4	-
Hispanic <sup>⊙</sup>	13.3	-1.9

Mortality<sup>❖</sup>

RACE/ETHNICITY	Rate 1990-1996	Trend 1990-1996
	Rate per 100,000 persons	EAPC (%)
	<u>Females</u>	<u>Females</u>
All Races	3.4	-0.6
White	3.2	-0.5
White Hispanic	2.6	0.9
White Non-Hispanic	3.2	-0.4
Black	5.8	-0.9
Asian/Pacific Islander <sup>‡</sup>	1.8	0.2
American Indian	2.0	2.1
Hispanic <sup>⊙</sup>	2.5	0.6

- The EAPC is the Estimated Annual Percent Change over the time interval.  
 - Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.  
 i Incidence data are from the 11 SEER areas (San Francisco, Connecticut, Detroit, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, and Los Angeles). Unless specified, other tables use data from 9 SEER areas (San Francisco, Connecticut, Detroit, Iowa, New Mexico, Seattle, Utah, and Atlanta).  
 ⊙ Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians. For incidence, all 11 SEER areas are included. For mortality, information is included for all states except Connecticut, Oklahoma, Louisiana, and New Hampshire.  
 ❖ Mortality data are analyzed from a public-use file provided by the National Center for Health Statistics (NCHS).  
 ‡ Population estimates for the Asian Pacific Islanders rates are adjusted Census Bureau estimates. The population estimates used in Wingo PA, Ries LAG, Rosenberg HM, Miller DS, Edwards BK. Cancer Incidence and mortality, 1973-1996. Cancer 82:1197-207, 1998 were unadjusted.  
 \* The EAPC is significantly different from zero (p<.05).

Table VII-8  
CORPUS AND UTERUS, NOS CANCER (Invasive)AGE-ADJUSTED SEER INCIDENCE RATES§ AND SEER MORTALITY RATES\*  
By Registry, Race and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES§, 1992-96</u>									
REGISTRY									
11 registries	21.1	3.5	75.3	22.3	3.4	80.5	15.1	2.1	55.1
9 registries	21.5	3.6	76.5	22.4	3.5	80.8	15.3	2.2	55.8
San Francisco-Oakland	20.9	3.3	75.4	23.0	3.1	84.4	14.8	2.1	53.8
Connecticut	24.5	4.0	87.5	25.0	4.2	89.4	17.1	2.8	61.4
Metropolitan Detroit	20.8	3.7	73.5	22.2	4.1	78.1	14.7	2.4	52.4
Hawaii	19.8	6.6	60.4	21.3	2.9	78.1	-	-	-
Iowa	22.9	3.5	82.9	23.0	3.5	83.0	-	-	-
New Mexico	16.0	3.1	55.6	16.4	2.9	58.0	-	-	-
Seattle-Puget Sound	23.3	3.4	84.4	24.0	3.5	87.0	11.4	0.2	45.8
Utah	21.6	3.4	77.6	21.5	3.3	77.8	-	-	-
Metropolitan Atlanta	18.3	2.8	66.0	18.8	3.1	67.2	17.3	1.9	64.7
San Jose-Monterey	21.0	2.8	77.4	22.7	2.7	84.6	9.1	2.6	29.3
Los Angeles	19.8	3.2	71.1	21.8	3.3	78.6	14.9	2.0	54.4
<u>SEER MORTALITY RATES*, 1992-96</u>									
REGISTRY									
11 registries	3.3	0.2	12.8	3.2	0.2	12.6	5.2	0.4	20.2
9 registries	3.3	0.2	12.7	3.2	0.2	12.4	5.2	0.4	19.9
San Francisco-Oakland	3.2	0.3	12.0	3.1	0.2	12.1	4.7	0.4	17.8
Connecticut	3.1	0.1	12.2	2.9	0.1	11.7	5.3	0.3	20.9
Metropolitan Detroit	3.6	0.3	13.7	3.2	0.2	12.4	4.9	0.5	18.3
Hawaii	2.5	0.2	9.7	3.2	0.2	12.6	-	-	-
Iowa	4.0	0.2	15.5	4.0	0.2	15.4	-	-	-
New Mexico	3.1	0.3	11.6	3.0	0.3	11.5	-	-	-
Seattle-Puget Sound	2.8	0.1	11.2	2.8	0.1	11.1	4.7	0.2	18.5
Utah	3.3	0.2	12.9	3.3	0.1	13.1	-	-	-
Metropolitan Atlanta	3.5	0.2	13.4	2.6	0.3	10.0	6.6	0.2	26.5
San Jose-Monterey	3.1	0.2	12.0	3.3	0.1	13.2	0.4	0.6	0.0
Los Angeles	3.5	0.4	13.2	3.5	0.4	13.1	5.6	0.5	21.4

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in tables where indicated.  
 § SEER Program. Incidence rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
 \* NCHS public use tape. Mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
 - Statistic could not be calculated or populations are inadequate.



Table VII-9  
CORPUS AND UTERUS, NOS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER MORTALITY\* RATES BY STATE, 1992-96

All Races, Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.3	0.02		
High Five States				
District of Columbia	5.3*	0.51	(01)	60.6
Vermont	4.1*	0.48	(02)	24.2
Delaware	4.1*	0.44	(03)	24.2
New Jersey	4.0⊕*	0.12	(04)	21.2
New York	4.0⊕*	0.08	(05)	21.2
Low Five States				
Arizona	2.6⊕*	0.14	(47)	-21.2
Idaho	2.6*	0.28	(48)	-21.2
Hawaii	2.5*	0.27	(49)	-24.2
North Dakota	2.4*	0.33	(50)	-27.3
Alaska	1.6*	0.44	(51)	-51.5

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.7*	0.14	(43)	-18.2	Montana	3.2	0.33	(31)	-3.0
Alaska	1.6*	0.44	(51)	-51.5	Nebraska	2.8*	0.22	(41)	-15.2
Arizona	2.6⊕*	0.14	(47)	-21.2	Nevada	2.7*	0.25	(45)	-18.2
Arkansas	3.0	0.19	(37)	-9.1	New Hampshire	3.4	0.32	(16)	3.0
California	3.2	0.06	(28)	-3.0	New Jersey	4.0⊕*	0.12	(04)	21.2
Colorado	2.7*	0.16	(42)	-18.2	New Mexico	3.1	0.26	(36)	-6.1
Connecticut	3.1	0.16	(35)	-6.1	New York	4.0⊕*	0.08	(05)	21.2
Delaware	4.1*	0.44	(03)	24.2	North Carolina	3.3	0.12	(25)	0.0
District of Columbia	5.3*	0.51	(01)	60.6	North Dakota	2.4*	0.33	(50)	-27.3
Florida	2.9⊕	0.07	(39)	-12.1	Ohio	3.9⊕*	0.10	(07)	18.2
Georgia	3.1	0.13	(32)	-6.1	Oklahoma	2.7*	0.16	(44)	-18.2
Hawaii	2.5*	0.27	(49)	-24.2	Oregon	3.3	0.17	(23)	0.0
Idaho	2.6*	0.28	(48)	-21.2	Pennsylvania	3.7	0.09	(08)	12.1
Illinois	3.5	0.10	(12)	6.1	Rhode Island	3.6	0.32	(11)	9.1
Indiana	3.5	0.14	(13)	6.1	South Carolina	3.4	0.17	(18)	3.0
Iowa	4.0*	0.20	(06)	21.2	South Dakota	3.3	0.37	(26)	0.0
Kansas	3.3	0.19	(22)	0.0	Tennessee	3.2	0.14	(29)	-3.0
Kentucky	3.2	0.16	(27)	-3.0	Texas	3.0	0.08	(38)	-9.1
Louisiana	3.2	0.16	(30)	-3.0	Utah	3.3	0.27	(24)	0.0
Maine	3.4	0.29	(20)	3.0	Vermont	4.1*	0.48	(02)	24.2
Maryland	3.4	0.15	(19)	3.0	Virginia	3.5	0.13	(15)	6.1
Massachusetts	3.3	0.13	(21)	0.0	Washington	2.9	0.13	(40)	-12.1
Michigan	3.6	0.11	(10)	9.1	West Virginia	3.1	0.21	(33)	-6.1
Minnesota	3.1	0.15	(34)	-6.1	Wisconsin	3.4	0.14	(17)	3.0
Mississippi	2.7*	0.18	(46)	-18.2	Wyoming	3.6	0.52	(09)	9.1
Missouri	3.5	0.14	(14)	6.1					

\* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
\* Absolute difference between state rate and total U.S. rate is 15% or more.  
⊕ Difference between state rate and total U.S. rate is statistically significant (p<=.0002).

# SEER Incidence and U.S. Mortality Corpus Uteri, Under 50 Years of Age By Race

## White

## Black





Figure VII-1

# SEER Incidence and U.S. Mortality Corpus Uteri, Ages 50 and Over By Race

## White

## Black





Figure VII-2