

Table VII-1
CORPUS AND UTERUS, NOS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1973-1999
WITH UP TO THREE JOINPOINTS BY RACE AND AGE

	All Races		Whites		Blacks	
	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>						
Age - All Ages	1973-75	6.4★	1973-75	6.9★	1973-99	0.0
	1975-79	-6.0★	1975-79	-6.2★		
	1979-88	-1.7★	1979-88	-1.6★		
	1988-99	0.6★	1988-99	0.5★		
Age - Under 50	1973-84	-3.6★	1973-83	-3.8★	1973-99	-0.7
	1984-99	0.4	1983-99	-0.1		
Age - 50 and Over	1973-75	8.1★	1973-75	8.6★	1973-99	0.1
	1975-79	-6.3★	1975-79	-6.3★		
	1979-88	-1.6★	1979-88	-1.6★		
	1988-99	0.5★	1988-99	0.6★		
<u>U.S. Cancer Mortality^b</u>						
Age - All Ages	1973-89	-1.6★	1973-91	-1.6★	1973-75	-7.0
	1989-99	-0.4★	1991-99	-0.2	1975-99	-0.7★
Age - Under 50	1973-85	-5.0★	1973-84	-5.0★	1973-99	-4.4★
	1985-99	-0.4	1984-99	-0.3		
Age - 50 and Over	1973-91	-1.4★	1973-91	-1.5★	1973-99	-0.6★
	1991-99	-0.3	1991-99	-0.3		

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

Trends are from the SEER 9 areas.

Trends are from the NCHS public use data file.

★ The APC is significantly different from zero (p<.05).

^a
^b
★

Table VII-2
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCES AND U.S. DEATH^â RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence [§]			US Death ^â		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-SPECIFIC RATES, 1995-99</u>						
AGE AT DIAGNOSIS/DEATH:						
0-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	1.2	1.0	-	0.1	-	-
30-34	2.8	2.4	-	0.2	0.2	-
35-39	6.0	5.7	-	0.4	0.4	0.5
40-44	12.1	11.8	8.6	0.8	0.7	1.0
45-49	23.9	24.1	10.3	1.7	1.7	1.8
50-54	50.6	53.4	26.3	3.3	3.2	4.3
55-59	69.7	74.5	41.1	6.1	5.7	9.5
60-64	85.6	91.1	67.7	10.8	9.8	20.1
65-69	92.8	97.8	86.7	15.5	14.0	31.1
70-74	101.9	107.1	79.6	20.9	19.5	39.5
75-79	101.9	108.1	69.4	25.0	23.8	42.9
80-84	98.1	103.8	74.8	30.7	29.7	46.8
85+	71.8	72.4	74.3	36.3	35.4	51.7
<u>AGE-ADJUSTED RATES, 1995-99</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	25.4	26.5	18.1	4.1	3.9	6.9
Under 65	15.3	15.9	9.4	1.3	1.2	2.0
65 and over	95.2	100.1	78.0	23.6	22.3	40.5
All ages (world std.) [⊕]	18.2	19.1	12.9	2.5	2.3	4.3

§ SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.
^â NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.
[⊕] Rates are per 100,000 and are age-adjusted to the world standard population.
 - Statistic not shown. Rate based on less than 25 cases for the time interval.

Table VII-3
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCES AND U.S. DEATH RATES, AGE-ADJUSTED RATES, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1973	31.7	7.1	96.1	32.9	7.2	100.1	18.2	-	57.9
1974	34.0	6.4	106.4	35.8	6.7	111.9	15.5	-	48.1
1975	35.4	6.5	111.2	37.2	6.8	117.0	19.9	-	62.6
1976	34.4	6.2	108.1	36.1	6.3	114.3	17.6	-	55.1
1977	31.6	5.7	99.5	33.1	5.9	104.3	20.1	-	63.4
1978	30.1	6.0	92.9	31.7	6.3	98.1	18.6	-	57.5
1979	27.9	5.3	87.0	29.1	5.3	91.5	16.8	-	50.5
1980	27.3	5.0	85.9	28.5	5.0	90.3	16.4	-	52.8
1981	27.3	5.6	84.0	28.3	5.4	88.2	17.4	-	55.5
1982	26.8	5.0	83.8	28.2	5.1	88.5	17.2	-	55.8
1983	26.6	4.8	83.4	27.8	4.9	87.9	18.8	-	59.2
1984	25.7	4.6	80.9	27.2	4.8	85.6	17.3	-	57.9
1985	25.2	4.6	79.2	26.5	4.8	83.3	17.5	-	57.5
1986	24.4	4.8	75.6	25.6	5.0	79.6	16.6	-	53.4
1987	24.7	5.0	76.4	26.1	5.3	80.7	15.5	-	51.7
1988	23.6	4.5	73.4	24.6	4.4	77.3	16.8	3.0	52.9
1989	24.4	4.5	76.7	25.6	4.8	79.9	19.3	-	65.1
1990	25.1	4.6	78.7	26.6	4.8	83.7	17.0	-	55.4
1991	24.6	4.4	77.5	25.9	4.4	82.3	17.5	3.3	54.6
1992	24.8	4.8	77.3	26.2	4.8	82.2	17.1	2.8	54.5
1993	24.3	5.2	74.5	25.7	5.1	79.6	17.2	3.6	52.9
1994	25.2	5.0	77.9	26.3	5.0	82.0	17.9	2.7	57.9
1995	25.4	4.7	79.4	26.4	4.6	83.5	18.7	2.5	61.2
1996	24.9	4.5	78.3	25.8	4.3	82.3	19.4	3.2	61.8
1997	25.9	5.0	80.8	27.2	5.0	85.5	16.5	2.4	53.3
1998	25.6	4.7	80.4	26.9	4.7	85.0	19.2	2.8	62.3
1999	25.1	5.2	77.3	26.3	5.0	82.1	16.9	2.8	53.9
1995-99	25.4	4.8	79.3	26.5	4.7	83.7	18.1	2.7	58.3
<u>U.S. DEATH RATES</u>									
<u>YEAR OF DEATH:</u>									
1973	5.6	0.6	18.8	5.3	0.5	17.8	9.5	1.4	30.7
1974	5.4	0.6	18.1	5.2	0.6	17.3	8.3	0.9	27.8
1975	5.3	0.5	17.7	5.1	0.5	17.1	7.8	0.9	25.8
1976	5.3	0.5	17.9	5.1	0.5	17.2	8.1	0.9	27.0
1977	5.2	0.5	17.5	5.0	0.5	16.7	8.4	0.8	28.0
1978	5.1	0.5	17.1	4.8	0.5	16.2	8.2	0.8	27.3
1979	5.0	0.4	17.0	4.7	0.4	16.2	7.9	0.7	26.7
1980	5.1	0.4	17.4	4.8	0.4	16.5	8.4	0.9	28.1
1981	4.9	0.4	16.6	4.7	0.4	15.9	7.3	0.6	25.1
1982	4.9	0.4	16.5	4.6	0.4	15.7	7.6	0.4	26.4
1983	4.8	0.4	16.4	4.6	0.3	15.8	7.3	0.6	24.8
1984	4.7	0.4	16.0	4.4	0.3	15.2	7.7	0.7	26.1
1985	4.6	0.4	15.7	4.4	0.3	15.0	7.4	0.5	25.3
1986	4.4	0.3	15.2	4.2	0.3	14.4	7.3	0.6	24.9
1987	4.4	0.3	15.1	4.2	0.3	14.3	7.3	0.5	25.1
1988	4.3	0.3	14.9	4.1	0.3	14.0	7.4	0.5	25.6
1989	4.3	0.3	14.5	4.0	0.3	13.6	7.3	0.5	25.0
1990	4.3	0.3	14.9	4.1	0.3	14.0	7.4	0.4	25.5
1991	4.2	0.3	14.3	3.9	0.3	13.3	7.3	0.5	25.1
1992	4.2	0.3	14.5	4.0	0.3	13.6	7.0	0.4	24.2
1993	4.2	0.3	14.3	3.9	0.3	13.4	7.3	0.4	25.3
1994	4.2	0.3	14.4	3.9	0.3	13.5	7.1	0.3	24.8
1995	4.2	0.3	14.2	4.0	0.3	13.6	6.5	0.3	22.6
1996	4.2	0.3	14.3	3.9	0.3	13.3	7.3	0.4	25.3
1997	4.0	0.3	13.8	3.8	0.3	13.0	6.8	0.3	23.9
1998	4.1	0.3	14.2	3.9	0.3	13.4	6.9	0.5	24.0
1999	4.1	0.4	14.1	3.9	0.3	13.3	6.9	0.4	24.1
1995-99	4.1	0.3	14.1	3.9	0.3	13.3	6.9	0.4	24.0

SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population.
NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population.
Statistic not shown. Rate based on less than 25 cases for the time interval.

Table VII-4
CORPUS AND UTERUS, NOS CANCER (Invasive)

SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>5-YR RELATIVE SURVIVAL RATES</u>									
YEAR OF DIAGNOSIS:									
1960-63 ^a	-	-	-	73	-	-	31	-	-
1970-73 ^a	-	-	-	81	-	-	44	-	-
1974-76 ^b	87.9	92.1	87.4	88.8	92.9	88.4	61.4	84.3↑	56.5
1977-79 ^b	85.0	92.1	84.1	86.3	92.9	85.5	57.9	81.0↑	52.6
1980-82 ^b	81.6	91.1	80.4	82.8	92.3	81.7	55.1	85.6↑	50.2
1983-85 ^b	82.8	92.5	81.6	84.5	93.3	83.4	54.2	81.4↑	50.1
1986-88 ^b	82.9	92.3	81.4	84.4	92.5	83.3	57.1	86.5	51.9
1989-91 ^b	83.7	88.4	83.0	85.6	90.8	84.8	57.5	71.7↑	55.0
1992-98 ^b	84.3Û	89.8	83.2Û	86.0Û	91.3	85.2Û	60.5	81.8	55.8
1992-98 ^b									
ALL STAGES	84.3	89.8	83.2	86.0	91.3	85.2	60.5	81.8	55.8
LOCALIZED	96.1	96.1	96.2	96.9	97.3	96.9	83.3	89.0	81.7
REGIONAL	63.5	79.1	60.7	65.7	79.8	63.4	44.8	80.1↑	38.2
DISTANT	26.1	44.8	23.2	27.8	48.1	24.8	14.9	-	11.9
UNSTAGED	52.6	80.4	48.8	51.6	78.4↑	48.4	49.8↑	-	45.9↑
<u>STAGE DISTRIBUTION (%) 1992-98^b</u>									
ALL STAGES									
Number of cases	19,442	2,771	16,671	16,949	2,164	14,785	1,212	195	1,017
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	73	77	72	75	77	74	53	69	50
REGIONAL	14	13	15	14	13	14	22	17	23
DISTANT	8	7	8	8	7	8	17	9	19
UNSTAGED	5	3	5	4	3	5	8	5	9
<u>5-YR RELATIVE SURVIVAL RATES, 1992-98^b</u>									
AGE AT DIAGNOSIS:									
<45	89.5	-	-	90.5	-	-	84.4	-	-
45-54	90.5	-	-	91.9	-	-	76.5	-	-
55-64	87.0	-	-	89.5	-	-	57.9	-	-
65-74	83.2	-	-	85.5	-	-	54.9	-	-
75+	74.3	-	-	76.0	-	-	42.5↑	-	-
Under 65	88.6	-	-	90.4	-	-	68.9	-	-
65 and over	79.9	-	-	81.8	-	-	50.8	-	-

^a Rates are based on End Results data from a series of hospital registries and one population-based registry.
^b Rates are from the SEER 9 areas. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients into 1999.
Û The difference in rates between 1974-76 and 1992-98 is statistically significant (p<.05).
↑ The standard error of the survival rate is between 5 and 10 percentage points.
↑ The standard error of the survival rate is greater than 10 percentage points.
- Statistic could not be calculated.

Table VII-5
CORPUS AND UTERUS, NOS CANCER (Invasive)

INCIDENCE, DEATH, AND SURVIVAL RATES*

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

	1975	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Incidence																					
SEER ^a	35.4	27.3	27.3	26.8	26.6	25.7	25.2	24.4	24.7	23.6	24.4	25.1	24.6	24.8	24.3	25.2	25.4	24.9	25.9	25.6	25.1
Mortality																					
U.S. ^b	5.3	5.1	4.9	4.9	4.8	4.7	4.6	4.4	4.4	4.3	4.3	4.3	4.2	4.2	4.2	4.2	4.2	4.2	4.0	4.1	4.1
SEER ^a	4.8	5.2	5.1	5.3	5.2	4.9	5.0	4.2	4.1	4.1	4.5	4.2	4.2	4.3	4.0	3.9	4.1	4.1	3.7	4.2	4.0
Relative Survival Rates (SEER ^a)																					
1-year	94.5	92.4	91.1	91.6	91.9	91.6	93.2	93.1	92.1	92.3	92.3	92.7	92.0	93.1	92.5	92.0	93.2	92.0	93.2	93.6	
2-year	91.9	86.3	86.4	86.4	86.9	87.5	88.2	87.1	87.4	88.4	88.0	88.2	88.1	89.7	88.6	87.4	88.7	88.5	89.2		
3-year	90.0	83.4	84.0	83.9	83.8	85.2	85.5	85.4	84.4	85.4	85.9	85.4	85.7	87.4	85.9	85.1	86.4	86.3			
4-year	88.6	81.8	82.7	83.2	82.5	84.3	83.8	84.0	83.5	83.9	84.9	84.0	84.2	86.1	84.7	83.6	85.0				
5-year	88.3	80.9	82.0	81.9	81.4	83.7	83.4	82.9	82.5	83.2	84.1	83.3	83.6	86.0	83.9	82.9					
6-year	88.1	80.7	81.4	81.5	80.9	83.4	82.9	82.6	81.7	82.9	83.7	82.1	82.5	85.3	83.0						
7-year	87.9	80.7	81.0	81.5	80.2	82.7	82.2	82.5	81.7	82.6	83.7	82.1	82.4	85.2							
8-year	87.9	80.1	80.4	81.0	79.7	82.7	82.1	82.5	81.5	82.6	82.7	81.9									
9-year	87.9	80.1	80.1	81.0	79.4	82.3	81.7	82.2	81.5	82.2	82.7	81.8									
10-year	87.9	80.1	79.9	81.0	79.4	81.8	81.7	82.2	81.5	81.7	82.7										
11-year	87.9	79.9	79.9	81.0	79.1	81.4	81.7	82.2	81.4	81.7											
12-year	87.9	79.5	79.9	81.0	79.1	81.3	81.7	82.0	81.2												
13-year	87.9	79.2	79.7	81.0	78.9	81.3	81.2	82.0													
14-year	87.9	79.2	79.7	81.0	77.7	81.3	80.8														
15-year	87.9	79.2	79.7	80.6	77.6	81.3															
16-year	87.9	78.9	79.2	80.6	76.9																
17-year	87.9	78.9	78.9	80.2																	
18-year	87.9	78.9	78.8																		
19-year	87.9	78.9																			
20-year	87.9																				
21-year	87.9																				
22-year	87.9																				
23-year	87.9																				
24-year	87.9																				

^a Rates are from the SEER 9 areas.

^b Rates are from the NCHS public use data file.

* Incidence and death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

Survival rates are relative rates expressed as percents.

Table VII-6
CORPUS AND UTERUS, NOS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1997-1999

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.69
10	0.00	0.01	0.05	2.72
20	0.01	0.05	0.23	2.72
30	0.05	0.22	0.78	2.73
40	0.18	0.74	1.52	2.70
50	0.58	1.37	2.10	2.58
60	0.84	1.60	2.03	2.12

Lifetime Risk of Being Diagnosed = 2.69%
 Lifetime Risk of Dying = 0.51%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.88
10	0.00	0.01	0.05	2.90
20	0.01	0.05	0.23	2.91
30	0.05	0.22	0.82	2.91
40	0.18	0.78	1.63	2.89
50	0.62	1.48	2.26	2.76
60	0.91	1.72	2.17	2.25

Lifetime Risk of Being Diagnosed = 2.88%
 Lifetime Risk of Dying = 0.49%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.72
10	0.00	0.01	0.04	1.75
20	0.01	0.03	0.12	1.76
30	0.02	0.11	0.41	1.76
40	0.08	0.39	1.03	1.77
50	0.32	0.99	1.45	1.76
60	0.74	1.25	1.52	1.58

Lifetime Risk of Being Diagnosed = 1.72%
 Lifetime Risk of Dying = 0.71%

Table VII-7
CORPUS AND UTERUS, NOS CANCER (Invasive)
AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidenceⁱ

RACE/ETHNICITY	Rate 1992-1999	Trend 1992-1999
	Rate per 100,000 persons	EAPC (%)
	Females	Females
All Races	24.5	0.3
White	26.0	0.1
White Hispanic	16.4	0.8
White Non-Hispanic	27.1	0.1
Black	17.7	0.8
Asian/Pacific Islander	17.0	2.6★
Amer Ind/Alaska Nat	10.6	-
Hispanic [⊗]	15.6	0.8

Mortality[❖]

RACE/ETHNICITY	Rate 1992-1999	Trend 1992-1999
	Rate per 100,000 persons	EAPC (%)
	Females	Females
All Races	4.2	-0.3
White	3.9	-0.3
White Hispanic	3.1	-0.2
White Non-Hispanic	3.9	-0.2
Black	7.0	-0.4
Asian/Pacific Islander	2.2	1.3
Amer Ind/Alaska Nat	2.3	-5.9★
Hispanic [⊗]	3.0	-0.4

- The EAPC is the Estimated Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ⁱ Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- [⊗] Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics does not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Oklahoma, New York, and New Hampshire.
- [❖] Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- [★] The EAPC is significantly different from zero (p<.05).

Table VII-8
CORPUS AND UTERUS, NOS CANCER (Invasive)AGE-ADJUSTED SEER INCIDENCE RATES§ AND SEER DEATH RATESä
By Registry, Race and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES§, 1995-99</u>									
REGISTRY									
11 registries	24.7	4.7	77.0	26.1	4.7	82.2	18.0	2.6	58.3
9 registries	25.4	4.8	79.3	26.5	4.7	83.7	18.1	2.7	58.3
San Francisco-Oakland	24.9	4.4	78.5	27.6	4.6	88.0	16.7	-	53.6
Connecticut	29.0	5.0	91.6	29.6	5.1	93.5	18.7	-	59.1
Metropolitan Detroit	24.9	4.8	77.6	26.7	5.2	83.1	18.6	3.3	58.5
Hawaii	25.1	10.5	63.3	26.0	4.5	82.2	-	-	-
Iowa	28.1	5.3	87.9	28.1	5.2	88.2	-	-	-
New Mexico	19.0	4.1	58.0	19.3	3.8	59.9	-	-	-
Seattle-Puget Sound	26.8	4.4	85.3	27.7	4.5	88.5	17.6	-	61.7
Utah	23.8	4.8	73.5	24.0	4.7	74.4	-	-	-
Metropolitan Atlanta	21.5	3.5	68.6	22.0	3.9	69.3	20.2	2.2	67.4
San Jose-Monterey	24.1	3.5	78.0	25.7	3.4	84.2	-	-	-
Los Angeles	22.7	4.7	69.8	24.7	5.0	76.1	18.1	2.5	59.1
<u>SEER DEATH RATESä, 1995-99</u>									
REGISTRY									
11 registries	4.0	0.3	13.7	4.0	0.3	13.7	5.8	0.5	19.6
9 registries	4.0	0.3	13.7	4.0	0.3	13.7	5.6	-	19.2
San Francisco-Oakland	3.8	0.4	12.8	3.9	-	13.3	4.9	-	17.2
Connecticut	4.2	-	14.4	4.1	-	13.9	6.7	-	22.4
Metropolitan Detroit	4.1	0.5	13.8	3.8	-	12.7	5.5	-	18.4
Hawaii	3.2	-	10.0	3.6	-	-	-	-	-
Iowa	4.6	-	15.9	4.6	-	15.8	-	-	-
New Mexico	3.8	-	12.9	3.8	-	13.1	-	-	-
Seattle-Puget Sound	4.0	-	13.9	4.0	-	14.0	-	-	-
Utah	3.6	-	12.6	3.6	-	12.8	-	-	-
Metropolitan Atlanta	4.0	-	13.9	3.4	-	11.9	6.4	-	22.4
San Jose-Monterey	3.8	-	13.3	4.0	-	14.0	-	-	-
Los Angeles	4.1	0.4	13.8	4.2	0.4	14.0	6.2	-	20.6

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in tables where indicated.
 § SEER 9 areas. Incidence rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.
 ä NCHS public use data file. Death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.
 - Statistic could not be calculated or populations are inadequate.

Table VII-9
CORPUS AND UTERUS, NOS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^ä RATES BY STATE, 1995-99

All Races, Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.1	0.02		
High Five States				
District of Columbia	6.7 \oplus *	0.63	(01)	63.4
New Jersey	5.0 \oplus *	0.14	(02)	22.0
New York	4.8 \oplus *	0.09	(03)	17.1
Vermont	4.8*	0.54	(04)	17.1
New Hampshire	4.8*	0.39	(05)	17.1
Low Five States				
Arizona	3.3 \oplus *	0.16	(47)	-19.5
Colorado	3.2 \oplus *	0.19	(48)	-22.0
Hawaii	3.2*	0.32	(49)	-22.0
Alaska	2.9*	0.69	(50)	-29.3
North Dakota	2.6 \oplus *	0.35	(51)	-36.6

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.6	0.17	(40)	-12.2	Montana	3.9	0.39	(30)	-4.9
Alaska	2.9*	0.69	(50)	-29.3	Nebraska	4.2	0.29	(19)	2.4
Arizona	3.3 \oplus *	0.16	(47)	-19.5	Nevada	3.3*	0.29	(46)	-19.5
Arkansas	3.8	0.22	(35)	-7.3	New Hampshire	4.8*	0.39	(05)	17.1
California	3.9 \oplus	0.07	(33)	-4.9	New Jersey	5.0 \oplus *	0.14	(02)	22.0
Colorado	3.2 \oplus *	0.19	(48)	-22.0	New Mexico	3.8	0.30	(37)	-7.3
Connecticut	4.2	0.20	(17)	2.4	New York	4.8 \oplus *	0.09	(03)	17.1
Delaware	4.6	0.47	(09)	12.2	North Carolina	4.1	0.14	(23)	0.0
District of Columbia	6.7 \oplus *	0.63	(01)	63.4	North Dakota	2.6 \oplus *	0.35	(51)	-36.6
Florida	3.6 \oplus	0.08	(42)	-12.2	Ohio	4.7 \oplus	0.12	(06)	14.6
Georgia	3.9	0.15	(32)	-4.9	Oklahoma	3.6	0.19	(44)	-12.2
Hawaii	3.2*	0.32	(49)	-22.0	Oregon	4.0	0.21	(28)	-2.4
Idaho	3.8	0.35	(38)	-7.3	Pennsylvania	4.7 \oplus	0.11	(07)	14.6
Illinois	4.5	0.11	(10)	9.8	Rhode Island	4.2	0.37	(22)	2.4
Indiana	4.4	0.16	(11)	7.3	South Carolina	4.3	0.20	(16)	4.9
Iowa	4.6	0.22	(08)	12.2	South Dakota	3.6	0.41	(41)	-12.2
Kansas	3.9	0.22	(29)	-4.9	Tennessee	3.8	0.16	(36)	-7.3
Kentucky	3.8	0.18	(34)	-7.3	Texas	3.9	0.09	(31)	-4.9
Louisiana	4.1	0.19	(27)	0.0	Utah	3.6	0.30	(43)	-12.2
Maine	4.1	0.33	(24)	0.0	Vermont	4.8*	0.54	(04)	17.1
Maryland	4.3	0.18	(14)	4.9	Virginia	4.2	0.15	(21)	2.4
Massachusetts	4.1	0.15	(26)	0.0	Washington	3.7	0.16	(39)	-9.8
Michigan	4.4	0.13	(12)	7.3	West Virginia	4.2	0.26	(18)	2.4
Minnesota	4.1	0.18	(25)	0.0	Wisconsin	4.2	0.17	(20)	2.4
Mississippi	3.4 \oplus *	0.21	(45)	-17.1	Wyoming	4.4	0.60	(13)	7.3
Missouri	4.3	0.16	(15)	4.9					

ä NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

SE Standard error of the rate.


PD Percent difference between state rate and total U.S. rate.

* Absolute percent difference between state rate and total U.S. rate is 15% or more.

\oplus Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).


Figure VII-1

SEER Incidence and U.S. Death Rates⁺ Corpus Uteri By Race



⁺ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 U.S. standard million population by 5-year age groups. Regression lines are calculated using the Joinpoint Regression Program.

SEER Incidence and U.S. Death Rates⁺ Corpus Uteri, By Age and Race



⁺ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 U.S. standard million population by 5-year age groups. Regression lines are calculated using the Joinpoint Regression Program.