

Table I-15

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site, Race and Sex

12 SEER Areas, 1998-2000

Site	All Races		Whites		Blacks	
	Males	Females	Males	Females	Males	Females
All Sites	45.19	38.67	45.40	39.99	42.45	32.09
Invasive and In Situ	46.40	41.98	46.70	43.44	42.83	34.32
Oral cavity and Pharynx	1.40	0.67	1.40	0.69	1.38	0.50
Esophagus	0.75	0.26	0.77	0.25	0.77	0.38
Stomach	1.25	0.79	1.09	0.66	1.32	1.03
Colon and Rectum	5.97	5.66	6.03	5.66	4.94	5.38
Invasive and In Situ	6.31	5.94	6.36	5.92	5.29	5.71
Liver and Intrahepatic bile duct	0.86	0.42	0.72	0.35	0.80	0.36
Pancreas	1.23	1.24	1.23	1.22	1.27	1.34
Larynx	0.65	0.16	0.65	0.17	0.86	0.23
Invasive and In Situ	0.70	0.18	0.71	0.18	0.89	0.24
Lung and Bronchus	7.75	5.79	7.75	6.08	8.29	5.37
Melanoma of skin	1.83	1.23	2.16	1.48	0.10	0.08
Invasive and In Situ	2.88	1.96	3.38	2.32	0.13	0.12
Breast	0.11	13.51	0.12	14.28	0.14	10.14
Invasive and In Situ	0.13	16.06	0.13	16.92	0.15	12.14
Cervix uteri	-	0.79	-	0.75	-	0.97
Corpus and Uterus, NOS	-	2.62	-	2.82	-	1.67
Invasive and In Situ	-	2.66	-	2.87	-	1.69
Ovary	-	1.72	-	1.85	-	1.10
Prostate	17.28	-	16.90	-	20.40	-
Testis	0.35	-	0.42	-	0.10	-
Urinary bladder(Invasive and In Situ)	3.52	1.13	3.91	1.22	1.42	0.78
Kidney and Renal pelvis	1.46	0.87	1.53	0.91	1.23	0.86
Brain and Other nervous system	0.66	0.53	0.75	0.59	0.32	0.30
Thyroid	0.30	0.84	0.32	0.87	0.14	0.46
Hodgkin lymphoma	0.23	0.20	0.26	0.22	0.19	0.15
Non-Hodgkin lymphoma	2.12	1.79	2.26	1.91	1.17	1.04
Multiple myeloma	0.66	0.54	0.65	0.49	0.89	0.93
Leukemia	1.45	1.03	1.55	1.09	0.89	0.74
Acute Lymphocytic Leukemia	0.12	0.11	0.13	0.12	0.06	0.06
Chronic Lymphocytic Leukemia	0.47	0.29	0.51	0.31	0.30	0.19
Acute Myeloid Leukemia	0.45	0.35	0.47	0.36	0.29	0.28
Chronic Myeloid Leukemia	0.19	0.14	0.20	0.14	0.13	0.10

Note: Invasive cancer only unless specified otherwise.

Table I-16

Lifetime Risk (Percent) of Dying from Cancer by Site, Race and Sex

Total U.S., 1998-2000

Site	All Races		Whites		Blacks	
	Males	Females	Males	Females	Males	Females
All Sites	23.60	19.99	23.66	20.22	24.23	19.47
Oral cavity and Pharynx	0.39	0.20	0.37	0.20	0.54	0.19
Esophagus	0.71	0.23	0.71	0.21	0.84	0.32
Stomach	0.61	0.41	0.56	0.37	0.92	0.69
Colon and Rectum	2.39	2.30	2.40	2.28	2.39	2.56
Liver and Intrahepatic bile duct	0.61	0.36	0.57	0.34	0.67	0.37
Pancreas	1.15	1.16	1.15	1.15	1.14	1.33
Larynx	0.24	0.06	0.22	0.06	0.39	0.09
Lung and Bronchus	7.35	4.81	7.41	4.99	7.40	3.93
Melanoma of skin	0.35	0.19	0.40	0.22	0.04	0.05
Breast	0.03	3.03	0.03	3.03	0.04	3.24
Cervix uteri	-	0.27	-	0.24	-	0.50
Corpus and Uterus, NOS	-	0.50	-	0.48	-	0.72
Ovary	-	1.02	-	1.07	-	0.72
Prostate	3.05	-	2.86	-	4.72	-
Testis	0.02	-	0.02	-	0.01	-
Urinary bladder	0.73	0.31	0.78	0.32	0.38	0.32
Kidney and Renal pelvis	0.56	0.33	0.58	0.35	0.44	0.27
Brain and Other nervous system	0.49	0.40	0.54	0.43	0.23	0.21
Thyroid	0.04	0.06	0.04	0.06	0.02	0.05
Hodgkin lymphoma	0.05	0.04	0.05	0.04	0.04	0.03
Non-Hodgkin lymphoma	0.96	0.85	1.03	0.91	0.50	0.44
Multiple myeloma	0.44	0.40	0.43	0.37	0.63	0.69
Leukemia	0.94	0.72	0.99	0.75	0.62	0.54
Acute Lymphocytic Leukemia	0.05	0.04	0.05	0.04	0.03	0.02
Chronic Lymphocytic Leukemia	0.22	0.15	0.23	0.16	0.15	0.11
Acute Myeloid Leukemia	0.30	0.24	0.31	0.25	0.18	0.17
Chronic Myeloid Leukemia	0.08	0.06	0.09	0.06	0.07	0.06

Table II-8
ALL SITES (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.17	0.33	0.67	45.19	0.16	0.32	0.75	38.67
10	0.16	0.51	1.21	45.88	0.16	0.61	1.79	39.14
20	0.35	1.06	2.89	46.02	0.45	1.63	4.60	39.12
30	0.72	2.57	8.57	46.34	1.20	4.19	10.05	38.99
40	1.90	8.03	21.49	46.64	3.05	9.01	18.28	38.49
50	6.43	20.55	36.53	46.92	6.23	15.91	26.89	37.02
60	15.91	33.92	43.70	45.63	10.59	22.61	31.30	33.70

Lifetime Risk of Being Diagnosed = 45.19%
 Lifetime Risk of Dying = 23.60%

Lifetime Risk of Being Diagnosed = 38.67%
 Lifetime Risk of Dying = 19.99%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.19	0.36	0.73	45.40	0.17	0.34	0.81	39.99
10	0.17	0.55	1.28	45.97	0.17	0.65	1.87	40.38
20	0.38	1.11	2.92	46.09	0.48	1.71	4.74	40.35
30	0.74	2.58	8.52	46.33	1.24	4.29	10.34	40.20
40	1.87	7.95	21.50	46.56	3.11	9.27	18.99	39.66
50	6.35	20.53	36.51	46.73	6.42	16.57	27.96	38.13
60	15.89	33.81	43.43	45.27	11.09	23.56	32.35	34.68

Lifetime Risk of Being Diagnosed = 45.40%
 Lifetime Risk of Dying = 23.66%

Lifetime Risk of Being Diagnosed = 39.99%
 Lifetime Risk of Dying = 20.22%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.10	0.22	0.45	42.45	0.11	0.23	0.57	32.09
10	0.12	0.36	1.00	43.90	0.12	0.47	1.52	32.97
20	0.24	0.89	3.28	44.21	0.35	1.41	4.19	32.98
30	0.67	3.12	11.21	45.17	1.07	3.88	9.32	32.98
40	2.55	10.94	25.95	46.17	2.88	8.45	16.81	32.70
50	9.18	25.60	40.30	47.72	5.91	14.79	24.02	31.65
60	20.33	38.53	46.24	47.71	10.00	20.40	27.08	29.01

Lifetime Risk of Being Diagnosed = 42.45%
 Lifetime Risk of Dying = 24.23%

Lifetime Risk of Being Diagnosed = 32.09%
 Lifetime Risk of Dying = 19.47%

Table III-8
BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.03	0.06	0.08	0.66	0.03	0.05	0.07	0.53
10	0.02	0.05	0.09	0.64	0.02	0.04	0.07	0.50
20	0.02	0.06	0.12	0.62	0.02	0.05	0.09	0.48
30	0.04	0.10	0.20	0.61	0.03	0.07	0.14	0.47
40	0.06	0.16	0.31	0.57	0.04	0.11	0.21	0.44
50	0.10	0.26	0.43	0.53	0.07	0.17	0.31	0.41
60	0.17	0.35	0.45	0.46	0.11	0.25	0.34	0.35

Lifetime Risk of Being Diagnosed = 0.66%
 Lifetime Risk of Dying = 0.49%

Lifetime Risk of Being Diagnosed = 0.53%
 Lifetime Risk of Dying = 0.40%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.04	0.06	0.09	0.75	0.03	0.06	0.08	0.59
10	0.03	0.05	0.10	0.72	0.02	0.04	0.08	0.56
20	0.03	0.07	0.13	0.70	0.02	0.06	0.10	0.54
30	0.04	0.11	0.22	0.68	0.03	0.08	0.15	0.52
40	0.07	0.18	0.35	0.64	0.05	0.12	0.24	0.49
50	0.12	0.30	0.48	0.60	0.08	0.19	0.35	0.45
60	0.20	0.39	0.50	0.51	0.12	0.28	0.37	0.39

Lifetime Risk of Being Diagnosed = 0.75%
 Lifetime Risk of Dying = 0.54%

Lifetime Risk of Being Diagnosed = 0.59%
 Lifetime Risk of Dying = 0.43%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.03	0.04	0.05	0.32	0.02	0.04	0.05	0.30
10	0.02	0.03	0.05	0.31	0.02	0.03	0.05	0.29
20	0.01	0.04	0.08	0.29	0.01	0.03	0.05	0.27
30	0.03	0.07	0.14	0.29	0.02	0.04	0.10	0.26
40	0.05	0.11	0.19	0.27	0.03	0.08	0.15	0.25
50	0.07	0.15	0.21	0.24	0.06	0.13	0.19	0.23
60	0.09	0.16	0.19	0.19	0.07	0.14	0.18	0.18

Lifetime Risk of Being Diagnosed = 0.32%
 Lifetime Risk of Dying = 0.23%

Lifetime Risk of Being Diagnosed = 0.30%
 Lifetime Risk of Dying = 0.21%

Table IV-11
BREAST CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.11	0.00	0.00	0.05	13.51
10	0.00	0.00	0.00	0.12	0.00	0.05	0.44	13.71
20	0.00	0.00	0.01	0.12	0.05	0.44	1.87	13.74
30	0.00	0.01	0.02	0.12	0.40	1.83	4.49	13.76
40	0.00	0.02	0.05	0.12	1.45	4.14	7.59	13.52
50	0.02	0.05	0.08	0.12	2.78	6.34	9.82	12.45
60	0.04	0.07	0.10	0.11	3.81	7.53	9.81	10.36

Lifetime Risk of Being Diagnosed = 0.11% Lifetime Risk of Being Diagnosed = 13.51%
 Lifetime Risk of Dying = 0.03% Lifetime Risk of Dying = 3.03%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.12	0.00	0.00	0.04	14.28
10	0.00	0.00	0.00	0.12	0.00	0.04	0.44	14.46
20	0.00	0.00	0.01	0.12	0.04	0.43	1.89	14.49
30	0.00	0.01	0.02	0.12	0.39	1.86	4.64	14.51
40	0.00	0.02	0.06	0.12	1.48	4.29	7.98	14.26
50	0.02	0.05	0.09	0.12	2.90	6.70	10.42	13.17
60	0.04	0.08	0.11	0.11	4.06	8.05	10.44	10.98

Lifetime Risk of Being Diagnosed = 0.12% Lifetime Risk of Being Diagnosed = 14.28%
 Lifetime Risk of Dying = 0.03% Lifetime Risk of Dying = 3.03%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.14	0.00	0.00	0.07	10.14
10	0.00	0.00	0.00	0.14	0.00	0.07	0.51	10.44
20	0.00	0.00	0.01	0.14	0.07	0.51	1.86	10.47
30	0.00	0.01	0.04	0.15	0.44	1.80	4.06	10.49
40	0.01	0.04	0.08	0.15	1.39	3.70	6.34	10.25
50	0.04	0.08	0.12	0.15	2.43	5.21	7.66	9.33
60	0.05	0.09	0.11	0.14	3.09	5.80	7.26	7.65

Lifetime Risk of Being Diagnosed = 0.14% Lifetime Risk of Being Diagnosed = 10.14%
 Lifetime Risk of Dying = 0.04% Lifetime Risk of Dying = 3.24%

Table V-6
CERVIX UTERI CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.04	0.79
10	0.00	0.04	0.16	0.80
20	0.04	0.16	0.32	0.80
30	0.12	0.28	0.43	0.76
40	0.16	0.31	0.46	0.65
50	0.15	0.30	0.41	0.50
60	0.16	0.27	0.34	0.36

Lifetime Risk of Being Diagnosed = 0.79%
 Lifetime Risk of Dying = 0.27%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.04	0.75
10	0.00	0.05	0.17	0.76
20	0.04	0.17	0.33	0.76
30	0.13	0.28	0.43	0.72
40	0.16	0.30	0.44	0.60
50	0.15	0.29	0.38	0.45
60	0.15	0.24	0.30	0.32

Lifetime Risk of Being Diagnosed = 0.75%
 Lifetime Risk of Dying = 0.24%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.03	0.97
10	0.00	0.03	0.15	1.00
20	0.03	0.15	0.34	1.00
30	0.12	0.31	0.46	0.98
40	0.20	0.35	0.55	0.88
50	0.16	0.36	0.55	0.71
60	0.22	0.42	0.54	0.60

Lifetime Risk of Being Diagnosed = 0.97%
 Lifetime Risk of Dying = 0.50%

Table VI-13
COLON AND RECTUM CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	5.97	0.00	0.00	0.01	5.66
10	0.00	0.01	0.06	6.07	0.00	0.01	0.06	5.75
20	0.01	0.06	0.25	6.10	0.01	0.06	0.23	5.76
30	0.05	0.25	0.90	6.17	0.05	0.22	0.71	5.77
40	0.20	0.86	2.33	6.22	0.17	0.67	1.79	5.77
50	0.68	2.21	4.33	6.24	0.50	1.65	3.41	5.71
60	1.65	3.94	5.62	6.00	1.20	3.05	4.87	5.47

Lifetime Risk of Being Diagnosed = 5.97%
 Lifetime Risk of Dying = 2.39%

Lifetime Risk of Being Diagnosed = 5.66%
 Lifetime Risk of Dying = 2.30%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	6.03	0.00	0.00	0.01	5.66
10	0.00	0.01	0.06	6.12	0.00	0.01	0.06	5.73
20	0.01	0.06	0.24	6.15	0.01	0.06	0.22	5.74
30	0.05	0.23	0.87	6.20	0.04	0.21	0.67	5.75
40	0.19	0.83	2.31	6.25	0.16	0.62	1.73	5.75
50	0.67	2.20	4.34	6.26	0.47	1.60	3.38	5.69
60	1.64	3.95	5.64	6.01	1.18	3.04	4.88	5.46

Lifetime Risk of Being Diagnosed = 6.03%
 Lifetime Risk of Dying = 2.40%

Lifetime Risk of Being Diagnosed = 5.66%
 Lifetime Risk of Dying = 2.28%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	4.94	0.00	0.00	0.02	5.38
10	0.00	0.01	0.06	5.12	0.00	0.02	0.08	5.55
20	0.01	0.06	0.31	5.16	0.02	0.08	0.30	5.56
30	0.05	0.31	1.08	5.28	0.07	0.29	0.96	5.59
40	0.26	1.06	2.48	5.40	0.22	0.91	2.26	5.62
50	0.86	2.39	4.25	5.54	0.71	2.12	3.84	5.62
60	1.79	3.95	5.23	5.46	1.54	3.42	4.88	5.37

Lifetime Risk of Being Diagnosed = 4.94%
 Lifetime Risk of Dying = 2.39%

Lifetime Risk of Being Diagnosed = 5.38%
 Lifetime Risk of Dying = 2.56%

Table VII-6
CORPUS AND UTERUS, NOS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.62
10	0.00	0.01	0.06	2.66
20	0.01	0.06	0.23	2.67
30	0.05	0.22	0.73	2.67
40	0.18	0.69	1.47	2.65
50	0.53	1.32	2.02	2.52
60	0.83	1.57	2.00	2.09

Lifetime Risk of Being Diagnosed = 2.62%
Lifetime Risk of Dying = 0.50%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.82
10	0.00	0.01	0.06	2.86
20	0.01	0.06	0.23	2.87
30	0.05	0.23	0.78	2.87
40	0.18	0.74	1.59	2.84
50	0.57	1.44	2.19	2.71
60	0.90	1.69	2.15	2.24

Lifetime Risk of Being Diagnosed = 2.82%
Lifetime Risk of Dying = 0.48%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.67
10	0.00	0.01	0.04	1.72
20	0.01	0.04	0.13	1.72
30	0.03	0.12	0.39	1.73
40	0.09	0.37	0.97	1.73
50	0.29	0.92	1.40	1.70
60	0.69	1.22	1.48	1.55

Lifetime Risk of Being Diagnosed = 1.67%
Lifetime Risk of Dying = 0.72%

Table VIII-8
ESOPHAGUS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.75	0.00	0.00	0.00	0.26
10	0.00	0.00	0.01	0.77	0.00	0.00	0.00	0.26
20	0.00	0.01	0.03	0.77	0.00	0.00	0.01	0.26
30	0.00	0.03	0.14	0.78	0.00	0.01	0.03	0.27
40	0.03	0.14	0.36	0.79	0.01	0.03	0.09	0.27
50	0.11	0.35	0.61	0.78	0.02	0.08	0.17	0.27
60	0.25	0.53	0.69	0.72	0.06	0.15	0.23	0.25

Lifetime Risk of Being Diagnosed = 0.75%
 Lifetime Risk of Dying = 0.71%

Lifetime Risk of Being Diagnosed = 0.26%
 Lifetime Risk of Dying = 0.23%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.77	0.00	0.00	0.00	0.25
10	0.00	0.00	0.01	0.78	0.00	0.00	0.00	0.26
20	0.00	0.01	0.03	0.78	0.00	0.00	0.01	0.26
30	0.01	0.03	0.14	0.79	0.00	0.00	0.02	0.26
40	0.03	0.14	0.36	0.80	0.00	0.02	0.08	0.26
50	0.11	0.34	0.61	0.79	0.02	0.08	0.16	0.26
60	0.25	0.53	0.70	0.73	0.06	0.15	0.22	0.25

Lifetime Risk of Being Diagnosed = 0.77%
 Lifetime Risk of Dying = 0.71%

Lifetime Risk of Being Diagnosed = 0.25%
 Lifetime Risk of Dying = 0.21%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.77	0.00	0.00	0.00	0.38
10	0.00	0.00	0.01	0.80	0.00	0.00	0.00	0.39
20	0.00	0.01	0.04	0.80	0.00	0.00	0.02	0.40
30	0.00	0.04	0.21	0.82	0.00	0.02	0.08	0.40
40	0.04	0.21	0.49	0.85	0.02	0.08	0.20	0.40
50	0.19	0.49	0.77	0.87	0.07	0.19	0.31	0.40
60	0.35	0.67	0.77	0.79	0.13	0.26	0.34	0.36

Lifetime Risk of Being Diagnosed = 0.77%
 Lifetime Risk of Dying = 0.84%

Lifetime Risk of Being Diagnosed = 0.38%
 Lifetime Risk of Dying = 0.32%

Table IX-8
HODGKIN LYMPHOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.06	0.23	0.00	0.02	0.06	0.20
10	0.02	0.06	0.09	0.23	0.02	0.06	0.09	0.20
20	0.04	0.08	0.11	0.22	0.04	0.07	0.09	0.18
30	0.04	0.07	0.10	0.18	0.03	0.05	0.07	0.14
40	0.03	0.06	0.09	0.14	0.02	0.04	0.06	0.11
50	0.03	0.06	0.10	0.12	0.02	0.05	0.07	0.09
60	0.04	0.07	0.09	0.09	0.03	0.05	0.07	0.07

Lifetime Risk of Being Diagnosed = 0.23%
 Lifetime Risk of Dying = 0.05%

Lifetime Risk of Being Diagnosed = 0.20%
 Lifetime Risk of Dying = 0.04%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.07	0.26	0.00	0.03	0.07	0.22
10	0.02	0.06	0.10	0.26	0.03	0.07	0.11	0.22
20	0.04	0.08	0.12	0.24	0.05	0.08	0.10	0.20
30	0.04	0.07	0.10	0.20	0.04	0.06	0.08	0.15
40	0.04	0.06	0.10	0.16	0.02	0.04	0.07	0.12
50	0.03	0.07	0.10	0.13	0.02	0.05	0.08	0.10
60	0.04	0.08	0.10	0.10	0.03	0.06	0.07	0.08

Lifetime Risk of Being Diagnosed = 0.26%
 Lifetime Risk of Dying = 0.05%

Lifetime Risk of Being Diagnosed = 0.22%
 Lifetime Risk of Dying = 0.04%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.05	0.19	0.00	0.01	0.05	0.15
10	0.01	0.05	0.09	0.19	0.01	0.05	0.08	0.15
20	0.04	0.08	0.12	0.18	0.04	0.07	0.09	0.14
30	0.04	0.08	0.11	0.15	0.03	0.05	0.07	0.10
40	0.04	0.07	0.09	0.11	0.02	0.04	0.05	0.07
50	0.03	0.05	0.07	0.07	0.02	0.03	0.05	0.05
60	0.02	0.04	0.04	0.04	0.01	0.03	0.03	0.04

Lifetime Risk of Being Diagnosed = 0.19%
 Lifetime Risk of Dying = 0.04%

Lifetime Risk of Being Diagnosed = 0.15%
 Lifetime Risk of Dying = 0.03%

Table XI-8

KIDNEY AND RENAL PELVIS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.01	0.02	1.46	0.01	0.02	0.02	0.87
10	0.00	0.01	0.03	1.47	0.00	0.00	0.02	0.87
20	0.00	0.02	0.11	1.48	0.00	0.02	0.07	0.87
30	0.02	0.11	0.35	1.49	0.02	0.06	0.18	0.87
40	0.09	0.34	0.76	1.50	0.05	0.16	0.39	0.86
50	0.26	0.69	1.15	1.46	0.12	0.35	0.61	0.83
60	0.47	0.97	1.25	1.29	0.24	0.51	0.70	0.75

Lifetime Risk of Being Diagnosed = 1.46%
Lifetime Risk of Dying = 0.56%

Lifetime Risk of Being Diagnosed = 0.87%
Lifetime Risk of Dying = 0.33%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.01	0.02	1.53	0.01	0.02	0.02	0.91
10	0.00	0.01	0.03	1.54	0.00	0.00	0.02	0.91
20	0.00	0.02	0.11	1.55	0.00	0.02	0.07	0.91
30	0.02	0.11	0.37	1.56	0.02	0.06	0.18	0.91
40	0.09	0.35	0.79	1.57	0.05	0.17	0.41	0.90
50	0.27	0.72	1.21	1.52	0.12	0.36	0.64	0.87
60	0.49	1.01	1.30	1.34	0.25	0.54	0.74	0.78

Lifetime Risk of Being Diagnosed = 1.53%
Lifetime Risk of Dying = 0.58%

Lifetime Risk of Being Diagnosed = 0.91%
Lifetime Risk of Dying = 0.35%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.01	0.02	1.23	0.02	0.02	0.03	0.86
10	0.00	0.01	0.04	1.27	0.00	0.01	0.03	0.86
20	0.01	0.03	0.14	1.28	0.01	0.03	0.09	0.86
30	0.03	0.13	0.40	1.30	0.02	0.09	0.23	0.87
40	0.11	0.38	0.77	1.32	0.06	0.21	0.44	0.85
50	0.29	0.71	1.09	1.30	0.15	0.40	0.65	0.83
60	0.49	0.93	1.13	1.17	0.27	0.54	0.69	0.73

Lifetime Risk of Being Diagnosed = 1.23%
Lifetime Risk of Dying = 0.44%

Lifetime Risk of Being Diagnosed = 0.86%
Lifetime Risk of Dying = 0.27%

Table XII-8
LARYNX CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.65	0.00	0.00	0.00	0.16
10	0.00	0.00	0.00	0.66	0.00	0.00	0.00	0.17
20	0.00	0.00	0.03	0.66	0.00	0.00	0.01	0.17
30	0.00	0.03	0.14	0.67	0.00	0.01	0.04	0.17
40	0.03	0.14	0.36	0.68	0.01	0.04	0.09	0.17
50	0.11	0.34	0.56	0.67	0.03	0.08	0.13	0.16
60	0.25	0.48	0.58	0.60	0.06	0.11	0.13	0.14

Lifetime Risk of Being Diagnosed = 0.65%
 Lifetime Risk of Dying = 0.24%

Lifetime Risk of Being Diagnosed = 0.16%
 Lifetime Risk of Dying = 0.06%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.65	0.00	0.00	0.00	0.17
10	0.00	0.00	0.00	0.66	0.00	0.00	0.00	0.17
20	0.00	0.00	0.03	0.66	0.00	0.00	0.01	0.17
30	0.00	0.03	0.13	0.67	0.00	0.01	0.04	0.17
40	0.03	0.13	0.36	0.68	0.01	0.04	0.09	0.17
50	0.11	0.34	0.56	0.67	0.03	0.08	0.14	0.16
60	0.25	0.48	0.58	0.60	0.06	0.12	0.14	0.14

Lifetime Risk of Being Diagnosed = 0.65%
 Lifetime Risk of Dying = 0.22%

Lifetime Risk of Being Diagnosed = 0.17%
 Lifetime Risk of Dying = 0.06%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.86	0.00	0.00	0.00	0.23
10	0.00	0.00	0.00	0.89	0.00	0.00	0.00	0.24
20	0.00	0.00	0.07	0.90	0.00	0.00	0.02	0.24
30	0.00	0.07	0.28	0.92	0.00	0.02	0.07	0.24
40	0.07	0.28	0.59	0.95	0.02	0.07	0.15	0.24
50	0.23	0.57	0.82	0.95	0.05	0.14	0.20	0.23
60	0.39	0.69	0.82	0.83	0.10	0.16	0.19	0.20

Lifetime Risk of Being Diagnosed = 0.86%
 Lifetime Risk of Dying = 0.39%

Lifetime Risk of Being Diagnosed = 0.23%
 Lifetime Risk of Dying = 0.09%

Table XIII-14

LEUKEMIA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.06	0.09	0.12	1.45	0.05	0.08	0.10	1.03
10	0.03	0.06	0.09	1.41	0.03	0.05	0.07	1.00
20	0.03	0.06	0.13	1.39	0.02	0.05	0.09	0.97
30	0.04	0.10	0.24	1.38	0.03	0.07	0.17	0.95
40	0.06	0.20	0.50	1.37	0.05	0.14	0.31	0.94
50	0.14	0.45	0.91	1.35	0.10	0.27	0.54	0.91
60	0.33	0.82	1.20	1.30	0.18	0.46	0.75	0.85

Lifetime Risk of Being Diagnosed = 1.45%
Lifetime Risk of Dying = 0.94%

Lifetime Risk of Being Diagnosed = 1.03%
Lifetime Risk of Dying = 0.72%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.07	0.10	0.13	1.55	0.06	0.09	0.11	1.09
10	0.03	0.06	0.10	1.50	0.03	0.05	0.08	1.05
20	0.03	0.06	0.13	1.48	0.02	0.05	0.10	1.02
30	0.04	0.10	0.24	1.46	0.03	0.07	0.17	1.00
40	0.06	0.21	0.52	1.45	0.05	0.14	0.33	0.98
50	0.15	0.48	0.97	1.43	0.10	0.28	0.57	0.95
60	0.35	0.88	1.27	1.37	0.20	0.49	0.79	0.89

Lifetime Risk of Being Diagnosed = 1.55%
Lifetime Risk of Dying = 0.99%

Lifetime Risk of Being Diagnosed = 1.09%
Lifetime Risk of Dying = 0.75%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.03	0.05	0.07	0.89	0.02	0.04	0.06	0.74
10	0.02	0.04	0.08	0.89	0.02	0.04	0.06	0.74
20	0.02	0.06	0.12	0.88	0.02	0.04	0.08	0.72
30	0.04	0.10	0.21	0.88	0.02	0.06	0.15	0.71
40	0.06	0.18	0.41	0.87	0.04	0.13	0.26	0.70
50	0.12	0.38	0.65	0.87	0.09	0.23	0.43	0.69
60	0.29	0.61	0.83	0.86	0.14	0.37	0.57	0.65

Lifetime Risk of Being Diagnosed = 0.89%
Lifetime Risk of Dying = 0.62%

Lifetime Risk of Being Diagnosed = 0.74%
Lifetime Risk of Dying = 0.54%

Table XIV-8
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.01	0.86	0.01	0.01	0.01	0.42
10	0.00	0.00	0.01	0.87	0.00	0.00	0.01	0.42
20	0.00	0.01	0.07	0.88	0.00	0.01	0.02	0.42
30	0.01	0.07	0.22	0.89	0.00	0.02	0.05	0.42
40	0.06	0.21	0.45	0.89	0.02	0.05	0.15	0.42
50	0.15	0.40	0.68	0.86	0.04	0.13	0.27	0.41
60	0.26	0.56	0.73	0.76	0.10	0.25	0.36	0.39

Lifetime Risk of Being Diagnosed = 0.86%
 Lifetime Risk of Dying = 0.61%

Lifetime Risk of Being Diagnosed = 0.42%
 Lifetime Risk of Dying = 0.36%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.01	0.72	0.01	0.01	0.01	0.35
10	0.00	0.00	0.01	0.72	0.00	0.00	0.01	0.35
20	0.00	0.01	0.05	0.73	0.00	0.00	0.02	0.35
30	0.01	0.05	0.16	0.73	0.00	0.01	0.04	0.35
40	0.05	0.16	0.35	0.74	0.01	0.04	0.11	0.35
50	0.11	0.31	0.55	0.72	0.03	0.10	0.22	0.35
60	0.21	0.47	0.61	0.64	0.08	0.20	0.30	0.33

Lifetime Risk of Being Diagnosed = 0.72%
 Lifetime Risk of Dying = 0.57%

Lifetime Risk of Being Diagnosed = 0.35%
 Lifetime Risk of Dying = 0.34%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	0.80	0.00	0.00	0.01	0.36
10	0.00	0.00	0.01	0.83	0.00	0.00	0.01	0.37
20	0.00	0.01	0.10	0.84	0.00	0.00	0.02	0.37
30	0.01	0.10	0.33	0.85	0.00	0.02	0.07	0.37
40	0.10	0.33	0.57	0.87	0.02	0.07	0.18	0.37
50	0.26	0.51	0.72	0.83	0.05	0.17	0.30	0.37
60	0.29	0.54	0.65	0.67	0.13	0.26	0.33	0.34

Lifetime Risk of Being Diagnosed = 0.80%
 Lifetime Risk of Dying = 0.67%

Lifetime Risk of Being Diagnosed = 0.36%
 Lifetime Risk of Dying = 0.37%

Table XV-13
LUNG AND BRONCHUS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	7.75	0.00	0.00	0.01	5.79
10	0.00	0.00	0.03	7.89	0.00	0.01	0.03	5.87
20	0.00	0.03	0.21	7.93	0.00	0.03	0.18	5.88
30	0.03	0.21	1.03	8.02	0.03	0.18	0.81	5.90
40	0.19	1.02	3.26	8.13	0.15	0.79	2.39	5.93
50	0.86	3.18	6.24	8.22	0.65	2.28	4.40	5.89
60	2.50	5.80	7.63	7.93	1.71	3.93	5.25	5.49

Lifetime Risk of Being Diagnosed = 7.75%
 Lifetime Risk of Dying = 7.35%

Lifetime Risk of Being Diagnosed = 5.79%
 Lifetime Risk of Dying = 4.81%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	7.75	0.00	0.00	0.01	6.08
10	0.00	0.00	0.03	7.86	0.00	0.01	0.03	6.15
20	0.00	0.03	0.19	7.90	0.00	0.03	0.18	6.16
30	0.02	0.19	0.96	7.98	0.03	0.17	0.83	6.18
40	0.17	0.95	3.20	8.08	0.15	0.81	2.51	6.20
50	0.81	3.13	6.20	8.17	0.67	2.41	4.65	6.16
60	2.49	5.78	7.60	7.89	1.81	4.16	5.51	5.73

Lifetime Risk of Being Diagnosed = 7.75%
 Lifetime Risk of Dying = 7.41%

Lifetime Risk of Being Diagnosed = 6.08%
 Lifetime Risk of Dying = 4.99%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	8.29	0.00	0.00	0.00	5.37
10	0.00	0.00	0.06	8.59	0.00	0.00	0.05	5.53
20	0.00	0.06	0.45	8.66	0.00	0.05	0.29	5.55
30	0.06	0.46	1.89	8.88	0.04	0.29	1.12	5.59
40	0.42	1.89	4.68	9.11	0.25	1.09	2.76	5.64
50	1.59	4.59	7.78	9.37	0.88	2.62	4.49	5.62
60	3.49	7.20	8.81	9.05	1.90	3.94	4.98	5.17

Lifetime Risk of Being Diagnosed = 8.29%
 Lifetime Risk of Dying = 7.40%

Lifetime Risk of Being Diagnosed = 5.37%
 Lifetime Risk of Dying = 3.93%

Table XVI-8
MELANOMA OF SKIN (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.04	1.83	0.00	0.01	0.07	1.23
10	0.01	0.04	0.12	1.86	0.01	0.07	0.19	1.25
20	0.03	0.12	0.29	1.86	0.06	0.18	0.35	1.24
30	0.09	0.26	0.57	1.85	0.12	0.29	0.51	1.19
40	0.18	0.49	0.93	1.79	0.18	0.39	0.63	1.08
50	0.32	0.77	1.27	1.67	0.22	0.46	0.71	0.92
60	0.49	1.03	1.38	1.46	0.25	0.51	0.68	0.73

Lifetime Risk of Being Diagnosed = 1.83%
 Lifetime Risk of Dying = 0.35%

Lifetime Risk of Being Diagnosed = 1.23%
 Lifetime Risk of Dying = 0.19%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.05	2.16	0.00	0.01	0.09	1.48
10	0.01	0.05	0.15	2.19	0.01	0.09	0.24	1.50
20	0.04	0.14	0.35	2.20	0.08	0.23	0.44	1.49
30	0.10	0.32	0.68	2.18	0.15	0.37	0.63	1.42
40	0.22	0.58	1.10	2.11	0.22	0.48	0.77	1.28
50	0.38	0.92	1.51	1.95	0.27	0.56	0.85	1.08
60	0.58	1.21	1.62	1.69	0.31	0.60	0.80	0.85

Lifetime Risk of Being Diagnosed = 2.16%
 Lifetime Risk of Dying = 0.40%

Lifetime Risk of Being Diagnosed = 1.48%
 Lifetime Risk of Dying = 0.22%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.10	0.00	0.00	0.00	0.08
10	0.00	0.00	0.01	0.11	0.00	0.00	0.01	0.08
20	0.00	0.01	0.01	0.11	0.00	0.01	0.02	0.08
30	0.01	0.01	0.02	0.11	0.01	0.01	0.03	0.08
40	0.01	0.02	0.05	0.10	0.01	0.03	0.04	0.08
50	0.01	0.05	0.07	0.11	0.02	0.03	0.05	0.07
60	0.04	0.07	0.10	0.11	0.02	0.03	0.05	0.06

Lifetime Risk of Being Diagnosed = 0.10%
 Lifetime Risk of Dying = 0.04%

Lifetime Risk of Being Diagnosed = 0.08%
 Lifetime Risk of Dying = 0.05%

Table XVII-8
MULTIPLE MYELOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.66	0.00	0.00	0.00	0.54
10	0.00	0.00	0.01	0.67	0.00	0.00	0.00	0.55
20	0.00	0.01	0.03	0.68	0.00	0.00	0.02	0.55
30	0.01	0.03	0.11	0.68	0.00	0.02	0.08	0.55
40	0.03	0.11	0.28	0.69	0.02	0.07	0.20	0.55
50	0.08	0.26	0.50	0.69	0.06	0.19	0.37	0.54
60	0.19	0.45	0.62	0.65	0.13	0.32	0.48	0.51

Lifetime Risk of Being Diagnosed = 0.66%
 Lifetime Risk of Dying = 0.44%

Lifetime Risk of Being Diagnosed = 0.54%
 Lifetime Risk of Dying = 0.40%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.65	0.00	0.00	0.00	0.49
10	0.00	0.00	0.01	0.66	0.00	0.00	0.00	0.50
20	0.00	0.01	0.03	0.66	0.00	0.00	0.02	0.50
30	0.01	0.03	0.10	0.67	0.00	0.02	0.06	0.50
40	0.02	0.10	0.26	0.67	0.01	0.06	0.18	0.50
50	0.08	0.25	0.48	0.67	0.05	0.16	0.33	0.50
60	0.18	0.43	0.60	0.63	0.12	0.29	0.44	0.47

Lifetime Risk of Being Diagnosed = 0.65%
 Lifetime Risk of Dying = 0.43%

Lifetime Risk of Being Diagnosed = 0.49%
 Lifetime Risk of Dying = 0.37%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.89	0.00	0.00	0.00	0.93
10	0.00	0.00	0.01	0.92	0.00	0.00	0.01	0.95
20	0.00	0.01	0.06	0.93	0.00	0.01	0.05	0.96
30	0.01	0.06	0.23	0.95	0.01	0.05	0.19	0.96
40	0.05	0.23	0.52	0.98	0.04	0.19	0.42	0.97
50	0.19	0.50	0.81	0.99	0.15	0.39	0.72	0.97
60	0.36	0.72	0.90	0.93	0.26	0.62	0.86	0.89

Lifetime Risk of Being Diagnosed = 0.89%
 Lifetime Risk of Dying = 0.63%

Lifetime Risk of Being Diagnosed = 0.93%
 Lifetime Risk of Dying = 0.69%

Table XVIII-13
NON-HODGKIN LYMPHOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.03	0.06	2.12	0.00	0.01	0.03	1.79
10	0.02	0.05	0.13	2.15	0.01	0.03	0.08	1.82
20	0.03	0.11	0.27	2.14	0.02	0.07	0.17	1.81
30	0.08	0.24	0.52	2.14	0.05	0.15	0.35	1.80
40	0.16	0.45	0.95	2.09	0.10	0.30	0.68	1.77
50	0.30	0.81	1.48	2.00	0.21	0.59	1.14	1.70
60	0.56	1.27	1.74	1.84	0.40	0.98	1.45	1.57

Lifetime Risk of Being Diagnosed = 2.12%
Lifetime Risk of Dying = 0.96%

Lifetime Risk of Being Diagnosed = 1.79%
Lifetime Risk of Dying = 0.85%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.03	0.06	2.26	0.00	0.01	0.03	1.91
10	0.02	0.05	0.13	2.28	0.01	0.03	0.08	1.93
20	0.03	0.11	0.27	2.28	0.02	0.07	0.17	1.92
30	0.08	0.24	0.54	2.27	0.05	0.15	0.36	1.91
40	0.16	0.47	1.01	2.22	0.10	0.32	0.73	1.88
50	0.32	0.87	1.59	2.13	0.22	0.64	1.23	1.81
60	0.60	1.36	1.85	1.94	0.44	1.06	1.54	1.66

Lifetime Risk of Being Diagnosed = 2.26%
Lifetime Risk of Dying = 1.03%

Lifetime Risk of Being Diagnosed = 1.91%
Lifetime Risk of Dying = 0.91%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.02	0.05	1.17	0.00	0.01	0.03	1.04
10	0.01	0.05	0.15	1.20	0.01	0.03	0.09	1.07
20	0.03	0.14	0.31	1.20	0.02	0.08	0.18	1.07
30	0.11	0.29	0.51	1.20	0.06	0.16	0.32	1.05
40	0.19	0.42	0.70	1.13	0.11	0.27	0.52	1.02
50	0.25	0.56	0.83	1.02	0.17	0.42	0.69	0.95
60	0.36	0.68	0.85	0.89	0.28	0.57	0.78	0.85

Lifetime Risk of Being Diagnosed = 1.17%
Lifetime Risk of Dying = 0.50%

Lifetime Risk of Being Diagnosed = 1.04%
Lifetime Risk of Dying = 0.44%

Table XIX-8
ORAL CAVITY AND PHARYNX CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.40	0.00	0.00	0.01	0.67
10	0.00	0.01	0.04	1.42	0.00	0.01	0.03	0.68
20	0.01	0.04	0.16	1.43	0.01	0.02	0.07	0.68
30	0.03	0.15	0.43	1.43	0.02	0.06	0.16	0.68
40	0.12	0.41	0.81	1.43	0.05	0.14	0.30	0.66
50	0.30	0.71	1.09	1.35	0.10	0.26	0.45	0.63
60	0.45	0.86	1.08	1.14	0.17	0.37	0.52	0.56

Lifetime Risk of Being Diagnosed = 1.40%
 Lifetime Risk of Dying = 0.39%

Lifetime Risk of Being Diagnosed = 0.67%
 Lifetime Risk of Dying = 0.20%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.40	0.00	0.00	0.01	0.69
10	0.00	0.01	0.04	1.42	0.00	0.01	0.02	0.70
20	0.01	0.04	0.15	1.43	0.01	0.02	0.06	0.69
30	0.03	0.14	0.42	1.44	0.01	0.05	0.14	0.69
40	0.11	0.39	0.80	1.43	0.04	0.13	0.30	0.68
50	0.29	0.71	1.10	1.36	0.09	0.26	0.47	0.65
60	0.45	0.87	1.09	1.15	0.18	0.39	0.54	0.58

Lifetime Risk of Being Diagnosed = 1.40%
 Lifetime Risk of Dying = 0.37%

Lifetime Risk of Being Diagnosed = 0.69%
 Lifetime Risk of Dying = 0.20%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.38	0.00	0.01	0.01	0.50
10	0.00	0.01	0.04	1.43	0.00	0.01	0.03	0.51
20	0.01	0.04	0.22	1.44	0.00	0.03	0.09	0.50
30	0.04	0.22	0.61	1.47	0.02	0.09	0.19	0.50
40	0.19	0.60	1.03	1.48	0.07	0.17	0.31	0.49
50	0.43	0.91	1.25	1.39	0.11	0.25	0.39	0.44
60	0.55	0.95	1.07	1.11	0.15	0.30	0.35	0.36

Lifetime Risk of Being Diagnosed = 1.38%
 Lifetime Risk of Dying = 0.54%

Lifetime Risk of Being Diagnosed = 0.50%
 Lifetime Risk of Dying = 0.19%

Table XX-6
OVARY CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.05	1.72
10	0.01	0.05	0.12	1.74
20	0.04	0.11	0.27	1.73
30	0.07	0.24	0.54	1.70
40	0.17	0.47	0.88	1.64
50	0.31	0.73	1.16	1.51
60	0.44	0.89	1.18	1.25

Lifetime Risk of Being Diagnosed = 1.72%

Lifetime Risk of Dying = 1.02%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.05	1.85
10	0.01	0.05	0.12	1.87
20	0.04	0.11	0.29	1.86
30	0.07	0.25	0.58	1.83
40	0.18	0.50	0.95	1.77
50	0.33	0.79	1.26	1.62
60	0.48	0.97	1.27	1.34

Lifetime Risk of Being Diagnosed = 1.85%

Lifetime Risk of Dying = 1.07%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.04	1.10
10	0.01	0.04	0.09	1.13
20	0.03	0.08	0.18	1.13
30	0.05	0.15	0.32	1.11
40	0.11	0.28	0.57	1.08
50	0.18	0.49	0.80	1.01
60	0.33	0.67	0.85	0.90

Lifetime Risk of Being Diagnosed = 1.10%

Lifetime Risk of Dying = 0.72%

Table XXI-8
PANCREAS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.23	0.00	0.00	0.00	1.24
10	0.00	0.00	0.01	1.25	0.00	0.00	0.01	1.26
20	0.00	0.01	0.05	1.26	0.00	0.01	0.03	1.26
30	0.01	0.05	0.19	1.27	0.01	0.03	0.13	1.26
40	0.04	0.19	0.51	1.29	0.03	0.12	0.37	1.27
50	0.15	0.49	0.91	1.29	0.10	0.35	0.76	1.26
60	0.36	0.82	1.16	1.22	0.26	0.69	1.09	1.22

Lifetime Risk of Being Diagnosed = 1.23%
 Lifetime Risk of Dying = 1.15%

Lifetime Risk of Being Diagnosed = 1.24%
 Lifetime Risk of Dying = 1.16%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.23	0.00	0.00	0.00	1.22
10	0.00	0.00	0.01	1.25	0.00	0.00	0.01	1.23
20	0.00	0.01	0.05	1.25	0.00	0.01	0.03	1.23
30	0.01	0.05	0.19	1.27	0.01	0.03	0.12	1.24
40	0.04	0.18	0.51	1.28	0.02	0.12	0.36	1.24
50	0.14	0.48	0.91	1.28	0.09	0.34	0.75	1.24
60	0.36	0.82	1.15	1.21	0.26	0.68	1.07	1.19

Lifetime Risk of Being Diagnosed = 1.23%
 Lifetime Risk of Dying = 1.15%

Lifetime Risk of Being Diagnosed = 1.22%
 Lifetime Risk of Dying = 1.15%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.27	0.00	0.00	0.00	1.34
10	0.00	0.00	0.02	1.32	0.00	0.00	0.01	1.38
20	0.00	0.02	0.08	1.33	0.00	0.01	0.05	1.38
30	0.01	0.08	0.29	1.36	0.01	0.05	0.18	1.39
40	0.07	0.29	0.69	1.39	0.04	0.18	0.48	1.41
50	0.23	0.67	1.15	1.42	0.14	0.45	0.93	1.42
60	0.50	1.07	1.30	1.38	0.34	0.86	1.26	1.39

Lifetime Risk of Being Diagnosed = 1.27%
 Lifetime Risk of Dying = 1.14%

Lifetime Risk of Being Diagnosed = 1.34%
 Lifetime Risk of Dying = 1.33%

Table XXII-6
PROSTATE CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	17.28
10	0.00	0.00	0.01	17.58
20	0.00	0.01	0.22	17.68
30	0.01	0.22	2.25	17.89
40	0.22	2.28	8.17	18.17
50	2.13	8.24	15.06	18.61
60	6.71	14.20	17.53	18.11

Lifetime Risk of Being Diagnosed = 17.28%
 Lifetime Risk of Dying = 3.05%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	16.90
10	0.00	0.00	0.01	17.16
20	0.00	0.01	0.20	17.24
30	0.01	0.20	2.17	17.42
40	0.20	2.20	8.00	17.68
50	2.06	8.06	14.71	18.07
60	6.55	13.82	16.95	17.48

Lifetime Risk of Being Diagnosed = 16.90%
 Lifetime Risk of Dying = 2.86%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	20.40
10	0.00	0.00	0.02	21.12
20	0.00	0.02	0.51	21.31
30	0.02	0.52	3.96	21.86
40	0.52	4.07	12.20	22.55
50	3.84	12.64	20.38	23.83
60	10.62	19.97	23.53	24.13

Lifetime Risk of Being Diagnosed = 20.40%
 Lifetime Risk of Dying = 4.72%

Table XXIII-8
STOMACH CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.25	0.00	0.00	0.00	0.79
10	0.00	0.00	0.02	1.27	0.00	0.00	0.01	0.81
20	0.00	0.02	0.06	1.28	0.00	0.01	0.04	0.81
30	0.02	0.06	0.19	1.29	0.01	0.04	0.10	0.81
40	0.04	0.17	0.47	1.30	0.03	0.09	0.23	0.80
50	0.13	0.44	0.88	1.30	0.06	0.21	0.45	0.79
60	0.33	0.81	1.16	1.25	0.15	0.40	0.66	0.76

Lifetime Risk of Being Diagnosed = 1.25%
 Lifetime Risk of Dying = 0.61%

Lifetime Risk of Being Diagnosed = 0.79%
 Lifetime Risk of Dying = 0.41%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.09	0.00	0.00	0.00	0.66
10	0.00	0.00	0.02	1.10	0.00	0.00	0.01	0.66
20	0.00	0.02	0.05	1.11	0.00	0.01	0.04	0.66
30	0.01	0.05	0.16	1.12	0.01	0.03	0.08	0.67
40	0.04	0.15	0.41	1.12	0.03	0.08	0.19	0.66
50	0.12	0.39	0.76	1.12	0.05	0.17	0.36	0.65
60	0.29	0.69	1.00	1.07	0.12	0.33	0.54	0.62

Lifetime Risk of Being Diagnosed = 1.09%
 Lifetime Risk of Dying = 0.56%

Lifetime Risk of Being Diagnosed = 0.66%
 Lifetime Risk of Dying = 0.37%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.32	0.00	0.00	0.00	1.03
10	0.00	0.01	0.03	1.37	0.00	0.00	0.02	1.06
20	0.01	0.03	0.09	1.38	0.00	0.02	0.05	1.07
30	0.02	0.09	0.24	1.41	0.01	0.05	0.13	1.07
40	0.07	0.23	0.56	1.43	0.04	0.12	0.30	1.08
50	0.17	0.53	1.05	1.47	0.09	0.27	0.58	1.08
60	0.42	1.03	1.38	1.51	0.20	0.53	0.89	1.08

Lifetime Risk of Being Diagnosed = 1.32%
 Lifetime Risk of Dying = 0.92%

Lifetime Risk of Being Diagnosed = 1.03%
 Lifetime Risk of Dying = 0.69%

Table XXIV-6
TESTIS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.11	0.35
10	0.02	0.11	0.23	0.36
20	0.10	0.21	0.29	0.34
30	0.12	0.20	0.23	0.25
40	0.08	0.11	0.12	0.13
50	0.03	0.04	0.05	0.05
60	0.01	0.02	0.02	0.02

Lifetime Risk of Being Diagnosed = 0.35%
 Lifetime Risk of Dying = 0.02%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.03	0.14	0.42
10	0.02	0.14	0.27	0.42
20	0.11	0.25	0.35	0.40
30	0.14	0.23	0.27	0.29
40	0.09	0.13	0.14	0.15
50	0.04	0.05	0.06	0.06
60	0.01	0.02	0.03	0.03

Lifetime Risk of Being Diagnosed = 0.42%
 Lifetime Risk of Dying = 0.02%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.03	0.10
10	0.01	0.03	0.06	0.10
20	0.02	0.05	0.07	0.09
30	0.03	0.05	0.07	0.07
40	0.02	0.04	0.04	0.04
50	0.01	0.02	0.02	0.02
60	0.01	0.01	0.01	0.01

Lifetime Risk of Being Diagnosed = 0.10%
 Lifetime Risk of Dying = 0.01%

Table XXV-8
THYROID CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.02	0.30	0.00	0.02	0.10	0.84
10	0.00	0.02	0.05	0.31	0.02	0.10	0.24	0.86
20	0.02	0.05	0.10	0.30	0.09	0.23	0.39	0.84
30	0.03	0.08	0.15	0.29	0.14	0.30	0.46	0.76
40	0.05	0.12	0.19	0.26	0.16	0.32	0.46	0.62
50	0.07	0.14	0.20	0.22	0.16	0.30	0.41	0.47
60	0.08	0.13	0.16	0.16	0.14	0.26	0.31	0.32

Lifetime Risk of Being Diagnosed = 0.30%
 Lifetime Risk of Dying = 0.04%

Lifetime Risk of Being Diagnosed = 0.84%
 Lifetime Risk of Dying = 0.06%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.02	0.32	0.00	0.02	0.12	0.87
10	0.00	0.02	0.06	0.32	0.02	0.12	0.27	0.88
20	0.02	0.05	0.10	0.32	0.09	0.25	0.42	0.86
30	0.03	0.09	0.16	0.30	0.15	0.32	0.48	0.77
40	0.05	0.12	0.20	0.27	0.17	0.33	0.46	0.62
50	0.07	0.15	0.20	0.23	0.16	0.30	0.40	0.46
60	0.08	0.14	0.16	0.16	0.14	0.25	0.30	0.31

Lifetime Risk of Being Diagnosed = 0.32%
 Lifetime Risk of Dying = 0.04%

Lifetime Risk of Being Diagnosed = 0.87%
 Lifetime Risk of Dying = 0.06%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	0.14	0.00	0.01	0.03	0.46
10	0.00	0.01	0.02	0.15	0.01	0.03	0.10	0.47
20	0.01	0.02	0.04	0.15	0.02	0.09	0.17	0.47
30	0.01	0.04	0.07	0.14	0.07	0.14	0.25	0.45
40	0.02	0.06	0.09	0.14	0.08	0.18	0.27	0.39
50	0.04	0.07	0.10	0.12	0.11	0.20	0.27	0.32
60	0.04	0.07	0.09	0.09	0.09	0.17	0.21	0.23

Lifetime Risk of Being Diagnosed = 0.14%
 Lifetime Risk of Dying = 0.02%

Lifetime Risk of Being Diagnosed = 0.46%
 Lifetime Risk of Dying = 0.05%

Table XXVI-8
URINARY BLADDER CANCER (Invasive and In Situ)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	3.52	0.00	0.00	0.00	1.13
10	0.00	0.01	0.02	3.58	0.00	0.00	0.01	1.15
20	0.00	0.02	0.11	3.60	0.00	0.01	0.04	1.15
30	0.02	0.10	0.41	3.64	0.01	0.03	0.13	1.15
40	0.09	0.40	1.21	3.68	0.03	0.12	0.35	1.16
50	0.32	1.16	2.51	3.72	0.10	0.33	0.70	1.15
60	0.90	2.36	3.43	3.66	0.25	0.64	0.99	1.10

Lifetime Risk of Being Diagnosed = 3.52%
 Lifetime Risk of Dying = 0.73%

Lifetime Risk of Being Diagnosed = 1.13%
 Lifetime Risk of Dying = 0.31%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	3.91	0.00	0.00	0.00	1.22
10	0.00	0.01	0.02	3.97	0.00	0.00	0.01	1.23
20	0.01	0.02	0.12	3.99	0.00	0.01	0.04	1.24
30	0.02	0.11	0.46	4.03	0.01	0.04	0.14	1.24
40	0.10	0.45	1.36	4.07	0.03	0.14	0.39	1.24
50	0.36	1.31	2.81	4.10	0.11	0.37	0.77	1.23
60	1.01	2.62	3.77	4.01	0.27	0.69	1.06	1.17

Lifetime Risk of Being Diagnosed = 3.91%
 Lifetime Risk of Dying = 0.78%

Lifetime Risk of Being Diagnosed = 1.22%
 Lifetime Risk of Dying = 0.32%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.42	0.00	0.00	0.00	0.78
10	0.00	0.00	0.01	1.47	0.00	0.00	0.00	0.80
20	0.00	0.01	0.06	1.48	0.00	0.00	0.02	0.80
30	0.01	0.06	0.21	1.52	0.00	0.02	0.09	0.81
40	0.05	0.21	0.59	1.56	0.02	0.09	0.26	0.82
50	0.17	0.57	1.14	1.62	0.08	0.25	0.50	0.83
60	0.47	1.13	1.61	1.68	0.18	0.46	0.74	0.82

Lifetime Risk of Being Diagnosed = 1.42%
 Lifetime Risk of Dying = 0.38%

Lifetime Risk of Being Diagnosed = 0.78%
 Lifetime Risk of Dying = 0.32%