

Table I-17  
US PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2002<sup>a</sup>  
USING DIFFERENT TUMOR INCLUSION CRITERIA<sup>b</sup>

Site/Sex	5-Year Limited Duration			27-year Limited Duration	
	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 27 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 27 Years <sup>d</sup>
All Sites	3,867,003	3,941,332	4,238,091	9,430,525	9,606,482
Male	1,957,153	1,983,555	2,125,418	4,356,626	4,406,570
Female	1,909,850	1,957,777	2,112,673	5,073,899	5,199,912
Oral Cavity & Pharynx	82,465	93,657	96,502	209,742	228,987
Male	54,587	61,351	63,058	134,000	144,924
Female	27,878	32,306	33,444	75,742	84,063
Esophagus	16,212	19,256	19,256	23,139	27,099
Male	12,545	14,664	14,664	17,520	20,164
Female	3,667	4,592	4,592	5,619	6,935
Stomach	29,294	33,978	34,082	56,255	63,243
Male	17,931	20,805	20,838	32,491	36,516
Female	11,363	13,173	13,244	23,764	26,727
Colon & Rectum	409,449	468,805	476,603	990,228	1,095,712
Male	205,302	235,516	238,820	482,617	530,469
Female	204,147	233,289	237,783	507,611	565,243
Liver & Intrahep	11,552	13,206	13,217	15,152	17,046
Male	7,508	8,594	8,605	9,437	10,650
Female	4,044	4,612	4,612	5,715	6,396
Pancreas	19,344	22,700	22,700	25,555	29,428
Male	9,438	11,214	11,214	12,274	14,270
Female	9,906	11,486	11,486	13,281	15,158
Larynx	32,997	37,965	38,182	91,754	100,304
Male	26,181	30,101	30,259	73,344	79,883
Female	6,816	7,864	7,923	18,410	20,421
Lung & Bronchus	202,674	246,254	250,392	335,447	395,971
Male	100,886	123,709	125,599	165,547	194,890
Female	101,788	122,545	124,793	169,900	201,081
Melanoma of the Skin	202,999	223,960	229,859	578,663	615,944
Male	107,851	120,689	124,506	286,244	307,023
Female	95,148	103,271	105,353	292,419	308,921
Breast					
Female	810,067	867,306	903,843	2,147,690	2,277,797
Cervix					
Female	44,613	46,898	46,925	179,134	184,371
Corpus & Uterus					
Female	142,978	160,405	160,468	485,208	522,668
Ovary					
Female	54,659	62,754	62,758	142,293	158,788

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER 9 registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census.

<sup>bcd</sup> (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion: (c) First invasive tumor ever; (d) First invasive tumor for each cancer site diagnosed during the previous 27 years (1975-2001); (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (1997-2001). For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 1997; Melanoma in 1998.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 27-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred 20 years ago. In method (d) the 1981 melanoma is counted for the melanoma and all sites 27-year limited duration prevalence. The 1997 breast cancer is counted for the breast 5-year and 27-year limited duration prevalence. In method (e) the 1997 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 1998 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table I-17 - continued  
US PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2002<sup>a</sup>  
USING DIFFERENT TUMOR INCLUSION CRITERIA<sup>b</sup>

Site/Sex	5-Year Limited Duration			27-year Limited Duration	
	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 27 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 27 Years <sup>d</sup>
Prostate					
Male	892,138	960,511	960,545	1,826,301	1,949,727
Testis					
Male	37,578	38,062	38,571	144,174	145,865
Urinary Bladder					
Male	184,483	221,675	223,454	471,823	535,752
Female	138,422	166,292	167,816	349,628	395,599
Female	46,061	55,383	55,638	122,195	140,153
Kidney & Renal Pelvis					
Male	89,323	107,038	107,849	207,723	237,885
Female	53,720	65,276	65,792	123,059	141,698
Female	35,603	41,762	42,057	84,664	96,187
Brain & Nervous System					
Male	35,503	37,509	37,641	91,881	94,763
Female	19,416	20,452	20,517	49,872	51,377
Female	16,087	17,057	17,124	42,009	43,386
Thyroid					
Male	88,240	94,590	94,806	279,685	292,893
Female	20,450	22,504	22,553	63,721	67,300
Female	67,790	72,086	72,253	215,964	225,593
Hodgkin Lymphoma					
Male	33,241	34,605	34,627	125,113	127,804
Female	17,560	18,312	18,312	65,330	66,779
Female	15,681	16,293	16,315	59,783	61,025
Non-Hodgkin Lymphoma					
Male	154,821	175,583	176,195	332,657	365,300
Female	81,328	92,287	92,613	172,701	188,781
Female	73,493	83,296	83,582	159,956	176,519
Myeloma					
Male	34,833	39,821	39,844	49,931	56,104
Female	18,832	21,683	21,706	27,208	30,707
Female	16,001	18,138	18,138	22,723	25,397
Leukemia					
Male	80,095	89,786	89,845	183,579	198,257
Female	46,350	52,154	52,201	103,787	112,084
Female	33,745	37,632	37,644	79,792	86,173
Acute Lymphocytic Leuk					
Male	13,598	13,731	13,731	44,730	44,955
Female	7,581	7,624	7,624	24,523	24,604
Female	6,017	6,107	6,107	20,207	20,351
Childhood (0-19)					
Male	55,930	56,015	56,307	215,915	216,360
Female	29,328	29,380	29,523	110,475	110,683
Female	26,602	26,635	26,784	105,440	105,677
Kaposi's Sarcoma					
Male	7,140	7,674	7,708	19,711	20,644
Female	6,545	7,024	7,058	18,332	19,151
Female	595	650	650	1,379	1,493
Mesothelioma					
Male	3,065	3,626	3,626	4,422	5,077
Female	2,144	2,643	2,643	2,686	3,256
Female	921	983	983	1,736	1,821

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER 9 registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census.

<sup>bcd</sup> (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion: (c) First invasive tumor ever; (d) First invasive tumor for each cancer site diagnosed during the previous 27 years (1975-2001); (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (1997-2001). For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 1997; Melanoma in 1998.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 27-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred 20 years ago. In method (d) the 1981 melanoma is counted for the melanoma and all sites 27-year limited duration prevalence. The 1997 breast cancer is counted for the breast 5-year and 27-year limited duration prevalence. In method (e) the 1997 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 1998 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table I-18  
US COMPLETE PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2002<sup>a</sup>  
BY AGE AT PREVALENCE

Age at Prevalence	Age Specific								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
<u>Site/Sex</u>									
All Sites									
Males	4,503,895	16,687	35,355	62,064	146,126	300,518	580,447	1,013,496	2,349,201
Females	5,642,429	14,040	30,938	66,090	210,795	556,663	970,592	1,154,712	2,638,599
Breast									
Males	11,780	0	11	0	153	502	1,913	3,066	6,135
Females	2,278,269	0	59	1,906	36,453	198,739	446,463	521,249	1,073,401
Colon & Rectum									
Males	505,267	0	80	644	4,721	19,026	56,080	112,718	311,998
Females	546,415	11	63	704	4,461	17,769	47,993	92,415	382,999
Corpus & Uterus, NOS									
Females	571,854	0	56	388	4,101	22,096	67,003	112,603	365,607
Esophagus									
Males	17,651	0	0	40	189	1,112	3,871	5,437	7,002
Females	5,751	0	11	11	71	208	629	1,349	3,471
Larynx									
Males	78,348	0	11	22	274	2,990	11,043	21,374	42,633
Females	19,555	0	0	33	210	1,335	2,433	5,714	9,829
Lung & Bronchus									
Males	174,384	34	77	269	976	6,943	24,344	50,180	91,561
Females	176,295	11	22	321	1,447	7,463	24,410	48,055	94,565
Melanoma of the Skin									
Males	304,097	78	552	4,111	16,149	45,683	69,092	66,751	101,680
Females	325,725	57	684	8,250	30,125	63,878	73,988	57,468	91,275
Non-Hodgkin Lymphoma									
Males	180,337	614	3,030	5,687	11,579	23,291	34,346	39,245	62,544
Females	166,702	307	1,354	2,960	7,638	16,193	27,250	34,245	76,756
Oral Cavity & Pharynx									
Males	147,272	56	468	1,229	3,579	13,645	30,185	36,114	61,995
Females	84,527	96	458	1,280	3,399	7,667	13,677	18,157	39,792
Ovary									
Females	169,875	56	1,014	2,956	7,766	21,466	34,390	36,707	65,519
Prostate									
Males	1,831,929	34	92	97	230	13,222	143,741	449,469	1,225,043
Stomach									
Males	34,258	3	11	107	471	2,113	4,669	8,130	18,753
Females	25,053	11	23	49	547	1,601	2,540	4,566	15,715
Urinary Bladder									
Males	367,550	28	171	623	2,641	13,157	39,326	82,579	229,024
Females	131,649	22	34	300	1,496	4,445	11,884	25,578	87,888
Myeloma									
Males	27,357	0	5	27	457	1,873	5,778	7,631	11,586
Females	23,127	0	11	33	259	1,184	4,161	5,316	12,162

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER 9 registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table II-12  
ALL SITES (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	3,867,003	2,460,330	1,504,635	872,558	553,193	7,039,730	9,430,525	715,799	10,146,324
Males	1,957,153	1,233,741	639,238	309,137	172,896	3,517,811	4,356,626	147,269	4,503,895
Females	1,909,850	1,226,589	865,397	563,421	380,297	3,521,919	5,073,899	568,530	5,642,429
White <sup>b</sup>	3,370,258	2,174,781	1,364,398	794,145	506,445	6,189,259	8,368,124	635,007	9,003,131
Males	1,693,011	1,082,602	582,459	282,797	159,821	3,072,611	3,841,476	136,108	3,977,584
Females	1,677,247	1,092,179	781,939	511,348	346,624	3,116,648	4,526,648	498,899	5,025,547
Black <sup>b</sup>	349,752	201,255	97,676	53,097	31,475	599,087	742,995	39,343	782,338
Males	190,761	111,002	40,814	17,758	8,479	323,678	371,112	4,700	375,812
Females	158,991	90,253	56,862	35,339	22,996	275,409	371,883	34,643	406,526
Asian/Pacific Islander <sup>c</sup>	85,392	45,835	+	+	+	143,180	+	+	+
Males	37,395	18,926	+	+	+	60,654	+	+	+
Females	47,997	26,909	+	+	+	82,526	+	+	+
Hispanic <sup>d</sup>	185,695	105,516	+	+	+	319,960	+	+	+
Males	90,068	48,833	+	+	+	150,533	+	+	+
Females	95,627	56,683	+	+	+	169,427	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	1.9552%	0.0765%	0.1025%	0.2064%	0.5256%	1.2483%	3.0137%	6.6073%	10.6061%	10.3405%	2.1145%
Males	1.9759%	0.0817%	0.1068%	0.1924%	0.4097%	0.8369%	2.5015%	7.6417%	14.6397%	15.1464%	2.4520%
Females	1.9350%	0.0710%	0.0979%	0.2211%	0.6448%	1.6542%	3.4996%	5.6874%	7.5704%	7.7514%	1.9148%
White <sup>c</sup>	2.1304%	0.0829%	0.1107%	0.2286%	0.5705%	1.3074%	3.1268%	6.9174%	11.0039%	10.5126%	2.1990%
Males	2.1231%	0.0884%	0.1151%	0.2136%	0.4516%	0.8832%	2.5559%	7.8285%	14.8780%	15.1483%	2.5028%
Females	2.1378%	0.0771%	0.1061%	0.2449%	0.6976%	1.7393%	3.6836%	6.0862%	8.0453%	8.0534%	2.0279%
Black <sup>c</sup>	1.4290%	0.0483%	0.0762%	0.1323%	0.3787%	1.0515%	2.9796%	6.9895%	10.6640%	9.0076%	2.0344%
Males	1.6402%	0.0492%	0.0800%	0.1181%	0.2628%	0.7536%	3.1344%	9.9353%	17.5275%	15.5995%	2.8308%
Females	1.2376%	0.0474%	0.0723%	0.1453%	0.4814%	1.3122%	2.8505%	4.6668%	6.1474%	5.9275%	1.5338%
Asian/Pacific Islander <sup>c</sup>	1.2329%	0.0657%	0.0792%	0.1298%	0.3354%	0.9396%	2.0542%	3.8217%	6.7362%	7.8294%	1.4017%
Males	1.1367%	0.0732%	0.0813%	0.1147%	0.2206%	0.4810%	1.3014%	3.8179%	9.1878%	11.1826%	1.4881%
Females	1.3230%	0.0578%	0.0770%	0.1445%	0.4437%	1.3521%	2.7151%	3.8247%	4.9208%	5.5284%	1.3704%
Hispanic <sup>d</sup>	0.7567%	0.0715%	0.0945%	0.1365%	0.3567%	0.8798%	1.9654%	4.3340%	7.5704%	7.5473%	1.4752%
Males	0.6890%	0.0757%	0.0991%	0.1278%	0.2583%	0.4818%	1.4325%	5.0228%	11.0553%	11.4012%	1.7291%
Females	0.8280%	0.0672%	0.0896%	0.1467%	0.4688%	1.2851%	2.4551%	3.7662%	5.0898%	5.4259%	1.3473%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table III-12  
BRAIN AND OTHER NERVOUS SYSTEM (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	35,503	21,327	16,177	9,926	6,728	64,276	91,881	14,079	105,960
Males	19,416	11,972	8,575	5,358	3,396	35,534	49,872	6,993 <sup>i</sup>	56,865
Females	16,087	9,355	7,602	4,568	3,332	28,742	42,009	7,086 <sup>i</sup>	49,095
White <sup>b</sup>	31,555	18,767	14,353	8,873	5,956	56,835	81,518	10,932	92,450
Males	17,290	10,601	7,506	4,757	3,012	31,496	44,225	5,691 <sup>i</sup>	49,916
Females	14,265	8,166	6,847	4,116	2,944	25,339	37,293	5,241 <sup>i</sup>	42,534
Black <sup>b</sup>	2,570	1,659	1,307	781	580	4,893	7,070	+	+
Males	1,460	838	753	454	294	2,687	3,876	+	+
Females	1,110	821	554	327	286	2,206	3,194	+	+
Asian/Pacific Islander <sup>c</sup>	867	505	+	+	+	1,518	+	+	+
Males	466	287	+	+	+	841	+	+	+
Females	401	218	+	+	+	677	+	+	+
Hispanic <sup>d</sup>	3,068	1,527	+	+	+	5,142	+	+	+
Males	1,665	815	+	+	+	2,781	+	+	+
Females	1,403	712	+	+	+	2,361	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0182%	0.0130%	0.0197%	0.0146%	0.0186%	0.0221%	0.0225%	0.0205%	0.0174%	0.0098%	0.0182%
Males	0.0203%	0.0142%	0.0212%	0.0163%	0.0204%	0.0246%	0.0256%	0.0241%	0.0210%	0.0121%	0.0204%
Females	0.0161%	0.0118%	0.0182%	0.0128%	0.0168%	0.0195%	0.0195%	0.0174%	0.0147%	0.0086%	0.0162%
White <sup>c</sup>	0.0202%	0.0145%	0.0213%	0.0159%	0.0213%	0.0252%	0.0253%	0.0229%	0.0190%	0.0093%	0.0203%
Males	0.0225%	0.0157%	0.0232%	0.0175%	0.0227%	0.0279%	0.0286%	0.0267%	0.0222%	0.0122%	0.0225%
Females	0.0180%	0.0132%	0.0194%	0.0142%	0.0197%	0.0225%	0.0221%	0.0194%	0.0165%	0.0078%	0.0181%
Black <sup>c</sup>	0.0114%	0.0092%	0.0150%	0.0099%	0.0099%	0.0104%	0.0144%	0.0111%	0.0128%	0.0114%	0.0114%
Males	0.0128%	0.0108%	0.0151%	0.0120%	0.0116%	0.0130%	0.0160%	0.0118%	0.0170%	-	0.0128%
Females	0.0100%	0.0077%	0.0149%	0.0080%	0.0084%	0.0082%	0.0131%	0.0106%	0.0101%	0.0148%	0.0102%
Asian/Pacific Islander <sup>c</sup>	0.0109%	0.0078%	0.0147%	0.0104%	0.0103%	0.0117%	0.0104%	0.0114%	0.0094%	0.0126%	0.0110%
Males	0.0124%	0.0091%	0.0150%	0.0119%	0.0122%	0.0117%	0.0128%	0.0130%	0.0158%	0.0167%	0.0126%
Females	0.0095%	0.0065%	0.0145%	0.0090%	0.0085%	0.0116%	0.0083%	0.0101%	0.0047%	0.0097%	0.0096%
Hispanic <sup>d</sup>	0.0123%	0.0103%	0.0132%	0.0076%	0.0128%	0.0170%	0.0164%	0.0178%	0.0143%	0.0090%	0.0132%
Males	0.0129%	0.0110%	0.0128%	0.0081%	0.0131%	0.0189%	0.0187%	0.0211%	0.0142%	-	0.0139%
Females	0.0117%	0.0095%	0.0136%	0.0070%	0.0124%	0.0152%	0.0143%	0.0152%	0.0144%	0.0120%	0.0124%

- <sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.
- <sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- <sup>g h i</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table IV-17  
BREAST CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	815,238	557,995	390,390	222,946	129,931	1,548,184	2,159,088	130,961	2,290,049
Males	5,171	2,750	1,674	971	627	8,769	11,398	382	11,780
Females	810,067	555,245	388,716	221,975	129,304	1,539,415	2,147,690	130,579	2,278,269
White <sup>b</sup>	718,104	498,369	355,168	203,033	119,018	1,374,678	1,932,860	116,667	2,049,527
Males	4,430	2,419	1,449	906	578	7,572	9,974	285 <sup>i</sup>	10,259
Females	713,674	495,950	353,719	202,127	118,440	1,367,106	1,922,886	116,382	2,039,268
Black <sup>b</sup>	67,474	42,592	25,613	14,511	7,975	122,140	160,589	+	+
Males	597	236	211	50	49	944	1,145	+	+
Females	66,877	42,356	25,402	14,461	7,926	121,196	159,444	8,928	168,372
Asian/Pacific Islander <sup>c</sup>	20,812	11,916	+	+	+	36,083	+	+	+
Males	54	74	+	+	+	132	+	+	+
Females	20,758	11,842	+	+	+	35,951	+	+	+
Hispanic <sup>d</sup>	36,401	22,488	+	+	+	65,375	+	+	+
Males	130	64	+	+	+	219	+	+	+
Females	36,271	22,424	+	+	+	65,156	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.4288%	-	0.0002%	0.0047%	0.0770%	0.3799%	0.9421%	1.4263%	1.8718%	2.0490%	0.4585%
Males	0.0050%	-	-	-	0.0006%	0.0023%	0.0088%	0.0211%	0.0312%	0.0395%	0.0062%
Females	0.8431%	-	0.0003%	0.0097%	0.1556%	0.7525%	1.8274%	2.6760%	3.2571%	3.1316%	0.8357%
White <sup>c</sup>	0.4714%	-	0.0002%	0.0046%	0.0754%	0.3868%	0.9833%	1.5196%	2.0033%	2.1706%	0.4822%
Males	0.0053%	-	-	-	0.0005%	0.0021%	0.0086%	0.0214%	0.0325%	0.0407%	0.0062%
Females	0.9350%	-	0.0003%	0.0096%	0.1553%	0.7786%	1.9338%	2.8863%	3.5082%	3.3005%	0.8867%
Black <sup>c</sup>	0.2915%	-	-	0.0061%	0.0935%	0.3580%	0.8203%	1.1928%	1.5573%	1.5962%	0.3950%
Males	0.0050%	-	-	-	-	0.0032%	0.0143%	0.0297%	0.0409%	0.0397%	0.0082%
Females	0.5511%	-	-	0.0118%	0.1754%	0.6685%	1.4923%	2.1099%	2.5552%	2.3235%	0.6813%
Asian/Pacific Islander <sup>c</sup>	0.2970%	-	-	0.0046%	0.0701%	0.3552%	0.7509%	0.9892%	1.1138%	1.1045%	0.3244%
Males	0.0026%	-	-	-	-	0.0021%	0.0038%	0.0095%	0.0152%	0.0284%	0.0033%
Females	0.5727%	-	-	0.0091%	0.1359%	0.6728%	1.4067%	1.7984%	1.9273%	1.8429%	0.5884%
Hispanic <sup>d</sup>	0.1555%	-	-	0.0034%	0.0495%	0.2660%	0.6169%	0.9047%	1.2065%	1.3197%	0.3010%
Males	0.0010%	-	-	-	-	0.0013%	0.0031%	0.0086%	0.0112%	0.0181%	0.0025%
Females	0.3182%	-	-	0.0075%	0.1058%	0.5357%	1.1810%	1.6433%	2.0574%	2.0362%	0.5431%

- <sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.
- <sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- <sup>g h i</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table V-10  
CERVIX UTERI CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	44,613	39,155	34,583	26,978	24,062	98,268	179,134	44,307 <sup>i</sup>	223,441
White <sup>b</sup>	34,944	31,782	28,654	22,132	19,696	78,681	145,291	31,811 <sup>i</sup>	177,102
Black <sup>b</sup>	6,795	4,686	4,128	3,623	3,363	13,289	23,999	+	+
Asian/Pacific Islander <sup>c</sup>	1,988	1,700	+	+	+	4,174	+	+	+
Hispanic <sup>d</sup>	7,695	6,102	+	+	+	16,006	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup> All Ages
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	0.0619%	-	0.0002%	0.0115%	0.0768%	0.1255%	0.1206%	0.1107%	0.0936%	0.0607%	0.0616%
White <sup>c</sup>	0.0620%	-	-	0.0127%	0.0852%	0.1304%	0.1153%	0.0985%	0.0792%	0.0472%	0.0611%
Black <sup>c</sup>	0.0577%	-	-	0.0100%	0.0631%	0.1093%	0.1220%	0.1485%	0.1337%	0.1510%	0.0653%
Asian/Pacific Islander <sup>c</sup>	0.0626%	-	-	0.0043%	0.0358%	0.1005%	0.1487%	0.1525%	0.1651%	0.1178%	0.0628%
Hispanic <sup>d</sup>	0.0759%	-	-	0.0119%	0.0882%	0.2152%	0.2253%	0.2178%	0.2271%	0.1216%	0.1071%

- <sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.
- <sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- <sup>g h i</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table VI-18  
COLON AND RECTUM CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	409,449	233,819	164,608	105,937	60,690	717,979	990,228	61,454	1,051,682
Males	205,302	115,014	81,422	49,292	25,427	357,337	482,617	22,650	505,267
Females	204,147	118,805	83,186	56,645	35,263	360,642	507,611	38,804	546,415
White <sup>b</sup>	354,178	205,555	146,219	94,980	54,726	625,996	869,716	55,060	924,776
Males	179,123	101,610	73,028	44,397	23,072	313,727	426,783	20,481	447,264
Females	175,055	103,945	73,191	50,583	31,654	312,269	442,933	34,579	477,512
Black <sup>b</sup>	38,946	19,764	13,345	7,960	4,126	64,804	85,307	4,222	89,529
Males	17,373	8,868	5,736	3,291	1,469	29,037	37,125	1,438	38,563
Females	21,573	10,896	7,609	4,669	2,657	35,767	48,182	2,784	50,966
Asian/Pacific Islander <sup>c</sup>	11,564	5,902	+	+	+	19,126	+	+	+
Males	6,084	3,083	+	+	+	10,002	+	+	+
Females	5,480	2,819	+	+	+	9,124	+	+	+
Hispanic <sup>d</sup>	17,876	8,640	+	+	+	29,325	+	+	+
Males	9,787	4,508	+	+	+	15,707	+	+	+
Females	8,089	4,132	+	+	+	13,618	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.1989%	-	0.0003%	0.0032%	0.0186%	0.0702%	0.2410%	0.6585%	1.2399%	1.6908%	0.2187%
Males	0.2019%	-	0.0003%	0.0032%	0.0192%	0.0735%	0.2748%	0.7965%	1.4955%	1.8970%	0.2560%
Females	0.1960%	-	0.0003%	0.0032%	0.0179%	0.0669%	0.2090%	0.5358%	1.0476%	1.5797%	0.1898%
White <sup>c</sup>	0.2110%	-	0.0004%	0.0030%	0.0181%	0.0663%	0.2321%	0.6572%	1.2564%	1.7179%	0.2186%
Males	0.2131%	-	0.0005%	0.0030%	0.0187%	0.0692%	0.2677%	0.7971%	1.5182%	1.9269%	0.2567%
Females	0.2089%	-	0.0003%	0.0030%	0.0175%	0.0633%	0.1974%	0.5296%	1.0564%	1.6070%	0.1886%
Black <sup>c</sup>	0.1524%	-	-	0.0034%	0.0188%	0.0858%	0.3050%	0.7552%	1.2340%	1.4591%	0.2281%
Males	0.1455%	-	-	0.0032%	0.0176%	0.0892%	0.3357%	0.8635%	1.3518%	1.5542%	0.2509%
Females	0.1587%	-	-	0.0035%	0.0199%	0.0827%	0.2794%	0.6699%	1.1566%	1.4147%	0.2123%
Asian/Pacific Islander <sup>c</sup>	0.1715%	-	-	0.0039%	0.0203%	0.0786%	0.2393%	0.5840%	1.1048%	1.5703%	0.2034%
Males	0.1856%	-	-	0.0037%	0.0227%	0.0867%	0.2643%	0.7168%	1.4009%	1.8144%	0.2437%
Females	0.1583%	-	-	0.0042%	0.0181%	0.0713%	0.2173%	0.4742%	0.8855%	1.4029%	0.1726%
Hispanic <sup>d</sup>	0.0680%	-	0.0004%	0.0025%	0.0130%	0.0490%	0.1720%	0.4679%	0.8932%	1.1132%	0.1542%
Males	0.0704%	-	-	0.0028%	0.0132%	0.0495%	0.1962%	0.6103%	1.1889%	1.2247%	0.1892%
Females	0.0655%	-	-	0.0021%	0.0128%	0.0484%	0.1497%	0.3505%	0.6828%	1.0519%	0.1281%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table VII-10  
CORPUS AND UTERUS, NOS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	142,978	107,011	82,698	66,457	59,049	286,056	485,208	86,646	571,854
White <sup>b</sup>	128,991	98,524	77,506	63,042	56,043	261,377	450,038	77,528	527,566
Black <sup>b</sup>	8,216	4,725	3,129	1,989	1,583	14,288	20,156	1,709	21,865
Asian/Pacific Islander <sup>c</sup>	3,446	2,161	+	+	+	6,165	+	+	+
Hispanic <sup>d</sup>	6,431	4,028	+	+	+	11,849	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	0-9	10-19	20-29	Age Specific (Crude)						Age-Adjusted <sup>f</sup>
					30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.1507%	-	0.0002%	0.0021%	0.0192%	0.0829%	0.2841%	0.5886%	0.6828%	0.5803%	0.1503%
White <sup>c</sup>	0.1726%	-	-	0.0020%	0.0189%	0.0832%	0.3064%	0.6600%	0.7572%	0.6340%	0.1641%
Black <sup>c</sup>	0.0647%	-	-	0.0018%	0.0164%	0.0471%	0.1289%	0.3418%	0.4219%	0.2676%	0.0836%
Asian/Pacific Islander <sup>c</sup>	0.0980%	-	-	0.0031%	0.0198%	0.1085%	0.2661%	0.3171%	0.3214%	0.2518%	0.1003%
Hispanic <sup>d</sup>	0.0566%	-	-	0.0029%	0.0237%	0.0722%	0.2003%	0.3630%	0.3734%	0.3087%	0.0965%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on <sup>b</sup> SEER 9 Areas <sup>c</sup> SEER 11 Areas and Rural Georgia <sup>d</sup> SEER 11 Areas excluding Hawaii and Detroit.  
<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VIII-12  
ESOPHAGUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	16,212	4,334	1,551	714	254	21,431	23,139	263	23,402
Males	12,545	3,143	1,099	476	214	16,288	17,520	131	17,651
Females	3,667	1,191	452	238	40	5,143	5,619	132	5,751
White <sup>b</sup>	14,207	3,742	1,261	547	232	18,662	20,033	221	20,254
Males	11,102	2,706	930	363	209	14,300	15,342	124	15,466
Females	3,105	1,036	331	184	23	4,362	4,691	97	4,788
Black <sup>b</sup>	1,576	512	245	164	14	2,231	2,532	63	2,595
Males	1,098	364	142	113	0	1,553	1,724	21	1,745
Females	478	148	103	51	14	678	808	42	850
Asian/Pacific Islander <sup>c</sup>	271	74	+	+	+	363	+	+	+
Males	208	59	+	+	+	282	+	+	+
Females	63	15	+	+	+	81	+	+	+
Hispanic <sup>d</sup>	583	176	+	+	+	800	+	+	+
Males	457	132	+	+	+	631	+	+	+
Females	126	44	+	+	+	169	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0059%	-	-	0.0001%	0.0005%	0.0025%	0.0112%	0.0253%	0.0344%	0.0256%	0.0065%
Males	0.0092%	-	-	0.0002%	0.0008%	0.0041%	0.0197%	0.0431%	0.0579%	0.0441%	0.0110%
Females	0.0028%	-	-	-	0.0002%	0.0009%	0.0032%	0.0096%	0.0167%	0.0156%	0.0028%
White <sup>c</sup>	0.0065%	-	-	-	0.0006%	0.0026%	0.0116%	0.0265%	0.0359%	0.0267%	0.0067%
Males	0.0100%	-	-	-	0.0009%	0.0043%	0.0208%	0.0449%	0.0599%	0.0465%	0.0115%
Females	0.0030%	-	-	-	0.0003%	0.0008%	0.0027%	0.0097%	0.0176%	0.0161%	0.0028%
Black <sup>c</sup>	0.0052%	-	-	-	-	0.0036%	0.0163%	0.0309%	0.0341%	0.0152%	0.0073%
Males	0.0076%	-	-	-	-	0.0058%	0.0253%	0.0499%	0.0580%	0.0198%	0.0116%
Females	0.0031%	-	-	-	-	0.0017%	0.0088%	0.0160%	0.0184%	0.0130%	0.0040%
Asian/Pacific Islander <sup>c</sup>	0.0033%	-	-	-	-	0.0011%	0.0047%	0.0135%	0.0236%	0.0214%	0.0039%
Males	0.0053%	-	-	-	-	0.0015%	0.0079%	0.0248%	0.0433%	0.0357%	0.0068%
Females	0.0015%	-	-	-	-	-	0.0019%	0.0041%	0.0090%	0.0117%	0.0016%
Hispanic <sup>d</sup>	0.0019%	-	-	-	-	0.0013%	0.0081%	0.0167%	0.0164%	0.0202%	0.0040%
Males	0.0029%	-	-	-	-	0.0021%	0.0138%	0.0283%	0.0309%	0.0291%	0.0069%
Females	0.0009%	-	-	-	-	-	0.0028%	0.0071%	0.0060%	0.0152%	0.0018%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table IX-12  
HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	33,241	28,119	25,148	19,628	14,505	71,954	125,113	20,388	145,501
Males	17,560	14,652	12,893	10,305	7,601	37,885	65,330	10,927	76,257
Females	15,681	13,467	12,255	9,323	6,904	34,069	59,783	9,461	69,244
White <sup>b</sup>	28,702	24,885	23,083	18,140	13,628	63,238	112,661	18,651	131,312
Males	15,097	13,046	11,884	9,405	7,036	33,371	58,587	9,685	68,272
Females	13,605	11,839	11,199	8,735	6,592	29,867	54,074	8,966	63,040
Black <sup>b</sup>	3,520	2,622	1,659	1,232	751	6,942	9,949	1,254	11,203
Males	1,857	1,306	790	765	488	3,533	5,361	873	6,234
Females	1,663	1,316	869	467	263	3,409	4,588	381	4,969
Asian/Pacific Islander <sup>c</sup>	523	298	+	+	+	913	+	+	+
Males	290	130	+	+	+	462	+	+	+
Females	233	168	+	+	+	451	+	+	+
Hispanic <sup>d</sup>	2,828	1,658	+	+	+	5,085	+	+	+
Males	1,625	933	+	+	+	2,885	+	+	+
Females	1,203	725	+	+	+	2,200	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0196%	0.0009%	0.0082%	0.0305%	0.0356%	0.0254%	0.0189%	0.0171%	0.0173%	0.0114%	0.0194%
Males	0.0211%	0.0014%	0.0081%	0.0287%	0.0361%	0.0286%	0.0230%	0.0222%	0.0215%	0.0142%	0.0209%
Females	0.0182%	0.0003%	0.0083%	0.0324%	0.0351%	0.0223%	0.0151%	0.0125%	0.0142%	0.0099%	0.0180%
White <sup>c</sup>	0.0221%	0.0010%	0.0095%	0.0346%	0.0400%	0.0287%	0.0209%	0.0188%	0.0192%	0.0124%	0.0218%
Males	0.0235%	0.0017%	0.0093%	0.0318%	0.0405%	0.0317%	0.0249%	0.0242%	0.0230%	0.0156%	0.0231%
Females	0.0206%	0.0004%	0.0097%	0.0375%	0.0396%	0.0256%	0.0170%	0.0139%	0.0162%	0.0107%	0.0206%
Black <sup>c</sup>	0.0168%	-	0.0051%	0.0280%	0.0327%	0.0218%	0.0190%	0.0155%	0.0133%	-	0.0170%
Males	0.0183%	-	0.0047%	0.0271%	0.0326%	0.0272%	0.0255%	0.0217%	0.0204%	-	0.0194%
Females	0.0155%	-	0.0056%	0.0288%	0.0327%	0.0171%	0.0136%	0.0106%	0.0086%	-	0.0151%
Asian/Pacific Islander <sup>c</sup>	0.0065%	-	0.0036%	0.0098%	0.0113%	0.0063%	0.0052%	0.0064%	0.0067%	0.0072%	0.0062%
Males	0.0069%	-	0.0045%	0.0096%	0.0102%	0.0068%	0.0064%	0.0088%	0.0109%	-	0.0068%
Females	0.0061%	-	0.0027%	0.0100%	0.0124%	0.0059%	0.0041%	0.0045%	-	-	0.0057%
Hispanic <sup>d</sup>	0.0120%	0.0015%	0.0085%	0.0166%	0.0173%	0.0136%	0.0165%	0.0196%	0.0209%	0.0138%	0.0133%
Males	0.0132%	0.0026%	0.0099%	0.0150%	0.0174%	0.0174%	0.0202%	0.0223%	0.0335%	0.0217%	0.0156%
Females	0.0107%	-	0.0069%	0.0185%	0.0171%	0.0097%	0.0130%	0.0174%	0.0119%	-	0.0113%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table X-12  
KAPOSI'S SARCOMA (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	7,140	8,705	2,599	822	288	17,179	19,711	+	+
Males	6,545	8,354	2,400	743	196	16,079	18,332	+	+
Females	595	351	199	79	92	1,100	1,379	+	+
White <sup>b</sup>	4,912	7,144	2,096	671	253	13,124	15,199	+	+
Males	4,484	6,882	1,945	603	161	12,316	14,170	+	+
Females	428	262	151	68	92	808	1,029	+	+
Black <sup>b</sup>	1,751	1,109	247	93	12	2,992	3,205	+	+
Males	1,633	1,035	210	93	12	2,776	2,976	+	+
Females	118	74	37	0	0	216	229	+	+
Asian/Pacific Islander <sup>c</sup>	89	97	+	+	+	192	+	+	+
Males	85	95	+	+	+	185	+	+	+
Females	4	2	+	+	+	7	+	+	+
Hispanic <sup>d</sup>	1,128	1,239	+	+	+	2,512	+	+	+
Males	1,051	1,206	+	+	+	2,387	+	+	+
Females	77	33	+	+	+	125	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0064%	-	-	0.0008%	0.0107%	0.0170%	0.0093%	0.0047%	0.0055%	0.0116%	0.0065%
Males	0.0124%	-	-	0.0014%	0.0206%	0.0337%	0.0186%	0.0089%	0.0106%	0.0209%	0.0126%
Females	0.0006%	-	-	-	0.0005%	0.0005%	0.0005%	0.0009%	0.0017%	0.0065%	0.0006%
White <sup>c</sup>	0.0065%	-	-	0.0005%	0.0105%	0.0168%	0.0095%	0.0048%	0.0058%	0.0119%	0.0064%
Males	0.0123%	-	-	0.0009%	0.0199%	0.0330%	0.0188%	0.0091%	0.0110%	0.0219%	0.0125%
Females	0.0006%	-	-	-	0.0005%	0.0003%	0.0004%	0.0008%	0.0019%	0.0065%	0.0006%
Black <sup>c</sup>	0.0086%	-	-	0.0026%	0.0177%	0.0245%	0.0111%	0.0051%	-	0.0114%	0.0090%
Males	0.0171%	-	-	0.0054%	0.0370%	0.0509%	0.0228%	0.0096%	-	-	0.0181%
Females	0.0008%	-	-	-	-	0.0014%	-	-	-	0.0111%	0.0010%
Asian/Pacific Islander <sup>c</sup>	0.0016%	-	-	-	0.0018%	0.0044%	0.0026%	-	-	0.0048%	0.0016%
Males	0.0031%	-	-	-	0.0037%	0.0092%	0.0051%	-	-	-	0.0032%
Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic <sup>d</sup>	0.0063%	-	-	0.0007%	0.0135%	0.0195%	0.0115%	0.0069%	0.0083%	0.0257%	0.0084%
Males	0.0117%	-	-	0.0014%	0.0246%	0.0382%	0.0231%	0.0129%	0.0187%	0.0362%	0.0158%
Females	0.0006%	-	-	-	0.0008%	-	-	-	-	0.0199%	0.0011%

- <sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.
- <sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- <sup>g h i</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table XI-12  
KIDNEY AND RENAL PELVIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	89,323	53,895	32,405	18,544	10,573	158,462	207,723	13,547	221,270
Males	53,720	32,010	18,695	11,071	5,894	94,880	123,059	7,595	130,654
Females	35,603	21,885	13,710	7,473	4,679	63,582	84,664	5,952	90,616
White <sup>b</sup>	76,701	46,650	28,943	16,667	9,425	136,918	181,029	13,474	194,503
Males	46,557	28,148	16,848	10,100	5,401	82,905	108,541	6,658	115,199
Females	30,144	18,502	12,095	6,567	4,024	54,013	72,488	6,816	79,304
Black <sup>b</sup>	9,741	5,706	2,849	1,546	956	16,798	21,078	2,062	23,140
Males	5,348	2,977	1,511	766	400	9,091	11,153	721	11,874
Females	4,393	2,729	1,338	780	556	7,707	9,925	1,341	11,266
Asian/Pacific Islander <sup>c</sup>	1,619	791	+	+	+	2,574	+	+	+
Males	990	512	+	+	+	1,593	+	+	+
Females	629	279	+	+	+	981	+	+	+
Hispanic <sup>d</sup>	6,039	3,029	+	+	+	9,952	+	+	+
Males	3,510	1,744	+	+	+	5,698	+	+	+
Females	2,529	1,285	+	+	+	4,254	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0434%	0.0074%	0.0040%	0.0015%	0.0074%	0.0291%	0.0825%	0.1611%	0.2064%	0.1824%	0.0466%
Males	0.0530%	0.0067%	0.0036%	0.0013%	0.0085%	0.0363%	0.1106%	0.2169%	0.2845%	0.2736%	0.0625%
Females	0.0340%	0.0082%	0.0043%	0.0018%	0.0062%	0.0220%	0.0558%	0.1115%	0.1477%	0.1333%	0.0339%
White <sup>c</sup>	0.0473%	0.0078%	0.0042%	0.0014%	0.0075%	0.0300%	0.0852%	0.1706%	0.2179%	0.1891%	0.0488%
Males	0.0577%	0.0074%	0.0039%	0.0009%	0.0086%	0.0377%	0.1141%	0.2275%	0.2986%	0.2886%	0.0653%
Females	0.0370%	0.0083%	0.0046%	0.0018%	0.0063%	0.0222%	0.0571%	0.1188%	0.1562%	0.1364%	0.0352%
Black <sup>c</sup>	0.0385%	0.0080%	0.0054%	0.0026%	0.0092%	0.0351%	0.1014%	0.1737%	0.2123%	0.1765%	0.0517%
Males	0.0453%	0.0054%	0.0051%	0.0026%	0.0098%	0.0428%	0.1357%	0.2363%	0.2946%	0.2460%	0.0681%
Females	0.0324%	0.0106%	0.0058%	0.0027%	0.0087%	0.0284%	0.0729%	0.1243%	0.1581%	0.1440%	0.0399%
Asian/Pacific Islander <sup>c</sup>	0.0223%	0.0040%	-	0.0013%	0.0048%	0.0164%	0.0444%	0.0791%	0.1116%	0.1108%	0.0251%
Males	0.0287%	0.0026%	-	0.0020%	0.0063%	0.0221%	0.0609%	0.1128%	0.1644%	0.1470%	0.0347%
Females	0.0164%	0.0055%	-	-	0.0034%	0.0114%	0.0298%	0.0512%	0.0724%	0.0860%	0.0173%
Hispanic <sup>d</sup>	0.0235%	0.0063%	0.0027%	0.0012%	0.0058%	0.0268%	0.0730%	0.1599%	0.2112%	0.1865%	0.0453%
Males	0.0262%	0.0053%	0.0017%	0.0008%	0.0065%	0.0288%	0.0992%	0.2221%	0.2679%	0.2820%	0.0595%
Females	0.0206%	0.0073%	0.0039%	0.0016%	0.0050%	0.0248%	0.0489%	0.1086%	0.1708%	0.1340%	0.0345%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XII-12  
LARYNX CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	32,997	22,778	16,877	10,713	6,720	63,365	91,754	6,149	97,903
Males	26,181	18,369	13,434	8,646	5,353	50,486	73,344	5,004	78,348
Females	6,816	4,409	3,443	2,067	1,367	12,879	18,410	1,145	19,555
White <sup>b</sup>	27,707	19,790	14,673	9,541	6,130	54,084	79,327	5,054	84,381
Males	22,100	15,986	11,762	7,751	4,875	43,201	63,682	4,097	67,779
Females	5,607	3,804	2,911	1,790	1,255	10,883	15,645	957	16,602
Black <sup>b</sup>	4,492	2,562	1,875	1,034	476	7,905	10,602	415	11,017
Males	3,413	2,053	1,440	785	388	6,152	8,217	348	8,565
Females	1,079	509	435	249	88	1,753	2,385	67	2,452
Asian/Pacific Islander <sup>c</sup>	445	275	+	+	+	792	+	+	+
Males	374	235	+	+	+	669	+	+	+
Females	71	40	+	+	+	123	+	+	+
Hispanic <sup>d</sup>	1,433	776	+	+	+	2,541	+	+	+
Males	1,188	656	+	+	+	2,123	+	+	+
Females	245	120	+	+	+	418	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0165%	-	-	0.0001%	0.0009%	0.0069%	0.0294%	0.0763%	0.0985%	0.0644%	0.0181%
Males	0.0267%	-	-	-	0.0010%	0.0101%	0.0497%	0.1313%	0.1846%	0.1457%	0.0327%
Females	0.0065%	-	-	0.0002%	0.0007%	0.0038%	0.0102%	0.0274%	0.0337%	0.0207%	0.0066%
White <sup>c</sup>	0.0178%	-	-	0.0001%	0.0010%	0.0069%	0.0298%	0.0790%	0.1017%	0.0663%	0.0185%
Males	0.0285%	-	-	-	0.0010%	0.0098%	0.0499%	0.1353%	0.1871%	0.1509%	0.0332%
Females	0.0071%	-	-	-	0.0009%	0.0040%	0.0101%	0.0276%	0.0365%	0.0213%	0.0068%
Black <sup>c</sup>	0.0183%	-	-	-	0.0009%	0.0122%	0.0489%	0.1150%	0.1279%	0.0695%	0.0258%
Males	0.0300%	-	-	-	-	0.0191%	0.0853%	0.1984%	0.2677%	0.1587%	0.0483%
Females	0.0077%	-	-	-	-	0.0062%	0.0186%	0.0492%	0.0359%	0.0278%	0.0098%
Asian/Pacific Islander <sup>c</sup>	0.0071%	-	-	-	-	0.0023%	0.0098%	0.0290%	0.0561%	0.0426%	0.0084%
Males	0.0124%	-	-	-	-	0.0039%	0.0181%	0.0512%	0.1144%	0.0928%	0.0164%
Females	0.0021%	-	-	-	-	-	0.0026%	0.0107%	0.0129%	0.0081%	0.0022%
Hispanic <sup>d</sup>	0.0056%	-	-	-	0.0006%	0.0031%	0.0153%	0.0517%	0.0769%	0.0462%	0.0122%
Males	0.0091%	-	-	-	-	0.0047%	0.0281%	0.0981%	0.1514%	0.1156%	0.0240%
Females	0.0020%	-	-	-	0.0008%	0.0014%	0.0035%	0.0135%	0.0238%	-	0.0035%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.  
<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.  
<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.  
<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.  
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
 + Not available.

Table XIII-18  
LEUKEMIA (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	80,095	47,653	27,123	16,503	9,867	140,112	183,579	6,286	189,865
Males	46,350	27,039	15,161	9,198	4,956	80,080	103,787	2,647	106,434
Females	33,745	20,614	11,962	7,305	4,911	60,032	79,792	3,639	83,431
White <sup>b</sup>	71,241	42,767	24,655	15,106	9,042	125,286	164,974	5,451	170,425
Males	41,371	24,359	14,002	8,416	4,606	71,939	93,791	2,275	96,066
Females	29,870	18,408	10,653	6,690	4,436	53,347	71,183	3,176	74,359
Black <sup>b</sup>	5,432	2,862	1,410	803	516	8,923	11,173	632	11,805
Males	2,761	1,445	612	464	208	4,444	5,540	299	5,839
Females	2,671	1,417	798	339	308	4,479	5,633	333	5,966
Asian/Pacific Islander <sup>c</sup>	1,632	1,002	+	+	+	2,929	+	+	+
Males	988	537	+	+	+	1,682	+	+	+
Females	644	465	+	+	+	1,247	+	+	+
Hispanic <sup>d</sup>	6,091	3,663	+	+	+	10,747	+	+	+
Males	3,389	2,129	+	+	+	6,110	+	+	+
Females	2,702	1,534	+	+	+	4,637	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0403%	0.0286%	0.0289%	0.0125%	0.0142%	0.0231%	0.0487%	0.0982%	0.1506%	0.1626%	0.0425%
Males	0.0472%	0.0310%	0.0303%	0.0136%	0.0172%	0.0287%	0.0629%	0.1325%	0.2032%	0.2159%	0.0535%
Females	0.0336%	0.0260%	0.0274%	0.0114%	0.0112%	0.0176%	0.0352%	0.0677%	0.1110%	0.1339%	0.0334%
White <sup>c</sup>	0.0453%	0.0319%	0.0317%	0.0139%	0.0149%	0.0242%	0.0525%	0.1087%	0.1662%	0.1774%	0.0464%
Males	0.0527%	0.0343%	0.0331%	0.0155%	0.0173%	0.0300%	0.0679%	0.1466%	0.2229%	0.2371%	0.0583%
Females	0.0379%	0.0294%	0.0302%	0.0121%	0.0123%	0.0183%	0.0375%	0.0740%	0.1228%	0.1458%	0.0365%
Black <sup>c</sup>	0.0218%	0.0113%	0.0178%	0.0069%	0.0108%	0.0172%	0.0351%	0.0645%	0.0947%	0.0891%	0.0266%
Males	0.0244%	0.0123%	0.0190%	0.0048%	0.0143%	0.0217%	0.0429%	0.0834%	0.1195%	0.0961%	0.0318%
Females	0.0195%	0.0103%	0.0165%	0.0088%	0.0077%	0.0134%	0.0286%	0.0496%	0.0784%	0.0858%	0.0226%
Asian/Pacific Islander <sup>c</sup>	0.0217%	0.0273%	0.0246%	0.0102%	0.0126%	0.0167%	0.0245%	0.0380%	0.0447%	0.0439%	0.0229%
Males	0.0260%	0.0319%	0.0261%	0.0102%	0.0178%	0.0208%	0.0274%	0.0476%	0.0634%	0.0689%	0.0280%
Females	0.0177%	0.0226%	0.0229%	0.0102%	0.0077%	0.0130%	0.0220%	0.0301%	0.0309%	0.0268%	0.0184%
Hispanic <sup>d</sup>	0.0261%	0.0315%	0.0353%	0.0126%	0.0151%	0.0189%	0.0305%	0.0488%	0.0698%	0.0785%	0.0303%
Males	0.0286%	0.0325%	0.0392%	0.0143%	0.0178%	0.0220%	0.0358%	0.0621%	0.0868%	0.0869%	0.0350%
Females	0.0234%	0.0305%	0.0313%	0.0105%	0.0121%	0.0157%	0.0257%	0.0378%	0.0578%	0.0738%	0.0261%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIII-19  
ACUTE LYMPHOCYTIC LEUKEMIA (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	13,598	10,017	8,594	6,594	4,892	27,251	44,730	+	+
Males	7,581	5,647	4,766	3,624	2,423	15,222	24,523	+	+
Females	6,017	4,370	3,828	2,970	2,469	12,029	20,207	+	+
White <sup>b</sup>	11,907	8,636	7,567	5,955	4,457	23,807	39,500	+	+
Males	6,596	4,922	4,283	3,284	2,249	13,350	21,789	+	+
Females	5,311	3,714	3,284	2,671	2,208	10,457	17,711	+	+
Black <sup>b</sup>	913	730	574	329	229	1,859	2,815	+	+
Males	508	372	248	197	86	968	1,439	+	+
Females	405	358	326	132	143	891	1,376	+	+
Asian/Pacific Islander <sup>c</sup>	510	441	+	+	+	1,083	+	+	+
Males	312	241	+	+	+	614	+	+	+
Females	198	200	+	+	+	469	+	+	+
Hispanic <sup>d</sup>	2,997	2,087	+	+	+	5,632	+	+	+
Males	1,647	1,255	+	+	+	3,202	+	+	+
Females	1,350	832	+	+	+	2,430	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0090%	0.0248%	0.0242%	0.0060%	0.0027%	0.0017%	0.0021%	0.0019%	0.0014%	0.0010%	0.0090%
Males	0.0103%	0.0269%	0.0256%	0.0075%	0.0035%	0.0018%	0.0025%	0.0024%	0.0018%	0.0012%	0.0099%
Females	0.0078%	0.0225%	0.0227%	0.0044%	0.0018%	0.0017%	0.0017%	0.0015%	0.0012%	0.0009%	0.0080%
White <sup>c</sup>	0.0099%	0.0280%	0.0271%	0.0071%	0.0029%	0.0019%	0.0022%	0.0020%	0.0015%	0.0010%	0.0101%
Males	0.0112%	0.0301%	0.0287%	0.0090%	0.0037%	0.0019%	0.0026%	0.0027%	0.0017%	-	0.0111%
Females	0.0087%	0.0259%	0.0255%	0.0051%	0.0020%	0.0019%	0.0019%	0.0014%	0.0013%	0.0010%	0.0091%
Black <sup>c</sup>	0.0045%	0.0089%	0.0128%	0.0020%	0.0015%	-	-	-	-	-	0.0039%
Males	0.0052%	0.0097%	0.0137%	-	0.0020%	-	-	-	-	-	0.0043%
Females	0.0039%	0.0080%	0.0120%	0.0024%	-	-	-	-	-	-	0.0036%
Asian/Pacific Islander <sup>c</sup>	0.0073%	0.0222%	0.0194%	0.0036%	0.0025%	0.0016%	0.0023%	0.0020%	-	-	0.0076%
Males	0.0087%	0.0266%	0.0204%	0.0041%	0.0038%	0.0017%	0.0030%	-	-	-	0.0087%
Females	0.0059%	0.0175%	0.0185%	0.0031%	0.0012%	0.0015%	-	0.0030%	-	-	0.0064%
Hispanic <sup>d</sup>	0.0138%	0.0284%	0.0305%	0.0067%	0.0037%	0.0030%	0.0033%	0.0018%	-	-	0.0110%
Males	0.0153%	0.0294%	0.0340%	0.0081%	0.0048%	0.0031%	0.0039%	-	-	-	0.0123%
Females	0.0122%	0.0272%	0.0268%	0.0050%	0.0023%	0.0028%	0.0028%	-	-	-	0.0097%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table XIV-12  
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	11,552	1,743	908	685	221	13,738	15,152	360	15,512
Males	7,508	1,082	401	300	85	8,836	9,437	61	9,498
Females	4,044	661	507	385	136	4,902	5,715	299	6,014
White <sup>b</sup>	8,299	1,281	722	568	150	9,922	11,049	182	11,231
Males	5,252	772	285	240	58	6,196	6,639	87 <sup>i</sup>	6,726
Females	3,047	509	437	328	92	3,726	4,410	95 <sup>i</sup>	4,505
Black <sup>b</sup>	1,550	151	108	73	47	1,743	1,922	+	+
Males	1,083	101	61	37	23	1,203	1,301	+	+
Females	467	50	47	36	24	540	621	+	+
Asian/Pacific Islander <sup>c</sup>	1,466	245	+	+	+	1,756	+	+	+
Males	1,014	157	+	+	+	1,199	+	+	+
Females	452	88	+	+	+	557	+	+	+
Hispanic <sup>d</sup>	1,480	267	+	+	+	1,779	+	+	+
Males	964	166	+	+	+	1,147	+	+	+
Females	516	101	+	+	+	632	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0057%	0.0016%	0.0005%	0.0005%	0.0010%	0.0040%	0.0114%	0.0221%	0.0262%	0.0172%	0.0061%
Males	0.0076%	0.0015%	0.0006%	0.0005%	0.0012%	0.0057%	0.0179%	0.0316%	0.0381%	0.0242%	0.0088%
Females	0.0039%	0.0018%	0.0004%	0.0006%	0.0007%	0.0023%	0.0052%	0.0137%	0.0172%	0.0134%	0.0039%
White <sup>c</sup>	0.0044%	0.0016%	0.0005%	0.0004%	0.0008%	0.0032%	0.0084%	0.0151%	0.0185%	0.0130%	0.0045%
Males	0.0057%	0.0014%	0.0007%	0.0004%	0.0008%	0.0044%	0.0130%	0.0214%	0.0261%	0.0174%	0.0063%
Females	0.0031%	0.0018%	0.0004%	0.0005%	0.0007%	0.0020%	0.0040%	0.0094%	0.0127%	0.0106%	0.0030%
Black <sup>c</sup>	0.0044%	-	-	-	-	0.0035%	0.0146%	0.0208%	0.0255%	0.0079%	0.0058%
Males	0.0065%	-	-	-	-	0.0056%	0.0249%	0.0347%	0.0414%	-	0.0093%
Females	0.0025%	-	-	-	-	0.0017%	0.0060%	0.0098%	0.0150%	-	0.0031%
Asian/Pacific Islander <sup>c</sup>	0.0158%	0.0029%	-	0.0011%	0.0026%	0.0102%	0.0282%	0.0704%	0.0824%	0.0619%	0.0176%
Males	0.0222%	0.0020%	-	-	0.0044%	0.0156%	0.0474%	0.1036%	0.1251%	0.0857%	0.0265%
Females	0.0099%	0.0037%	-	-	-	0.0053%	0.0113%	0.0429%	0.0507%	0.0456%	0.0104%
Hispanic <sup>d</sup>	0.0045%	0.0015%	0.0007%	0.0004%	0.0008%	0.0051%	0.0142%	0.0340%	0.0380%	0.0241%	0.0085%
Males	0.0056%	0.0014%	0.0011%	-	0.0010%	0.0072%	0.0235%	0.0445%	0.0551%	0.0259%	0.0117%
Females	0.0033%	0.0016%	-	-	-	0.0030%	0.0055%	0.0253%	0.0259%	0.0231%	0.0057%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XV-17  
LUNG AND BRONCHUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	202,674	66,449	34,847	19,044	9,992	285,709	335,447	15,232	350,679
Males	100,886	31,832	17,196	9,088	5,225	140,466	165,547	8,837	174,384
Females	101,788	34,617	17,651	9,956	4,767	145,243	169,900	6,395	176,295
White <sup>b</sup>	175,678	58,481	31,066	16,790	8,911	248,962	293,060	7,873	300,933
Males	85,877	27,627	15,105	7,794	4,621	120,327	142,175	3,983	146,158
Females	89,801	30,854	15,961	8,996	4,290	128,635	150,885	3,890	154,775
Black <sup>b</sup>	21,046	6,293	2,977	1,846	823	28,719	33,207	795	34,002
Males	11,548	3,238	1,612	1,039	465	15,490	18,045	458	18,503
Females	9,498	3,055	1,365	807	358	13,229	15,162	337	15,499
Asian/Pacific Islander <sup>c</sup>	4,370	1,128	+	+	+	5,785	+	+	+
Males	2,541	661	+	+	+	3,384	+	+	+
Females	1,829	467	+	+	+	2,401	+	+	+
Hispanic <sup>d</sup>	5,121	1,563	+	+	+	7,119	+	+	+
Males	2,639	797	+	+	+	3,629	+	+	+
Females	2,482	766	+	+	+	3,490	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0788%	-	0.0002%	0.0011%	0.0039%	0.0254%	0.1142%	0.3380%	0.5493%	0.3730%	0.0870%
Males	0.0796%	-	0.0003%	0.0010%	0.0035%	0.0250%	0.1195%	0.3709%	0.6454%	0.4743%	0.0987%
Females	0.0781%	-	0.0012%	0.0012%	0.0044%	0.0258%	0.1091%	0.3086%	0.4770%	0.3184%	0.0784%
White <sup>c</sup>	0.0849%	-	0.0002%	0.0012%	0.0040%	0.0249%	0.1132%	0.3509%	0.5701%	0.3766%	0.0892%
Males	0.0827%	-	0.0004%	0.0010%	0.0034%	0.0237%	0.1126%	0.3718%	0.6482%	0.4690%	0.0978%
Females	0.0871%	-	-	0.0014%	0.0047%	0.0260%	0.1138%	0.3318%	0.5104%	0.3276%	0.0830%
Black <sup>c</sup>	0.0681%	-	-	0.0009%	0.0044%	0.0393%	0.1649%	0.4007%	0.5499%	0.3391%	0.0987%
Males	0.0772%	-	-	-	0.0040%	0.0425%	0.2121%	0.5055%	0.7107%	0.5122%	0.1271%
Females	0.0598%	-	-	-	0.0046%	0.0365%	0.1255%	0.3181%	0.4441%	0.2582%	0.0789%
Asian/Pacific Islander <sup>c</sup>	0.0539%	-	-	-	0.0032%	0.0176%	0.0792%	0.2116%	0.4017%	0.3550%	0.0632%
Males	0.0651%	-	-	-	0.0039%	0.0202%	0.0939%	0.2675%	0.5873%	0.4924%	0.0850%
Females	0.0433%	-	-	-	0.0025%	0.0153%	0.0663%	0.1654%	0.2643%	0.2607%	0.0466%
Hispanic <sup>d</sup>	0.0171%	-	-	0.0008%	0.0021%	0.0115%	0.0364%	0.1341%	0.2587%	0.2180%	0.0388%
Males	0.0168%	-	-	0.0007%	0.0021%	0.0115%	0.0406%	0.1435%	0.3305%	0.2732%	0.0463%
Females	0.0174%	-	-	0.0009%	0.0020%	0.0115%	0.0326%	0.1264%	0.2077%	0.1877%	0.0337%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XVI-12  
MELANOMA OF THE SKIN (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	202,999	140,468	102,284	72,580	47,934	388,400	578,663	51,159	629,822
Males	107,851	71,956	49,016	32,680	19,675	202,121	286,244	17,853	304,097
Females	95,148	68,512	53,268	39,900	28,259	186,279	292,419	33,306	325,725
White <sup>b</sup>	195,440	134,001	97,729	68,677	46,323	372,536	553,759	48,322	602,081
Males	103,825	68,668	47,201	30,945	18,921	194,063	274,314	16,994	291,308
Females	91,615	65,333	50,528	37,732	27,402	178,473	279,445	31,328	310,773
Black <sup>b</sup>	821	518	446	223	37	1,523	2,154	149	2,303
Males	387	279	188	49	11	759	937	61	998
Females	434	239	258	174	26	764	1,217	88	1,305
Asian/Pacific Islander <sup>c</sup>	578	297	+	+	+	937	+	+	+
Males	261	150	+	+	+	425	+	+	+
Females	317	147	+	+	+	512	+	+	+
Hispanic <sup>d</sup>	3,296	1,906	+	+	+	5,862	+	+	+
Males	1,310	705	+	+	+	2,277	+	+	+
Females	1,986	1,201	+	+	+	3,585	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.1013%	0.0003%	0.0023%	0.0237%	0.0688%	0.1337%	0.2001%	0.2640%	0.3297%	0.3086%	0.1058%
Males	0.1080%	0.0003%	0.0019%	0.0161%	0.0500%	0.1253%	0.2258%	0.3470%	0.4650%	0.4857%	0.1236%
Females	0.0947%	0.0003%	0.0027%	0.0317%	0.0881%	0.1419%	0.1757%	0.1901%	0.2278%	0.2132%	0.0937%
White <sup>c</sup>	0.1249%	0.0003%	0.0028%	0.0290%	0.0848%	0.1632%	0.2398%	0.3168%	0.3851%	0.3473%	0.1264%
Males	0.1322%	0.0004%	0.0022%	0.0190%	0.0599%	0.1506%	0.2668%	0.4115%	0.5397%	0.5558%	0.1454%
Females	0.1176%	0.0003%	0.0034%	0.0399%	0.1114%	0.1761%	0.2135%	0.2305%	0.2670%	0.2366%	0.1140%
Black <sup>c</sup>	0.0037%	-	-	0.0011%	0.0025%	0.0035%	0.0079%	0.0173%	0.0169%	0.0168%	0.0049%
Males	0.0039%	-	-	-	0.0014%	0.0042%	0.0082%	0.0249%	0.0221%	0.0198%	0.0059%
Females	0.0035%	-	-	-	0.0035%	0.0028%	0.0076%	0.0113%	0.0135%	0.0154%	0.0042%
Asian/Pacific Islander <sup>c</sup>	0.0078%	-	-	0.0024%	0.0056%	0.0075%	0.0151%	0.0204%	0.0282%	0.0321%	0.0084%
Males	0.0077%	-	-	0.0029%	0.0029%	0.0060%	0.0145%	0.0246%	0.0423%	0.0389%	0.0091%
Females	0.0078%	-	-	0.0020%	0.0082%	0.0089%	0.0157%	0.0169%	0.0177%	0.0274%	0.0079%
Hispanic <sup>d</sup>	0.0137%	-	-	0.0031%	0.0125%	0.0260%	0.0452%	0.0509%	0.0872%	0.0958%	0.0236%
Males	0.0101%	-	-	0.0019%	0.0050%	0.0181%	0.0358%	0.0520%	0.1085%	0.1234%	0.0222%
Females	0.0175%	-	-	0.0045%	0.0211%	0.0340%	0.0539%	0.0501%	0.0720%	0.0806%	0.0258%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XVII-10  
MESOTHELIOMA (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	3,065	641	267	213	175	3,825	4,422	+	+
Males	2,144	286	73	117	59	2,465	2,686	+	+
Females	921	355	194	96	116	1,360	1,736	+	+
White <sup>b</sup>	2,868	561	261	193	152	3,545	4,088	+	+
Males	2,017	249	70	97	47	2,301	2,480	+	+
Females	851	312	191	96	105	1,244	1,608	+	+
Black <sup>b</sup>	106	63	0	17	24	169	215	+	+
Males	58	37	0	17	13	95	130	+	+
Females	48	26	0	0	11	74	85	+	+
Asian/Pacific Islander <sup>c</sup>	26	9	+	+	+	36	+	+	+
Males	18	5	+	+	+	23	+	+	+
Females	8	4	+	+	+	13	+	+	+
Hispanic <sup>d</sup>	192	19	+	+	+	217	+	+	+
Males	150	5	+	+	+	155	+	+	+
Females	42	14	+	+	+	62	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0010%	-	-	-	0.0002%	0.0005%	0.0017%	0.0033%	0.0067%	0.0060%	0.0011%
Males	0.0014%	-	-	-	-	0.0005%	0.0021%	0.0045%	0.0128%	0.0119%	0.0018%
Females	0.0007%	-	-	-	0.0003%	0.0005%	0.0013%	0.0022%	0.0022%	0.0028%	0.0007%
White <sup>c</sup>	0.0012%	-	-	-	0.0002%	0.0006%	0.0018%	0.0039%	0.0076%	0.0064%	0.0013%
Males	0.0017%	-	-	-	-	0.0005%	0.0023%	0.0056%	0.0139%	0.0132%	0.0020%
Females	0.0008%	-	-	-	0.0004%	0.0006%	0.0014%	0.0024%	0.0027%	0.0028%	0.0008%
Black <sup>c</sup>	0.0005%	-	-	-	-	-	0.0012%	0.0021%	0.0034%	-	0.0007%
Males	0.0006%	-	-	-	-	-	-	-	0.0085%	-	0.0011%
Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0004%
Asian/Pacific Islander <sup>c</sup>	0.0003%	-	-	-	-	-	-	-	0.0030%	-	0.0004%
Males	0.0005%	-	-	-	-	-	-	-	0.0070%	-	0.0007%
Females	0.0002%	-	-	-	-	-	-	-	-	-	0.0002%
Hispanic <sup>d</sup>	0.0005%	-	-	-	-	0.0006%	0.0013%	0.0028%	0.0070%	0.0088%	0.0012%
Males	0.0008%	-	-	-	-	-	0.0020%	0.0063%	0.0126%	-	0.0021%
Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0006%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.  
<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.  
<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.  
<sup>g h i</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.  
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
 + Not available.

Table XVIII-12  
MYELOMA (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	34,833	9,855	3,462	1,168	488	46,736	49,931	553	50,484
Males	18,832	5,529	1,951	650	207	25,527	27,208	149	27,357
Females	16,001	4,326	1,511	518	281	21,209	22,723	404	23,127
White <sup>b</sup>	27,527	7,684	2,587	834	354	36,762	39,056	149	39,205
Males	15,422	4,405	1,492	444	151	20,714	21,926	0 <sup>i</sup>	21,926
Females	12,105	3,279	1,095	390	203	16,048	17,130	149 <sup>i</sup>	17,279
Black <sup>b</sup>	6,158	1,824	778	271	117	8,409	9,187	+	+
Males	2,797	923	432	173	52	3,962	4,388	+	+
Females	3,361	901	346	98	65	4,447	4,799	+	+
Asian/Pacific Islander <sup>c</sup>	660	190	+	+	+	882	+	+	+
Males	332	114	+	+	+	459	+	+	+
Females	328	76	+	+	+	423	+	+	+
Hispanic <sup>d</sup>	1,959	552	+	+	+	2,622	+	+	+
Males	1,077	333	+	+	+	1,481	+	+	+
Females	882	219	+	+	+	1,141	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0139%	-	-	0.0002%	0.0015%	0.0067%	0.0257%	0.0519%	0.0797%	0.0747%	0.0152%
Males	0.0153%	-	-	0.0002%	0.0021%	0.0082%	0.0308%	0.0641%	0.0986%	0.0889%	0.0185%
Females	0.0126%	-	-	0.0002%	0.0010%	0.0053%	0.0208%	0.0411%	0.0655%	0.0671%	0.0124%
White <sup>c</sup>	0.0138%	-	-	0.0002%	0.0016%	0.0064%	0.0237%	0.0492%	0.0753%	0.0694%	0.0142%
Males	0.0155%	-	-	-	0.0024%	0.0079%	0.0297%	0.0612%	0.0958%	0.0853%	0.0179%
Females	0.0120%	-	-	-	0.0008%	0.0048%	0.0178%	0.0382%	0.0597%	0.0609%	0.0112%
Black <sup>c</sup>	0.0203%	-	-	-	0.0022%	0.0116%	0.0527%	0.0954%	0.1682%	0.1410%	0.0297%
Males	0.0201%	-	-	-	-	0.0136%	0.0559%	0.1219%	0.1909%	0.1230%	0.0330%
Females	0.0205%	-	-	-	0.0031%	0.0098%	0.0499%	0.0745%	0.1533%	0.1493%	0.0272%
Asian/Pacific Islander <sup>c</sup>	0.0082%	-	-	-	0.0006%	0.0046%	0.0147%	0.0297%	0.0468%	0.0610%	0.0094%
Males	0.0088%	-	-	-	0.0013%	0.0049%	0.0172%	0.0337%	0.0530%	0.0779%	0.0112%
Females	0.0076%	-	-	-	-	0.0043%	0.0126%	0.0264%	0.0422%	0.0494%	0.0081%
Hispanic <sup>d</sup>	0.0065%	-	-	-	0.0019%	0.0068%	0.0212%	0.0426%	0.0769%	0.0592%	0.0135%
Males	0.0070%	-	-	-	0.0033%	0.0076%	0.0262%	0.0442%	0.0948%	0.1033%	0.0171%
Females	0.0059%	-	-	-	-	0.0060%	0.0166%	0.0413%	0.0641%	0.0349%	0.0109%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIX-16  
NON-HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	154,821	85,244	47,766	26,952	14,294	262,289	332,657	14,382	347,039
Males	81,328	44,005	24,757	13,815	7,109	136,827	172,701	7,636	180,337
Females	73,493	41,239	23,009	13,137	7,185	125,462	159,956	6,746	166,702
White <sup>b</sup>	138,315	76,197	43,410	24,625	13,166	234,688	299,055	14,461	313,516
Males	72,757	39,068	22,584	12,599	6,587	122,246	155,210	8,088	163,298
Females	65,558	37,129	20,826	12,026	6,579	112,442	143,845	6,373	150,218
Black <sup>b</sup>	10,554	5,631	2,955	1,554	808	17,628	21,686	1,032	22,718
Males	5,479	3,112	1,453	821	334	9,311	11,262	470	11,732
Females	5,075	2,519	1,502	733	474	8,317	10,424	562	10,986
Asian/Pacific Islander <sup>c</sup>	3,478	1,791	+	+	+	5,722	+	+	+
Males	1,832	897	+	+	+	2,977	+	+	+
Females	1,646	894	+	+	+	2,745	+	+	+
Hispanic <sup>d</sup>	8,241	4,261	+	+	+	13,531	+	+	+
Males	4,400	2,247	+	+	+	7,149	+	+	+
Females	3,841	2,014	+	+	+	6,382	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0749%	0.0022%	0.0084%	0.0140%	0.0293%	0.0626%	0.1243%	0.2312%	0.3411%	0.3502%	0.0800%
Males	0.0795%	0.0030%	0.0112%	0.0169%	0.0342%	0.0750%	0.1439%	0.2624%	0.3842%	0.3897%	0.0918%
Females	0.0705%	0.0013%	0.0055%	0.0109%	0.0244%	0.0503%	0.1057%	0.2035%	0.3086%	0.3289%	0.0696%
White <sup>c</sup>	0.0831%	0.0023%	0.0086%	0.0144%	0.0307%	0.0659%	0.1323%	0.2521%	0.3649%	0.3682%	0.0852%
Males	0.0874%	0.0031%	0.0116%	0.0172%	0.0357%	0.0789%	0.1522%	0.2833%	0.4110%	0.4093%	0.0974%
Females	0.0788%	0.0015%	0.0054%	0.0113%	0.0254%	0.0527%	0.1129%	0.2237%	0.3296%	0.3464%	0.0743%
Black <sup>c</sup>	0.0425%	0.0014%	0.0063%	0.0136%	0.0285%	0.0571%	0.1038%	0.1395%	0.1735%	0.1623%	0.0536%
Males	0.0478%	0.0021%	0.0088%	0.0166%	0.0333%	0.0702%	0.1318%	0.1653%	0.1880%	0.1961%	0.0640%
Females	0.0376%	-	0.0037%	0.0109%	0.0242%	0.0456%	0.0804%	0.1192%	0.1640%	0.1465%	0.0452%
Asian/Pacific Islander <sup>c</sup>	0.0484%	0.0023%	0.0100%	0.0115%	0.0195%	0.0392%	0.0733%	0.1412%	0.2451%	0.2634%	0.0541%
Males	0.0511%	0.0036%	0.0111%	0.0150%	0.0226%	0.0450%	0.0809%	0.1603%	0.2682%	0.2961%	0.0609%
Females	0.0459%	-	0.0088%	0.0080%	0.0165%	0.0340%	0.0667%	0.1254%	0.2280%	0.2410%	0.0483%
Hispanic <sup>d</sup>	0.0327%	0.0020%	0.0067%	0.0088%	0.0212%	0.0457%	0.0848%	0.1699%	0.2637%	0.2537%	0.0589%
Males	0.0336%	0.0030%	0.0090%	0.0102%	0.0252%	0.0516%	0.0960%	0.1825%	0.2939%	0.2198%	0.0637%
Females	0.0317%	0.0009%	0.0043%	0.0071%	0.0167%	0.0396%	0.0746%	0.1595%	0.2423%	0.2724%	0.0540%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XX-12  
 ORAL CAVITY AND PHARYNX CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	82,465	51,737	33,393	22,747	14,821	149,578	209,742	22,057	231,799
Males	54,587	33,348	20,442	13,862	8,841	97,579	134,000	13,272	147,272
Females	27,878	18,389	12,951	8,885	5,980	51,999	75,742	8,785	84,527
White <sup>b</sup>	71,152	45,325	29,834	20,540	13,178	130,229	184,232	18,813	203,045
Males	47,417	29,496	18,557	12,706	8,083	85,704	119,000	11,440	130,440
Females	23,735	15,829	11,277	7,834	5,095	44,525	65,232	7,373	72,605
Black <sup>b</sup>	7,268	3,919	2,152	1,392	1,019	12,179	15,988	1,037	17,025
Males	4,690	2,319	1,103	720	412	7,518	9,340	429	9,769
Females	2,578	1,600	1,049	672	607	4,661	6,648	608	7,256
Asian/Pacific Islander <sup>c</sup>	2,689	1,501	+	+	+	4,625	+	+	+
Males	1,675	928	+	+	+	2,877	+	+	+
Females	1,014	573	+	+	+	1,748	+	+	+
Hispanic <sup>d</sup>	3,350	1,862	+	+	+	5,669	+	+	+
Males	2,203	1,091	+	+	+	3,545	+	+	+
Females	1,147	771	+	+	+	2,124	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0424%	0.0003%	0.0017%	0.0046%	0.0120%	0.0358%	0.0908%	0.1511%	0.1782%	0.1583%	0.0451%
Males	0.0562%	0.0002%	0.0016%	0.0046%	0.0128%	0.0491%	0.1349%	0.2246%	0.2602%	0.2364%	0.0650%
Females	0.0288%	0.0004%	0.0017%	0.0046%	0.0112%	0.0226%	0.0489%	0.0857%	0.1165%	0.1163%	0.0285%
White <sup>c</sup>	0.0449%	0.0003%	0.0017%	0.0042%	0.0109%	0.0333%	0.0914%	0.1593%	0.1876%	0.1652%	0.0460%
Males	0.0597%	-	0.0013%	0.0041%	0.0119%	0.0467%	0.1360%	0.2356%	0.2725%	0.2511%	0.0665%
Females	0.0301%	0.0003%	0.0022%	0.0044%	0.0100%	0.0196%	0.0479%	0.0896%	0.1227%	0.1196%	0.0286%
Black <sup>c</sup>	0.0296%	-	0.0021%	0.0048%	0.0098%	0.0373%	0.0913%	0.1328%	0.1237%	0.0768%	0.0381%
Males	0.0396%	-	0.0036%	0.0051%	0.0088%	0.0469%	0.1457%	0.2059%	0.1874%	0.1033%	0.0558%
Females	0.0205%	-	-	0.0044%	0.0107%	0.0289%	0.0459%	0.0753%	0.0818%	0.0644%	0.0245%
Asian/Pacific Islander <sup>c</sup>	0.0374%	-	0.0008%	0.0061%	0.0180%	0.0480%	0.0811%	0.1044%	0.1402%	0.1302%	0.0404%
Males	0.0479%	-	-	0.0073%	0.0194%	0.0648%	0.1122%	0.1527%	0.2031%	0.1623%	0.0552%
Females	0.0276%	-	-	0.0050%	0.0166%	0.0330%	0.0537%	0.0645%	0.0937%	0.1082%	0.0282%
Hispanic <sup>d</sup>	0.0135%	-	0.0012%	0.0024%	0.0061%	0.0175%	0.0458%	0.0841%	0.1142%	0.0986%	0.0256%
Males	0.0164%	-	-	0.0028%	0.0058%	0.0220%	0.0640%	0.1288%	0.1766%	0.1199%	0.0359%
Females	0.0104%	-	0.0019%	0.0019%	0.0065%	0.0128%	0.0291%	0.0473%	0.0698%	0.0869%	0.0173%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on <sup>b</sup> SEER 9 Areas <sup>c</sup> SEER 11 Areas and Rural Georgia <sup>d</sup> SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXI-10  
OVARY CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
 By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	54,659	28,651	21,938	18,448	13,976	92,773	142,293	27,582	169,875
White <sup>b</sup>	48,713	25,490	20,168	16,592	12,544	82,993	127,668	24,101	151,769
Black <sup>b</sup>	3,644	1,942	1,108	1,251	1,000	5,937	9,296	2,334	11,630
Asian/Pacific Islander <sup>c</sup>	1,622	854	+	+	+	2,737	+	+	+
Hispanic <sup>d</sup>	3,201	1,748	+	+	+	5,487	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>												
All Races <sup>c</sup>	0.0529%	0.0004%	0.0050%	0.0120%	0.0203%	0.0538%	0.1158%	0.1613%	0.1770%	0.1184%	0.0528%	
White <sup>c</sup>	0.0586%	0.0005%	0.0049%	0.0122%	0.0206%	0.0572%	0.1251%	0.1763%	0.1928%	0.1249%	0.0567%	
Black <sup>c</sup>	0.0280%	-	0.0045%	0.0092%	0.0168%	0.0338%	0.0640%	0.0981%	0.1073%	0.0890%	0.0332%	
Asian/Pacific Islander <sup>c</sup>	0.0418%	-	0.0064%	0.0138%	0.0216%	0.0497%	0.0976%	0.1134%	0.1132%	0.0727%	0.0419%	
Hispanic <sup>d</sup>	0.0270%	0.0006%	0.0055%	0.0116%	0.0160%	0.0418%	0.0854%	0.1173%	0.1452%	0.0953%	0.0413%	

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on <sup>b</sup> SEER 9 Areas <sup>c</sup> SEER 11 Areas and Rural Georgia <sup>d</sup> SEER 11 Areas excluding Hawaii and Detroit.  
<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table XXII-12  
PANCREAS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	19,344	3,085	1,717	801	459	23,263	25,555	524	26,079
Males	9,438	1,458	766	336	191	11,294	12,274	217	12,491
Females	9,906	1,627	951	465	268	11,969	13,281	307	13,588
White <sup>b</sup>	16,499	2,496	1,353	555	389	19,740	21,415	428	21,843
Males	8,296	1,174	628	251	177	9,838	10,593	193	10,786
Females	8,203	1,322	725	304	212	9,902	10,822	235	11,057
Black <sup>b</sup>	1,964	402	303	172	67	2,434	2,934	213	3,147
Males	779	172	131	51	11	976	1,161	84	1,245
Females	1,185	230	172	121	56	1,458	1,773	129	1,902
Asian/Pacific Islander <sup>c</sup>	580	81	+	+	+	684	+	+	+
Males	241	37	+	+	+	287	+	+	+
Females	339	44	+	+	+	397	+	+	+
Hispanic <sup>d</sup>	879	266	+	+	+	1,175	+	+	+
Males	406	120	+	+	+	547	+	+	+
Females	473	146	+	+	+	628	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0070%	-	0.0002%	0.0003%	0.0011%	0.0042%	0.0118%	0.0255%	0.0397%	0.0369%	0.0077%
Males	0.0068%	-	-	-	0.0010%	0.0045%	0.0132%	0.0279%	0.0412%	0.0413%	0.0082%
Females	0.0073%	-	0.0003%	0.0005%	0.0013%	0.0039%	0.0105%	0.0233%	0.0386%	0.0346%	0.0072%
White <sup>c</sup>	0.0073%	-	0.0002%	0.0004%	0.0011%	0.0040%	0.0119%	0.0251%	0.0398%	0.0353%	0.0076%
Males	0.0073%	-	-	-	0.0010%	0.0047%	0.0133%	0.0286%	0.0423%	0.0409%	0.0084%
Females	0.0073%	-	0.0002%	0.0006%	0.0012%	0.0034%	0.0105%	0.0220%	0.0378%	0.0324%	0.0069%
Black <sup>c</sup>	0.0062%	-	-	-	0.0015%	0.0055%	0.0138%	0.0324%	0.0383%	0.0393%	0.0087%
Males	0.0051%	-	-	-	0.0020%	0.0049%	0.0148%	0.0302%	0.0290%	0.0278%	0.0077%
Females	0.0071%	-	-	-	-	0.0060%	0.0130%	0.0341%	0.0444%	0.0447%	0.0092%
Asian/Pacific Islander <sup>c</sup>	0.0064%	-	-	-	0.0009%	0.0039%	0.0090%	0.0226%	0.0396%	0.0503%	0.0075%
Males	0.0056%	-	-	-	-	0.0024%	0.0110%	0.0217%	0.0383%	0.0522%	0.0073%
Females	0.0072%	-	-	-	0.0015%	0.0052%	0.0072%	0.0233%	0.0406%	0.0490%	0.0077%
Hispanic <sup>d</sup>	0.0030%	-	-	0.0003%	0.0007%	0.0027%	0.0069%	0.0191%	0.0406%	0.0339%	0.0064%
Males	0.0026%	-	-	-	-	0.0024%	0.0074%	0.0202%	0.0488%	0.0299%	0.0066%
Females	0.0033%	-	-	0.0007%	0.0009%	0.0030%	0.0066%	0.0182%	0.0348%	0.0361%	0.0061%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on <sup>b</sup> SEER 9 Areas <sup>c</sup> SEER 11 Areas and Rural Georgia <sup>d</sup> SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXIII-10  
PROSTATE CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	892,138	620,579	237,807	54,128	18,778	1,657,793	1,826,301	5,628	1,831,929
White <sup>b</sup>	747,669	531,325	215,173	49,002	17,015	1,410,102	1,562,680	5,936	1,568,616
Black <sup>b</sup>	112,639	72,176	18,784	4,448	1,575	196,364	209,959	488	210,447
Asian/Pacific Islander <sup>c</sup>	13,698	7,345	+	+	+	22,302	+	+	+
Hispanic <sup>d</sup>	38,785	21,955	+	+	+	64,707	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.9191%	-	-	0.0002%	0.0008%	0.0547%	0.8044%	4.0236%	8.7542%	8.6619%	1.1987%
White <sup>c</sup>	0.9602%	-	-	0.0003%	0.0006%	0.0483%	0.7831%	4.0093%	8.7100%	8.3757%	1.1813%
Black <sup>c</sup>	0.9919%	-	-	-	0.0020%	0.1391%	1.5347%	6.7798%	13.0251%	11.4908%	1.8469%
Asian/Pacific Islander <sup>c</sup>	0.4562%	-	-	-	-	0.0139%	0.2549%	1.5471%	4.9142%	6.2639%	0.6446%
Hispanic <sup>d</sup>	0.2955%	-	-	-	-	0.0227%	0.4460%	2.7384%	7.1107%	7.2745%	0.9243%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXIV-12  
STOMACH CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	29,294	11,591	7,306	4,872	2,565	44,043	56,255	3,056	59,311
Males	17,931	6,643	3,532	2,634	1,379	25,993	32,491	1,767	34,258
Females	11,363	4,948	3,774	2,238	1,186	18,050	23,764	1,289	25,053
White <sup>b</sup>	22,398	8,790	5,374	3,777	2,128	33,423	42,983	2,473	45,456
Males	14,067	5,010	2,595	2,093	1,172	20,059	25,238	1,366	26,604
Females	8,331	3,780	2,779	1,684	956	13,364	17,745	1,107	18,852
Black <sup>b</sup>	4,291	1,649	1,131	735	226	6,471	8,109	408	8,517
Males	2,352	970	573	364	123	3,592	4,431	239	4,670
Females	1,939	679	558	371	103	2,879	3,678	169	3,847
Asian/Pacific Islander <sup>c</sup>	2,313	1,018	+	+	+	3,647	+	+	+
Males	1,285	596	+	+	+	2,054	+	+	+
Females	1,028	422	+	+	+	1,593	+	+	+
Hispanic <sup>d</sup>	3,175	1,173	+	+	+	4,639	+	+	+
Males	1,876	565	+	+	+	2,573	+	+	+
Females	1,299	608	+	+	+	2,066	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0158%	-	0.0001%	0.0004%	0.0027%	0.0083%	0.0212%	0.0500%	0.0964%	0.1133%	0.0172%
Males	0.0187%	-	-	0.0005%	0.0027%	0.0094%	0.0279%	0.0701%	0.1343%	0.1554%	0.0234%
Females	0.0129%	-	-	0.0003%	0.0027%	0.0072%	0.0148%	0.0321%	0.0679%	0.0906%	0.0126%
White <sup>c</sup>	0.0133%	-	0.0001%	0.0002%	0.0024%	0.0068%	0.0171%	0.0417%	0.0733%	0.0887%	0.0137%
Males	0.0160%	-	-	0.0003%	0.0024%	0.0076%	0.0231%	0.0603%	0.1040%	0.1245%	0.0189%
Females	0.0105%	-	-	-	0.0023%	0.0060%	0.0113%	0.0247%	0.0499%	0.0697%	0.0096%
Black <sup>c</sup>	0.0155%	-	-	0.0011%	0.0024%	0.0131%	0.0288%	0.0641%	0.1378%	0.1292%	0.0229%
Males	0.0178%	-	-	-	0.0027%	0.0161%	0.0380%	0.0887%	0.1958%	0.1280%	0.0300%
Females	0.0134%	-	-	-	0.0021%	0.0104%	0.0212%	0.0446%	0.0997%	0.1298%	0.0178%
Asian/Pacific Islander <sup>c</sup>	0.0331%	-	-	0.0008%	0.0049%	0.0134%	0.0419%	0.0939%	0.2359%	0.3335%	0.0396%
Males	0.0385%	-	-	-	0.0045%	0.0151%	0.0541%	0.1241%	0.3204%	0.4163%	0.0513%
Females	0.0280%	-	-	-	0.0053%	0.0119%	0.0311%	0.0689%	0.1733%	0.2768%	0.0307%
Hispanic <sup>d</sup>	0.0112%	-	-	0.0004%	0.0049%	0.0112%	0.0317%	0.0635%	0.1244%	0.1717%	0.0240%
Males	0.0121%	-	-	0.0006%	0.0049%	0.0114%	0.0361%	0.0872%	0.1710%	0.2330%	0.0313%
Females	0.0103%	-	-	-	0.0049%	0.0110%	0.0276%	0.0441%	0.0912%	0.1379%	0.0189%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXV-10  
TESTIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	37,578	32,462	29,752	22,954	16,896	82,309	144,174	19,835	164,009
White <sup>b</sup>	35,220	30,808	28,513	22,189	16,326	77,733	137,335	19,072	156,407
Black <sup>b</sup>	1,138	649	438	361	259	1,951	2,967	192	3,159
Asian/Pacific Islander <sup>c</sup>	610	468	+	+	+	1,261	+	+	+
Hispanic <sup>d</sup>	3,411	2,416	+	+	+	6,739	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0471%	0.0021%	0.0042%	0.0512%	0.1084%	0.0935%	0.0446%	0.0169%	0.0077%	0.0059%	0.0454%
White <sup>c</sup>	0.0565%	0.0020%	0.0048%	0.0626%	0.1310%	0.1113%	0.0526%	0.0197%	0.0090%	0.0067%	0.0543%
Black <sup>c</sup>	0.0106%	-	0.0031%	0.0086%	0.0248%	0.0240%	0.0097%	-	-	-	0.0106%
Asian/Pacific Islander <sup>c</sup>	0.0173%	0.0036%	0.0016%	0.0183%	0.0370%	0.0328%	0.0162%	0.0059%	-	-	0.0163%
Hispanic <sup>d</sup>	0.0305%	0.0024%	0.0041%	0.0448%	0.0678%	0.0531%	0.0219%	0.0100%	-	-	0.0288%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on <sup>b</sup> SEER 9 Areas <sup>c</sup> SEER 11 Areas and Rural Georgia <sup>d</sup> SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVI-12  
 THYROID CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	88,240	64,212	47,035	37,303	31,523	172,922	279,685	47,718	327,403
Males	20,450	15,144	10,315	8,315	6,831	40,148	63,721	11,264	74,985
Females	67,790	49,068	36,720	28,988	24,692	132,774	215,964	36,454	252,418
White <sup>b</sup>	77,398	56,657	42,256	33,568	28,399	152,440	248,569	43,144	291,713
Males	18,433	13,747	9,572	7,494	6,283	36,396	58,010	9,572	67,582
Females	58,965	42,910	32,684	26,074	22,116	116,044	190,559	33,572	224,131
Black <sup>b</sup>	5,217	3,670	2,179	1,773	1,532	9,923	14,933	1,979	16,912
Males	902	727	373	370	194	1,818	2,648	212	2,860
Females	4,315	2,943	1,806	1,403	1,338	8,105	12,285	1,767	14,052
Asian/Pacific Islander <sup>c</sup>	3,906	2,703	+	+	+	7,382	+	+	+
Males	754	472	+	+	+	1,351	+	+	+
Females	3,152	2,231	+	+	+	6,031	+	+	+
Hispanic <sup>d</sup>	7,297	5,077	+	+	+	13,737	+	+	+
Males	1,341	960	+	+	+	2,564	+	+	+
Females	5,956	4,117	+	+	+	11,173	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0514%	0.0001%	0.0035%	0.0279%	0.0656%	0.0890%	0.0971%	0.0934%	0.0784%	0.0553%	0.0519%
Males	0.0237%	-	0.0014%	0.0080%	0.0232%	0.0367%	0.0529%	0.0591%	0.0554%	0.0405%	0.0252%
Females	0.0784%	-	0.0057%	0.0489%	0.1093%	0.1406%	0.1391%	0.1240%	0.0956%	0.0632%	0.0778%
White <sup>c</sup>	0.0548%	0.0001%	0.0038%	0.0311%	0.0713%	0.0955%	0.1001%	0.0950%	0.0772%	0.0529%	0.0546%
Males	0.0261%	-	0.0015%	0.0090%	0.0253%	0.0403%	0.0563%	0.0617%	0.0571%	0.0394%	0.0269%
Females	0.0835%	-	0.0062%	0.0550%	0.1205%	0.1517%	0.1429%	0.1254%	0.0925%	0.0601%	0.0823%
Black <sup>c</sup>	0.0233%	-	0.0013%	0.0090%	0.0263%	0.0404%	0.0595%	0.0587%	0.0591%	0.0341%	0.0271%
Males	0.0096%	-	-	0.0022%	0.0084%	0.0139%	0.0290%	0.0316%	0.0460%	0.0198%	0.0127%
Females	0.0356%	-	0.0019%	0.0153%	0.0423%	0.0635%	0.0849%	0.0801%	0.0677%	0.0408%	0.0391%
Asian/Pacific Islander <sup>c</sup>	0.0554%	-	0.0041%	0.0263%	0.0646%	0.0893%	0.1062%	0.1057%	0.0965%	0.0946%	0.0559%
Males	0.0216%	-	-	0.0067%	0.0208%	0.0325%	0.0461%	0.0593%	0.0477%	0.0617%	0.0234%
Females	0.0871%	-	0.0078%	0.0456%	0.1060%	0.1403%	0.1589%	0.1441%	0.1326%	0.1171%	0.0844%
Hispanic <sup>d</sup>	0.0332%	-	0.0031%	0.0199%	0.0466%	0.0735%	0.0913%	0.0870%	0.0849%	0.0693%	0.0456%
Males	0.0118%	-	0.0010%	0.0053%	0.0134%	0.0235%	0.0382%	0.0505%	0.0530%	0.0398%	0.0192%
Females	0.0557%	-	0.0054%	0.0372%	0.0844%	0.1244%	0.1401%	0.1172%	0.1077%	0.0855%	0.0712%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVII-12  
URINARY BLADDER CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	184,483	122,644	79,318	48,761	29,222	342,732	471,823	27,376	499,199
Males	138,422	91,050	59,229	35,167	20,705	255,871	349,628	17,922	367,550
Females	46,061	31,594	20,089	13,594	8,517	86,861	122,195	9,454	131,649
White <sup>b</sup>	172,521	115,474	75,754	46,434	27,816	321,782	444,983	21,738	466,721
Males	130,313	85,835	56,672	33,456	19,791	241,279	330,853	13,986	344,839
Females	42,208	29,639	19,082	12,978	8,025	80,503	114,130	7,752	121,882
Black <sup>b</sup>	7,282	4,453	2,406	1,453	851	12,963	16,661	450	17,111
Males	4,533	3,155	1,668	1,075	560	8,502	11,147	357	11,504
Females	2,749	1,298	738	378	291	4,461	5,514	93	5,607
Asian/Pacific Islander <sup>c</sup>	2,241	1,364	+	+	+	3,946	+	+	+
Males	1,620	1,050	+	+	+	2,919	+	+	+
Females	621	314	+	+	+	1,027	+	+	+
Hispanic <sup>d</sup>	4,836	3,097	+	+	+	8,760	+	+	+
Males	3,653	2,339	+	+	+	6,600	+	+	+
Females	1,183	758	+	+	+	2,160	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2002, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0897%	0.0002%	0.0002%	0.0016%	0.0062%	0.0266%	0.0991%	0.3107%	0.6121%	0.7181%	0.0991%
Males	0.1366%	0.0002%	0.0003%	0.0021%	0.0083%	0.0410%	0.1577%	0.5145%	1.0893%	1.4417%	0.1758%
Females	0.0439%	-	-	0.0011%	0.0040%	0.0124%	0.0436%	0.1295%	0.2530%	0.3282%	0.0428%
White <sup>c</sup>	0.1068%	0.0002%	0.0002%	0.0020%	0.0071%	0.0303%	0.1135%	0.3581%	0.6879%	0.7743%	0.1112%
Males	0.1620%	0.0002%	0.0003%	0.0027%	0.0092%	0.0460%	0.1783%	0.5865%	1.2217%	1.5817%	0.1968%
Females	0.0520%	-	-	0.0013%	0.0048%	0.0143%	0.0503%	0.1498%	0.2803%	0.3459%	0.0476%
Black <sup>c</sup>	0.0301%	-	-	-	0.0036%	0.0144%	0.0525%	0.1427%	0.2537%	0.3490%	0.0460%
Males	0.0425%	-	-	-	0.0060%	0.0230%	0.0859%	0.2354%	0.4261%	0.5770%	0.0765%
Females	0.0189%	-	-	-	0.0015%	0.0068%	0.0246%	0.0696%	0.1403%	0.2424%	0.0257%
Asian/Pacific Islander <sup>c</sup>	0.0366%	-	-	-	0.0025%	0.0105%	0.0361%	0.1116%	0.2801%	0.4141%	0.0446%
Males	0.0559%	-	-	-	0.0036%	0.0173%	0.0593%	0.1903%	0.4778%	0.7221%	0.0762%
Females	0.0186%	-	-	-	0.0015%	0.0043%	0.0158%	0.0466%	0.1336%	0.2028%	0.0208%
Hispanic <sup>d</sup>	0.0202%	-	-	0.0007%	0.0035%	0.0109%	0.0417%	0.1412%	0.2810%	0.3988%	0.0476%
Males	0.0295%	-	-	0.0007%	0.0039%	0.0153%	0.0656%	0.2502%	0.5216%	0.7709%	0.0862%
Females	0.0104%	-	-	-	0.0032%	0.0064%	0.0197%	0.0514%	0.1098%	0.1940%	0.0206%

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVIII-12  
 CHILDHOOD CANCER (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2002  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <12 <sup>e</sup>	0 to <27 <sup>e</sup>	>=27 <sup>f</sup>	Complete <sup>g</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	55,930	46,645	39,770	34,275	29,318	119,335	215,915	+	+
Males	29,328	24,549	20,805	17,308	13,920	62,959	110,475	+	+
Females	26,602	22,096	18,965	16,967	15,398	56,376	105,440	+	+
White <sup>b</sup>	47,142	38,768	33,956	30,073	25,780	100,231	184,521	+	+
Males	24,620	20,501	17,881	15,311	12,424	53,000	94,774	+	+
Females	22,522	18,267	16,075	14,762	13,356	47,231	89,747	+	+
Black <sup>b</sup>	5,776	5,116	3,932	2,820	2,544	12,511	21,023	+	+
Males	3,047	2,566	1,944	1,331	1,028	6,408	10,321	+	+
Females	2,729	2,550	1,988	1,489	1,516	6,103	10,702	+	+
Asian/Pacific Islander <sup>c</sup>	1,846	1,614	+	+	+	3,973	+	+	+
Males	1,000	822	+	+	+	2,085	+	+	+
Females	846	792	+	+	+	1,888	+	+	+
Hispanic <sup>d</sup>	8,371	6,138	+	+	+	16,498	+	+	+
Males	4,428	3,347	+	+	+	8,911	+	+	+
Females	3,943	2,791	+	+	+	7,587	+	+	+

<sup>a</sup> US 2002 cancer prevalence counts are based on 2002 cancer prevalence proportions from the SEER registries and 1/1/2002 US population estimates based on the average of 2001 and 2002 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on b) SEER 9 Areas c) SEER 11 Areas and Rural Georgia d) SEER 11 Areas excluding Hawaii and Detroit.

<sup>e</sup> Maximum limited-duration prevalence: 27 years for 1975-2002 SEER 9 data; 12 years for 1990-2002 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f g</sup> (f) Cases diagnosed more than 27 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <27 and >=27.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.