

Table II-16  
ALL SITES (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2003  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <13 <sup>e</sup>	0 to <28 <sup>e</sup>	>=28 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	3,962,252	2,511,031	1,605,669	924,744	564,549	7,559,653	9,822,273	673,712	10,495,985
Males	2,012,274	1,244,498	717,207	332,858	179,942	3,770,369	4,554,140	138,242	4,692,382
Females	1,949,978	1,266,533	888,462	591,886	384,607	3,789,284	5,268,133	535,470	5,803,603
White <sup>b</sup>	3,442,564	2,213,728	1,450,941	842,073	516,600	6,635,621	8,698,301	596,996	9,295,297
Males	1,736,715	1,087,998	650,645	304,446	166,160	3,288,860	4,008,076	127,969	4,136,045
Females	1,705,849	1,125,730	800,296	537,627	350,440	3,346,761	4,690,225	469,027	5,159,252
Black <sup>b</sup>	361,605	208,112	107,160	55,249	32,825	643,940	779,028	37,221	816,249
Males	196,730	114,039	47,712	18,925	9,172	346,414	389,648	4,256	393,904
Females	164,875	94,073	59,448	36,324	23,653	297,526	389,380	32,965	422,345
Asian/Pacific Islander <sup>c</sup>	89,637	48,904	+	+	+	157,208	+	+	+
Males	39,159	20,104	+	+	+	66,122	+	+	+
Females	50,478	28,800	+	+	+	91,086	+	+	+
Hispanic <sup>d</sup>	197,915	110,621	+	+	+	354,861	+	+	+
Males	96,036	52,006	+	+	+	167,076	+	+	+
Females	101,879	58,615	+	+	+	187,785	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2003, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	1.9814%	0.0768%	0.1062%	0.2104%	0.5283%	1.2572%	3.0482%	6.6866%	10.6455%	10.0716%	2.1199%
Males	1.9956%	0.0805%	0.1135%	0.1943%	0.4104%	0.8423%	2.5543%	7.7010%	14.6841%	14.4892%	2.4448%
Females	1.9676%	0.0729%	0.0985%	0.2273%	0.6497%	1.6678%	3.5165%	5.7819%	7.6049%	7.6709%	1.9260%
White <sup>c</sup>	2.1570%	0.0830%	0.1160%	0.2329%	0.5747%	1.3185%	3.1557%	6.9973%	11.0680%	10.2418%	2.2061%
Males	2.1418%	0.0868%	0.1238%	0.2155%	0.4544%	0.8883%	2.6042%	7.8891%	14.9232%	14.4624%	2.4951%
Females	2.1721%	0.0790%	0.1078%	0.2517%	0.7036%	1.7581%	3.6939%	6.1814%	8.1122%	7.9806%	2.0416%
Black <sup>c</sup>	1.4627%	0.0494%	0.0718%	0.1362%	0.3876%	1.0606%	3.0732%	7.0856%	10.6798%	8.7713%	2.0473%
Males	1.6751%	0.0478%	0.0767%	0.1203%	0.2747%	0.7684%	3.2879%	10.0411%	17.5713%	14.6300%	2.8295%
Females	1.2704%	0.0512%	0.0668%	0.1510%	0.4875%	1.3167%	2.8952%	4.7686%	6.1511%	5.9829%	1.5501%
Asian/Pacific Islander <sup>c</sup>	1.2518%	0.0626%	0.0804%	0.1312%	0.3326%	0.9321%	2.0757%	3.8570%	6.7205%	7.6464%	1.3969%
Males	1.1453%	0.0694%	0.0867%	0.1168%	0.2069%	0.4809%	1.2944%	3.8179%	9.2331%	10.9872%	1.4822%
Females	1.3516%	0.0555%	0.0736%	0.1455%	0.4515%	1.3396%	2.7596%	3.8896%	4.9006%	5.3938%	1.3718%
Hispanic <sup>d</sup>	0.7866%	0.0716%	0.0954%	0.1401%	0.3580%	0.8862%	2.0074%	4.3784%	7.5976%	7.2520%	1.4755%
Males	0.7218%	0.0740%	0.1034%	0.1292%	0.2591%	0.4908%	1.4776%	5.0627%	11.1673%	11.1041%	1.7353%
Females	0.8547%	0.0692%	0.0869%	0.1529%	0.4705%	1.2922%	2.4962%	3.8135%	5.0495%	5.0865%	1.3425%

- <sup>a</sup> US 2003 cancer prevalence counts are based on 2003 cancer prevalence proportions from the SEER registries and 1/1/2003 US population estimates based on the average of 2002 and 2003 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia excluding Hawaii and Seattle.
- <sup>e</sup> Maximum limited-duration prevalence: 28 years for 1975-2003 SEER 9 data; 13 years for 1990-2003 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- <sup>g h</sup> (g) Cases diagnosed more than 28 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <28 and >=28.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.