

Table X-9
KAPOSI SARCOMA (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002
Relative Survival Rates (SEER ^b)																				
1-year	93.5	91.9	85.1	100.0	79.3	80.8	66.9	77.6	89.9	74.7	70.9	82.3	88.0	79.5	70.6	86.8	73.7	79.3	87.0	90.7
2-year	89.7	86.7	85.1	77.7	79.3	80.8	54.2	50.4	78.9	53.5	70.9	56.5	86.1	79.5	65.0	86.8	67.2	79.3	68.1	
3-year	89.7	76.8	85.1	69.8	79.3	78.6	40.7	50.4	74.4	43.2	50.7	48.3	73.8	66.9	65.0	86.8	62.6	69.2		
4-year	83.8	73.7	84.9	69.8	79.3	75.5	40.7	43.4	74.4	31.9	50.7	43.6	68.5	66.9	64.5	86.8	62.6			
5-year	81.7	68.4	84.9	69.8	79.3	69.0	37.4	40.1	73.4	31.9	50.7	43.6	49.4	57.6	50.0	86.8				
6-year	81.7	64.3	84.9	69.8	79.3	69.0	29.5	34.6	73.4	26.9	50.7	40.5	49.4	51.4	50.0					
7-year	81.7	64.3	84.9	46.3	79.3	65.0	29.5	34.6	73.4	26.9	44.4	35.2	49.4	51.4						
8-year	81.7	58.1	84.9	46.3	66.0	57.3	29.5	33.3	73.4	22.0	44.4	35.2	49.4							
9-year	81.7	55.9	84.9	46.3	53.2	57.3	29.5	33.3	73.4	22.0	40.4	30.7								
11-year	81.7	50.6	84.9	46.3	19.1	52.4	29.5	27.0	73.4	22.0	40.4									
10-year	81.7	48.4	82.1	41.7	0.0	52.4	25.6	19.7	73.4	22.0										
12-year	81.7	48.4	61.9	41.7	0.0	44.2	25.6	19.7	73.4											
13-year	81.7	48.4	50.2	41.7	0.0	44.2	14.3	19.7												
14-year	81.7	48.4	19.2	41.7	0.0	34.2	14.3													
15-year	81.7	48.4	19.2	41.7	0.0	17.5														
16-year	81.7	48.4	19.2	41.7	0.0															
17-year	81.7	39.6	0.0	41.7																
18-year	81.7	22.6	0.0																	
19-year	81.7	22.6																		
20-year	81.7																			

^a Survival rates are relative rates expressed as percents.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).