

Table XVII-11
MESOTHELIOMA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2003
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <13 ^e	0 to <28 ^e	>=28 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	2,802	598	315	192	151	3,619	4,182	+	+
Males	1,903	269	87	96	72	2,232	2,438	+	+
Females	899	329	228	96	79	1,387	1,744	+	+
White ^b	2,609	523	279	167	138	3,319	3,828	+	+
Males	1,770	233	70	71	59	2,051	2,214	+	+
Females	839	290	209	96	79	1,268	1,614	+	+
Black ^b	116	61	26	22	13	202	249	+	+
Males	81	36	13	22	13	130	165	+	+
Females	35	25	13	0	0	72	84	+	+
Asian/Pacific Islander ^c	19	9	+	+	+	32	+	+	+
Males	14	7	+	+	+	21	+	+	+
Females	5	2	+	+	+	11	+	+	+
Hispanic ^d	213	20	+	+	+	243	+	+	+
Males	167	10	+	+	+	177	+	+	+
Females	46	10	+	+	+	66	+	+	+

Estimated prevalence percent^a on January 1, 2003, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0010%	-	-	-	0.0001%	0.0006%	0.0017%	0.0036%	0.0050%	0.0063%	0.0011%
Males	0.0014%	-	-	-	-	0.0007%	0.0018%	0.0050%	0.0098%	0.0127%	0.0017%
Females	0.0007%	-	-	-	0.0002%	0.0006%	0.0015%	0.0023%	0.0014%	0.0028%	0.0007%
White ^c	0.0012%	-	-	-	0.0002%	0.0007%	0.0018%	0.0041%	0.0059%	0.0068%	0.0012%
Males	0.0016%	-	-	-	-	0.0007%	0.0020%	0.0058%	0.0113%	0.0136%	0.0019%
Females	0.0008%	-	-	-	0.0003%	0.0007%	0.0017%	0.0026%	0.0017%	0.0031%	0.0008%
Black ^c	0.0005%	-	-	-	-	-	0.0011%	0.0021%	-	-	0.0006%
Males	0.0007%	-	-	-	-	-	-	-	-	-	0.0012%
Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0003%
Asian/Pacific Islander ^c	0.0003%	-	-	-	-	-	-	-	-	-	0.0003%
Males	0.0004%	-	-	-	-	-	-	-	-	-	0.0005%
Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic ^d	0.0006%	-	-	-	-	0.0008%	0.0016%	0.0034%	0.0067%	0.0077%	0.0012%
Males	0.0009%	-	-	-	-	-	0.0025%	0.0060%	0.0135%	-	0.0022%
Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0005%

^a US 2003 cancer prevalence counts are based on 2003 cancer prevalence proportions from the SEER registries and 1/1/2003 US population estimates based on the average of 2002 and 2003 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia excluding Hawaii and Seattle.

^e Maximum limited-duration prevalence: 28 years for 1975-2003 SEER 9 data; 13 years for 1990-2003 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 28 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <28 and >=28. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.