

Table I-21  
US PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2004<sup>a</sup>  
USING DIFFERENT TUMOR INCLUSION CRITERIA<sup>b</sup>

Site/Sex	5-Year Limited Duration			29-year Limited Duration	
	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 29 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 29 Years <sup>d</sup>
All Sites	4,000,969	4,077,054	4,404,865	10,133,875	10,325,806
Male	2,045,700	2,073,736	2,232,399	4,719,465	4,775,763
Female	1,955,269	2,003,318	2,172,466	5,414,410	5,550,043
Oral Cavity & Pharynx	84,371	96,606	99,624	217,906	238,970
Male	56,621	64,295	65,983	140,286	152,405
Female	27,750	32,311	33,641	77,620	86,565
Esophagus	17,196	20,284	20,296	24,565	28,664
Male	13,312	15,498	15,510	18,653	21,473
Female	3,884	4,786	4,786	5,912	7,191
Stomach	29,647	35,113	35,259	57,826	65,836
Male	17,872	21,205	21,244	33,280	37,946
Female	11,775	13,908	14,015	24,546	27,890
Colon & Rectum	413,172	475,209	483,080	1,026,682	1,141,407
Male	208,481	240,519	244,040	503,545	556,381
Female	204,691	234,690	239,040	523,137	585,026
Liver & Intrahep	14,336	16,338	16,338	19,061	21,427
Male	9,723	11,009	11,009	12,344	13,826
Female	4,613	5,329	5,329	6,717	7,601
Pancreas	21,231	25,021	25,021	27,987	32,353
Male	10,156	12,184	12,184	13,235	15,435
Female	11,075	12,837	12,837	14,752	16,918
Larynx	31,966	36,772	37,057	90,186	98,875
Male	25,659	29,412	29,650	72,417	79,020
Female	6,307	7,360	7,407	17,769	19,855
Lung & Bronchus	206,480	254,806	259,381	345,236	412,115
Male	101,519	126,137	128,221	167,413	199,420
Female	104,961	128,669	131,160	177,823	212,695
Melanoma of the Skin	214,760	238,166	246,012	642,642	686,025
Male	113,145	127,550	132,610	317,120	341,495
Female	101,615	110,616	113,402	325,522	344,530
Breast					
Female	813,699	871,649	910,855	2,299,047	2,438,866
Cervix					
Female	41,612	43,636	43,691	186,488	191,924
Corpus & Uterus					
Female	144,321	161,682	161,696	500,188	540,981
Ovary <sup>f</sup>					
Female	54,533	63,083	63,116	148,412	166,361

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER 9 registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census.

<sup>bcd</sup> (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion: (c) First invasive tumor ever; (d) First invasive tumor for each cancer site diagnosed during the previous 29 years (1975-2003); (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (1999-2003). For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 1999; Melanoma in 2000.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 29-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2004.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 29-year limited duration prevalence. The 1999 breast cancer is counted for the breast 5-year and 29-year limited duration prevalence.

In method (e) the 1999 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2000 melanoma is counted for 5-year limited duration prevalence for melanoma.

<sup>f</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table I-21 - continued  
US PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2004<sup>a</sup>  
USING DIFFERENT TUMOR INCLUSION CRITERIA<sup>b</sup>

Site/Sex	5-Year Limited Duration			29-year Limited Duration	
	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 29 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 29 Years <sup>d</sup>
Prostate					
Male	942,369	1,015,465	1,015,501	2,020,336	2,155,814
Testis					
Male	38,541	39,085	39,519	158,516	160,522
Urinary Bladder					
Male	185,870	226,468	229,260	489,252	560,822
Female	139,001	169,421	171,644	362,809	414,423
Female	46,869	57,047	57,616	126,443	146,399
Kidney & Renal Pelvis					
Male	98,835	119,954	120,885	228,714	264,104
Female	59,318	72,826	73,444	135,506	157,273
Female	39,517	47,128	47,441	93,208	106,831
Brain & Nervous System					
Male	38,349	40,291	40,487	100,542	103,478
Female	20,364	21,337	21,448	53,857	55,282
Female	17,985	18,954	19,039	46,685	48,196
Thyroid					
Male	103,273	111,414	111,642	321,227	337,609
Female	23,545	26,152	26,179	72,214	76,826
Female	79,728	85,262	85,463	249,013	260,783
Hodgkin Lymphoma					
Male	33,763	35,330	35,354	135,338	138,313
Female	17,936	18,787	18,798	70,330	71,915
Female	15,827	16,543	16,556	65,008	66,398
Non-Hodgkin Lymphoma					
Male	165,166	188,724	189,721	368,283	405,953
Female	86,010	98,988	99,481	190,649	209,512
Female	79,156	89,736	90,240	177,634	196,441
Myeloma					
Male	36,315	41,836	41,906	53,382	60,424
Female	20,046	23,478	23,548	29,616	33,879
Female	16,269	18,358	18,358	23,766	26,545
Leukemia					
Male	86,548	97,217	97,309	202,266	218,659
Female	50,266	56,859	56,917	114,783	124,335
Female	36,282	40,358	40,392	87,483	94,324
Acute Lymphocytic Leuk					
Male	13,978	14,185	14,185	49,880	50,189
Female	7,914	7,981	7,981	27,421	27,513
Female	6,064	6,204	6,204	22,459	22,676
Childhood (0-19)					
Male	58,085	58,205	58,498	237,628	238,095
Female	30,670	30,716	30,847	122,159	122,351
Female	27,415	27,489	27,651	115,469	115,744
Kaposi Sarcoma					
Male	6,370	6,864	6,864	20,160	21,169
Female	5,838	6,252	6,252	18,784	19,642
Female	532	612	612	1,376	1,527
Mesothelioma					
Male	2,668	3,333	3,333	4,161	4,920
Female	1,880	2,386	2,386	2,501	3,078
Female	788	947	947	1,660	1,842

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER 9 registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census.

<sup>bcde</sup> (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion: (c) First invasive tumor ever; (d) First invasive tumor for each cancer site diagnosed during the previous 29 years (1975-2003); (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (1999-2003). For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 1999; Melanoma in 2000.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 29-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2004.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 29-year limited duration prevalence. The 1999 breast cancer is counted for the breast 5-year and 29-year limited duration prevalence.

In method (e) the 1999 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2000 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table I-22  
 US COMPLETE PREVALENCE COUNTS, INVASIVE CANCERS ONLY, January 1, 2004<sup>a</sup>  
 BY AGE AT PREVALENCE

Age at Prevalence	Age Specific								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
<u>Site/Sex</u>									
All Sites									
Males	4,848,429	16,120	38,774	67,794	144,693	318,859	645,699	1,116,039	2,500,451
Females	5,913,785	14,522	31,876	73,689	212,191	581,836	1,039,847	1,258,340	2,701,485
Oral Cavity & Pharynx									
Males	150,946	46	527	1,324	3,270	14,521	32,689	37,725	60,844
Females	84,910	85	511	1,519	3,268	8,168	14,433	18,310	38,616
Esophagus									
Males	18,763	0	0	32	255	973	3,750	5,978	7,775
Females	6,022	0	12	23	33	226	854	1,378	3,497
Stomach									
Males	34,708	0	17	119	607	2,209	4,775	8,337	18,643
Females	25,592	33	23	105	446	1,708	3,234	4,649	15,394
Colon & Rectum									
Males	521,676	0	46	822	4,958	20,770	62,456	115,774	316,850
Females	554,659	11	75	819	4,801	19,051	52,318	94,583	383,002
Liver & Intrahep									
Males	12,411	263	389	319	267	1,237	4,028	2,656	3,252
Females	7,018	389	369	267	359	624	1,127	1,435	2,448
Pancreas									
Males	13,423	0	47	46	250	1,182	2,659	3,674	5,567
Females	15,024	0	63	147	356	1,206	2,256	3,699	7,297
Larynx									
Males	76,247	0	0	45	244	2,757	10,911	21,016	41,274
Females	18,600	0	0	46	230	1,255	2,655	5,229	9,185
Lung & Bronchus									
Males	174,880	23	73	268	1,109	6,906	24,315	50,502	91,686
Females	183,248	0	33	329	1,478	8,138	24,765	50,219	98,287
Melanoma of the Skin									
Males	333,330	55	769	4,499	16,117	46,145	75,025	76,142	114,578
Females	356,691	68	870	9,662	30,498	66,872	82,644	64,932	101,145

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table I-22 - continued  
US COMPLETE PREVALENCE COUNTS, INVASIVE CANCERS ONLY, January 1, 2004<sup>a</sup>  
 BY AGE AT PREVALENCE

Age at Prevalence	Age Specific								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
<u>Site/Sex</u>									
Breast									
Males	12,270	0	12	0	91	532	1,950	3,278	6,407
Females	2,407,943	0	82	2,097	35,151	203,932	471,756	578,108	1,116,817
Cervix									
Females	250,726	0	23	2,300	19,359	45,393	54,883	49,076	79,691
Corpus & Uterus, NOS									
Females	568,407	0	33	550	4,497	22,806	71,239	117,700	351,581
Ovary <sup>d</sup>									
Females	172,765	57	990	3,128	7,443	21,361	35,354	39,174	65,259
Prostate									
Males	2,024,489	34	58	103	282	15,786	167,564	509,996	1,330,665
Urinary Bladder									
Males	377,523	39	109	680	2,499	12,816	40,436	84,367	236,576
Females	134,267	22	45	375	1,333	4,592	12,214	26,715	88,970
Kidney & Renal Pelvis									
Males	141,899	1,550	2,415	2,125	3,869	12,865	26,944	36,480	55,651
Females	98,367	1,636	2,499	2,462	3,637	9,204	15,277	21,466	42,186
Hodgkin Lymphoma									
Males	78,455	175	2,171	8,440	16,776	21,227	16,175	8,444	5,048
Females	72,752	78	1,567	8,936	17,183	20,266	13,657	6,072	4,995
Non-Hodgkin Lymphoma									
Males	197,416	704	3,407	6,025	11,723	24,843	38,664	44,245	67,805
Females	183,713	308	1,403	3,404	8,069	17,550	30,736	39,111	83,133
Myeloma									
Males	29,745	0	5	12	574	2,166	6,246	8,590	12,152
Females	23,967	0	0	33	248	1,346	4,192	6,198	11,950
Leukemia									
Males	117,771	5,715	10,976	9,236	8,274	10,308	15,154	21,104	37,003
Females	90,849	4,882	9,186	8,526	6,528	6,998	9,832	13,426	31,471
Acute Lymphocytic Leuk									
Males	29,334	4,841	9,521	7,192	4,745	1,602	640	513	279
Females	24,073	4,257	7,787	6,202	3,388	1,218	725	253	242

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

<sup>d</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table II-16  
ALL CANCER SITES (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	4,000,969	2,575,551	1,684,932	967,733	576,385	8,006,111	10,133,875	628,339	10,762,214
Males	2,045,700	1,263,135	783,314	350,307	188,472	3,991,427	4,719,465	128,964	4,848,429
Females	1,955,269	1,312,416	901,618	617,426	387,913	4,014,684	5,414,410	499,375	5,913,785
White <sup>b</sup>	3,470,077	2,268,586	1,516,326	880,613	527,020	7,022,890	8,963,343	556,535	9,519,878
Males	1,763,511	1,103,097	705,971	320,832	174,241	3,480,290	4,149,428	119,653	4,269,081
Females	1,706,566	1,165,489	810,355	559,781	352,779	3,542,600	4,813,915	436,882	5,250,797
Black <sup>b</sup>	369,267	212,712	116,988	58,155	33,350	683,128	808,407	34,714	843,121
Males	201,699	116,049	55,791	19,784	9,246	367,667	406,546	3,856	410,402
Females	167,568	96,663	61,197	38,371	24,104	315,461	401,861	30,858	432,719
Asian/Pacific Islander <sup>c</sup>	94,884	52,855	+	+	+	173,402	+	+	+
Males	41,766	21,367	+	+	+	72,822	+	+	+
Females	53,118	31,488	+	+	+	100,580	+	+	+
Hispanic <sup>d</sup>	209,405	119,228	+	+	+	391,682	+	+	+
Males	101,410	55,591	+	+	+	183,756	+	+	+
Females	107,995	63,637	+	+	+	207,926	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	2.0010%	0.0753%	0.1068%	0.2153%	0.5353%	1.2656%	3.0498%	6.7185%	10.6190%	9.8820%	2.1180%
Males	2.0133%	0.0786%	0.1146%	0.1977%	0.4107%	0.8480%	2.5748%	7.7294%	14.6025%	14.1535%	2.4355%
Females	1.9890%	0.0719%	0.0985%	0.2338%	0.6640%	1.6800%	3.5000%	5.8134%	7.6124%	7.5419%	1.9274%
White <sup>c</sup>	2.1765%	0.0809%	0.1167%	0.2401%	0.5853%	1.3279%	3.1515%	7.0255%	11.0647%	10.0733%	2.2067%
Males	2.1593%	0.0845%	0.1254%	0.2202%	0.4568%	0.8954%	2.6178%	7.9086%	14.8755%	14.1521%	2.4884%
Females	2.1935%	0.0772%	0.1076%	0.2617%	0.7236%	1.7711%	3.6727%	6.2142%	8.1276%	7.8683%	2.0450%
Black <sup>c</sup>	1.4862%	0.0519%	0.0703%	0.1327%	0.3877%	1.0599%	3.1068%	7.1388%	10.6235%	8.6047%	2.0451%
Males	1.7028%	0.0498%	0.0736%	0.1132%	0.2737%	0.7699%	3.3913%	10.1198%	17.3533%	14.1910%	2.8194%
Females	1.2902%	0.0541%	0.0669%	0.1508%	0.4882%	1.3139%	2.8712%	4.8051%	6.2123%	5.9111%	1.5508%
Asian/Pacific Islander <sup>c</sup>	1.2855%	0.0600%	0.0833%	0.1311%	0.3360%	0.9499%	2.1151%	3.9201%	6.7560%	7.4756%	1.4050%
Males	1.1713%	0.0670%	0.0914%	0.1202%	0.2086%	0.4836%	1.3102%	3.9184%	9.2258%	10.8071%	1.4858%
Females	1.3923%	0.0527%	0.0748%	0.1419%	0.4568%	1.3730%	2.8196%	3.9214%	4.9839%	5.2597%	1.3862%
Hispanic <sup>d</sup>	0.8010%	0.0694%	0.0963%	0.1459%	0.3653%	0.8955%	2.0241%	4.4695%	7.5576%	7.0980%	1.4796%
Males	0.7304%	0.0717%	0.1015%	0.1340%	0.2682%	0.4921%	1.5016%	5.1135%	11.0515%	10.7381%	1.7246%
Females	0.8756%	0.0670%	0.0908%	0.1603%	0.4771%	1.3147%	2.5110%	3.9338%	5.0612%	5.0177%	1.3566%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table III-16  
BRAIN AND OTHER NERVOUS SYSTEM (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	38,349	21,518	17,187	12,032	6,865	74,262	100,542	14,540	115,082
Males	20,364	11,819	9,672	6,098	3,571	40,225	53,857	6,630	60,487
Females	17,985	9,699	7,515	5,934	3,294	34,037	46,685	7,910	54,595
White <sup>b</sup>	33,840	18,924	15,184	10,752	6,095	65,469	88,966	8,080	97,046
Males	18,168	10,465	8,516	5,376	3,150	35,692	47,836	4,447 <sup>i</sup>	52,283
Females	15,672	8,459	6,668	5,376	2,945	29,777	41,130	3,633 <sup>i</sup>	44,763
Black <sup>b</sup>	2,901	1,647	1,433	914	596	5,731	7,799	+	+
Males	1,450	875	796	495	339	2,984	4,070	+	+
Females	1,451	772	637	419	257	2,747	3,729	+	+
Asian/Pacific Islander <sup>c</sup>	982	532	+	+	+	1,821	+	+	+
Males	494	251	+	+	+	939	+	+	+
Females	488	281	+	+	+	882	+	+	+
Hispanic <sup>d</sup>	3,546	1,772	+	+	+	6,452	+	+	+
Males	1,871	892	+	+	+	3,363	+	+	+
Females	1,675	880	+	+	+	3,089	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0187%	0.0135%	0.0201%	0.0156%	0.0185%	0.0226%	0.0224%	0.0212%	0.0189%	0.0119%	0.0187%
Males	0.0204%	0.0133%	0.0219%	0.0163%	0.0198%	0.0246%	0.0254%	0.0242%	0.0233%	0.0154%	0.0205%
Females	0.0171%	0.0137%	0.0183%	0.0148%	0.0171%	0.0206%	0.0196%	0.0185%	0.0156%	0.0101%	0.0171%
White <sup>c</sup>	0.0209%	0.0147%	0.0223%	0.0171%	0.0211%	0.0255%	0.0251%	0.0238%	0.0206%	0.0120%	0.0209%
Males	0.0228%	0.0145%	0.0247%	0.0179%	0.0225%	0.0275%	0.0281%	0.0275%	0.0249%	0.0160%	0.0228%
Females	0.0189%	0.0149%	0.0198%	0.0161%	0.0195%	0.0235%	0.0221%	0.0204%	0.0172%	0.0099%	0.0191%
Black <sup>c</sup>	0.0117%	0.0108%	0.0131%	0.0114%	0.0101%	0.0119%	0.0134%	0.0127%	0.0107%	0.0092%	0.0116%
Males	0.0125%	0.0107%	0.0137%	0.0116%	0.0112%	0.0140%	0.0153%	0.0127%	0.0169%	-	0.0125%
Females	0.0109%	0.0109%	0.0125%	0.0111%	0.0091%	0.0101%	0.0120%	0.0127%	0.0067%	0.0137%	0.0108%
Asian/Pacific Islander <sup>c</sup>	0.0113%	0.0075%	0.0149%	0.0100%	0.0111%	0.0130%	0.0113%	0.0100%	0.0123%	0.0106%	0.0113%
Males	0.0115%	0.0077%	0.0146%	0.0097%	0.0107%	0.0127%	0.0130%	0.0091%	0.0157%	0.0163%	0.0117%
Females	0.0111%	0.0073%	0.0153%	0.0102%	0.0114%	0.0134%	0.0097%	0.0108%	0.0099%	0.0068%	0.0111%
Hispanic <sup>d</sup>	0.0132%	0.0104%	0.0146%	0.0084%	0.0131%	0.0186%	0.0169%	0.0192%	0.0174%	0.0094%	0.0143%
Males	0.0133%	0.0099%	0.0140%	0.0086%	0.0133%	0.0198%	0.0187%	0.0198%	0.0145%	0.0144%	0.0146%
Females	0.0131%	0.0108%	0.0152%	0.0080%	0.0129%	0.0175%	0.0152%	0.0187%	0.0195%	-	0.0140%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table IV-21  
BREAST CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	819,033	604,268	407,714	261,657	138,936	1,763,214	2,311,009	109,204	2,420,213
Males	5,334	2,860	1,694	1,016	758	9,673	11,962	308	12,270
Females	813,699	601,408	406,020	260,641	138,178	1,753,541	2,299,047	108,896	2,407,943
White <sup>b</sup>	716,788	538,074	367,895	239,127	126,620	1,561,045	2,061,456	97,253	2,158,709
Males	4,528	2,533	1,463	904	724	8,338	10,440	181 <sup>i</sup>	10,621
Females	712,260	535,541	366,432	238,223	125,896	1,552,707	2,051,016	97,072	2,148,088
Black <sup>b</sup>	70,732	45,895	28,262	16,372	9,016	140,495	174,874	+	+
Males	664	217	197	96	30	1,053	1,217	+	+
Females	70,068	45,678	28,065	16,276	8,986	139,442	173,657	7,527	181,184
Asian/Pacific Islander <sup>c</sup>	22,676	14,429	+	+	+	44,218	+	+	+
Males	77	55	+	+	+	163	+	+	+
Females	22,599	14,374	+	+	+	44,055	+	+	+
Hispanic <sup>d</sup>	40,609	25,430	+	+	+	79,851	+	+	+
Males	157	76	+	+	+	266	+	+	+
Females	40,452	25,354	+	+	+	79,585	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.4389%	-	0.0002%	0.0051%	0.0761%	0.3795%	0.9238%	1.4648%	1.8751%	1.9770%	0.4564%
Males	0.0050%	-	-	-	0.0005%	0.0024%	0.0082%	0.0205%	0.0314%	0.0398%	0.0061%
Females	0.8634%	-	0.0004%	0.0103%	0.1542%	0.7536%	1.7918%	2.7579%	3.2667%	3.0382%	0.8342%
White <sup>c</sup>	0.4810%	-	0.0002%	0.0050%	0.0750%	0.3852%	0.9594%	1.5602%	2.0095%	2.1014%	0.4797%
Males	0.0054%	-	-	-	0.0005%	0.0023%	0.0082%	0.0206%	0.0334%	0.0413%	0.0062%
Females	0.9550%	-	0.0003%	0.0103%	0.1552%	0.7777%	1.8884%	2.9745%	3.5326%	3.2150%	0.8856%
Black <sup>c</sup>	0.3044%	-	-	0.0064%	0.0894%	0.3602%	0.8268%	1.2201%	1.5701%	1.6334%	0.3988%
Males	0.0052%	-	-	-	-	0.0040%	0.0131%	0.0369%	0.0298%	0.0426%	0.0081%
Females	0.5752%	-	-	0.0123%	0.1677%	0.6722%	1.5005%	2.1463%	2.5797%	2.4005%	0.6876%
Asian/Pacific Islander <sup>c</sup>	0.3141%	-	-	0.0046%	0.0703%	0.3597%	0.7675%	1.0208%	1.1634%	1.0529%	0.3303%
Males	0.0025%	-	-	-	-	-	0.0042%	0.0080%	0.0169%	0.0242%	0.0031%
Females	0.6056%	-	-	0.0091%	0.1366%	0.6852%	1.4356%	1.8696%	1.9860%	1.7371%	0.5982%
Hispanic <sup>d</sup>	0.1640%	-	-	0.0043%	0.0490%	0.2696%	0.6173%	0.9534%	1.1581%	1.1752%	0.2969%
Males	0.0011%	-	-	-	-	0.0011%	0.0040%	0.0096%	0.0089%	0.0144%	0.0024%
Females	0.3357%	-	-	0.0093%	0.1051%	0.5485%	1.1887%	1.7384%	1.9791%	1.8386%	0.5409%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table V-12  
CERVIX UTERI CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	41,612	39,781	35,196	28,613	23,332	109,686	186,488	64,238	250,726
White <sup>b</sup>	32,718	31,459	29,292	23,868	19,049	87,662	151,179	53,612	204,791
Black <sup>b</sup>	6,245	5,289	3,997	3,322	3,255	14,783	24,669	+	+
Asian/Pacific Islander <sup>c</sup>	1,937	1,806	+	+	+	4,789	+	+	+
Hispanic <sup>d</sup>	8,304	6,184	+	+	+	18,932	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	0-9	10-19	20-29	Age Specific (Crude)						Age-Adjusted <sup>f</sup> All Ages
					30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	0.0599%	-	0.0002%	0.0105%	0.0754%	0.1201%	0.1141%	0.1034%	0.0905%	0.0565%	0.0590%
White <sup>c</sup>	0.0600%	-	-	0.0113%	0.0845%	0.1250%	0.1100%	0.0916%	0.0792%	0.0423%	0.0588%
Black <sup>c</sup>	0.0563%	-	-	0.0090%	0.0611%	0.1033%	0.1165%	0.1402%	0.1374%	0.1396%	0.0625%
Asian/Pacific Islander <sup>c</sup>	0.0591%	-	-	0.0037%	0.0332%	0.0985%	0.1316%	0.1440%	0.1301%	0.1166%	0.0574%
Hispanic <sup>d</sup>	0.0745%	-	-	0.0126%	0.0841%	0.1980%	0.2202%	0.2084%	0.2148%	0.1120%	0.1016%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VI-23  
COLON AND RECTUM CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	413,172	245,282	166,300	110,399	61,991	796,433	1,026,682	49,653	1,076,335
Males	208,481	120,903	81,877	53,796	26,796	397,719	503,545	18,131	521,676
Females	204,691	124,379	84,423	56,603	35,195	398,714	523,137	31,522	554,659
White <sup>b</sup>	354,407	214,315	146,951	98,437	55,723	690,690	896,513	44,382	940,895
Males	180,121	106,272	72,849	48,403	24,252	347,063	442,470	16,341	458,811
Females	174,286	108,043	74,102	50,034	31,471	343,627	454,043	28,041	482,084
Black <sup>b</sup>	40,623	21,272	13,602	8,557	4,435	73,035	90,401	3,421	93,822
Males	18,483	9,395	6,019	3,564	1,599	32,992	39,740	1,159	40,899
Females	22,140	11,877	7,583	4,993	2,836	40,043	50,661	2,262	52,923
Asian/Pacific Islander <sup>c</sup>	12,676	6,902	+	+	+	22,945	+	+	+
Males	6,655	3,687	+	+	+	12,057	+	+	+
Females	6,021	3,215	+	+	+	10,888	+	+	+
Hispanic <sup>d</sup>	19,880	10,031	+	+	+	35,328	+	+	+
Males	10,684	5,381	+	+	+	18,809	+	+	+
Females	9,196	4,650	+	+	+	16,519	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.2012%	-	0.0002%	0.0035%	0.0201%	0.0736%	0.2526%	0.6378%	1.2236%	1.6114%	0.2156%
Males	0.2045%	-	0.0002%	0.0037%	0.0209%	0.0765%	0.2884%	0.7661%	1.4730%	1.8018%	0.2514%
Females	0.1979%	-	0.0003%	0.0033%	0.0193%	0.0706%	0.2187%	0.5230%	1.0354%	1.5071%	0.1876%
White <sup>c</sup>	0.2121%	-	0.0003%	0.0036%	0.0196%	0.0709%	0.2414%	0.6307%	1.2394%	1.6405%	0.2152%
Males	0.2143%	-	0.0003%	0.0037%	0.0201%	0.0732%	0.2780%	0.7605%	1.4851%	1.8358%	0.2509%
Females	0.2098%	-	0.0003%	0.0034%	0.0190%	0.0684%	0.2057%	0.5116%	1.0500%	1.5350%	0.1866%
Black <sup>c</sup>	0.1583%	-	-	0.0030%	0.0197%	0.0872%	0.3209%	0.7625%	1.2256%	1.3858%	0.2279%
Males	0.1524%	-	-	0.0032%	0.0194%	0.0927%	0.3664%	0.8529%	1.3561%	1.4399%	0.2511%
Females	0.1636%	-	-	0.0029%	0.0199%	0.0824%	0.2833%	0.6918%	1.1401%	1.3597%	0.2113%
Asian/Pacific Islander <sup>c</sup>	0.1777%	-	-	0.0036%	0.0215%	0.0747%	0.2569%	0.5774%	1.0972%	1.4880%	0.2009%
Males	0.1924%	-	-	0.0048%	0.0242%	0.0795%	0.2781%	0.7198%	1.4367%	1.7137%	0.2433%
Females	0.1640%	-	-	0.0024%	0.0190%	0.0704%	0.2384%	0.4580%	0.8537%	1.3379%	0.1692%
Hispanic <sup>d</sup>	0.0724%	-	-	0.0029%	0.0155%	0.0535%	0.1767%	0.4627%	0.8854%	1.0642%	0.1533%
Males	0.0746%	-	-	0.0033%	0.0155%	0.0501%	0.2039%	0.5792%	1.1795%	1.2354%	0.1877%
Females	0.0701%	-	-	0.0024%	0.0156%	0.0570%	0.1513%	0.3658%	0.6752%	0.9663%	0.1278%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VII-12  
CORPUS AND UTERUS, NOS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	144,321	111,240	84,444	64,539	52,023	325,042	500,188	68,219	568,407
White <sup>b</sup>	128,644	101,702	79,124	60,661	49,320	295,404	461,176	60,537	521,713
Black <sup>b</sup>	9,355	5,220	3,091	2,276	1,376	17,120	22,275	1,425	23,700
Asian/Pacific Islander <sup>c</sup>	3,867	2,512	+	+	+	7,637	+	+	+
Hispanic <sup>d</sup>	7,272	4,561	+	+	+	14,687	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	0-9	10-19	20-29	Age Specific (Crude)						Age-Adjusted <sup>f</sup>
					30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.1522%	-	-	0.0026%	0.0207%	0.0848%	0.2844%	0.5902%	0.6642%	0.5303%	0.1480%
White <sup>c</sup>	0.1730%	-	-	0.0023%	0.0210%	0.0850%	0.3045%	0.6534%	0.7410%	0.5792%	0.1610%
Black <sup>c</sup>	0.0712%	-	-	0.0032%	0.0173%	0.0504%	0.1358%	0.3784%	0.4291%	0.2874%	0.0890%
Asian/Pacific Islander <sup>c</sup>	0.1039%	-	-	0.0028%	0.0199%	0.1070%	0.2752%	0.3438%	0.3162%	0.2382%	0.1020%
Hispanic <sup>d</sup>	0.0602%	-	-	0.0030%	0.0266%	0.0769%	0.2041%	0.3760%	0.3559%	0.2709%	0.0969%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VIII-16  
ESOPHAGUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	17,196	4,268	1,939	764	288	23,242	24,565	220	24,785
Males	13,312	3,238	1,367	449	216	17,785	18,653	110	18,763
Females	3,884	1,030	572	315	72	5,457	5,912	110	6,022
White <sup>b</sup>	15,061	3,783	1,578	642	256	20,301	21,406	182	21,588
Males	11,838	2,941	1,095	385	198	15,765	16,529	103	16,632
Females	3,223	842	483	257	58	4,536	4,877	79	4,956
Black <sup>b</sup>	1,725	409	308	112	29	2,407	2,609	56	2,665
Males	1,151	236	231	64	15	1,596	1,697	18	1,715
Females	574	173	77	48	14	811	912	38	950
Asian/Pacific Islander <sup>c</sup>	266	76	+	+	+	386	+	+	+
Males	201	57	+	+	+	292	+	+	+
Females	65	19	+	+	+	94	+	+	+
Hispanic <sup>d</sup>	551	154	+	+	+	804	+	+	+
Males	462	110	+	+	+	656	+	+	+
Females	89	44	+	+	+	148	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0061%	-	-	0.0001%	0.0005%	0.0022%	0.0104%	0.0254%	0.0376%	0.0263%	0.0066%
Males	0.0094%	-	-	-	0.0008%	0.0035%	0.0172%	0.0436%	0.0642%	0.0465%	0.0111%
Females	0.0029%	-	-	-	-	0.0010%	0.0040%	0.0091%	0.0175%	0.0153%	0.0028%
White <sup>c</sup>	0.0068%	-	-	0.0001%	0.0006%	0.0023%	0.0108%	0.0269%	0.0407%	0.0270%	0.0069%
Males	0.0105%	-	-	-	0.0010%	0.0037%	0.0185%	0.0464%	0.0699%	0.0473%	0.0119%
Females	0.0031%	-	-	-	-	0.0009%	0.0033%	0.0090%	0.0182%	0.0161%	0.0028%
Black <sup>c</sup>	0.0053%	-	-	-	-	0.0029%	0.0149%	0.0317%	0.0322%	0.0242%	0.0072%
Males	0.0069%	-	-	-	-	0.0037%	0.0197%	0.0467%	0.0526%	0.0390%	0.0108%
Females	0.0038%	-	-	-	-	0.0022%	0.0109%	0.0199%	0.0189%	0.0171%	0.0047%
Asian/Pacific Islander <sup>c</sup>	0.0031%	-	-	-	-	0.0009%	0.0043%	0.0105%	0.0204%	0.0229%	0.0034%
Males	0.0048%	-	-	-	-	0.0017%	0.0058%	0.0209%	0.0328%	0.0470%	0.0060%
Females	0.0015%	-	-	-	-	-	0.0030%	-	0.0115%	0.0069%	0.0015%
Hispanic <sup>d</sup>	0.0017%	-	-	-	0.0004%	0.0010%	0.0048%	0.0124%	0.0220%	0.0142%	0.0035%
Males	0.0027%	-	-	-	0.0006%	0.0015%	0.0079%	0.0229%	0.0446%	0.0246%	0.0064%
Females	0.0007%	-	-	-	-	-	0.0018%	0.0037%	0.0059%	0.0083%	0.0012%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table IX-16  
HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	33,763	29,194	25,949	21,398	15,732	83,961	135,338	15,869	151,207
Males	17,936	15,240	13,243	10,919	8,529	43,928	70,330	8,125	78,455
Females	15,827	13,954	12,706	10,479	7,203	40,033	65,008	7,744	72,752
White <sup>b</sup>	28,830	25,990	23,389	19,824	14,615	73,653	121,430	17,719	139,149
Males	15,387	13,648	12,006	10,017	7,794	38,751	62,968	9,182	72,150
Females	13,443	12,342	11,383	9,807	6,821	34,902	58,462	8,537	66,999
Black <sup>b</sup>	3,628	2,620	2,133	1,248	927	8,088	10,971	1,259	12,230
Males	1,778	1,295	1,023	724	636	3,942	5,732	877	6,609
Females	1,850	1,325	1,110	524	291	4,146	5,239	382	5,621
Asian/Pacific Islander <sup>c</sup>	677	316	+	+	+	1,203	+	+	+
Males	368	165	+	+	+	626	+	+	+
Females	309	151	+	+	+	577	+	+	+
Hispanic <sup>d</sup>	2,862	1,946	+	+	+	6,134	+	+	+
Males	1,731	1,029	+	+	+	3,538	+	+	+
Females	1,131	917	+	+	+	2,596	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0196%	0.0006%	0.0083%	0.0310%	0.0349%	0.0253%	0.0191%	0.0186%	0.0174%	0.0109%	0.0195%
Males	0.0212%	0.0010%	0.0097%	0.0288%	0.0352%	0.0289%	0.0228%	0.0228%	0.0211%	0.0143%	0.0211%
Females	0.0181%	0.0003%	0.0069%	0.0333%	0.0347%	0.0218%	0.0157%	0.0149%	0.0145%	0.0091%	0.0180%
White <sup>c</sup>	0.0220%	0.0007%	0.0096%	0.0353%	0.0398%	0.0280%	0.0211%	0.0206%	0.0194%	0.0116%	0.0219%
Males	0.0237%	0.0012%	0.0110%	0.0324%	0.0396%	0.0318%	0.0249%	0.0249%	0.0228%	0.0152%	0.0234%
Females	0.0204%	-	0.0081%	0.0384%	0.0400%	0.0241%	0.0174%	0.0166%	0.0169%	0.0096%	0.0205%
Black <sup>c</sup>	0.0169%	-	0.0053%	0.0257%	0.0309%	0.0251%	0.0181%	0.0174%	0.0130%	0.0115%	0.0172%
Males	0.0176%	-	0.0055%	0.0233%	0.0302%	0.0288%	0.0206%	0.0225%	0.0215%	0.0213%	0.0188%
Females	0.0162%	-	0.0051%	0.0279%	0.0314%	0.0218%	0.0159%	0.0134%	0.0074%	-	0.0160%
Asian/Pacific Islander <sup>c</sup>	0.0072%	-	0.0040%	0.0123%	0.0116%	0.0082%	0.0057%	0.0069%	0.0060%	0.0055%	0.0069%
Males	0.0081%	-	0.0060%	0.0118%	0.0118%	0.0091%	0.0073%	0.0094%	0.0095%	-	0.0078%
Females	0.0065%	-	0.0020%	0.0128%	0.0114%	0.0074%	0.0044%	0.0048%	-	-	0.0061%
Hispanic <sup>d</sup>	0.0119%	0.0009%	0.0081%	0.0166%	0.0169%	0.0147%	0.0144%	0.0241%	0.0175%	0.0117%	0.0131%
Males	0.0132%	0.0017%	0.0108%	0.0151%	0.0169%	0.0175%	0.0181%	0.0284%	0.0267%	0.0172%	0.0154%
Females	0.0105%	-	0.0053%	0.0185%	0.0168%	0.0118%	0.0109%	0.0204%	0.0110%	0.0085%	0.0112%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table X-13  
KAPOSI SARCOMAEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	6,370	8,539	3,456	1,174	374	17,922	20,160	+	+
Males	5,838	8,148	3,207	1,146	300	16,760	18,784	+	+
Females	532	391	249	28	74	1,162	1,376	+	+
White <sup>b</sup>	4,522	6,768	2,906	1,000	302	13,822	15,699	+	+
Males	4,080	6,516	2,704	972	239	12,937	14,633	+	+
Females	442	252	202	28	63	885	1,066	+	+
Black <sup>b</sup>	1,531	1,225	321	71	39	3,043	3,198	+	+
Males	1,452	1,131	285	71	39	2,834	2,989	+	+
Females	79	94	36	0	0	209	209	+	+
Asian/Pacific Islander <sup>c</sup>	87	92	+	+	+	190	+	+	+
Males	84	92	+	+	+	187	+	+	+
Females	3	0	+	+	+	3	+	+	+
Hispanic <sup>d</sup>	1,057	1,330	+	+	+	2,845	+	+	+
Males	988	1,288	+	+	+	2,705	+	+	+
Females	69	42	+	+	+	140	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0059%	-	-	0.0007%	0.0083%	0.0161%	0.0090%	0.0050%	0.0053%	0.0104%	0.0059%
Males	0.0113%	-	-	0.0013%	0.0160%	0.0318%	0.0180%	0.0097%	0.0096%	0.0189%	0.0115%
Females	0.0006%	-	-	-	0.0003%	0.0004%	0.0004%	0.0009%	0.0020%	0.0058%	0.0005%
White <sup>c</sup>	0.0059%	-	-	0.0005%	0.0076%	0.0162%	0.0088%	0.0052%	0.0058%	0.0108%	0.0058%
Males	0.0112%	-	-	0.0008%	0.0144%	0.0317%	0.0177%	0.0098%	0.0104%	0.0195%	0.0112%
Females	0.0006%	-	-	-	0.0003%	0.0004%	-	0.0010%	0.0023%	0.0061%	0.0005%
Black <sup>c</sup>	0.0080%	-	-	0.0026%	0.0170%	0.0210%	0.0126%	0.0044%	-	0.0081%	0.0083%
Males	0.0161%	-	-	0.0050%	0.0356%	0.0437%	0.0264%	0.0090%	-	-	0.0170%
Females	0.0007%	-	-	-	-	-	-	-	-	0.0086%	0.0008%
Asian/Pacific Islander <sup>c</sup>	0.0014%	-	-	-	0.0011%	0.0036%	0.0028%	0.0021%	-	-	0.0014%
Males	0.0028%	-	-	-	0.0023%	0.0075%	0.0061%	0.0040%	-	-	0.0029%
Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic <sup>d</sup>	0.0059%	-	-	0.0008%	0.0113%	0.0189%	0.0091%	0.0075%	0.0105%	0.0240%	0.0078%
Males	0.0110%	-	-	0.0014%	0.0208%	0.0366%	0.0186%	0.0145%	0.0213%	0.0345%	0.0147%
Females	0.0006%	-	-	-	-	-	-	-	-	0.0181%	0.0011%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XI-16  
KIDNEY AND RENAL PELVIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	98,835	57,255	35,468	20,124	11,497	185,832	228,714	11,552	240,266
Males	59,318	34,237	20,735	11,573	6,711	111,099	135,506	6,393	141,899
Females	39,517	23,018	14,733	8,551	4,786	74,733	93,208	5,159	98,367
White <sup>b</sup>	84,557	49,158	31,637	18,003	10,132	160,099	198,486	13,137	211,623
Males	51,165	29,695	18,718	10,551	6,081	96,614	118,884	6,222	125,106
Females	33,392	19,463	12,919	7,452	4,051	63,485	79,602	6,915	86,517
Black <sup>b</sup>	10,517	6,218	3,033	1,727	1,110	19,361	23,041	1,983	25,024
Males	5,832	3,406	1,534	791	504	10,588	12,257	697	12,954
Females	4,685	2,812	1,499	936	606	8,773	10,784	1,286	12,070
Asian/Pacific Islander <sup>c</sup>	1,983	969	+	+	+	3,314	+	+	+
Males	1,213	614	+	+	+	2,059	+	+	+
Females	770	355	+	+	+	1,255	+	+	+
Hispanic <sup>d</sup>	7,378	3,517	+	+	+	12,673	+	+	+
Males	4,355	1,943	+	+	+	7,304	+	+	+
Females	3,023	1,574	+	+	+	5,369	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0470%	0.0069%	0.0042%	0.0015%	0.0082%	0.0325%	0.0860%	0.1703%	0.2227%	0.1836%	0.0493%
Males	0.0573%	0.0065%	0.0035%	0.0016%	0.0090%	0.0399%	0.1156%	0.2313%	0.3084%	0.2594%	0.0657%
Females	0.0368%	0.0074%	0.0050%	0.0015%	0.0074%	0.0251%	0.0580%	0.1157%	0.1581%	0.1421%	0.0360%
White <sup>c</sup>	0.0510%	0.0072%	0.0045%	0.0014%	0.0084%	0.0338%	0.0897%	0.1786%	0.2334%	0.1902%	0.0515%
Males	0.0622%	0.0072%	0.0039%	0.0014%	0.0093%	0.0413%	0.1196%	0.2401%	0.3221%	0.2739%	0.0685%
Females	0.0399%	0.0073%	0.0051%	0.0015%	0.0074%	0.0261%	0.0604%	0.1221%	0.1650%	0.1450%	0.0374%
Black <sup>c</sup>	0.0412%	0.0076%	0.0048%	0.0027%	0.0106%	0.0368%	0.0997%	0.1835%	0.2294%	0.1901%	0.0538%
Males	0.0488%	0.0055%	0.0032%	0.0028%	0.0109%	0.0445%	0.1417%	0.2453%	0.3317%	0.2341%	0.0711%
Females	0.0343%	0.0098%	0.0064%	0.0026%	0.0104%	0.0300%	0.0649%	0.1350%	0.1623%	0.1689%	0.0412%
Asian/Pacific Islander <sup>c</sup>	0.0255%	0.0031%	0.0015%	-	0.0046%	0.0172%	0.0474%	0.0931%	0.1322%	0.1138%	0.0278%
Males	0.0325%	0.0022%	-	-	0.0053%	0.0234%	0.0652%	0.1388%	0.1820%	0.1530%	0.0384%
Females	0.0191%	0.0040%	0.0020%	-	0.0045%	0.0115%	0.0318%	0.0549%	0.0964%	0.0878%	0.0194%
Hispanic <sup>d</sup>	0.0265%	0.0059%	0.0027%	0.0010%	0.0073%	0.0291%	0.0821%	0.1611%	0.2531%	0.1696%	0.0487%
Males	0.0295%	0.0050%	0.0016%	0.0010%	0.0081%	0.0327%	0.1088%	0.2082%	0.3445%	0.2143%	0.0624%
Females	0.0233%	0.0069%	0.0038%	0.0010%	0.0064%	0.0254%	0.0572%	0.1219%	0.1877%	0.1441%	0.0379%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XII-16  
LARYNX CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	31,966	21,902	15,704	10,618	6,945	66,967	90,186	4,661	94,847
Males	25,659	17,456	12,484	8,756	5,583	53,605	72,417	3,830	76,247
Females	6,307	4,446	3,220	1,862	1,362	13,362	17,769	831	18,600
White <sup>b</sup>	26,814	18,834	13,547	9,366	6,312	57,012	77,676	3,790	81,466
Males	21,623	14,991	10,813	7,824	5,091	45,716	62,632	3,107	65,739
Females	5,191	3,843	2,734	1,542	1,221	11,296	15,044	683	15,727
Black <sup>b</sup>	4,298	2,561	1,829	1,082	538	8,345	10,501	320	10,821
Males	3,287	2,044	1,422	813	411	6,509	8,121	268	8,389
Females	1,011	517	407	269	127	1,836	2,380	52	2,432
Asian/Pacific Islander <sup>c</sup>	505	298	+	+	+	943	+	+	+
Males	447	264	+	+	+	832	+	+	+
Females	58	34	+	+	+	111	+	+	+
Hispanic <sup>d</sup>	1,470	856	+	+	+	2,834	+	+	+
Males	1,253	725	+	+	+	2,410	+	+	+
Females	217	131	+	+	+	424	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0156%	-	-	0.0002%	0.0009%	0.0065%	0.0267%	0.0708%	0.0908%	0.0606%	0.0167%
Males	0.0255%	-	-	0.0002%	0.0010%	0.0092%	0.0450%	0.1221%	0.1706%	0.1398%	0.0303%
Females	0.0060%	-	-	0.0002%	0.0008%	0.0038%	0.0093%	0.0248%	0.0306%	0.0173%	0.0060%
White <sup>c</sup>	0.0167%	-	-	0.0002%	0.0009%	0.0067%	0.0268%	0.0721%	0.0937%	0.0620%	0.0171%
Males	0.0269%	-	-	-	0.0009%	0.0094%	0.0446%	0.1230%	0.1729%	0.1425%	0.0305%
Females	0.0066%	-	-	0.0003%	0.0009%	0.0040%	0.0095%	0.0255%	0.0327%	0.0185%	0.0063%
Black <sup>c</sup>	0.0176%	-	-	-	0.0012%	0.0100%	0.0447%	0.1127%	0.1174%	0.0635%	0.0240%
Males	0.0291%	-	-	-	-	0.0146%	0.0798%	0.2031%	0.2359%	0.1560%	0.0453%
Females	0.0072%	-	-	-	-	0.0060%	0.0156%	0.0419%	0.0396%	0.0188%	0.0089%
Asian/Pacific Islander <sup>c</sup>	0.0073%	-	-	-	-	0.0017%	0.0104%	0.0300%	0.0505%	0.0462%	0.0082%
Males	0.0133%	-	-	-	-	0.0034%	0.0195%	0.0576%	0.1055%	0.1065%	0.0170%
Females	0.0016%	-	-	-	-	-	0.0023%	0.0069%	0.0111%	-	0.0017%
Hispanic <sup>d</sup>	0.0056%	-	-	-	0.0007%	0.0037%	0.0142%	0.0441%	0.0756%	0.0460%	0.0116%
Males	0.0092%	-	-	-	0.0006%	0.0052%	0.0247%	0.0848%	0.1609%	0.1091%	0.0233%
Females	0.0018%	-	-	-	0.0007%	0.0021%	0.0044%	0.0103%	0.0146%	0.0099%	0.0029%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIII-24  
LEUKEMIA (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	86,548	50,210	30,629	18,629	10,673	162,426	202,266	6,354	208,620
Males	50,266	28,357	17,336	10,330	5,879	93,201	114,783	2,988	117,771
Females	36,282	21,853	13,293	8,299	4,794	69,225	87,483	3,366	90,849
White <sup>b</sup>	76,866	45,030	27,828	16,850	9,713	145,205	181,450	5,401	186,851
Males	44,967	25,258	16,033	9,439	5,360	83,717	103,537	2,196	105,733
Females	31,899	19,772	11,795	7,411	4,353	61,488	77,913	3,205	81,118
Black <sup>b</sup>	5,588	2,964	1,544	1,023	580	9,906	11,995	627	12,622
Males	2,851	1,625	700	494	323	5,083	6,092	299	6,391
Females	2,737	1,339	844	529	257	4,823	5,903	328	6,231
Asian/Pacific Islander <sup>c</sup>	1,898	1,097	+	+	+	3,582	+	+	+
Males	1,118	617	+	+	+	2,049	+	+	+
Females	780	480	+	+	+	1,533	+	+	+
Hispanic <sup>d</sup>	6,815	4,066	+	+	+	13,223	+	+	+
Males	3,814	2,334	+	+	+	7,508	+	+	+
Females	3,001	1,732	+	+	+	5,715	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0424%	0.0276%	0.0310%	0.0130%	0.0149%	0.0245%	0.0478%	0.1067%	0.1545%	0.1706%	0.0441%
Males	0.0495%	0.0289%	0.0341%	0.0144%	0.0171%	0.0304%	0.0609%	0.1401%	0.2106%	0.2318%	0.0555%
Females	0.0354%	0.0262%	0.0278%	0.0116%	0.0126%	0.0186%	0.0355%	0.0768%	0.1121%	0.1371%	0.0348%
White <sup>c</sup>	0.0476%	0.0307%	0.0346%	0.0146%	0.0157%	0.0259%	0.0510%	0.1176%	0.1708%	0.1861%	0.0483%
Males	0.0552%	0.0321%	0.0375%	0.0158%	0.0178%	0.0319%	0.0646%	0.1542%	0.2311%	0.2545%	0.0604%
Females	0.0399%	0.0292%	0.0315%	0.0133%	0.0135%	0.0197%	0.0376%	0.0839%	0.1243%	0.1491%	0.0382%
Black <sup>c</sup>	0.0223%	0.0106%	0.0172%	0.0072%	0.0111%	0.0164%	0.0358%	0.0638%	0.1056%	0.0873%	0.0270%
Males	0.0254%	0.0095%	0.0189%	0.0081%	0.0132%	0.0218%	0.0459%	0.0800%	0.1440%	0.0929%	0.0332%
Females	0.0196%	0.0117%	0.0155%	0.0065%	0.0093%	0.0117%	0.0273%	0.0511%	0.0803%	0.0846%	0.0224%
Asian/Pacific Islander <sup>c</sup>	0.0232%	0.0261%	0.0243%	0.0099%	0.0138%	0.0186%	0.0250%	0.0473%	0.0483%	0.0520%	0.0244%
Males	0.0278%	0.0291%	0.0289%	0.0126%	0.0168%	0.0222%	0.0287%	0.0576%	0.0637%	0.0835%	0.0300%
Females	0.0190%	0.0228%	0.0194%	0.0073%	0.0109%	0.0153%	0.0217%	0.0386%	0.0372%	0.0311%	0.0195%
Hispanic <sup>d</sup>	0.0269%	0.0296%	0.0365%	0.0139%	0.0150%	0.0193%	0.0311%	0.0592%	0.0739%	0.0789%	0.0315%
Males	0.0295%	0.0305%	0.0391%	0.0161%	0.0174%	0.0212%	0.0351%	0.0708%	0.1083%	0.1089%	0.0374%
Females	0.0243%	0.0287%	0.0337%	0.0113%	0.0122%	0.0173%	0.0275%	0.0496%	0.0493%	0.0617%	0.0267%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIII-25  
ACUTE LYMPHOCYTIC LEUKEMIA (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	13,978	11,128	9,163	7,176	5,427	32,415	49,880	3,527	53,407
Males	7,914	6,233	5,014	3,943	2,826	18,126	27,421	1,913	29,334
Females	6,064	4,895	4,149	3,233	2,601	14,289	22,459	1,614	24,073
White <sup>b</sup>	12,171	9,655	8,009	6,310	4,948	28,208	43,888	+	+
Males	6,920	5,368	4,522	3,503	2,595	15,899	24,331	+	+
Females	5,251	4,287	3,487	2,807	2,353	12,309	19,557	+	+
Black <sup>b</sup>	870	767	652	416	244	2,162	3,093	+	+
Males	436	451	284	205	120	1,091	1,549	+	+
Females	434	316	368	211	124	1,071	1,544	+	+
Asian/Pacific Islander <sup>c</sup>	573	456	+	+	+	1,314	+	+	+
Males	344	274	+	+	+	750	+	+	+
Females	229	182	+	+	+	564	+	+	+
Hispanic <sup>d</sup>	3,210	2,295	+	+	+	6,802	+	+	+
Males	1,753	1,319	+	+	+	3,825	+	+	+
Females	1,457	976	+	+	+	2,977	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0092%	0.0237%	0.0262%	0.0064%	0.0028%	0.0020%	0.0023%	0.0024%	0.0012%	0.0011%	0.0093%
Males	0.0105%	0.0248%	0.0291%	0.0081%	0.0034%	0.0022%	0.0023%	0.0029%	0.0015%	0.0011%	0.0103%
Females	0.0080%	0.0226%	0.0231%	0.0046%	0.0022%	0.0018%	0.0023%	0.0020%	0.0011%	0.0011%	0.0083%
White <sup>c</sup>	0.0102%	0.0267%	0.0298%	0.0077%	0.0030%	0.0022%	0.0024%	0.0025%	0.0014%	0.0010%	0.0105%
Males	0.0115%	0.0279%	0.0327%	0.0093%	0.0037%	0.0023%	0.0025%	0.0031%	0.0015%	-	0.0115%
Females	0.0090%	0.0255%	0.0268%	0.0060%	0.0023%	0.0020%	0.0023%	0.0020%	0.0013%	0.0011%	0.0095%
Black <sup>c</sup>	0.0041%	0.0078%	0.0119%	0.0022%	0.0011%	-	-	-	-	-	0.0036%
Males	0.0046%	0.0069%	0.0129%	0.0034%	-	-	-	-	-	-	0.0039%
Females	0.0036%	0.0087%	0.0109%	-	0.0013%	-	-	-	-	-	0.0033%
Asian/Pacific Islander <sup>c</sup>	0.0073%	0.0216%	0.0199%	0.0034%	0.0028%	0.0022%	0.0027%	0.0028%	-	-	0.0079%
Males	0.0091%	0.0237%	0.0251%	0.0060%	0.0031%	0.0026%	0.0027%	-	-	-	0.0093%
Females	0.0057%	0.0193%	0.0144%	-	0.0026%	0.0020%	0.0027%	0.0039%	-	-	0.0064%
Hispanic <sup>d</sup>	0.0138%	0.0261%	0.0316%	0.0077%	0.0036%	0.0035%	0.0037%	0.0048%	-	-	0.0114%
Males	0.0149%	0.0268%	0.0344%	0.0095%	0.0043%	0.0038%	0.0032%	0.0048%	-	-	0.0122%
Females	0.0126%	0.0255%	0.0286%	0.0054%	0.0027%	0.0031%	0.0041%	0.0048%	-	-	0.0105%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIV-16  
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	14,336	2,580	953	769	296	17,763	19,061	368	19,429
Males	9,723	1,564	524	321	152	11,787	12,344	67	12,411
Females	4,613	1,016	429	448	144	5,976	6,717	301	7,018
White <sup>b</sup>	10,603	2,007	746	610	239	13,261	14,290	338	14,628
Males	7,099	1,194	407	219	122	8,675	9,085	117 <sup>i</sup>	9,202
Females	3,504	813	339	391	117	4,586	5,205	221 <sup>i</sup>	5,426
Black <sup>b</sup>	1,653	173	117	84	38	1,931	2,098	+	+
Males	1,155	124	58	50	26	1,337	1,423	+	+
Females	498	49	59	34	12	594	675	+	+
Asian/Pacific Islander <sup>c</sup>	1,801	342	+	+	+	2,234	+	+	+
Males	1,270	227	+	+	+	1,559	+	+	+
Females	531	115	+	+	+	675	+	+	+
Hispanic <sup>d</sup>	1,942	336	+	+	+	2,388	+	+	+
Males	1,321	191	+	+	+	1,595	+	+	+
Females	621	145	+	+	+	793	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0070%	0.0017%	0.0006%	0.0006%	0.0010%	0.0043%	0.0165%	0.0244%	0.0306%	0.0203%	0.0073%
Males	0.0096%	0.0016%	0.0005%	0.0006%	0.0013%	0.0064%	0.0269%	0.0347%	0.0449%	0.0290%	0.0107%
Females	0.0045%	0.0018%	0.0006%	0.0006%	0.0008%	0.0022%	0.0066%	0.0153%	0.0198%	0.0156%	0.0044%
White <sup>c</sup>	0.0056%	0.0018%	0.0006%	0.0006%	0.0007%	0.0034%	0.0133%	0.0172%	0.0212%	0.0156%	0.0055%
Males	0.0075%	0.0015%	0.0007%	0.0006%	0.0006%	0.0047%	0.0217%	0.0244%	0.0301%	0.0216%	0.0079%
Females	0.0037%	0.0020%	0.0005%	0.0006%	0.0008%	0.0020%	0.0051%	0.0106%	0.0143%	0.0123%	0.0035%
Black <sup>c</sup>	0.0048%	-	-	-	0.0012%	0.0029%	0.0182%	0.0198%	0.0232%	0.0093%	0.0060%
Males	0.0071%	-	-	-	0.0018%	0.0044%	0.0317%	0.0317%	0.0396%	-	0.0096%
Females	0.0027%	-	-	-	-	0.0015%	0.0070%	0.0105%	0.0125%	0.0086%	0.0032%
Asian/Pacific Islander <sup>c</sup>	0.0185%	0.0028%	-	-	0.0025%	0.0114%	0.0344%	0.0741%	0.0981%	0.0694%	0.0200%
Males	0.0263%	0.0029%	-	-	0.0046%	0.0194%	0.0571%	0.1078%	0.1517%	0.0979%	0.0307%
Females	0.0111%	0.0027%	-	-	-	0.0043%	0.0146%	0.0459%	0.0597%	0.0505%	0.0114%
Hispanic <sup>d</sup>	0.0055%	0.0015%	0.0005%	0.0008%	0.0007%	0.0053%	0.0196%	0.0338%	0.0460%	0.0403%	0.0101%
Males	0.0071%	0.0017%	-	0.0007%	0.0008%	0.0082%	0.0327%	0.0472%	0.0626%	0.0487%	0.0143%
Females	0.0039%	0.0013%	-	0.0009%	-	0.0023%	0.0074%	0.0227%	0.0341%	0.0354%	0.0065%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XV-22  
LUNG AND BRONCHUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	206,480	68,243	36,705	19,051	9,818	305,761	345,236	12,892	358,128
Males	101,519	31,773	17,638	9,160	4,783	147,917	167,413	7,467	174,880
Females	104,961	36,470	19,067	9,891	5,035	157,844	177,823	5,425	183,248
White <sup>b</sup>	178,187	59,851	32,779	16,905	8,674	265,677	300,763	6,351	307,114
Males	86,324	27,305	15,553	7,948	4,198	126,476	143,550	3,172	146,722
Females	91,863	32,546	17,226	8,957	4,476	139,201	157,213	3,179	160,392
Black <sup>b</sup>	21,464	6,451	3,102	1,692	903	30,595	34,037	610	34,647
Males	11,492	3,328	1,621	923	466	16,198	18,085	344	18,429
Females	9,972	3,123	1,481	769	437	14,397	15,952	266	16,218
Asian/Pacific Islander <sup>c</sup>	4,900	1,321	+	+	+	6,712	+	+	+
Males	2,691	779	+	+	+	3,749	+	+	+
Females	2,209	542	+	+	+	2,963	+	+	+
Hispanic <sup>d</sup>	5,693	1,636	+	+	+	8,142	+	+	+
Males	3,027	845	+	+	+	4,236	+	+	+
Females	2,666	791	+	+	+	3,906	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0792%	-	0.0002%	0.0011%	0.0044%	0.0252%	0.1052%	0.3251%	0.5545%	0.3823%	0.0858%
Males	0.0782%	-	0.0002%	0.0010%	0.0039%	0.0242%	0.1081%	0.3469%	0.6361%	0.4780%	0.0954%
Females	0.0801%	-	0.0002%	0.0013%	0.0049%	0.0262%	0.1024%	0.3056%	0.4929%	0.3298%	0.0789%
White <sup>c</sup>	0.0851%	-	0.0002%	0.0013%	0.0045%	0.0245%	0.1033%	0.3373%	0.5764%	0.3868%	0.0879%
Males	0.0812%	-	-	0.0010%	0.0040%	0.0221%	0.1006%	0.3466%	0.6430%	0.4744%	0.0945%
Females	0.0890%	-	-	0.0015%	0.0051%	0.0270%	0.1060%	0.3287%	0.5251%	0.3394%	0.0834%
Black <sup>c</sup>	0.0687%	-	-	0.0009%	0.0040%	0.0364%	0.1615%	0.3793%	0.5540%	0.3534%	0.0968%
Males	0.0765%	-	-	-	0.0039%	0.0431%	0.2087%	0.4645%	0.6748%	0.5144%	0.1218%
Females	0.0617%	-	-	-	0.0042%	0.0306%	0.1225%	0.3126%	0.4748%	0.2757%	0.0795%
Asian/Pacific Islander <sup>c</sup>	0.0563%	-	-	-	0.0043%	0.0195%	0.0725%	0.2138%	0.4131%	0.3543%	0.0636%
Males	0.0649%	-	-	-	0.0040%	0.0217%	0.0798%	0.2701%	0.5702%	0.4822%	0.0824%
Females	0.0483%	-	-	-	0.0047%	0.0176%	0.0661%	0.1667%	0.3004%	0.2692%	0.0497%
Hispanic <sup>d</sup>	0.0177%	-	-	0.0010%	0.0026%	0.0110%	0.0392%	0.1280%	0.2530%	0.1970%	0.0377%
Males	0.0179%	-	-	-	0.0025%	0.0109%	0.0424%	0.1423%	0.3339%	0.2731%	0.0464%
Females	0.0174%	-	-	0.0015%	0.0027%	0.0112%	0.0362%	0.1161%	0.1952%	0.1534%	0.0317%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XVI-16  
MELANOMA OF THE SKIN (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	214,760	156,160	109,126	81,224	54,484	460,827	642,642	47,379	690,021
Males	113,145	80,114	53,672	36,196	22,987	237,858	317,120	16,210	333,330
Females	101,615	76,046	55,454	45,028	31,497	222,969	325,522	31,169	356,691
White <sup>b</sup>	206,665	149,731	104,603	77,045	52,848	442,601	616,365	44,770	661,135
Males	109,025	76,878	51,691	34,499	22,257	228,799	304,746	15,451	320,197
Females	97,640	72,853	52,912	42,546	30,591	213,802	311,619	29,319	340,938
Black <sup>b</sup>	737	628	374	200	109	1,623	2,159	122	2,281
Males	342	317	148	48	35	774	916	46	962
Females	395	311	226	152	74	849	1,243	76	1,319
Asian/Pacific Islander <sup>c</sup>	549	323	+	+	+	1,004	+	+	+
Males	246	143	+	+	+	437	+	+	+
Females	303	180	+	+	+	567	+	+	+
Hispanic <sup>d</sup>	3,695	2,365	+	+	+	7,478	+	+	+
Males	1,330	893	+	+	+	2,746	+	+	+
Females	2,365	1,472	+	+	+	4,732	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.1065%	0.0003%	0.0029%	0.0256%	0.0703%	0.1334%	0.2032%	0.2762%	0.3511%	0.3229%	0.1092%
Males	0.1134%	0.0002%	0.0026%	0.0165%	0.0506%	0.1213%	0.2262%	0.3602%	0.5036%	0.5107%	0.1272%
Females	0.0998%	0.0003%	0.0031%	0.0351%	0.0906%	0.1455%	0.1815%	0.2011%	0.2360%	0.2199%	0.0973%
White <sup>c</sup>	0.1329%	0.0003%	0.0036%	0.0322%	0.0882%	0.1644%	0.2477%	0.3331%	0.4191%	0.3684%	0.1323%
Males	0.1404%	0.0003%	0.0032%	0.0198%	0.0619%	0.1469%	0.2726%	0.4277%	0.5946%	0.5921%	0.1515%
Females	0.1254%	0.0003%	0.0040%	0.0457%	0.1165%	0.1823%	0.2233%	0.2461%	0.2838%	0.2474%	0.1201%
Black <sup>c</sup>	0.0037%	-	-	0.0008%	0.0033%	0.0038%	0.0055%	0.0177%	0.0180%	0.0202%	0.0048%
Males	0.0037%	-	-	-	0.0027%	0.0040%	0.0052%	0.0223%	0.0220%	0.0284%	0.0055%
Females	0.0038%	-	-	-	0.0038%	0.0035%	0.0058%	0.0141%	0.0154%	0.0162%	0.0044%
Asian/Pacific Islander <sup>c</sup>	0.0072%	-	0.0009%	0.0014%	0.0062%	0.0075%	0.0117%	0.0173%	0.0246%	0.0306%	0.0075%
Males	0.0067%	-	-	-	0.0035%	0.0054%	0.0118%	0.0201%	0.0402%	0.0244%	0.0077%
Females	0.0076%	-	-	0.0014%	0.0088%	0.0095%	0.0116%	0.0150%	0.0135%	0.0348%	0.0074%
Hispanic <sup>d</sup>	0.0150%	-	-	0.0035%	0.0140%	0.0281%	0.0450%	0.0531%	0.0837%	0.1009%	0.0243%
Males	0.0105%	-	-	0.0017%	0.0057%	0.0176%	0.0325%	0.0558%	0.0922%	0.1349%	0.0217%
Females	0.0197%	-	-	0.0056%	0.0235%	0.0390%	0.0565%	0.0508%	0.0777%	0.0814%	0.0278%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XVII-11  
MESOTHELIOMAEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	2,668	713	298	203	165	3,631	4,161	+	+
Males	1,880	347	97	63	91	2,299	2,501	+	+
Females	788	366	201	140	74	1,332	1,660	+	+
White <sup>b</sup>	2,425	637	266	188	153	3,283	3,760	+	+
Males	1,736	306	81	48	79	2,100	2,260	+	+
Females	689	331	185	140	74	1,183	1,500	+	+
Black <sup>b</sup>	153	60	25	12	12	238	285	+	+
Males	82	36	13	12	12	131	167	+	+
Females	71	24	12	0	0	107	118	+	+
Asian/Pacific Islander <sup>c</sup>	26	9	+	+	+	36	+	+	+
Males	22	9	+	+	+	31	+	+	+
Females	4	0	+	+	+	5	+	+	+
Hispanic <sup>d</sup>	218	29	+	+	+	267	+	+	+
Males	137	24	+	+	+	166	+	+	+
Females	81	5	+	+	+	101	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0010%	-	-	-	0.0001%	0.0005%	0.0017%	0.0033%	0.0058%	0.0058%	0.0011%
Males	0.0014%	-	-	-	-	0.0005%	0.0020%	0.0047%	0.0103%	0.0122%	0.0017%
Females	0.0007%	-	-	-	0.0002%	0.0005%	0.0013%	0.0019%	0.0024%	0.0023%	0.0006%
White <sup>c</sup>	0.0012%	-	-	-	0.0002%	0.0005%	0.0018%	0.0037%	0.0065%	0.0066%	0.0012%
Males	0.0016%	-	-	-	-	0.0005%	0.0021%	0.0054%	0.0114%	0.0137%	0.0018%
Females	0.0008%	-	-	-	0.0002%	0.0006%	0.0015%	0.0021%	0.0028%	0.0027%	0.0007%
Black <sup>c</sup>	0.0005%	-	-	-	-	-	0.0015%	0.0028%	0.0034%	-	0.0007%
Males	0.0006%	-	-	-	-	-	-	-	-	-	0.0010%
Females	0.0004%	-	-	-	-	-	-	0.0035%	-	-	0.0005%
Asian/Pacific Islander <sup>c</sup>	0.0003%	-	-	-	-	-	-	-	-	-	0.0003%
Males	0.0006%	-	-	-	-	-	-	-	-	-	0.0007%
Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic <sup>d</sup>	0.0006%	-	-	-	-	0.0006%	0.0025%	0.0027%	0.0068%	0.0056%	0.0012%
Males	0.0008%	-	-	-	-	-	0.0034%	0.0048%	0.0100%	-	0.0017%
Females	0.0004%	-	-	-	-	-	0.0016%	-	0.0046%	-	0.0008%

- <sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
- <sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
- <sup>g h i</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table XVIII-16  
MYELOMA (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	36,315	11,045	3,880	1,484	494	50,825	53,382	330	53,712
Males	20,046	6,178	2,255	854	231	28,226	29,616	129	29,745
Females	16,269	4,867	1,625	630	263	22,599	23,766	201	23,967
White <sup>b</sup>	28,905	8,630	2,903	1,085	345	40,179	42,003	62	42,065
Males	16,372	5,025	1,752	628	162	22,975	23,988	0 <sup>i</sup>	23,988
Females	12,533	3,605	1,151	457	183	17,204	18,015	62 <sup>i</sup>	18,077
Black <sup>b</sup>	6,139	2,008	827	328	133	8,840	9,460	+	+
Males	3,040	927	437	192	65	4,330	4,661	+	+
Females	3,099	1,081	390	136	68	4,510	4,799	+	+
Asian/Pacific Islander <sup>c</sup>	765	229	+	+	+	1,070	+	+	+
Males	365	133	+	+	+	539	+	+	+
Females	400	96	+	+	+	531	+	+	+
Hispanic <sup>d</sup>	2,310	610	+	+	+	3,127	+	+	+
Males	1,256	341	+	+	+	1,722	+	+	+
Females	1,054	269	+	+	+	1,405	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0146%	-	-	0.0002%	0.0017%	0.0072%	0.0256%	0.0548%	0.0830%	0.0705%	0.0155%
Males	0.0162%	-	-	-	0.0024%	0.0086%	0.0313%	0.0673%	0.1039%	0.0836%	0.0190%
Females	0.0130%	-	-	0.0002%	0.0011%	0.0058%	0.0203%	0.0437%	0.0672%	0.0633%	0.0126%
White <sup>c</sup>	0.0144%	-	-	0.0001%	0.0016%	0.0069%	0.0235%	0.0525%	0.0785%	0.0660%	0.0146%
Males	0.0163%	-	-	-	0.0023%	0.0082%	0.0294%	0.0664%	0.1002%	0.0789%	0.0183%
Females	0.0125%	-	-	-	0.0008%	0.0055%	0.0178%	0.0397%	0.0617%	0.0591%	0.0115%
Black <sup>c</sup>	0.0211%	-	-	-	0.0040%	0.0122%	0.0510%	0.1047%	0.1585%	0.1241%	0.0294%
Males	0.0224%	-	-	-	0.0041%	0.0156%	0.0595%	0.1274%	0.1953%	0.1304%	0.0350%
Females	0.0200%	-	-	-	0.0039%	0.0093%	0.0439%	0.0869%	0.1344%	0.1211%	0.0253%
Asian/Pacific Islander <sup>c</sup>	0.0089%	-	-	-	0.0006%	0.0044%	0.0163%	0.0287%	0.0541%	0.0608%	0.0099%
Males	0.0092%	-	-	-	0.0013%	0.0048%	0.0184%	0.0264%	0.0631%	0.0807%	0.0114%
Females	0.0087%	-	-	-	-	0.0041%	0.0145%	0.0305%	0.0476%	0.0476%	0.0089%
Hispanic <sup>d</sup>	0.0071%	-	-	-	0.0018%	0.0067%	0.0207%	0.0484%	0.0814%	0.0619%	0.0141%
Males	0.0075%	-	-	-	0.0029%	0.0076%	0.0251%	0.0540%	0.0908%	0.0962%	0.0169%
Females	0.0067%	-	-	-	-	0.0057%	0.0166%	0.0438%	0.0747%	0.0424%	0.0119%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIX-20  
NON-HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	165,166	95,646	52,508	31,037	16,904	305,401	368,283	12,846	381,129
Males	86,010	49,465	27,027	16,066	8,691	158,561	190,649	6,767	197,416
Females	79,156	46,181	25,481	14,971	8,213	146,840	177,634	6,079	183,713
White <sup>b</sup>	146,983	85,106	47,343	28,235	15,426	272,139	329,596	13,018	342,614
Males	76,940	43,608	24,348	14,636	7,954	141,247	170,662	7,257	177,919
Females	70,043	41,498	22,995	13,599	7,472	130,892	158,934	5,761	164,695
Black <sup>b</sup>	11,589	6,714	3,331	1,839	1,007	21,259	24,849	941	25,790
Males	5,669	3,797	1,668	971	466	10,971	12,722	418	13,140
Females	5,920	2,917	1,663	868	541	10,288	12,127	523	12,650
Asian/Pacific Islander <sup>c</sup>	3,888	2,083	+	+	+	6,899	+	+	+
Males	2,004	1,076	+	+	+	3,539	+	+	+
Females	1,884	1,007	+	+	+	3,360	+	+	+
Hispanic <sup>d</sup>	9,816	4,920	+	+	+	16,972	+	+	+
Males	5,206	2,559	+	+	+	8,922	+	+	+
Females	4,610	2,361	+	+	+	8,050	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0801%	0.0024%	0.0092%	0.0148%	0.0302%	0.0644%	0.1319%	0.2425%	0.3634%	0.3565%	0.0838%
Males	0.0845%	0.0032%	0.0126%	0.0177%	0.0346%	0.0764%	0.1515%	0.2709%	0.4050%	0.4004%	0.0954%
Females	0.0759%	0.0015%	0.0056%	0.0117%	0.0256%	0.0524%	0.1133%	0.2171%	0.3320%	0.3325%	0.0737%
White <sup>c</sup>	0.0889%	0.0025%	0.0096%	0.0149%	0.0314%	0.0672%	0.1394%	0.2656%	0.3935%	0.3770%	0.0895%
Males	0.0931%	0.0034%	0.0131%	0.0176%	0.0361%	0.0796%	0.1606%	0.2961%	0.4330%	0.4225%	0.1014%
Females	0.0847%	0.0017%	0.0059%	0.0119%	0.0263%	0.0546%	0.1187%	0.2376%	0.3630%	0.3524%	0.0788%
Black <sup>c</sup>	0.0473%	0.0015%	0.0065%	0.0162%	0.0320%	0.0616%	0.1141%	0.1517%	0.1858%	0.1590%	0.0578%
Males	0.0518%	0.0021%	0.0098%	0.0189%	0.0366%	0.0752%	0.1372%	0.1618%	0.2163%	0.1809%	0.0673%
Females	0.0432%	-	0.0031%	0.0137%	0.0279%	0.0496%	0.0949%	0.1439%	0.1659%	0.1485%	0.0502%
Asian/Pacific Islander <sup>c</sup>	0.0509%	0.0019%	0.0103%	0.0121%	0.0184%	0.0408%	0.0825%	0.1359%	0.2449%	0.2698%	0.0549%
Males	0.0535%	0.0025%	0.0130%	0.0164%	0.0209%	0.0490%	0.0832%	0.1475%	0.2888%	0.3038%	0.0623%
Females	0.0485%	-	0.0073%	0.0079%	0.0161%	0.0334%	0.0819%	0.1262%	0.2133%	0.2472%	0.0489%
Hispanic <sup>d</sup>	0.0361%	0.0022%	0.0075%	0.0092%	0.0225%	0.0465%	0.0970%	0.1764%	0.2815%	0.2474%	0.0621%
Males	0.0367%	0.0027%	0.0105%	0.0112%	0.0261%	0.0532%	0.1083%	0.1813%	0.2936%	0.2430%	0.0665%
Females	0.0355%	0.0017%	0.0044%	0.0069%	0.0184%	0.0395%	0.0865%	0.1724%	0.2728%	0.2499%	0.0576%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XX-16  
 CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	84,371	53,189	34,017	23,121	14,774	166,010	217,906	17,950	235,856
Males	56,621	34,320	21,542	13,940	8,760	109,190	140,286	10,660	150,946
Females	27,750	18,869	12,475	9,181	6,014	56,820	77,620	7,290	84,910
White <sup>b</sup>	72,469	46,784	29,976	20,744	13,156	144,341	190,725	15,254	205,979
Males	49,197	30,423	19,313	12,735	8,049	96,013	124,432	9,152	133,584
Females	23,272	16,361	10,663	8,009	5,107	48,328	66,293	6,102	72,395
Black <sup>b</sup>	7,507	3,608	2,398	1,515	942	13,096	16,436	896	17,332
Males	4,671	2,189	1,250	759	348	7,876	9,384	346	9,730
Females	2,836	1,419	1,148	756	594	5,220	7,052	550	7,602
Asian/Pacific Islander <sup>c</sup>	2,873	1,819	+	+	+	5,550	+	+	+
Males	1,783	1,129	+	+	+	3,443	+	+	+
Females	1,090	690	+	+	+	2,107	+	+	+
Hispanic <sup>d</sup>	3,534	2,119	+	+	+	6,739	+	+	+
Males	2,286	1,297	+	+	+	4,172	+	+	+
Females	1,248	822	+	+	+	2,567	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0428%	0.0003%	0.0017%	0.0047%	0.0115%	0.0366%	0.0924%	0.1449%	0.1754%	0.1474%	0.0443%
Males	0.0573%	0.0002%	0.0017%	0.0044%	0.0117%	0.0499%	0.1379%	0.2164%	0.2605%	0.2236%	0.0641%
Females	0.0286%	0.0003%	0.0018%	0.0051%	0.0113%	0.0234%	0.0493%	0.0810%	0.1111%	0.1056%	0.0277%
White <sup>c</sup>	0.0453%	0.0003%	0.0017%	0.0045%	0.0104%	0.0348%	0.0927%	0.1522%	0.1847%	0.1551%	0.0452%
Males	0.0610%	0.0002%	0.0015%	0.0038%	0.0112%	0.0481%	0.1395%	0.2264%	0.2728%	0.2391%	0.0658%
Females	0.0296%	0.0003%	0.0020%	0.0053%	0.0097%	0.0212%	0.0471%	0.0839%	0.1168%	0.1097%	0.0276%
Black <sup>c</sup>	0.0293%	-	0.0019%	0.0044%	0.0094%	0.0338%	0.0901%	0.1240%	0.1236%	0.0810%	0.0366%
Males	0.0391%	-	0.0027%	0.0050%	0.0079%	0.0420%	0.1409%	0.1976%	0.1841%	0.1164%	0.0534%
Females	0.0204%	-	-	0.0038%	0.0106%	0.0266%	0.0481%	0.0663%	0.0840%	0.0640%	0.0238%
Asian/Pacific Islander <sup>c</sup>	0.0388%	-	0.0016%	0.0063%	0.0176%	0.0473%	0.0860%	0.1067%	0.1394%	0.1127%	0.0403%
Males	0.0498%	-	-	0.0075%	0.0166%	0.0655%	0.1170%	0.1547%	0.2084%	0.1551%	0.0554%
Females	0.0285%	-	0.0020%	0.0051%	0.0185%	0.0308%	0.0588%	0.0665%	0.0900%	0.0846%	0.0281%
Hispanic <sup>d</sup>	0.0138%	-	0.0009%	0.0024%	0.0061%	0.0184%	0.0451%	0.0746%	0.1097%	0.1038%	0.0247%
Males	0.0168%	-	0.0008%	0.0020%	0.0067%	0.0236%	0.0635%	0.1116%	0.1566%	0.1477%	0.0345%
Females	0.0105%	-	0.0011%	0.0028%	0.0055%	0.0129%	0.0280%	0.0439%	0.0761%	0.0786%	0.0169%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXI-12  
OVARY CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
 By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	54,533	29,287	22,242	18,878	14,397	102,197	148,412	24,353	172,765
White <sup>b</sup>	48,314	25,867	20,104	17,113	12,929	90,767	132,464	21,146	153,610
Black <sup>b</sup>	3,648	1,984	1,309	1,196	950	6,721	9,791	2,123	11,914
Asian/Pacific Islander <sup>c</sup>	1,871	978	+	+	+	3,408	+	+	+
Hispanic <sup>d</sup>	3,637	1,880	+	+	+	6,565	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup> All Ages	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0529%	0.0005%	0.0050%	0.0128%	0.0208%	0.0513%	0.1091%	0.1646%	0.1695%	0.1184%	0.0516%	
White <sup>c</sup>	0.0582%	0.0006%	0.0051%	0.0134%	0.0207%	0.0540%	0.1167%	0.1807%	0.1824%	0.1250%	0.0551%	
Black <sup>c</sup>	0.0277%	-	0.0041%	0.0085%	0.0171%	0.0336%	0.0572%	0.0936%	0.1044%	0.0952%	0.0321%	
Asian/Pacific Islander <sup>c</sup>	0.0449%	-	0.0056%	0.0137%	0.0238%	0.0497%	0.1019%	0.1202%	0.1290%	0.0677%	0.0438%	
Hispanic <sup>d</sup>	0.0281%	0.0007%	0.0058%	0.0133%	0.0170%	0.0365%	0.0870%	0.1209%	0.1401%	0.1065%	0.0414%	

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.  
 US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
- <sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
- <sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table XXII-16  
PANCREAS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	21,231	3,252	1,826	865	653	26,083	27,987	460	28,447
Males	10,156	1,570	792	353	253	12,431	13,235	188	13,423
Females	11,075	1,682	1,034	512	400	13,652	14,752	272	15,024
White <sup>b</sup>	17,925	2,607	1,485	588	555	21,870	23,289	373	23,662
Males	8,905	1,334	653	271	224	10,843	11,468	168	11,636
Females	9,020	1,273	832	317	331	11,027	11,821	205	12,026
Black <sup>b</sup>	2,456	380	227	225	67	2,990	3,381	216	3,597
Males	858	120	86	79	13	1,027	1,182	77	1,259
Females	1,598	260	141	146	54	1,963	2,199	139	2,338
Asian/Pacific Islander <sup>c</sup>	624	131	+	+	+	790	+	+	+
Males	285	51	+	+	+	347	+	+	+
Females	339	80	+	+	+	443	+	+	+
Hispanic <sup>d</sup>	988	231	+	+	+	1,354	+	+	+
Males	470	75	+	+	+	610	+	+	+
Females	518	156	+	+	+	744	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0075%	-	0.0002%	0.0004%	0.0012%	0.0046%	0.0120%	0.0275%	0.0411%	0.0366%	0.0080%
Males	0.0074%	-	-	0.0002%	0.0010%	0.0048%	0.0136%	0.0302%	0.0459%	0.0389%	0.0086%
Females	0.0076%	-	0.0003%	0.0006%	0.0014%	0.0044%	0.0106%	0.0251%	0.0375%	0.0353%	0.0074%
White <sup>c</sup>	0.0078%	-	0.0002%	0.0005%	0.0012%	0.0042%	0.0128%	0.0272%	0.0401%	0.0344%	0.0078%
Males	0.0079%	-	-	0.0003%	0.0010%	0.0050%	0.0148%	0.0308%	0.0455%	0.0363%	0.0088%
Females	0.0076%	-	-	0.0007%	0.0015%	0.0034%	0.0108%	0.0239%	0.0360%	0.0334%	0.0071%
Black <sup>c</sup>	0.0068%	-	-	-	0.0015%	0.0061%	0.0113%	0.0336%	0.0503%	0.0474%	0.0096%
Males	0.0054%	-	-	-	0.0021%	0.0040%	0.0113%	0.0298%	0.0549%	0.0213%	0.0083%
Females	0.0081%	-	-	-	-	0.0080%	0.0113%	0.0366%	0.0474%	0.0600%	0.0103%
Asian/Pacific Islander <sup>c</sup>	0.0068%	-	-	-	0.0009%	0.0046%	0.0077%	0.0251%	0.0400%	0.0499%	0.0076%
Males	0.0063%	-	-	-	-	0.0028%	0.0066%	0.0272%	0.0417%	0.0718%	0.0080%
Females	0.0072%	-	-	-	0.0012%	0.0061%	0.0086%	0.0233%	0.0388%	0.0354%	0.0074%
Hispanic <sup>d</sup>	0.0030%	-	0.0003%	-	0.0008%	0.0020%	0.0093%	0.0202%	0.0303%	0.0296%	0.0058%
Males	0.0025%	-	-	-	-	0.0016%	0.0100%	0.0197%	0.0325%	0.0317%	0.0059%
Females	0.0034%	-	-	-	0.0012%	0.0024%	0.0087%	0.0207%	0.0287%	0.0283%	0.0058%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXIII-12  
PROSTATE CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
 By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	942,369	618,932	358,847	73,754	20,826	1,889,390	2,020,336	4,153	2,024,489
White <sup>b</sup>	786,788	524,932	319,277	67,060	19,019	1,603,095	1,722,113	4,475	1,726,588
Black <sup>b</sup>	120,665	75,526	32,102	5,690	1,569	225,930	236,062	363	236,425
Asian/Pacific Islander <sup>c</sup>	15,620	8,169	+	+	+	27,161	+	+	+
Hispanic <sup>d</sup>	43,918	25,606	+	+	+	79,891	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.9320%	-	-	0.0002%	0.0011%	0.0630%	0.8682%	4.1465%	8.6580%	7.7814%	1.1823%
White <sup>c</sup>	0.9691%	-	-	-	0.0010%	0.0561%	0.8380%	4.1273%	8.6232%	7.4719%	1.1639%
Black <sup>c</sup>	1.0358%	-	-	-	0.0018%	0.1570%	1.7295%	7.0469%	12.8348%	10.2270%	1.8386%
Asian/Pacific Islander <sup>c</sup>	0.4695%	-	-	-	-	0.0170%	0.2626%	1.6149%	4.9266%	5.9176%	0.6398%
Hispanic <sup>d</sup>	0.3185%	-	-	-	-	0.0303%	0.4931%	2.9006%	7.0766%	6.7356%	0.9225%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXIV-16  
STOMACH CANCER (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	29,647	12,158	7,142	5,104	2,646	47,696	57,826	2,474	60,300
Males	17,872	7,150	3,489	2,648	1,492	27,931	33,280	1,428	34,708
Females	11,775	5,008	3,653	2,456	1,154	19,765	24,546	1,046	25,592
White <sup>b</sup>	22,358	9,101	5,322	3,884	2,129	35,811	43,675	1,996	45,671
Males	13,840	5,411	2,561	2,020	1,244	21,350	25,563	1,101	26,664
Females	8,518	3,690	2,761	1,864	885	14,461	18,112	895	19,007
Black <sup>b</sup>	4,441	1,842	921	792	266	7,056	8,407	325	8,732
Males	2,405	961	485	406	154	3,791	4,505	190	4,695
Females	2,036	881	436	386	112	3,265	3,902	135	4,037
Asian/Pacific Islander <sup>c</sup>	2,452	1,134	+	+	+	4,222	+	+	+
Males	1,371	664	+	+	+	2,390	+	+	+
Females	1,081	470	+	+	+	1,832	+	+	+
Hispanic <sup>d</sup>	3,503	1,340	+	+	+	5,461	+	+	+
Males	1,921	643	+	+	+	2,854	+	+	+
Females	1,582	697	+	+	+	2,607	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.0161%	-	0.0001%	0.0006%	0.0025%	0.0088%	0.0216%	0.0490%	0.0947%	0.1116%	0.0171%
Males	0.0188%	-	-	0.0006%	0.0028%	0.0098%	0.0263%	0.0687%	0.1308%	0.1529%	0.0229%
Females	0.0134%	-	0.0002%	0.0005%	0.0022%	0.0078%	0.0170%	0.0313%	0.0675%	0.0890%	0.0128%
White <sup>c</sup>	0.0135%	-	0.0001%	0.0005%	0.0023%	0.0070%	0.0169%	0.0412%	0.0734%	0.0880%	0.0137%
Males	0.0161%	-	-	0.0006%	0.0025%	0.0080%	0.0207%	0.0590%	0.1047%	0.1223%	0.0186%
Females	0.0109%	-	0.0002%	0.0005%	0.0020%	0.0060%	0.0132%	0.0248%	0.0493%	0.0694%	0.0098%
Black <sup>c</sup>	0.0156%	-	-	-	0.0026%	0.0140%	0.0349%	0.0576%	0.1140%	0.1283%	0.0219%
Males	0.0172%	-	-	-	0.0035%	0.0166%	0.0418%	0.0767%	0.1478%	0.1560%	0.0276%
Females	0.0141%	-	-	-	0.0018%	0.0117%	0.0292%	0.0427%	0.0918%	0.1149%	0.0178%
Asian/Pacific Islander <sup>c</sup>	0.0329%	-	-	-	0.0040%	0.0149%	0.0400%	0.0935%	0.2250%	0.2999%	0.0376%
Males	0.0383%	-	-	-	0.0043%	0.0156%	0.0509%	0.1301%	0.3022%	0.3789%	0.0490%
Females	0.0279%	-	-	-	0.0038%	0.0144%	0.0305%	0.0628%	0.1696%	0.2474%	0.0290%
Hispanic <sup>d</sup>	0.0118%	-	0.0003%	0.0007%	0.0036%	0.0123%	0.0306%	0.0601%	0.1305%	0.1742%	0.0240%
Males	0.0120%	-	-	0.0007%	0.0041%	0.0117%	0.0340%	0.0780%	0.1598%	0.2170%	0.0290%
Females	0.0116%	-	-	0.0007%	0.0030%	0.0129%	0.0274%	0.0452%	0.1096%	0.1498%	0.0205%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXV-12  
TESTIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>									
All Races <sup>b</sup>	38,541	34,082	31,208	25,262	19,329	97,489	158,516	19,002	177,518
White <sup>b</sup>	36,147	32,157	29,918	24,402	18,691	92,143	150,982	18,259	169,241
Black <sup>b</sup>	1,061	845	425	373	310	2,220	3,213	179	3,392
Asian/Pacific Islander <sup>c</sup>	617	481	+	+	+	1,443	+	+	+
Hispanic <sup>d</sup>	3,881	2,780	+	+	+	8,541	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0476%	0.0018%	0.0044%	0.0532%	0.1092%	0.0947%	0.0453%	0.0179%	0.0086%	0.0054%	0.0462%
White <sup>c</sup>	0.0573%	0.0019%	0.0048%	0.0652%	0.1342%	0.1129%	0.0529%	0.0210%	0.0102%	0.0062%	0.0556%
Black <sup>c</sup>	0.0108%	-	0.0030%	0.0082%	0.0232%	0.0252%	0.0150%	-	-	-	0.0109%
Asian/Pacific Islander <sup>c</sup>	0.0164%	0.0025%	0.0035%	0.0182%	0.0306%	0.0329%	0.0168%	0.0060%	-	-	0.0155%
Hispanic <sup>d</sup>	0.0319%	0.0023%	0.0041%	0.0468%	0.0751%	0.0516%	0.0211%	0.0096%	-	-	0.0297%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVI-16  
 THYROID CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	103,273	71,568	51,794	39,835	30,644	217,395	321,227	45,239	366,466
Males	23,545	15,823	11,984	8,661	6,597	49,430	72,214	10,668	82,882
Females	79,728	55,745	39,810	31,174	24,047	167,965	249,013	34,571	283,584
White <sup>b</sup>	90,228	63,138	46,255	35,635	27,746	191,304	284,802	40,931	325,733
Males	21,193	14,353	10,974	7,886	5,972	44,723	65,584	9,062	74,646
Females	69,035	48,785	35,281	27,749	21,774	146,581	219,218	31,869	251,087
Black <sup>b</sup>	6,167	3,912	2,770	1,832	1,484	12,450	17,305	1,864	19,169
Males	1,132	648	581	336	281	2,292	3,103	202	3,305
Females	5,035	3,264	2,189	1,496	1,203	10,158	14,202	1,662	15,864
Asian/Pacific Islander <sup>c</sup>	4,594	3,044	+	+	+	9,337	+	+	+
Males	884	523	+	+	+	1,703	+	+	+
Females	3,710	2,521	+	+	+	7,634	+	+	+
Hispanic <sup>d</sup>	8,933	5,698	+	+	+	17,898	+	+	+
Males	1,623	1,005	+	+	+	3,224	+	+	+
Females	7,310	4,693	+	+	+	14,674	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0572%	-	0.0036%	0.0301%	0.0729%	0.0975%	0.1063%	0.1050%	0.0872%	0.0612%	0.0573%
Males	0.0256%	-	0.0014%	0.0085%	0.0259%	0.0379%	0.0525%	0.0673%	0.0592%	0.0435%	0.0268%
Females	0.0882%	-	0.0060%	0.0528%	0.1215%	0.1566%	0.1572%	0.1387%	0.1083%	0.0709%	0.0868%
White <sup>c</sup>	0.0614%	-	0.0038%	0.0336%	0.0800%	0.1058%	0.1102%	0.1061%	0.0865%	0.0606%	0.0606%
Males	0.0282%	-	0.0015%	0.0094%	0.0283%	0.0419%	0.0571%	0.0695%	0.0614%	0.0454%	0.0287%
Females	0.0944%	-	0.0062%	0.0599%	0.1356%	0.1713%	0.1620%	0.1398%	0.1058%	0.0688%	0.0926%
Black <sup>c</sup>	0.0257%	-	0.0015%	0.0097%	0.0316%	0.0416%	0.0597%	0.0701%	0.0602%	0.0409%	0.0294%
Males	0.0100%	-	-	0.0022%	0.0117%	0.0131%	0.0258%	0.0379%	0.0390%	0.0177%	0.0129%
Females	0.0399%	-	0.0020%	0.0167%	0.0491%	0.0666%	0.0879%	0.0953%	0.0740%	0.0521%	0.0431%
Asian/Pacific Islander <sup>c</sup>	0.0596%	-	0.0040%	0.0282%	0.0668%	0.0927%	0.1140%	0.1189%	0.1042%	0.0795%	0.0588%
Males	0.0230%	-	-	0.0089%	0.0228%	0.0323%	0.0394%	0.0731%	0.0568%	0.0421%	0.0242%
Females	0.0939%	-	0.0077%	0.0473%	0.1084%	0.1475%	0.1794%	0.1573%	0.1382%	0.1045%	0.0894%
Hispanic <sup>d</sup>	0.0364%	-	0.0028%	0.0221%	0.0490%	0.0786%	0.0954%	0.1004%	0.0874%	0.0759%	0.0486%
Males	0.0125%	-	0.0009%	0.0056%	0.0141%	0.0225%	0.0409%	0.0588%	0.0433%	0.0421%	0.0194%
Females	0.0617%	-	0.0049%	0.0419%	0.0892%	0.1368%	0.1463%	0.1349%	0.1190%	0.0952%	0.0773%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVII-16  
URINARY BLADDER CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>g</sup>	Complete <sup>h</sup>
<b>Race/Sex</b>									
All Races <sup>b</sup>	185,870	124,783	82,465	51,924	30,594	379,619	489,252	22,538	511,790
Males	139,001	92,860	61,148	38,627	21,832	283,235	362,809	14,714	377,523
Females	46,869	31,923	21,317	13,297	8,762	96,384	126,443	7,824	134,267
White <sup>b</sup>	173,651	117,485	78,353	49,437	29,399	356,461	461,231	17,762	478,993
Males	130,508	87,561	58,097	36,808	21,016	266,730	342,878	11,379	354,257
Females	43,143	29,924	20,256	12,629	8,383	89,731	118,353	6,383	124,736
Black <sup>b</sup>	7,484	4,210	2,596	1,670	701	14,017	17,041	349	17,390
Males	4,824	2,963	1,874	1,244	438	9,473	11,601	281	11,882
Females	2,660	1,247	722	426	263	4,544	5,440	68	5,508
Asian/Pacific Islander <sup>c</sup>	2,461	1,424	+	+	+	4,623	+	+	+
Males	1,889	1,026	+	+	+	3,480	+	+	+
Females	572	398	+	+	+	1,143	+	+	+
Hispanic <sup>d</sup>	5,360	3,161	+	+	+	10,351	+	+	+
Males	4,080	2,432	+	+	+	7,858	+	+	+
Females	1,280	729	+	+	+	2,493	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<b>Race/Sex</b>											
All Races <sup>c</sup>	0.0894%	0.0002%	0.0002%	0.0017%	0.0060%	0.0247%	0.0963%	0.2968%	0.5997%	0.7140%	0.0967%
Males	0.1359%	0.0002%	0.0002%	0.0021%	0.0083%	0.0377%	0.1530%	0.4886%	1.0702%	1.4124%	0.1710%
Females	0.0439%	-	-	0.0012%	0.0036%	0.0118%	0.0425%	0.1250%	0.2446%	0.3314%	0.0418%
White <sup>c</sup>	0.1066%	0.0002%	-	0.0021%	0.0072%	0.0280%	0.1101%	0.3432%	0.6805%	0.7731%	0.1090%
Males	0.1611%	0.0003%	-	0.0026%	0.0097%	0.0417%	0.1721%	0.5592%	1.2062%	1.5513%	0.1917%
Females	0.0523%	-	-	0.0015%	0.0045%	0.0139%	0.0497%	0.1447%	0.2753%	0.3524%	0.0470%
Black <sup>c</sup>	0.0309%	-	-	-	0.0025%	0.0148%	0.0569%	0.1370%	0.2589%	0.3251%	0.0455%
Males	0.0448%	-	-	-	0.0035%	0.0255%	0.1003%	0.2209%	0.4541%	0.5256%	0.0772%
Females	0.0183%	-	-	-	0.0016%	0.0054%	0.0210%	0.0712%	0.1309%	0.2284%	0.0243%
Asian/Pacific Islander <sup>c</sup>	0.0367%	-	-	-	0.0022%	0.0106%	0.0321%	0.1001%	0.2567%	0.4424%	0.0430%
Males	0.0566%	-	-	-	0.0034%	0.0183%	0.0512%	0.1722%	0.4524%	0.7962%	0.0755%
Females	0.0181%	-	-	-	-	0.0036%	0.0153%	0.0398%	0.1163%	0.2072%	0.0191%
Hispanic <sup>d</sup>	0.0205%	-	-	0.0005%	0.0035%	0.0109%	0.0419%	0.1312%	0.2756%	0.3623%	0.0453%
Males	0.0302%	-	-	-	0.0043%	0.0161%	0.0662%	0.2320%	0.5135%	0.6914%	0.0823%
Females	0.0102%	-	-	-	0.0027%	0.0056%	0.0192%	0.0473%	0.1056%	0.1742%	0.0190%

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h</sup> (g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVIII-12  
 CHILDHOOD CANCER (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2004  
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 <sup>e</sup>	0 to <29 <sup>e</sup>	>=29 <sup>f</sup>	Complete <sup>g</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	58,085	48,407	42,984	36,298	30,462	141,268	237,628	+	+
Males	30,670	25,794	22,695	18,102	15,221	74,840	122,159	+	+
Females	27,415	22,613	20,289	18,196	15,241	66,428	115,469	+	+
White <sup>b</sup>	48,916	40,601	36,378	31,566	26,790	118,905	203,160	+	+
Males	26,047	21,547	19,463	15,744	13,526	63,344	105,005	+	+
Females	22,869	19,054	16,915	15,822	13,264	55,561	98,155	+	+
Black <sup>b</sup>	5,891	4,940	4,317	3,133	2,577	14,309	22,638	+	+
Males	2,885	2,625	2,151	1,500	1,184	7,228	11,032	+	+
Females	3,006	2,315	2,166	1,633	1,393	7,081	11,606	+	+
Asian/Pacific Islander <sup>c</sup>	1,915	1,614	+	+	+	4,627	+	+	+
Males	1,065	881	+	+	+	2,474	+	+	+
Females	850	733	+	+	+	2,153	+	+	+
Hispanic <sup>d</sup>	9,087	6,640	+	+	+	20,187	+	+	+
Males	4,949	3,445	+	+	+	10,934	+	+	+
Females	4,138	3,195	+	+	+	9,253	+	+	+

<sup>a</sup> US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f g</sup> (f) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <29 and >=29.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.