

Table XVII-9
MESOTHELIOMA

SEER INCIDENCE
AGE-ADJUSTED RATES AND TRENDS^a
By Race/Ethnicity and Sex

RACE/ETHNICITY	Rate 2000-2004 ^b Rate per 100,000 persons			Trend 1995-2004 ^c APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	1.1	2.1	0.4	-1.3*	-1.5*	-0.9
White	1.2	2.3	0.5	-0.8	-1.1	-0.3
White Hispanic ^d	1.1	1.9	0.5	-2.6	-4.1	-
White Non-Hispanic ^d	1.3	2.4	0.5	-0.7	-0.8	-0.9
Black	0.6	1.1	0.2	-5.0	-	-
Asian/Pacific Islander	0.4	0.6	0.2	-3.4	-	-
Amer Ind/Alaska Nat ^e	-	-	-	-	-	-
Hispanic ^d	1.1	1.8	0.5	-2.3	-3.9	-

- The APC is the Annual Percent Change over the time interval.
Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).
- ^c Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).
- ^d Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky.
- ^e Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.
- * The APC is significantly different from zero ($p < .05$).