

Table IV-5
FEMALE BREAST CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE^a RATES BY YEAR, RACE AND AGE

	<u>All Races, Females</u>			<u>White Females</u>			<u>Black Females</u>		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	105.1	40.6	273.9	107.4	40.7	281.9	93.7	42.6	227.5
1976	101.9	40.1	263.9	104.8	40.9	272.3	85.3	37.8	209.8
1977	100.8	39.1	262.4	103.4	39.4	271.0	87.0	37.6	216.4
1978	100.6	38.8	262.3	103.6	39.3	272.0	86.1	40.2	206.1
1979	102.1	38.0	270.0	104.7	38.6	278.0	86.8	37.7	215.3
1980	102.2	37.7	270.9	105.1	38.3	280.0	89.2	37.4	224.9
1981	106.3	39.2	282.0	109.9	40.1	292.6	94.2	38.9	239.3
1982	106.4	40.5	279.0	110.0	41.3	289.8	93.7	38.1	239.2
1983	111.1	40.3	296.7	114.5	40.9	307.2	103.4	41.2	266.0
1984	115.9	42.3	308.6	119.9	43.1	321.0	102.2	44.8	252.5
1985	124.2	44.3	333.2	127.9	44.6	346.0	111.5	45.4	284.4
1986	126.8	44.4	342.4	130.4	45.3	353.4	115.0	45.5	296.8
1987	134.4	45.5	367.3	140.3	46.5	385.7	109.3	43.1	282.7
1988	131.2	44.0	359.7	136.3	45.1	375.4	118.6	42.9	316.6
1989	127.1	43.4	346.4	132.3	44.4	362.3	104.8	42.2	268.9
1990	131.7	45.6	357.4	136.2	45.4	374.1	118.1	48.7	299.7
1991	133.7	47.1	360.3	138.7	47.1	378.6	117.4	50.9	291.4
1992	131.9	43.4	363.6	135.7	43.6	376.8	123.6	46.9	324.4
1993	129.1	42.8	355.1	133.1	43.3	368.4	119.5	45.2	314.2
1994	130.8	42.3	362.6	135.6	42.7	379.1	122.1	47.2	318.2
1995	132.4	43.1	366.3	137.4	43.7	382.9	123.8	45.4	329.0
1996	133.4	43.4	369.1	137.9	43.9	384.3	123.2	43.8	331.1
1997	137.6	43.4	384.2	142.5	44.0	400.3	124.2	44.0	334.0
1998	141.1	45.4	391.6	146.5	45.7	410.5	122.6	46.1	323.0
1999	140.9	43.6	395.8	146.6	44.9	412.9	124.9	43.4	338.6
2000	136.0	43.3	378.8	142.6	43.8	401.2	116.3	44.1	305.5
2001	137.7	43.4	384.6	143.9	44.3	404.7	113.8	39.1	309.5
2002	134.4	42.2	375.6	140.0	43.0	394.1	120.4	40.1	330.8
2003	125.2	42.6	341.4	129.9	43.4	356.3	121.4	42.3	328.4
2004	125.7	43.9	340.0	129.8	43.9	354.6	120.0	45.1	316.0
2005	123.7	42.8	335.5	128.6	43.4	351.5	113.3	43.1	297.0
1975-2005	124.6	42.8	338.7	128.6	43.4	351.9	112.2	43.4	292.2

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.