

Table XXVIII-3

AGE-ADJUSTED CANCER U.S. DEATH RATES^a, 1975-2005

By Primary Cancer Site

All Races, Males and Females

Year of Death

Age/Site	1975	1980	1985	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<u>Ages 0-14</u>																			
All Sites	4.9	4.3	3.6	3.1	3.1	3.0	3.0	2.8	2.7	2.6	2.6	2.4	2.5	2.5	2.5	2.6	2.5	2.5	2.4
Bone & Joint	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Brain & Other nervous	1.0	0.9	0.8	0.9	0.9	0.8	0.8	0.8	0.7	0.8	0.7	0.7	0.7	0.7	0.7	0.8	0.8	0.7	0.7
Hodgkin lymphoma	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Kidney & Renal pelvis	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Leukemia	2.1	1.7	1.4	1.1	1.1	1.1	1.0	1.0	1.0	0.9	0.9	0.8	0.8	0.8	0.8	0.7	0.8	0.7	0.7
Acute lymphocytic	1.2	0.9	0.8	0.5	0.5	0.5	0.4	0.4	0.5	0.4	0.4	0.4	0.3	0.4	0.3	0.4	0.4	0.3	0.3
Non-Hodgkin lymphoma	0.4	0.3	0.2	0.2	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Soft tissue	0.1	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.1
<u>Ages 0-19</u>																			
All Sites	5.1	4.6	3.8	3.4	3.4	3.3	3.3	3.1	3.0	2.9	2.9	2.7	2.8	2.8	2.8	2.8	2.7	2.7	2.7
Bone & Joint	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.3
Brain & Other nervous	0.9	0.8	0.8	0.8	0.8	0.7	0.8	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.7	0.7	0.7	0.7
Hodgkin lymphoma	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Kidney & Renal pelvis	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Leukemia	2.0	1.8	1.4	1.2	1.2	1.2	1.1	1.1	1.0	0.9	1.0	0.8	0.8	0.9	0.9	0.8	0.8	0.8	0.8
Acute lymphocytic	1.0	0.9	0.7	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3
Non-Hodgkin lymphoma	0.4	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Soft tissue	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic could not be calculated. Rate based on less than 16 cases for the time interval.