

Table 1.21  
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2006<sup>a</sup>  
Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			31-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 31 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 31 Years <sup>d</sup>
All Sites	Both Sexes	4,077,488	4,154,826	4,519,388	10,831,324	11,027,782
	Male	2,080,191	2,108,988	2,284,134	5,056,502	5,116,252
	Female	1,997,297	2,045,838	2,235,254	5,774,822	5,911,530
Oral Cavity & Pharynx	Both Sexes	88,149	100,786	104,275	229,332	251,715
	Male	59,730	67,492	69,596	148,389	161,197
	Female	28,419	33,294	34,679	80,943	90,518
Esophagus	Both Sexes	18,231	21,903	21,915	27,407	32,201
	Male	13,976	16,810	16,810	20,647	24,147
	Female	4,255	5,093	5,105	6,760	8,054
Stomach	Both Sexes	31,763	38,156	38,279	62,024	71,142
	Male	19,112	22,988	23,004	35,526	40,842
	Female	12,651	15,168	15,275	26,498	30,300
Colon & Rectum	Both Sexes	410,878	473,926	481,943	1,063,937	1,184,011
	Male	207,419	239,595	243,087	522,508	578,045
	Female	203,459	234,331	238,856	541,429	605,966
Liver & Intrahepatic Bile Duct	Both Sexes	18,570	20,914	20,914	24,536	27,242
	Male	13,017	14,616	14,616	16,411	18,199
	Female	5,553	6,298	6,298	8,125	9,043
Pancreas	Both Sexes	23,089	27,435	27,446	30,784	35,776
	Male	11,266	13,503	13,514	14,813	17,277
	Female	11,823	13,932	13,932	15,971	18,499
Larynx	Both Sexes	30,174	35,697	35,972	88,041	97,768
	Male	24,109	28,434	28,686	70,364	77,761
	Female	6,065	7,263	7,286	17,677	20,007
Lung & Bronchus	Both Sexes	209,050	260,966	266,149	354,138	426,254
	Male	98,835	124,555	126,720	165,363	199,378
	Female	110,215	136,411	139,429	188,775	226,876
Melanoma of the Skin	Both Sexes	233,089	260,265	269,851	714,792	765,882
	Male	122,409	139,106	145,296	353,155	381,997
	Female	110,680	121,159	124,555	361,637	383,885
Breast	Female	801,058	861,011	905,789	2,442,116	2,589,991
Cervix	Female	39,451	41,186	41,269	192,354	197,813
Corpus & Uterus, NOS	Female	146,768	165,629	165,699	518,293	563,072
Ovary <sup>f</sup>	Female	54,346	63,586	63,631	154,428	173,886

<sup>a</sup> U.S. 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER 9 registries and 1/1/2006 U.S. population estimates based on the average of 2005 and 2006 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 31 years (1975-2005)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2001-2005)  
For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2001; Melanoma in 2002.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 31-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2006.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 31-year limited duration prevalence. The 2001 breast cancer is counted for the breast 5-year and 31-year limited duration prevalence.

In method (e) the 2001 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2002 melanoma is counted for 5-year limited duration prevalence for melanoma.

<sup>f</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.21 - continued  
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2006<sup>a</sup>  
 Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			31-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 31 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 31 Years <sup>d</sup>
Prostate	Male	930,125	1,003,839	1,003,911	2,174,982	2,320,295
Testis	Male	39,842	40,565	41,075	172,591	174,841
Urinary Bladder	Both Sexes	188,227	231,571	235,384	508,818	586,875
	Male	140,439	173,578	176,622	376,871	433,882
	Female	47,788	57,993	58,762	131,947	152,993
Kidney & Renal Pelvis	Both Sexes	111,278	134,837	135,889	254,925	295,615
	Male	66,844	81,868	82,613	150,001	175,152
	Female	44,434	52,969	53,276	104,924	120,463
Brain & Nervous System	Both Sexes	39,632	41,635	41,953	108,661	111,711
	Male	21,221	22,407	22,579	58,031	59,668
	Female	18,411	19,228	19,374	50,630	52,043
Thyroid	Both Sexes	120,479	130,868	131,194	367,333	387,569
	Male	26,876	30,552	30,586	81,388	87,358
	Female	93,603	100,316	100,608	285,945	300,211
Hodgkin Lymphoma	Both Sexes	35,015	36,820	36,831	145,147	148,461
	Male	18,420	19,309	19,320	74,925	76,593
	Female	16,595	17,511	17,511	70,222	71,868
Non-Hodgkin Lymphoma	Both Sexes	178,216	206,582	208,073	407,962	452,723
	Male	92,687	108,381	109,127	211,075	233,951
	Female	85,529	98,201	98,946	196,887	218,772
Myeloma	Both Sexes	39,615	45,364	45,401	58,905	66,529
	Male	21,534	24,854	24,878	32,370	36,858
	Female	18,081	20,510	20,523	26,535	29,671
Leukemia	Both Sexes	96,231	108,820	108,981	225,790	245,225
	Male	55,777	63,265	63,346	127,972	139,195
	Female	40,454	45,555	45,635	97,818	106,030
Acute Lymphocytic Leukemia	Both Sexes	15,118	15,404	15,404	55,306	55,720
	Male	8,660	8,708	8,708	30,447	30,533
	Female	6,458	6,696	6,696	24,859	25,187
Childhood (Ages 0-19)	Both Sexes	60,231	60,330	60,634	259,480	259,966
	Male	32,416	32,472	32,581	133,367	133,580
	Female	27,815	27,858	28,053	126,113	126,386
Kaposi Sarcoma	Both Sexes	6,695	7,138	7,138	21,168	22,209
	Male	6,215	6,544	6,544	19,850	20,703
	Female	480	594	594	1,318	1,506
Mesothelioma	Both Sexes	2,774	3,453	3,453	4,194	5,000
	Male	2,022	2,510	2,510	2,603	3,185
	Female	752	943	943	1,591	1,815

<sup>a</sup> U.S. 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER 9 registries and 1/1/2006 U.S. population estimates based on the average of 2005 and 2006 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 31 years (1975-2005)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2001-2005)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2001; Melanoma in 2002.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 31-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2006.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 31-year limited duration prevalence. The 2001 breast cancer is counted for the breast 5-year and 31-year limited duration prevalence.

In method (e) the 2001 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2002 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.22  
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2006<sup>a</sup>  
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	5,168,889	17,479	39,370	75,164	144,948	331,732	720,142	1,194,219	2,645,836
Females	6,216,003	14,832	33,518	81,003	213,538	597,268	1,130,077	1,333,585	2,812,181
Oral Cavity & Pharynx									
Males	157,250	23	492	1,318	3,276	14,407	37,003	40,095	60,638
Females	87,223	101	501	1,668	3,489	8,222	15,686	18,110	39,446
Esophagus									
Males	20,742	0	0	17	130	1,166	4,085	6,436	8,907
Females	6,861	0	0	12	50	303	909	1,550	4,037
Stomach									
Males	36,725	0	5	86	649	2,426	5,764	8,519	19,274
Females	27,397	11	23	91	491	1,896	3,966	4,776	16,143
Colon & Rectum									
Males	536,944	11	46	961	5,116	22,614	69,355	116,858	321,983
Females	567,158	0	115	1,190	4,881	20,121	58,565	96,908	385,378
Liver & Intrahep									
Males	16,484	415	403	334	382	1,391	5,714	3,821	4,025
Females	8,445	424	329	318	328	843	1,678	1,630	2,896
Pancreas									
Males	14,979	0	35	69	255	1,190	3,251	4,246	5,933
Females	16,201	0	44	114	434	1,327	2,738	4,093	7,452
Larynx									
Males	73,244	0	0	11	232	2,513	10,523	19,979	39,986
Females	18,326	0	0	46	132	1,174	3,010	4,737	9,227
Lung & Bronchus									
Males	171,522	45	46	345	1,118	6,174	24,269	49,093	90,433
Females	193,474	0	32	372	1,441	8,500	25,665	51,689	105,775
Melanoma of the Skin									
Males	367,925	72	692	5,217	16,598	46,626	83,425	86,050	129,247
Females	390,763	84	980	10,591	32,264	68,250	93,268	73,818	111,509

<sup>a</sup> U.S. 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2006 U.S. population estimates based on the average of 2005 and 2006 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.22 - continued  
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2006<sup>a</sup>  
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	13,132	0	0	23	75	542	2,078	3,322	7,093
Females	2,533,193	0	104	2,335	35,060	206,965	500,267	614,770	1,173,691
Cervix									
Females	248,166	11	92	2,502	18,006	43,602	57,740	48,934	77,278
Corpus & Uterus, NOS									
Females	572,603	0	55	543	4,924	23,372	76,468	124,512	342,730
Ovary <sup>d</sup>									
Females	176,007	45	891	3,473	7,019	20,547	37,874	39,554	66,605
Prostate									
Males	2,177,975	45	92	69	349	16,618	187,672	552,329	1,420,801
Urinary Bladder									
Males	388,965	23	143	717	2,390	12,796	41,822	85,952	245,121
Females	138,531	33	68	414	1,120	4,555	13,137	26,664	92,539
Kidney & Renal Pelvis									
Males	156,166	1,489	2,424	2,363	4,140	13,996	31,742	40,436	59,577
Females	109,902	1,565	2,655	2,595	3,885	10,133	17,631	24,085	47,353
Hodgkin Lymphoma									
Males	82,450	129	2,058	8,923	16,247	22,575	18,055	9,050	5,412
Females	77,557	101	1,477	10,135	16,381	21,291	15,813	6,811	5,549
Non-Hodgkin Lymphoma									
Males	217,143	663	3,555	6,616	12,138	26,624	42,988	48,865	75,694
Females	202,390	360	1,529	3,938	7,562	18,662	35,213	43,070	92,057
Myeloma									
Males	32,483	0	0	51	549	2,188	6,759	9,771	13,165
Females	26,720	0	0	23	222	1,661	4,882	7,103	12,830
Leukemia									
Males	130,746	6,418	11,449	10,565	9,207	10,991	17,345	23,135	41,637
Females	101,110	5,023	9,672	9,353	7,857	7,821	11,507	14,611	35,265
Acute Lymphocytic Leuk									
Males	32,255	5,310	9,965	8,164	5,276	1,976	781	520	263
Females	26,601	4,202	8,131	6,920	4,273	1,699	654	442	281

<sup>a</sup> U.S. 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2006 U.S. population estimates based on the average of 2005 and 2006 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.18  
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	4,077,488	2,756,143	1,827,790	1,067,357	628,995	408,422	8,921,919	10,831,324	553,568	11,384,892
	Males	2,080,191	1,358,523	873,389	402,083	208,914	117,265	4,419,347	5,056,502	112,387	5,168,889
	Females	1,997,297	1,397,620	954,401	665,274	420,081	291,157	4,502,572	5,774,822	441,181	6,216,003
White <sup>b</sup>	Both Sexes	3,520,272	2,420,229	1,629,494	969,935	573,796	373,186	7,806,111	9,546,459	489,074	10,035,533
	Males	1,788,398	1,184,235	776,817	368,293	192,182	108,601	3,847,393	4,433,351	104,730	4,538,081
	Females	1,731,874	1,235,994	852,677	601,642	381,614	264,585	3,958,718	5,113,108	384,344	5,497,452
Black <sup>b</sup>	Both Sexes	383,691	229,026	136,791	64,711	36,625	22,580	765,969	877,013	31,078	908,091
	Males	205,824	124,429	70,194	22,917	10,825	5,090	406,933	440,037	3,170	443,207
	Females	177,867	104,597	66,597	41,794	25,800	17,490	359,036	436,976	27,908	464,884
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	106,876	61,997	36,247	+	+	+	210,102	+	+	+
	Males	46,735	25,186	13,666	+	+	+	87,178	+	+	+
	Females	60,141	36,811	22,581	+	+	+	122,924	+	+	+
Hispanic <sup>d</sup>	Both Sexes	235,971	137,977	84,667	+	+	+	471,035	+	+	+
	Males	113,967	64,417	36,535	+	+	+	219,442	+	+	+
	Females	122,004	73,560	48,132	+	+	+	251,593	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	2.6125%	0.0793%	0.1565%	0.2705%	0.6644%	1.5625%	3.7265%	8.1929%	13.5686%	14.5035%	2.7072%
	Males	2.6003%	0.0849%	0.1644%	0.2517%	0.5169%	1.0685%	3.0327%	8.9806%	18.2203%	21.3291%	3.0961%
	Females	2.6243%	0.0734%	0.1482%	0.2906%	0.8161%	2.0556%	4.3885%	7.4821%	9.9786%	10.7168%	2.4852%
White <sup>c</sup>	Both Sexes	2.8537%	0.0852%	0.1709%	0.3002%	0.7370%	1.6543%	3.8634%	8.5564%	14.2545%	14.9548%	2.8361%
	Males	2.8095%	0.0914%	0.1792%	0.2805%	0.5835%	1.1472%	3.1082%	9.1880%	18.7633%	21.6733%	3.1916%
	Females	2.8979%	0.0787%	0.1621%	0.3221%	0.9016%	2.1772%	4.6067%	7.9711%	10.6939%	11.2707%	2.6436%
Black <sup>c</sup>	Both Sexes	1.8858%	0.0545%	0.1071%	0.1695%	0.4686%	1.2564%	3.6290%	8.5309%	13.3964%	12.4397%	2.5470%
	Males	2.1113%	0.0529%	0.1120%	0.1484%	0.3284%	0.8707%	3.6866%	11.5962%	21.5351%	21.3968%	3.4812%
	Females	1.6811%	0.0562%	0.1020%	0.1897%	0.5912%	1.5952%	3.5812%	6.1444%	7.9816%	8.1016%	1.9708%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1.6845%	0.0639%	0.1182%	0.1711%	0.3946%	1.1504%	2.6637%	4.9710%	8.2974%	10.3437%	1.7781%
	Males	1.4974%	0.0723%	0.1278%	0.1540%	0.2500%	0.5948%	1.6187%	4.6846%	10.5931%	14.9192%	1.8270%
	Females	1.8603%	0.0550%	0.1079%	0.1882%	0.5316%	1.6602%	3.5809%	5.2137%	6.6146%	7.3284%	1.7941%
Hispanic <sup>d</sup>	Both Sexes	1.0593%	0.0750%	0.1452%	0.1889%	0.4534%	1.1045%	2.5183%	5.5502%	9.4464%	10.0358%	1.8825%
	Males	0.9496%	0.0811%	0.1499%	0.1800%	0.3352%	0.6388%	1.7880%	5.9158%	13.2025%	15.1482%	2.1375%
	Females	1.1753%	0.0686%	0.1402%	0.1995%	0.5903%	1.5988%	3.2110%	5.2405%	6.7129%	7.0643%	1.7685%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 3.18  
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	39,632	22,564	18,270	13,267	8,008	5,917	83,594	108,661	14,026	122,687
	Males	21,221	11,940	10,182	6,986	4,240	2,979	45,088	58,031	6,401	64,432
	Females	18,411	10,624	8,088	6,281	3,768	2,938	38,506	50,630	7,625	58,255
White <sup>b</sup>	Both Sexes	35,067	19,983	16,065	11,742	7,089	5,361	73,757	96,220	7,226	103,446
	Males	19,035	10,677	8,903	6,072	3,772	2,712	40,080	51,619	4,000 <sup>i</sup>	55,619
	Females	16,032	9,306	7,162	5,670	3,317	2,649	33,677	44,601	3,226 <sup>i</sup>	47,827
Black <sup>b</sup>	Both Sexes	2,947	1,678	1,409	1,105	687	419	6,413	8,325	+	+
	Males	1,396	851	811	658	363	190	3,282	4,300	+	+
	Females	1,551	827	598	447	324	229	3,131	4,025	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1,062	568	427	+	+	+	2,121	+	+	+
	Males	527	262	237	+	+	+	1,062	+	+	+
	Females	535	306	190	+	+	+	1,059	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,873	2,236	1,413	+	+	+	7,807	+	+	+
	Males	2,145	1,120	714	+	+	+	4,140	+	+	+
	Females	1,728	1,116	699	+	+	+	3,667	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0249%	0.0137%	0.0279%	0.0239%	0.0244%	0.0303%	0.0303%	0.0281%	0.0241%	0.0138%	0.0248%
	Males	0.0270%	0.0143%	0.0299%	0.0249%	0.0260%	0.0327%	0.0341%	0.0315%	0.0266%	0.0155%	0.0268%
	Females	0.0229%	0.0131%	0.0259%	0.0228%	0.0228%	0.0279%	0.0267%	0.0250%	0.0223%	0.0128%	0.0229%
White <sup>c</sup>	Both Sexes	0.0279%	0.0149%	0.0311%	0.0261%	0.0283%	0.0346%	0.0341%	0.0312%	0.0264%	0.0142%	0.0278%
	Males	0.0303%	0.0159%	0.0333%	0.0270%	0.0300%	0.0369%	0.0385%	0.0357%	0.0287%	0.0168%	0.0300%
	Females	0.0256%	0.0139%	0.0288%	0.0251%	0.0264%	0.0323%	0.0298%	0.0270%	0.0246%	0.0128%	0.0256%
Black <sup>c</sup>	Both Sexes	0.0156%	0.0094%	0.0184%	0.0183%	0.0150%	0.0142%	0.0185%	0.0160%	0.0181%	0.0133%	0.0156%
	Males	0.0162%	0.0085%	0.0202%	0.0188%	0.0165%	0.0144%	0.0200%	0.0152%	0.0227%	-	0.0162%
	Females	0.0150%	0.0104%	0.0164%	0.0177%	0.0136%	0.0140%	0.0173%	0.0165%	0.0151%	0.0181%	0.0151%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0145%	0.0094%	0.0182%	0.0148%	0.0122%	0.0175%	0.0153%	0.0159%	0.0141%	0.0088%	0.0145%
	Males	0.0148%	0.0101%	0.0185%	0.0156%	0.0119%	0.0197%	0.0149%	0.0116%	0.0164%	0.0111%	0.0149%
	Females	0.0141%	0.0087%	0.0179%	0.0141%	0.0125%	0.0155%	0.0157%	0.0195%	0.0124%	0.0072%	0.0142%
Hispanic <sup>d</sup>	Both Sexes	0.0176%	0.0112%	0.0202%	0.0129%	0.0160%	0.0247%	0.0256%	0.0235%	0.0215%	0.0131%	0.0188%
	Males	0.0180%	0.0117%	0.0208%	0.0130%	0.0173%	0.0248%	0.0264%	0.0236%	0.0207%	0.0175%	0.0194%
	Females	0.0171%	0.0108%	0.0196%	0.0129%	0.0145%	0.0247%	0.0247%	0.0235%	0.0221%	0.0106%	0.0183%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 4.23  
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	806,524	648,228	436,782	291,728	158,753	96,802	1,959,764	2,454,992	91,333	2,546,325
	Males	5,466	3,352	1,709	1,182	736	406	10,895	12,876	256	13,132
	Females	801,058	644,876	435,073	290,546	158,017	96,396	1,948,869	2,442,116	91,077	2,533,193
White <sup>b</sup>	Both Sexes	700,075	574,925	391,645	266,134	144,496	88,457	1,728,591	2,180,598	81,191	2,261,789
	Males	4,614	2,984	1,464	1,052	674	393	9,379	11,206	95 <sup>i</sup>	11,301
	Females	695,461	571,941	390,181	265,082	143,822	88,064	1,719,212	2,169,392	81,096	2,250,488
Black <sup>b</sup>	Both Sexes	73,021	49,094	31,491	18,074	10,411	5,834	157,985	188,832	+	+
	Males	681	270	161	120	50	13	1,160	1,295	+	+
	Females	72,340	48,824	31,330	17,954	10,361	5,821	156,825	187,537	6,396	193,933
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	25,304	17,399	9,956	+	+	+	54,134	+	+	+
	Males	106	45	48	+	+	+	203	+	+	+
	Females	25,198	17,354	9,908	+	+	+	53,931	+	+	+
Hispanic <sup>d</sup>	Both Sexes	45,325	29,708	18,423	+	+	+	96,279	+	+	+
	Males	176	101	27	+	+	+	320	+	+	+
	Females	45,149	29,607	18,396	+	+	+	95,959	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.5773%	-	0.0002%	0.0057%	0.0814%	0.4260%	1.1244%	1.9183%	2.4904%	2.8998%	0.5857%
	Males	0.0063%	-	-	-	0.0004%	0.0023%	0.0093%	0.0242%	0.0412%	0.0550%	0.0075%
	Females	1.1379%	-	0.0004%	0.0116%	0.1647%	0.8490%	2.1882%	3.6274%	4.3807%	4.4781%	1.0696%
White <sup>c</sup>	Both Sexes	0.6330%	-	0.0002%	0.0053%	0.0808%	0.4294%	1.1649%	2.0317%	2.6780%	3.0991%	0.6162%
	Males	0.0068%	-	-	-	0.0002%	0.0022%	0.0091%	0.0241%	0.0435%	0.0591%	0.0077%
	Females	1.2596%	-	0.0004%	0.0110%	0.1671%	0.8699%	2.3024%	3.8921%	4.7584%	4.7661%	1.1365%
Black <sup>c</sup>	Both Sexes	0.3941%	-	-	0.0078%	0.0962%	0.4214%	0.9915%	1.6239%	2.0368%	2.3114%	0.5063%
	Males	0.0061%	-	-	-	-	0.0036%	0.0132%	0.0415%	0.0448%	0.0473%	0.0093%
	Females	0.7463%	-	-	0.0150%	0.1797%	0.7883%	1.8049%	2.8559%	3.3621%	3.4078%	0.8709%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.4222%	-	-	0.0059%	0.0726%	0.4106%	0.9699%	1.3902%	1.5756%	1.6294%	0.4308%
	Males	0.0034%	-	-	-	-	-	0.0067%	0.0106%	0.0230%	0.0266%	0.0040%
	Females	0.8155%	-	-	0.0118%	0.1404%	0.7865%	1.8153%	2.5596%	2.7136%	2.6856%	0.7801%
Hispanic <sup>d</sup>	Both Sexes	0.2191%	-	-	0.0046%	0.0554%	0.2939%	0.7737%	1.2684%	1.6101%	1.7243%	0.3870%
	Males	0.0013%	-	-	-	-	0.0010%	0.0048%	0.0081%	0.0152%	0.0200%	0.0029%
	Females	0.4495%	-	-	0.0100%	0.1191%	0.6046%	1.5030%	2.3360%	2.7707%	2.7149%	0.7060%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 5.14  
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	39,451	38,535	34,731	30,783	23,101	21,728	119,712	192,354	55,812	248,166
White <sup>b</sup>	31,048	30,600	28,499	25,641	19,137	17,691	95,858	155,925	46,577	202,502
Black <sup>b</sup>	5,677	5,244	4,021	3,443	3,036	3,056	15,832	25,053	+	+
Asian/Pacific Islander <sup>c</sup>	2,055	1,752	1,476	+	+	+	5,523	+	+	+
Hispanic <sup>d</sup>	8,083	6,813	5,682	+	+	+	21,602	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0822%	-	0.0004%	0.0108%	0.0835%	0.1594%	0.1706%	0.1513%	0.1363%	0.0801%	0.0795%	
White <sup>c</sup>	0.0833%	-	0.0004%	0.0115%	0.0951%	0.1698%	0.1681%	0.1396%	0.1191%	0.0614%	0.0802%	
Black <sup>c</sup>	0.0731%	-	-	0.0111%	0.0647%	0.1336%	0.1645%	0.1878%	0.1745%	0.1595%	0.0795%	
Asian/Pacific Islander <sup>c</sup>	0.0797%	-	-	0.0024%	0.0337%	0.1155%	0.1800%	0.1909%	0.2144%	0.1769%	0.0756%	
Hispanic <sup>d</sup>	0.0999%	-	-	0.0102%	0.0926%	0.2510%	0.3054%	0.3142%	0.2999%	0.1769%	0.1352%	

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 6.25  
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	410,878	263,947	164,750	114,290	67,504	37,785	866,951	1,063,937	40,165	1,104,102
	Males	207,419	131,746	80,079	56,122	30,135	15,092	432,913	522,508	14,436	536,944
	Females	203,459	132,201	84,671	58,168	37,369	22,693	434,038	541,429	25,729	567,158
White <sup>b</sup>	Both Sexes	348,874	230,188	144,309	100,824	60,586	33,731	747,384	922,890	35,761	958,651
	Males	178,455	115,778	70,634	50,227	27,060	13,523	376,890	457,468	12,995	470,463
	Females	170,419	114,410	73,675	50,597	33,526	20,208	370,494	465,422	22,766	488,188
Black <sup>b</sup>	Both Sexes	42,135	22,725	13,952	9,736	4,877	2,702	81,186	96,390	2,818	99,208
	Males	18,615	9,998	6,066	3,967	1,949	942	35,792	41,606	925	42,531
	Females	23,520	12,727	7,886	5,769	2,928	1,760	45,394	54,784	1,893	56,677
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	14,085	7,932	4,648	+	+	+	27,330	+	+	+
	Males	7,267	4,156	2,376	+	+	+	14,143	+	+	+
	Females	6,818	3,776	2,272	+	+	+	13,187	+	+	+
Hispanic <sup>d</sup>	Both Sexes	22,853	11,699	6,712	+	+	+	42,336	+	+	+
	Males	12,138	6,353	3,420	+	+	+	22,391	+	+	+
	Females	10,715	5,346	3,292	+	+	+	19,945	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Race	Sex	Age Specific (Crude)								Age-Adjusted <sup>f</sup>		
			All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
	All Races <sup>c</sup>	Both Sexes	0.2543%	-	0.0003%	0.0045%	0.0223%	0.0872%	0.3008%	0.7479%	1.5006%	2.1764%	0.2665%
		Males	0.2585%	-	0.0002%	0.0042%	0.0228%	0.0915%	0.3376%	0.8840%	1.8231%	2.4866%	0.3115%
		Females	0.2502%	-	0.0004%	0.0048%	0.0218%	0.0828%	0.2658%	0.6250%	1.2517%	2.0043%	0.2318%
	White <sup>c</sup>	Both Sexes	0.2675%	-	0.0004%	0.0048%	0.0221%	0.0845%	0.2877%	0.7362%	1.5137%	2.2204%	0.2659%
		Males	0.2714%	-	0.0003%	0.0044%	0.0227%	0.0883%	0.3280%	0.8782%	1.8385%	2.5317%	0.3118%
		Females	0.2636%	-	0.0005%	0.0052%	0.0215%	0.0806%	0.2481%	0.6046%	1.2572%	2.0496%	0.2296%
	Black <sup>c</sup>	Both Sexes	0.1990%	-	-	0.0031%	0.0228%	0.0971%	0.3738%	0.8986%	1.5369%	1.9071%	0.2811%
		Males	0.1888%	-	-	0.0036%	0.0227%	0.1013%	0.3968%	0.9702%	1.7753%	2.0846%	0.3102%
		Females	0.2083%	-	-	0.0026%	0.0229%	0.0935%	0.3546%	0.8428%	1.3782%	1.8211%	0.2619%
	Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.2283%	-	-	0.0035%	0.0218%	0.0903%	0.3087%	0.6857%	1.3575%	1.9678%	0.2479%
		Males	0.2425%	-	-	0.0038%	0.0228%	0.0975%	0.3321%	0.8247%	1.6999%	2.3356%	0.2950%
		Females	0.2148%	-	-	0.0032%	0.0208%	0.0836%	0.2880%	0.5679%	1.1064%	1.7255%	0.2125%
	Hispanic <sup>d</sup>	Both Sexes	0.0948%	-	0.0004%	0.0033%	0.0170%	0.0651%	0.2203%	0.5743%	1.0688%	1.4087%	0.1909%
		Males	0.0967%	-	-	0.0035%	0.0168%	0.0655%	0.2493%	0.6907%	1.4129%	1.5978%	0.2301%
		Females	0.0928%	-	-	0.0032%	0.0172%	0.0647%	0.1927%	0.4756%	0.8184%	1.2988%	0.1614%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 7.14  
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	146,768	115,083	87,371	64,913	49,514	44,765	363,887	518,293	54,310	572,603
White <sup>b</sup>	129,587	104,717	81,008	60,777	46,939	42,510	329,098	475,036	47,688	522,724
Black <sup>b</sup>	10,229	5,402	3,518	2,365	1,326	1,061	19,600	24,112	1,222	25,334
Asian/Pacific Islander <sup>c</sup>	4,567	2,923	1,827	+	+	+	9,575	+	+	+
Hispanic <sup>d</sup>	8,788	5,171	3,534	+	+	+	18,162	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.2066%	-	0.0002%	0.0029%	0.0249%	0.0992%	0.3384%	0.7640%	0.9301%	0.8471%	0.1956%	
White <sup>c</sup>	0.2358%	-	-	0.0027%	0.0248%	0.0992%	0.3606%	0.8414%	1.0452%	0.9392%	0.2135%	
Black <sup>c</sup>	0.0918%	-	-	0.0029%	0.0208%	0.0622%	0.1588%	0.4604%	0.5839%	0.3942%	0.1123%	
Asian/Pacific Islander <sup>c</sup>	0.1436%	-	-	0.0036%	0.0269%	0.1246%	0.3501%	0.4895%	0.4485%	0.3630%	0.1363%	
Hispanic <sup>d</sup>	0.0842%	-	-	0.0037%	0.0325%	0.0994%	0.2531%	0.5176%	0.5162%	0.4307%	0.1319%	

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 8.18  
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	18,231	5,609	2,279	802	338	148	26,395	27,407	196	27,603
	Males	13,976	4,100	1,704	560	212	96	19,977	20,647	95	20,742
	Females	4,255	1,509	575	242	126	52	6,418	6,760	101	6,861
White <sup>b</sup>	Both Sexes	16,146	4,919	2,016	621	274	121	23,280	24,096	160	24,256
	Males	12,668	3,681	1,497	441	184	96	17,998	18,568	89	18,657
	Females	3,478	1,238	519	180	90	25	5,282	5,528	71	5,599
Black <sup>b</sup>	Both Sexes	1,583	539	203	151	65	27	2,391	2,568	51	2,619
	Males	906	312	154	101	28	0	1,412	1,501	13	1,514
	Females	677	227	49	50	37	27	979	1,067	38	1,105
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	330	105	52	+	+	+	490	+	+	+
	Males	258	71	42	+	+	+	372	+	+	+
	Females	72	34	10	+	+	+	118	+	+	+
Hispanic <sup>d</sup>	Both Sexes	710	170	111	+	+	+	1,003	+	+	+
	Males	585	130	91	+	+	+	818	+	+	+
	Females	125	40	20	+	+	+	185	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0073%	-	-	-	0.0004%	0.0027%	0.0111%	0.0290%	0.0452%	0.0348%	0.0077%
	Males	0.0111%	-	-	-	0.0007%	0.0043%	0.0186%	0.0492%	0.0743%	0.0611%	0.0128%
	Females	0.0036%	-	-	-	0.0002%	0.0011%	0.0039%	0.0108%	0.0227%	0.0201%	0.0034%
White <sup>c</sup>	Both Sexes	0.0082%	-	-	-	0.0005%	0.0030%	0.0119%	0.0306%	0.0483%	0.0360%	0.0081%
	Males	0.0126%	-	-	-	0.0009%	0.0048%	0.0203%	0.0527%	0.0807%	0.0625%	0.0138%
	Females	0.0038%	-	-	-	-	0.0011%	0.0035%	0.0101%	0.0227%	0.0215%	0.0034%
Black <sup>c</sup>	Both Sexes	0.0055%	-	-	-	-	0.0024%	0.0130%	0.0323%	0.0437%	0.0214%	0.0075%
	Males	0.0068%	-	-	-	-	0.0030%	0.0166%	0.0458%	0.0590%	0.0374%	0.0105%
	Females	0.0044%	-	-	-	-	0.0018%	0.0100%	0.0218%	0.0336%	0.0136%	0.0054%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0042%	-	-	-	-	0.0013%	0.0046%	0.0153%	0.0261%	0.0347%	0.0046%
	Males	0.0065%	-	-	-	-	0.0024%	0.0088%	0.0256%	0.0432%	0.0652%	0.0080%
	Females	0.0020%	-	-	-	-	-	-	0.0065%	0.0137%	0.0146%	0.0020%
Hispanic <sup>d</sup>	Both Sexes	0.0023%	-	-	-	0.0005%	0.0015%	0.0051%	0.0161%	0.0254%	0.0263%	0.0044%
	Males	0.0036%	-	-	-	0.0009%	0.0025%	0.0091%	0.0293%	0.0471%	0.0491%	0.0081%
	Females	0.0009%	-	-	-	-	-	-	0.0048%	0.0096%	0.0131%	0.0016%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 9.18  
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	35,015	30,332	25,848	22,424	17,530	12,078	96,298	145,147	14,860	160,007
	Males	18,420	15,663	13,204	11,289	9,268	6,129	50,028	74,925	7,525	82,450
	Females	16,595	14,669	12,644	11,135	8,262	5,949	46,270	70,222	7,335	77,557
White <sup>b</sup>	Both Sexes	29,840	26,587	23,228	20,651	16,308	11,385	84,252	129,866	16,583	146,449
	Males	15,842	13,727	11,920	10,455	8,474	5,674	44,025	66,993	8,573	75,566
	Females	13,998	12,860	11,308	10,196	7,834	5,711	40,227	62,873	8,010	70,883
Black <sup>b</sup>	Both Sexes	3,876	2,969	2,202	1,366	1,020	568	9,465	12,016	1,094	13,110
	Males	1,834	1,507	1,086	624	696	375	4,589	6,143	742	6,885
	Females	2,042	1,462	1,116	742	324	193	4,876	5,873	352	6,225
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	752	431	277	+	+	+	1,498	+	+	+
	Males	399	241	124	+	+	+	778	+	+	+
	Females	353	190	153	+	+	+	720	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,143	2,344	1,584	+	+	+	7,435	+	+	+
	Males	1,767	1,338	904	+	+	+	4,187	+	+	+
	Females	1,376	1,006	680	+	+	+	3,248	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0280%	0.0006%	0.0089%	0.0382%	0.0515%	0.0419%	0.0296%	0.0248%	0.0247%	0.0166%	0.0278%
	Males	0.0296%	0.0008%	0.0104%	0.0347%	0.0508%	0.0458%	0.0346%	0.0306%	0.0303%	0.0199%	0.0295%
	Females	0.0263%	0.0004%	0.0074%	0.0421%	0.0522%	0.0380%	0.0249%	0.0195%	0.0204%	0.0148%	0.0264%
White <sup>c</sup>	Both Sexes	0.0315%	0.0007%	0.0097%	0.0435%	0.0593%	0.0471%	0.0329%	0.0275%	0.0278%	0.0184%	0.0314%
	Males	0.0333%	0.0009%	0.0111%	0.0388%	0.0579%	0.0510%	0.0383%	0.0339%	0.0340%	0.0222%	0.0330%
	Females	0.0297%	0.0006%	0.0082%	0.0486%	0.0607%	0.0430%	0.0276%	0.0216%	0.0228%	0.0163%	0.0301%
Black <sup>c</sup>	Both Sexes	0.0238%	-	0.0076%	0.0290%	0.0448%	0.0394%	0.0291%	0.0246%	0.0189%	0.0133%	0.0244%
	Males	0.0244%	-	0.0080%	0.0266%	0.0420%	0.0446%	0.0345%	0.0280%	0.0221%	0.0204%	0.0258%
	Females	0.0232%	-	0.0072%	0.0314%	0.0472%	0.0348%	0.0245%	0.0218%	0.0167%	0.0099%	0.0234%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0101%	-	0.0051%	0.0173%	0.0167%	0.0124%	0.0085%	0.0076%	0.0081%	0.0041%	0.0096%
	Males	0.0109%	-	0.0080%	0.0171%	0.0172%	0.0128%	0.0081%	0.0112%	0.0101%	-	0.0105%
	Females	0.0093%	-	0.0020%	0.0174%	0.0162%	0.0120%	0.0089%	0.0046%	0.0066%	-	0.0087%
Hispanic <sup>d</sup>	Both Sexes	0.0164%	0.0010%	0.0090%	0.0221%	0.0250%	0.0232%	0.0212%	0.0268%	0.0265%	0.0150%	0.0181%
	Males	0.0180%	0.0013%	0.0114%	0.0211%	0.0252%	0.0263%	0.0265%	0.0307%	0.0371%	0.0276%	0.0209%
	Females	0.0147%	0.0007%	0.0065%	0.0233%	0.0248%	0.0198%	0.0162%	0.0236%	0.0187%	0.0078%	0.0158%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 10.13  
Kaposi SarcomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	6,695	6,311	5,712	1,642	545	231	19,214	21,168	+	+
	Males	6,215	5,970	5,428	1,578	511	115	18,087	19,850	+	+
	Females	480	341	284	64	34	116	1,127	1,318	+	+
White <sup>b</sup>	Both Sexes	4,701	4,681	4,824	1,351	447	172	14,600	16,208	+	+
	Males	4,338	4,457	4,599	1,310	424	79	13,789	15,240	+	+
	Females	363	224	225	41	23	93	811	968	+	+
Black <sup>b</sup>	Both Sexes	1,662	1,254	601	112	52	12	3,541	3,694	+	+
	Males	1,563	1,181	542	112	52	12	3,310	3,463	+	+
	Females	99	73	59	0	0	0	231	231	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	105	83	48	+	+	+	238	+	+	+
	Males	100	80	48	+	+	+	230	+	+	+
	Females	5	3	0	+	+	+	8	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,235	1,094	871	+	+	+	3,254	+	+	+
	Males	1,169	1,050	837	+	+	+	3,105	+	+	+
	Females	66	44	34	+	+	+	149	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0074%	-	-	0.0008%	0.0080%	0.0197%	0.0132%	0.0070%	0.0078%	0.0136%	0.0073%
	Males	0.0142%	-	-	0.0014%	0.0156%	0.0389%	0.0265%	0.0136%	0.0149%	0.0259%	0.0143%
	Females	0.0007%	-	-	-	0.0002%	0.0006%	0.0004%	0.0012%	0.0023%	0.0067%	0.0006%
White <sup>c</sup>	Both Sexes	0.0074%	-	-	0.0006%	0.0070%	0.0201%	0.0132%	0.0072%	0.0085%	0.0139%	0.0072%
	Males	0.0141%	-	-	0.0011%	0.0134%	0.0390%	0.0263%	0.0136%	0.0158%	0.0266%	0.0140%
	Females	0.0007%	-	-	-	-	0.0005%	0.0003%	0.0012%	0.0028%	0.0070%	0.0006%
Black <sup>c</sup>	Both Sexes	0.0098%	-	-	0.0019%	0.0187%	0.0272%	0.0160%	0.0085%	0.0045%	0.0111%	0.0103%
	Males	0.0197%	-	-	0.0036%	0.0397%	0.0567%	0.0335%	0.0177%	0.0097%	-	0.0212%
	Females	0.0009%	-	-	-	-	0.0013%	-	-	-	0.0115%	0.0010%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0018%	-	-	-	0.0014%	0.0033%	0.0045%	0.0023%	0.0026%	0.0044%	0.0017%
	Males	0.0035%	-	-	-	0.0029%	0.0067%	0.0096%	0.0044%	0.0060%	0.0091%	0.0036%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic <sup>d</sup>	Both Sexes	0.0075%	-	-	0.0009%	0.0108%	0.0233%	0.0148%	0.0114%	0.0145%	0.0387%	0.0101%
	Males	0.0139%	-	-	0.0017%	0.0199%	0.0449%	0.0298%	0.0218%	0.0297%	0.0702%	0.0193%
	Females	0.0007%	-	-	-	-	-	-	0.0026%	-	0.0204%	0.0012%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 11.18  
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	111,278	61,228	38,793	22,513	12,707	7,477	216,737	254,925	11,143	266,068
	Males	66,844	35,728	22,828	12,604	7,573	3,858	128,580	150,001	6,165	156,166
	Females	44,434	25,500	15,965	9,909	5,134	3,619	88,157	104,924	4,978	109,902
White <sup>b</sup>	Both Sexes	95,119	52,432	33,890	20,212	11,285	6,680	186,269	220,503	12,716	233,219
	Males	57,212	31,202	20,208	11,467	6,887	3,527	111,481	131,035	5,866	136,901
	Females	37,907	21,230	13,682	8,745	4,398	3,153	74,788	89,468	6,850	96,318
Black <sup>b</sup>	Both Sexes	11,588	6,692	3,755	1,875	1,183	638	22,510	25,768	1,931	27,699
	Males	6,809	3,303	1,956	897	545	248	12,317	13,791	686	14,477
	Females	4,779	3,389	1,799	978	638	390	10,193	11,977	1,245	13,222
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,432	1,132	592	+	+	+	4,221	+	+	+
	Males	1,531	682	356	+	+	+	2,609	+	+	+
	Females	901	450	236	+	+	+	1,612	+	+	+
Hispanic <sup>d</sup>	Both Sexes	8,659	4,167	2,352	+	+	+	15,590	+	+	+
	Males	5,105	2,367	1,323	+	+	+	8,951	+	+	+
	Females	3,554	1,800	1,029	+	+	+	6,639	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0628%	0.0067%	0.0088%	0.0024%	0.0102%	0.0391%	0.1085%	0.2188%	0.2962%	0.2651%	0.0644%
	Males	0.0762%	0.0064%	0.0079%	0.0023%	0.0107%	0.0473%	0.1446%	0.2971%	0.4003%	0.3777%	0.0852%
	Females	0.0497%	0.0070%	0.0098%	0.0024%	0.0096%	0.0309%	0.0740%	0.1482%	0.2159%	0.2025%	0.0475%
White <sup>c</sup>	Both Sexes	0.0686%	0.0070%	0.0094%	0.0023%	0.0108%	0.0411%	0.1119%	0.2321%	0.3129%	0.2762%	0.0677%
	Males	0.0829%	0.0070%	0.0085%	0.0023%	0.0112%	0.0492%	0.1478%	0.3113%	0.4234%	0.4001%	0.0892%
	Females	0.0542%	0.0071%	0.0103%	0.0024%	0.0105%	0.0328%	0.0767%	0.1587%	0.2257%	0.2083%	0.0498%
Black <sup>c</sup>	Both Sexes	0.0534%	0.0075%	0.0101%	0.0035%	0.0113%	0.0426%	0.1295%	0.2161%	0.2921%	0.2801%	0.0680%
	Males	0.0633%	0.0060%	0.0084%	0.0038%	0.0128%	0.0518%	0.1827%	0.3005%	0.3764%	0.3793%	0.0895%
	Females	0.0443%	0.0090%	0.0120%	0.0032%	0.0099%	0.0345%	0.0852%	0.1504%	0.2360%	0.2320%	0.0524%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0339%	0.0029%	0.0028%	-	0.0050%	0.0213%	0.0600%	0.1168%	0.1759%	0.1576%	0.0357%
	Males	0.0431%	0.0019%	0.0027%	-	0.0049%	0.0286%	0.0832%	0.1722%	0.2444%	0.2101%	0.0490%
	Females	0.0254%	0.0040%	0.0030%	-	0.0050%	0.0145%	0.0397%	0.0698%	0.1257%	0.1230%	0.0251%
Hispanic <sup>d</sup>	Both Sexes	0.0350%	0.0059%	0.0064%	0.0016%	0.0092%	0.0327%	0.0980%	0.2062%	0.3084%	0.2791%	0.0621%
	Males	0.0391%	0.0059%	0.0044%	0.0014%	0.0093%	0.0365%	0.1236%	0.2709%	0.4336%	0.3727%	0.0803%
	Females	0.0307%	0.0059%	0.0084%	0.0018%	0.0091%	0.0287%	0.0738%	0.1513%	0.2172%	0.2247%	0.0479%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 12.18  
Cancer of the Larynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	30,174	21,452	15,138	10,127	7,016	3,697	69,307	88,041	3,529	91,570
	Males	24,109	16,972	12,146	8,124	5,592	3,040	55,235	70,364	2,880	73,244
	Females	6,065	4,480	2,992	2,003	1,424	657	14,072	17,677	649	18,326
White <sup>b</sup>	Both Sexes	25,135	18,490	13,143	8,652	6,406	3,347	58,950	75,507	2,841	78,348
	Males	20,026	14,728	10,534	7,076	5,162	2,730	47,017	60,553	2,310	62,863
	Females	5,109	3,762	2,609	1,576	1,244	617	11,933	14,954	531	15,485
Black <sup>b</sup>	Both Sexes	4,254	2,395	1,708	1,223	528	253	8,650	10,460	244	10,704
	Males	3,391	1,760	1,396	885	368	229	6,764	8,108	208	8,316
	Females	863	635	312	338	160	24	1,886	2,352	36	2,388
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	505	320	201	+	+	+	1,050	+	+	+
	Males	451	278	174	+	+	+	923	+	+	+
	Females	54	42	27	+	+	+	127	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,457	960	591	+	+	+	3,137	+	+	+
	Males	1,232	798	493	+	+	+	2,630	+	+	+
	Females	225	162	98	+	+	+	507	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0191%	-	-	0.0002%	0.0009%	0.0064%	0.0284%	0.0805%	0.1178%	0.0869%	0.0200%
	Males	0.0311%	-	-	-	0.0010%	0.0091%	0.0464%	0.1401%	0.2174%	0.2001%	0.0365%
	Females	0.0074%	-	-	0.0003%	0.0007%	0.0037%	0.0112%	0.0267%	0.0409%	0.0241%	0.0071%
White <sup>c</sup>	Both Sexes	0.0207%	-	-	0.0002%	0.0010%	0.0069%	0.0289%	0.0828%	0.1225%	0.0884%	0.0207%
	Males	0.0332%	-	-	-	0.0011%	0.0094%	0.0465%	0.1428%	0.2214%	0.2032%	0.0371%
	Females	0.0082%	-	-	0.0004%	0.0009%	0.0042%	0.0116%	0.0271%	0.0443%	0.0254%	0.0076%
Black <sup>c</sup>	Both Sexes	0.0205%	-	-	-	0.0009%	0.0075%	0.0473%	0.1231%	0.1533%	0.0972%	0.0278%
	Males	0.0341%	-	-	-	-	0.0114%	0.0816%	0.2201%	0.3131%	0.2470%	0.0535%
	Females	0.0082%	-	-	-	-	0.0041%	0.0187%	0.0477%	0.0470%	0.0247%	0.0100%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0088%	-	-	-	-	0.0025%	0.0093%	0.0310%	0.0631%	0.0681%	0.0097%
	Males	0.0160%	-	-	-	-	0.0049%	0.0168%	0.0610%	0.1295%	0.1506%	0.0198%
	Females	0.0021%	-	-	-	-	-	0.0028%	0.0056%	0.0144%	0.0138%	0.0021%
Hispanic <sup>d</sup>	Both Sexes	0.0068%	-	-	-	0.0009%	0.0038%	0.0143%	0.0536%	0.0862%	0.0701%	0.0138%
	Males	0.0111%	-	-	-	0.0010%	0.0047%	0.0249%	0.1049%	0.1731%	0.1557%	0.0271%
	Females	0.0023%	-	-	-	-	0.0028%	0.0043%	0.0100%	0.0230%	0.0204%	0.0039%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.26  
LeukemiaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	96,231	52,982	34,771	20,681	12,556	7,615	188,645	225,790	6,066	231,856
	Males	55,777	30,114	19,599	11,630	6,635	3,703	108,018	127,972	2,774	130,746
	Females	40,454	22,868	15,172	9,051	5,921	3,912	80,627	97,818	3,292	101,110
White <sup>b</sup>	Both Sexes	84,636	47,391	31,462	18,751	11,467	7,015	167,739	201,609	5,119	206,728
	Males	49,605	26,911	17,900	10,695	6,088	3,475	96,770	115,154	2,014	117,168
	Females	35,031	20,480	13,562	8,056	5,379	3,540	70,969	86,455	3,105	89,560
Black <sup>b</sup>	Both Sexes	6,915	3,168	1,861	1,018	585	356	12,148	13,959	677	14,636
	Males	3,571	1,635	894	463	323	114	6,180	7,033	308	7,341
	Females	3,344	1,533	967	555	262	242	5,968	6,926	369	7,295
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,180	1,258	820	+	+	+	4,422	+	+	+
	Males	1,200	736	428	+	+	+	2,451	+	+	+
	Females	980	522	392	+	+	+	1,971	+	+	+
Hispanic <sup>d</sup>	Both Sexes	8,258	4,586	3,263	+	+	+	16,552	+	+	+
	Males	4,751	2,481	1,852	+	+	+	9,380	+	+	+
	Females	3,507	2,105	1,411	+	+	+	7,172	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0565%	0.0294%	0.0501%	0.0216%	0.0210%	0.0307%	0.0604%	0.1297%	0.2050%	0.2279%	0.0582%
	Males	0.0656%	0.0321%	0.0537%	0.0236%	0.0238%	0.0369%	0.0761%	0.1700%	0.2779%	0.3075%	0.0724%
	Females	0.0475%	0.0266%	0.0464%	0.0195%	0.0181%	0.0245%	0.0453%	0.0933%	0.1488%	0.1837%	0.0464%
White <sup>c</sup>	Both Sexes	0.0634%	0.0322%	0.0562%	0.0241%	0.0222%	0.0321%	0.0644%	0.1437%	0.2277%	0.2500%	0.0637%
	Males	0.0735%	0.0353%	0.0597%	0.0264%	0.0252%	0.0388%	0.0809%	0.1877%	0.3083%	0.3389%	0.0793%
	Females	0.0533%	0.0291%	0.0524%	0.0217%	0.0190%	0.0252%	0.0481%	0.1029%	0.1640%	0.2012%	0.0507%
Black <sup>c</sup>	Both Sexes	0.0304%	0.0136%	0.0235%	0.0146%	0.0170%	0.0223%	0.0459%	0.0734%	0.1338%	0.1220%	0.0357%
	Males	0.0339%	0.0142%	0.0244%	0.0154%	0.0177%	0.0272%	0.0579%	0.0962%	0.1654%	0.1690%	0.0433%
	Females	0.0272%	0.0129%	0.0226%	0.0138%	0.0165%	0.0180%	0.0360%	0.0556%	0.1128%	0.0993%	0.0302%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0312%	0.0286%	0.0428%	0.0137%	0.0179%	0.0249%	0.0331%	0.0548%	0.0675%	0.0610%	0.0325%
	Males	0.0357%	0.0314%	0.0484%	0.0150%	0.0213%	0.0278%	0.0390%	0.0636%	0.0855%	0.0859%	0.0381%
	Females	0.0269%	0.0257%	0.0368%	0.0124%	0.0147%	0.0223%	0.0279%	0.0474%	0.0543%	0.0446%	0.0276%
Hispanic <sup>d</sup>	Both Sexes	0.0375%	0.0310%	0.0590%	0.0233%	0.0208%	0.0268%	0.0404%	0.0801%	0.0942%	0.1097%	0.0432%
	Males	0.0409%	0.0340%	0.0614%	0.0274%	0.0233%	0.0312%	0.0472%	0.0879%	0.1254%	0.1374%	0.0498%
	Females	0.0339%	0.0279%	0.0565%	0.0185%	0.0180%	0.0222%	0.0339%	0.0735%	0.0715%	0.0936%	0.0375%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 13.27  
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	15,118	11,509	9,569	8,046	6,316	4,302	37,886	55,306	3,550	58,856
	Males	8,660	6,303	5,361	4,490	3,194	2,206	21,220	30,447	1,808	32,255
	Females	6,458	5,206	4,208	3,556	3,122	2,096	16,666	24,859	1,742	26,601
White <sup>b</sup>	Both Sexes	13,060	10,055	8,321	7,067	5,743	3,997	32,962	48,661	+	+
	Males	7,590	5,471	4,777	3,984	2,928	2,086	18,662	27,047	+	+
	Females	5,470	4,584	3,544	3,083	2,815	1,911	14,300	21,614	+	+
Black <sup>b</sup>	Both Sexes	1,025	778	696	497	278	143	2,565	3,439	+	+
	Males	517	432	291	243	147	55	1,274	1,708	+	+
	Females	508	346	405	254	131	88	1,291	1,731	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	691	455	370	+	+	+	1,604	+	+	+
	Males	384	288	180	+	+	+	896	+	+	+
	Females	307	167	190	+	+	+	708	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,777	2,536	1,937	+	+	+	8,519	+	+	+
	Males	2,150	1,358	1,121	+	+	+	4,795	+	+	+
	Females	1,627	1,178	816	+	+	+	3,724	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0132%	0.0249%	0.0429%	0.0128%	0.0051%	0.0029%	0.0028%	0.0028%	0.0018%	0.0014%	0.0134%
	Males	0.0149%	0.0270%	0.0465%	0.0148%	0.0063%	0.0033%	0.0030%	0.0029%	0.0019%	0.0019%	0.0148%
	Females	0.0114%	0.0226%	0.0392%	0.0108%	0.0038%	0.0025%	0.0026%	0.0027%	0.0016%	0.0012%	0.0120%
White <sup>c</sup>	Both Sexes	0.0146%	0.0276%	0.0490%	0.0149%	0.0056%	0.0032%	0.0030%	0.0031%	0.0018%	0.0015%	0.0152%
	Males	0.0165%	0.0302%	0.0528%	0.0171%	0.0072%	0.0035%	0.0032%	0.0033%	0.0021%	0.0020%	0.0168%
	Females	0.0127%	0.0249%	0.0451%	0.0124%	0.0039%	0.0029%	0.0027%	0.0028%	0.0015%	0.0013%	0.0135%
Black <sup>c</sup>	Both Sexes	0.0064%	0.0098%	0.0174%	0.0070%	0.0029%	0.0013%	-	-	-	-	0.0058%
	Males	0.0068%	0.0103%	0.0176%	0.0075%	0.0021%	0.0015%	-	-	-	-	0.0058%
	Females	0.0061%	0.0093%	0.0172%	0.0066%	0.0035%	-	-	-	-	-	0.0058%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0102%	0.0240%	0.0342%	0.0070%	0.0042%	0.0028%	0.0025%	0.0031%	-	-	0.0111%
	Males	0.0119%	0.0248%	0.0398%	0.0085%	0.0047%	0.0033%	0.0028%	-	-	-	0.0123%
	Females	0.0087%	0.0231%	0.0283%	0.0055%	0.0037%	0.0023%	0.0023%	0.0042%	-	-	0.0098%
Hispanic <sup>d</sup>	Both Sexes	0.0193%	0.0269%	0.0521%	0.0144%	0.0064%	0.0043%	0.0049%	0.0051%	0.0038%	-	0.0162%
	Males	0.0210%	0.0294%	0.0548%	0.0172%	0.0071%	0.0055%	0.0041%	-	-	-	0.0175%
	Females	0.0175%	0.0243%	0.0494%	0.0110%	0.0057%	0.0031%	0.0055%	0.0068%	-	-	0.0148%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 14.18  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	18,570	3,431	1,169	708	477	181	23,311	24,536	393	24,929
	Males	13,017	2,146	674	252	240	71	15,903	16,411	73	16,484
	Females	5,553	1,285	495	456	237	110	7,408	8,125	320	8,445
White <sup>b</sup>	Both Sexes	13,534	2,580	884	562	373	119	17,088	18,053	374	18,427
	Males	9,546	1,546	516	171	174	46	11,640	12,010	139 <sup>i</sup>	12,149
	Females	3,988	1,034	368	391	199	73	5,448	6,043	235 <sup>i</sup>	6,278
Black <sup>b</sup>	Both Sexes	2,552	244	83	95	48	45	2,915	3,068	+	+
	Males	1,707	162	26	47	26	22	1,918	1,990	+	+
	Females	845	82	57	48	22	23	997	1,078	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,247	536	143	+	+	+	2,938	+	+	+
	Males	1,590	382	100	+	+	+	2,078	+	+	+
	Females	657	154	43	+	+	+	860	+	+	+
Hispanic <sup>d</sup>	Both Sexes	2,501	403	199	+	+	+	3,124	+	+	+
	Males	1,742	236	136	+	+	+	2,130	+	+	+
	Females	759	167	63	+	+	+	994	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0093%	0.0022%	0.0013%	0.0007%	0.0014%	0.0054%	0.0218%	0.0306%	0.0401%	0.0243%	0.0094%
	Males	0.0129%	0.0023%	0.0013%	0.0007%	0.0019%	0.0079%	0.0349%	0.0451%	0.0574%	0.0367%	0.0138%
	Females	0.0058%	0.0020%	0.0012%	0.0006%	0.0010%	0.0030%	0.0093%	0.0176%	0.0267%	0.0174%	0.0056%
White <sup>c</sup>	Both Sexes	0.0073%	0.0023%	0.0014%	0.0006%	0.0011%	0.0038%	0.0179%	0.0217%	0.0272%	0.0176%	0.0071%
	Males	0.0100%	0.0023%	0.0016%	0.0006%	0.0013%	0.0051%	0.0281%	0.0324%	0.0392%	0.0254%	0.0102%
	Females	0.0046%	0.0022%	0.0013%	0.0005%	0.0010%	0.0025%	0.0078%	0.0118%	0.0176%	0.0133%	0.0043%
Black <sup>c</sup>	Both Sexes	0.0072%	-	-	0.0012%	0.0015%	0.0049%	0.0251%	0.0321%	0.0296%	0.0190%	0.0087%
	Males	0.0101%	-	-	-	0.0023%	0.0071%	0.0432%	0.0467%	0.0397%	0.0275%	0.0131%
	Females	0.0046%	-	-	0.0017%	-	0.0030%	0.0101%	0.0208%	0.0228%	0.0149%	0.0054%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0236%	0.0030%	0.0010%	-	0.0030%	0.0158%	0.0427%	0.0865%	0.1279%	0.0800%	0.0248%
	Males	0.0342%	0.0032%	-	-	0.0050%	0.0264%	0.0726%	0.1295%	0.1899%	0.1236%	0.0383%
	Females	0.0137%	0.0027%	-	-	-	0.0061%	0.0165%	0.0501%	0.0825%	0.0512%	0.0138%
Hispanic <sup>d</sup>	Both Sexes	0.0071%	0.0021%	0.0013%	0.0006%	0.0008%	0.0052%	0.0279%	0.0436%	0.0512%	0.0337%	0.0120%
	Males	0.0094%	0.0025%	0.0014%	0.0006%	0.0011%	0.0083%	0.0442%	0.0660%	0.0706%	0.0372%	0.0173%
	Females	0.0047%	0.0018%	0.0013%	-	-	0.0020%	0.0124%	0.0247%	0.0371%	0.0317%	0.0074%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 15.24  
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	209,050	69,333	38,216	20,390	10,614	5,704	321,698	354,138	10,858	364,996
	Males	98,835	31,266	17,584	9,575	4,868	2,761	149,826	165,363	6,159	171,522
	Females	110,215	38,067	20,632	10,815	5,746	2,943	171,872	188,775	4,699	193,474
White <sup>b</sup>	Both Sexes	179,953	60,907	33,688	18,225	9,281	5,065	279,134	307,813	5,180	312,993
	Males	84,397	26,889	15,254	8,362	4,190	2,419	128,428	141,881	2,489	144,370
	Females	95,556	34,018	18,434	9,863	5,091	2,646	150,706	165,932	2,691	168,623
Black <sup>b</sup>	Both Sexes	21,763	6,180	3,492	1,697	1,076	472	31,842	34,803	482	35,285
	Males	10,526	3,132	1,745	974	524	266	15,613	17,265	259	17,524
	Females	11,237	3,048	1,747	723	552	206	16,229	17,538	223	17,761
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	5,727	1,557	660	+	+	+	8,044	+	+	+
	Males	3,052	859	379	+	+	+	4,342	+	+	+
	Females	2,675	698	281	+	+	+	3,702	+	+	+
Hispanic <sup>d</sup>	Both Sexes	6,284	1,751	965	+	+	+	9,167	+	+	+
	Males	3,314	906	457	+	+	+	4,751	+	+	+
	Females	2,970	845	508	+	+	+	4,416	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Race	Sex	Age Specific (Crude)								Age-Adjusted <sup>f</sup>		
			All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
	All Races <sup>c</sup>	Both Sexes	0.0907%	0.0001%	0.0002%	0.0015%	0.0051%	0.0261%	0.1095%	0.3490%	0.6279%	0.4910%	0.0962%
		Males	0.0865%	0.0002%	0.0002%	0.0014%	0.0045%	0.0226%	0.1105%	0.3619%	0.6871%	0.5888%	0.1033%
		Females	0.0949%	-	0.0002%	0.0016%	0.0056%	0.0295%	0.1085%	0.3373%	0.5822%	0.4368%	0.0914%
	White <sup>c</sup>	Both Sexes	0.0973%	0.0001%	0.0002%	0.0017%	0.0052%	0.0258%	0.1059%	0.3595%	0.6531%	0.5031%	0.0986%
		Males	0.0900%	0.0002%	0.0002%	0.0015%	0.0048%	0.0222%	0.1029%	0.3580%	0.6950%	0.5911%	0.1027%
		Females	0.1047%	-	-	0.0018%	0.0058%	0.0296%	0.1089%	0.3609%	0.6200%	0.4547%	0.0960%
	Black <sup>c</sup>	Both Sexes	0.0772%	-	-	0.0009%	0.0053%	0.0346%	0.1682%	0.4036%	0.6451%	0.4113%	0.1062%
		Males	0.0795%	-	-	-	0.0043%	0.0293%	0.1950%	0.4835%	0.7630%	0.5325%	0.1251%
		Females	0.0751%	-	-	-	0.0062%	0.0392%	0.1459%	0.3413%	0.5667%	0.3526%	0.0935%
	Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0676%	-	-	0.0010%	0.0039%	0.0200%	0.0850%	0.2482%	0.4628%	0.4443%	0.0735%
		Males	0.0754%	-	-	-	0.0032%	0.0191%	0.0963%	0.3094%	0.5893%	0.5998%	0.0917%
		Females	0.0603%	-	-	-	0.0046%	0.0208%	0.0751%	0.1963%	0.3701%	0.3418%	0.0599%
	Hispanic <sup>d</sup>	Both Sexes	0.0206%	-	-	0.0012%	0.0023%	0.0116%	0.0399%	0.1367%	0.2779%	0.2697%	0.0424%
		Males	0.0205%	-	-	0.0009%	0.0028%	0.0114%	0.0427%	0.1480%	0.3501%	0.3602%	0.0510%
		Females	0.0207%	-	-	0.0016%	0.0017%	0.0118%	0.0372%	0.1272%	0.2253%	0.2171%	0.0365%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 16.18  
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	233,089	170,371	118,667	88,895	60,159	38,309	541,829	714,792	43,896	758,688
	Males	122,409	87,895	58,917	40,417	25,909	15,628	278,623	353,155	14,770	367,925
	Females	110,680	82,476	59,750	48,478	34,250	22,681	263,206	361,637	29,126	390,763
White <sup>b</sup>	Both Sexes	225,347	164,016	113,738	85,054	57,712	36,868	522,176	687,691	41,522	729,213
	Males	118,610	84,648	56,570	38,946	24,872	14,965	268,979	340,488	14,095	354,583
	Females	106,737	79,368	57,168	46,108	32,840	21,903	253,197	347,203	27,427	374,630
Black <sup>b</sup>	Both Sexes	811	635	431	298	142	86	1,942	2,451	119	2,570
	Males	335	274	232	93	19	26	866	990	46	1,036
	Females	476	361	199	205	123	60	1,076	1,461	73	1,534
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	618	410	177	+	+	+	1,240	+	+	+
	Males	270	177	75	+	+	+	531	+	+	+
	Females	348	233	102	+	+	+	709	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,093	2,799	1,907	+	+	+	9,141	+	+	+
	Males	1,543	1,067	709	+	+	+	3,424	+	+	+
	Females	2,550	1,732	1,198	+	+	+	5,717	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1476%	0.0004%	0.0033%	0.0293%	0.0892%	0.1721%	0.2762%	0.3912%	0.4918%	0.4840%	0.1485%
	Males	0.1555%	0.0003%	0.0028%	0.0194%	0.0624%	0.1510%	0.2964%	0.4969%	0.6976%	0.7540%	0.1711%
	Females	0.1399%	0.0004%	0.0038%	0.0399%	0.1167%	0.1932%	0.2570%	0.2959%	0.3330%	0.3342%	0.1340%
White <sup>c</sup>	Both Sexes	0.1856%	0.0004%	0.0042%	0.0367%	0.1137%	0.2142%	0.3395%	0.4729%	0.5940%	0.5619%	0.1814%
	Males	0.1941%	0.0004%	0.0035%	0.0238%	0.0773%	0.1849%	0.3596%	0.5933%	0.8333%	0.8856%	0.2055%
	Females	0.1771%	0.0004%	0.0048%	0.0510%	0.1528%	0.2445%	0.3197%	0.3614%	0.4049%	0.3844%	0.1666%
Black <sup>c</sup>	Both Sexes	0.0047%	-	-	0.0015%	0.0038%	0.0038%	0.0074%	0.0196%	0.0250%	0.0269%	0.0061%
	Males	0.0045%	-	-	-	0.0031%	0.0030%	0.0068%	0.0227%	0.0387%	0.0340%	0.0070%
	Females	0.0049%	-	-	0.0023%	0.0044%	0.0045%	0.0078%	0.0172%	0.0158%	0.0235%	0.0056%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0093%	-	-	0.0025%	0.0070%	0.0103%	0.0149%	0.0266%	0.0253%	0.0342%	0.0094%
	Males	0.0084%	-	-	0.0032%	0.0041%	0.0074%	0.0155%	0.0266%	0.0319%	0.0347%	0.0091%
	Females	0.0102%	-	-	0.0018%	0.0097%	0.0130%	0.0144%	0.0266%	0.0204%	0.0339%	0.0097%
Hispanic <sup>d</sup>	Both Sexes	0.0205%	-	0.0006%	0.0046%	0.0168%	0.0372%	0.0565%	0.0795%	0.1050%	0.1464%	0.0323%
	Males	0.0148%	-	0.0006%	0.0027%	0.0069%	0.0230%	0.0439%	0.0737%	0.1125%	0.2174%	0.0294%
	Females	0.0266%	-	-	0.0069%	0.0283%	0.0523%	0.0684%	0.0844%	0.0995%	0.1051%	0.0367%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 17.11  
MesotheliomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	2,774	552	349	226	161	123	3,725	4,194	+	+
	Males	2,022	314	80	62	89	36	2,429	2,603	+	+
	Females	752	238	269	164	72	87	1,296	1,591	+	+
White <sup>b</sup>	Both Sexes	2,585	485	297	211	150	99	3,414	3,834	+	+
	Males	1,893	281	68	47	78	23	2,253	2,389	+	+
	Females	692	204	229	164	72	76	1,161	1,445	+	+
Black <sup>b</sup>	Both Sexes	132	36	38	11	11	24	205	252	+	+
	Males	99	12	13	11	11	13	123	159	+	+
	Females	33	24	25	0	0	11	82	93	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	30	12	5	+	+	+	47	+	+	+
	Males	18	9	3	+	+	+	30	+	+	+
	Females	12	3	2	+	+	+	17	+	+	+
Hispanic <sup>d</sup>	Both Sexes	210	55	16	+	+	+	280	+	+	+
	Males	133	35	0	+	+	+	168	+	+	+
	Females	77	20	16	+	+	+	112	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0011%	-	-	-	0.0002%	0.0005%	0.0017%	0.0033%	0.0067%	0.0056%	0.0011%
	Males	0.0014%	-	-	-	-	0.0004%	0.0019%	0.0046%	0.0121%	0.0119%	0.0017%
	Females	0.0007%	-	-	-	0.0003%	0.0006%	0.0014%	0.0022%	0.0026%	0.0022%	0.0007%
White <sup>c</sup>	Both Sexes	0.0013%	-	-	-	0.0003%	0.0006%	0.0018%	0.0038%	0.0081%	0.0059%	0.0013%
	Males	0.0017%	-	-	-	-	0.0004%	0.0021%	0.0053%	0.0142%	0.0124%	0.0020%
	Females	0.0009%	-	-	-	0.0003%	0.0007%	0.0016%	0.0025%	0.0033%	0.0024%	0.0008%
Black <sup>c</sup>	Both Sexes	0.0005%	-	-	-	-	-	0.0015%	0.0022%	-	-	0.0006%
	Males	0.0007%	-	-	-	-	-	-	-	-	-	0.0010%
	Females	0.0004%	-	-	-	-	-	-	-	-	-	0.0004%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0004%	-	-	-	-	-	-	-	-	0.0051%	0.0004%
	Males	0.0006%	-	-	-	-	-	-	-	-	0.0110%	0.0007%
	Females	0.0002%	-	-	-	-	-	-	-	-	-	0.0002%
Hispanic <sup>d</sup>	Both Sexes	0.0006%	-	-	-	-	0.0007%	0.0019%	0.0026%	0.0094%	0.0046%	0.0012%
	Males	0.0007%	-	-	-	-	-	0.0023%	0.0037%	0.0131%	-	0.0017%
	Females	0.0005%	-	-	-	-	-	0.0014%	-	0.0066%	-	0.0009%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 18.18  
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	39,615	12,048	4,647	1,740	622	195	56,829	58,905	298	59,203	
	Males	21,534	6,777	2,616	1,054	301	88	31,232	32,370	113	32,483	
	Females	18,081	5,271	2,031	686	321	107	25,597	26,535	185	26,720	
White <sup>b</sup>	Both Sexes	30,902	9,472	3,556	1,265	363	182	44,323	45,751	87	45,838	
	Males	17,260	5,545	2,024	783	163	84	25,068	25,859	0 <sup>i</sup>	25,859	
	Females	13,642	3,927	1,532	482	200	98	19,255	19,892	87 <sup>i</sup>	19,979	
Black <sup>b</sup>	Both Sexes	7,133	2,118	911	403	229	0	10,275	10,819	+	+	
	Males	3,467	995	510	241	126	0	5,034	5,338	+	+	
	Females	3,666	1,123	401	162	103	0	5,241	5,481	+	+	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	862	291	110	+	+	+	1,270	+	+	+	
	Males	433	156	56	+	+	+	647	+	+	+	
	Females	429	135	54	+	+	+	623	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	2,700	823	275	+	+	+	3,833	+	+	+	
	Males	1,399	470	161	+	+	+	2,054	+	+	+	
	Females	1,301	353	114	+	+	+	1,779	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Race	Sex	Age Specific (Crude)								Age-Adjusted <sup>f</sup>	
			All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+
All Races <sup>c</sup>	Both Sexes	0.0173%	-	-	0.0002%	0.0020%	0.0082%	0.0288%	0.0655%	0.0926%	0.0852%	0.0178%
	Males	0.0190%	-	-	0.0003%	0.0027%	0.0092%	0.0338%	0.0802%	0.1140%	0.1044%	0.0216%
	Females	0.0156%	-	-	-	0.0013%	0.0071%	0.0240%	0.0523%	0.0760%	0.0745%	0.0147%
White <sup>c</sup>	Both Sexes	0.0168%	-	-	-	0.0019%	0.0077%	0.0260%	0.0616%	0.0880%	0.0780%	0.0166%
	Males	0.0188%	-	-	-	0.0028%	0.0087%	0.0308%	0.0769%	0.1113%	0.0971%	0.0206%
	Females	0.0148%	-	-	-	0.0010%	0.0067%	0.0213%	0.0474%	0.0696%	0.0675%	0.0134%
Black <sup>c</sup>	Both Sexes	0.0259%	-	-	0.0007%	0.0039%	0.0155%	0.0611%	0.1252%	0.1703%	0.1732%	0.0348%
	Males	0.0270%	-	-	-	0.0040%	0.0175%	0.0698%	0.1557%	0.1944%	0.2041%	0.0407%
	Females	0.0249%	-	-	-	0.0037%	0.0138%	0.0540%	0.1014%	0.1542%	0.1583%	0.0305%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0106%	-	-	-	0.0007%	0.0037%	0.0176%	0.0407%	0.0565%	0.0713%	0.0113%
	Males	0.0112%	-	-	-	-	0.0043%	0.0217%	0.0417%	0.0652%	0.0926%	0.0131%
	Females	0.0101%	-	-	-	-	0.0033%	0.0140%	0.0399%	0.0501%	0.0572%	0.0099%
Hispanic <sup>d</sup>	Both Sexes	0.0088%	-	-	-	0.0025%	0.0080%	0.0224%	0.0599%	0.0931%	0.0755%	0.0166%
	Males	0.0090%	-	-	-	0.0035%	0.0080%	0.0272%	0.0676%	0.1061%	0.0955%	0.0193%
	Females	0.0085%	-	-	-	0.0013%	0.0080%	0.0178%	0.0533%	0.0836%	0.0638%	0.0144%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 19.22  
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	178,216	105,133	59,771	33,931	19,256	10,268	351,503	407,962	11,571	419,533	
	Males	92,687	54,051	31,316	17,469	9,896	4,966	182,379	211,075	6,068	217,143	
	Females	85,529	51,082	28,455	16,462	9,360	5,302	169,124	196,887	5,503	202,390	
White <sup>b</sup>	Both Sexes	157,198	93,731	53,538	30,738	17,606	9,442	311,965	363,533	11,814	375,347	
	Males	81,948	48,226	28,007	15,917	9,036	4,577	162,065	188,361	6,585	194,946	
	Females	75,250	45,505	25,531	14,821	8,570	4,865	149,900	175,172	5,229	180,401	
Black <sup>b</sup>	Both Sexes	13,856	7,121	3,838	2,144	1,022	584	25,436	28,629	891	29,520	
	Males	7,011	3,626	2,008	1,016	522	249	12,931	14,455	389	14,844	
	Females	6,845	3,495	1,830	1,128	500	335	12,505	14,174	502	14,676	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	4,455	2,442	1,394	+	+	+	8,471	+	+	+	
	Males	2,324	1,300	678	+	+	+	4,393	+	+	+	
	Females	2,131	1,142	716	+	+	+	4,078	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	10,965	6,225	3,051	+	+	+	20,706	+	+	+	
	Males	5,675	3,329	1,531	+	+	+	10,753	+	+	+	
	Females	5,290	2,896	1,520	+	+	+	9,953	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.1041%	0.0023%	0.0113%	0.0196%	0.0365%	0.0802%	0.1623%	0.3054%	0.4639%	0.4992%	0.1067%
	Males	0.1095%	0.0029%	0.0153%	0.0240%	0.0422%	0.0954%	0.1834%	0.3442%	0.5188%	0.5578%	0.1215%
	Females	0.0988%	0.0016%	0.0071%	0.0150%	0.0307%	0.0651%	0.1421%	0.2704%	0.4215%	0.4667%	0.0938%
White <sup>c</sup>	Both Sexes	0.1154%	0.0025%	0.0116%	0.0203%	0.0379%	0.0846%	0.1726%	0.3285%	0.5057%	0.5300%	0.1141%
	Males	0.1206%	0.0032%	0.0155%	0.0247%	0.0444%	0.0994%	0.1950%	0.3678%	0.5619%	0.5904%	0.1292%
	Females	0.1102%	0.0017%	0.0075%	0.0153%	0.0308%	0.0693%	0.1506%	0.2921%	0.4613%	0.4969%	0.1005%
Black <sup>c</sup>	Both Sexes	0.0627%	0.0019%	0.0092%	0.0202%	0.0361%	0.0751%	0.1381%	0.2211%	0.2588%	0.2355%	0.0758%
	Males	0.0682%	0.0019%	0.0138%	0.0242%	0.0415%	0.0961%	0.1606%	0.2480%	0.2880%	0.2446%	0.0873%
	Females	0.0577%	0.0019%	0.0045%	0.0164%	0.0315%	0.0567%	0.1194%	0.2001%	0.2393%	0.2311%	0.0663%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0667%	0.0015%	0.0116%	0.0150%	0.0247%	0.0522%	0.1004%	0.1840%	0.2931%	0.3649%	0.0696%
	Males	0.0706%	0.0022%	0.0151%	0.0197%	0.0256%	0.0628%	0.1060%	0.2096%	0.3343%	0.4239%	0.0796%
	Females	0.0630%	-	0.0079%	0.0103%	0.0239%	0.0424%	0.0955%	0.1623%	0.2629%	0.3261%	0.0615%
Hispanic <sup>d</sup>	Both Sexes	0.0469%	0.0021%	0.0089%	0.0132%	0.0267%	0.0570%	0.1181%	0.2170%	0.3611%	0.3425%	0.0782%
	Males	0.0471%	0.0025%	0.0120%	0.0161%	0.0301%	0.0640%	0.1294%	0.2226%	0.3749%	0.3464%	0.0833%
	Females	0.0467%	0.0017%	0.0058%	0.0098%	0.0228%	0.0496%	0.1074%	0.2122%	0.3511%	0.3403%	0.0732%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 20.18  
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	88,149	54,740	35,668	23,399	15,340	10,487	184,526	229,332	15,141	244,473	
	Males	59,730	35,629	22,720	14,107	9,060	6,161	121,846	148,389	8,861	157,250	
	Females	28,419	19,111	12,948	9,292	6,280	4,326	62,680	80,943	6,280	87,223	
White <sup>b</sup>	Both Sexes	75,611	47,666	31,443	20,922	13,658	9,255	160,023	199,915	12,831	212,746	
	Males	52,030	31,379	20,320	12,839	8,269	5,595	107,162	131,318	7,583	138,901	
	Females	23,581	16,287	11,123	8,083	5,389	3,660	52,861	68,597	5,248	73,845	
Black <sup>b</sup>	Both Sexes	7,560	3,920	2,478	1,453	1,008	707	14,343	17,246	776	18,022	
	Males	4,627	2,322	1,335	743	429	270	8,459	9,780	308	10,088	
	Females	2,933	1,598	1,143	710	579	437	5,884	7,466	468	7,934	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,186	2,038	1,172	+	+	+	6,552	+	+	+	
	Males	1,981	1,254	704	+	+	+	4,030	+	+	+	
	Females	1,205	784	468	+	+	+	2,522	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	4,089	2,379	1,449	+	+	+	8,131	+	+	+	
	Males	2,676	1,463	806	+	+	+	5,062	+	+	+	
	Females	1,413	916	643	+	+	+	3,069	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0550%	0.0003%	0.0020%	0.0058%	0.0143%	0.0419%	0.1125%	0.1847%	0.2236%	0.2066%	0.0556%
	Males	0.0732%	-	0.0018%	0.0050%	0.0144%	0.0548%	0.1667%	0.2775%	0.3295%	0.3072%	0.0797%
	Females	0.0372%	0.0004%	0.0022%	0.0066%	0.0142%	0.0289%	0.0607%	0.1009%	0.1419%	0.1507%	0.0353%
White <sup>c</sup>	Both Sexes	0.0584%	0.0003%	0.0020%	0.0056%	0.0129%	0.0399%	0.1135%	0.1940%	0.2374%	0.2150%	0.0569%
	Males	0.0784%	-	0.0015%	0.0046%	0.0131%	0.0534%	0.1696%	0.2910%	0.3511%	0.3253%	0.0824%
	Females	0.0384%	0.0004%	0.0024%	0.0068%	0.0127%	0.0259%	0.0583%	0.1040%	0.1476%	0.1545%	0.0351%
Black <sup>c</sup>	Both Sexes	0.0361%	-	0.0026%	0.0055%	0.0128%	0.0365%	0.1023%	0.1498%	0.1617%	0.1093%	0.0442%
	Males	0.0461%	-	0.0039%	0.0059%	0.0106%	0.0427%	0.1514%	0.2297%	0.2431%	0.1323%	0.0617%
	Females	0.0271%	-	0.0014%	0.0052%	0.0147%	0.0310%	0.0615%	0.0877%	0.1076%	0.0982%	0.0309%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0502%	-	0.0014%	0.0066%	0.0214%	0.0560%	0.1085%	0.1402%	0.1643%	0.1727%	0.0506%
	Males	0.0637%	-	-	0.0068%	0.0231%	0.0714%	0.1515%	0.2072%	0.2200%	0.2423%	0.0685%
	Females	0.0375%	-	0.0016%	0.0064%	0.0198%	0.0418%	0.0707%	0.0834%	0.1234%	0.1268%	0.0360%
Hispanic <sup>d</sup>	Both Sexes	0.0182%	-	0.0011%	0.0025%	0.0079%	0.0204%	0.0569%	0.1057%	0.1366%	0.1340%	0.0315%
	Males	0.0220%	-	0.0010%	0.0021%	0.0074%	0.0249%	0.0783%	0.1559%	0.2005%	0.1786%	0.0434%
	Females	0.0143%	-	0.0013%	0.0029%	0.0084%	0.0157%	0.0366%	0.0632%	0.0901%	0.1081%	0.0219%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 21.14  
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	54,346	30,384	22,476	18,636	15,590	11,063	111,375	154,428	21,579	176,007
White <sup>b</sup>	47,888	26,827	19,849	17,082	13,988	9,915	98,499	137,232	18,682	155,914
Black <sup>b</sup>	3,790	1,969	1,531	993	1,040	845	7,424	10,332	1,896	12,228
Asian/Pacific Islander <sup>c</sup>	2,033	1,128	790	+	+	+	4,062	+	+	+
Hispanic <sup>d</sup>	4,070	2,028	1,431	+	+	+	7,776	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	0.0675%	0.0003%	0.0053%	0.0169%	0.0267%	0.0637%	0.1345%	0.2007%	0.2134%	0.1618%	0.0643%
White <sup>c</sup>	0.0741%	0.0004%	0.0057%	0.0170%	0.0266%	0.0669%	0.1430%	0.2176%	0.2337%	0.1726%	0.0686%
Black <sup>c</sup>	0.0355%	-	0.0040%	0.0112%	0.0251%	0.0402%	0.0779%	0.1208%	0.1137%	0.1023%	0.0397%
Asian/Pacific Islander <sup>c</sup>	0.0589%	-	0.0049%	0.0226%	0.0280%	0.0633%	0.1310%	0.1525%	0.1526%	0.1081%	0.0557%
Hispanic <sup>d</sup>	0.0363%	-	0.0069%	0.0173%	0.0223%	0.0474%	0.0999%	0.1593%	0.1794%	0.1254%	0.0516%

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.  
<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census.  
<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.  
<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.  
<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.  
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
 + Not available.

Table 22.18  
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	23,089	3,805	1,959	1,044	483	372	29,138	30,784	396	31,180	
	Males	11,266	1,847	846	434	192	201	14,094	14,813	166	14,979	
	Females	11,823	1,958	1,113	610	291	171	15,044	15,971	230	16,201	
White <sup>b</sup>	Both Sexes	19,466	3,007	1,550	785	382	312	24,264	25,531	321	25,852	
	Males	9,931	1,588	684	355	155	173	12,325	12,910	150	13,060	
	Females	9,535	1,419	866	430	227	139	11,939	12,621	171	12,792	
Black <sup>b</sup>	Both Sexes	2,665	502	249	217	62	55	3,452	3,753	180	3,933	
	Males	883	146	74	76	26	23	1,116	1,231	54	1,285	
	Females	1,782	356	175	141	36	32	2,336	2,522	126	2,648	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	698	152	56	+	+	+	913	+	+	+	
	Males	340	65	27	+	+	+	433	+	+	+	
	Females	358	87	29	+	+	+	480	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	1,438	229	181	+	+	+	1,859	+	+	+	
	Males	603	83	65	+	+	+	762	+	+	+	
	Females	835	146	116	+	+	+	1,097	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0089%	-	0.0002%	0.0005%	0.0016%	0.0050%	0.0149%	0.0315%	0.0447%	0.0448%	0.0092%
	Males	0.0087%	-	-	0.0003%	0.0012%	0.0047%	0.0161%	0.0351%	0.0498%	0.0480%	0.0098%
	Females	0.0091%	-	0.0002%	0.0006%	0.0019%	0.0053%	0.0137%	0.0282%	0.0407%	0.0431%	0.0086%
White <sup>c</sup>	Both Sexes	0.0092%	-	0.0002%	0.0004%	0.0017%	0.0046%	0.0149%	0.0311%	0.0449%	0.0436%	0.0091%
	Males	0.0092%	-	-	0.0003%	0.0012%	0.0045%	0.0168%	0.0355%	0.0513%	0.0477%	0.0100%
	Females	0.0092%	-	0.0003%	0.0005%	0.0021%	0.0047%	0.0131%	0.0270%	0.0399%	0.0414%	0.0083%
Black <sup>c</sup>	Both Sexes	0.0084%	-	-	-	0.0018%	0.0059%	0.0183%	0.0415%	0.0521%	0.0553%	0.0113%
	Males	0.0064%	-	-	-	0.0021%	0.0051%	0.0170%	0.0369%	0.0422%	0.0239%	0.0090%
	Females	0.0103%	-	-	-	0.0016%	0.0066%	0.0194%	0.0450%	0.0586%	0.0705%	0.0127%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0075%	-	-	-	0.0010%	0.0055%	0.0117%	0.0261%	0.0349%	0.0472%	0.0079%
	Males	0.0075%	-	-	-	-	0.0043%	0.0103%	0.0332%	0.0401%	0.0621%	0.0088%
	Females	0.0075%	-	-	-	0.0012%	0.0065%	0.0129%	0.0200%	0.0310%	0.0373%	0.0074%
Hispanic <sup>d</sup>	Both Sexes	0.0043%	-	-	0.0005%	0.0014%	0.0035%	0.0114%	0.0261%	0.0419%	0.0428%	0.0080%
	Males	0.0033%	-	-	-	0.0011%	0.0027%	0.0087%	0.0277%	0.0400%	0.0354%	0.0073%
	Females	0.0053%	-	-	-	0.0017%	0.0045%	0.0139%	0.0248%	0.0432%	0.0471%	0.0086%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 23.14  
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	930,125	678,835	425,357	107,135	25,089	7,820	2,071,562	2,174,982	2,993	2,177,975
White <sup>b</sup>	771,984	573,491	371,231	97,545	22,772	7,134	1,750,649	1,844,754	3,289	1,848,043
Black <sup>b</sup>	120,753	83,385	44,242	7,854	2,057	592	251,034	258,895	267	259,162
Asian/Pacific Islander <sup>c</sup>	17,274	10,081	4,918	+	+	+	32,638	+	+	+
Hispanic <sup>d</sup>	48,519	29,822	15,122	+	+	+	94,658	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	1.2060%	0.0002%	0.0002%	-	0.0013%	0.0671%	0.9276%	4.6199%	10.8389%	12.5805%	1.5140%
White <sup>c</sup>	1.2659%	-	0.0003%	-	0.0010%	0.0597%	0.8924%	4.5828%	10.9325%	12.4498%	1.5071%
Black <sup>c</sup>	1.2893%	-	-	-	0.0036%	0.1619%	1.8218%	7.9219%	16.0258%	15.8494%	2.2927%
Asian/Pacific Islander <sup>c</sup>	0.6012%	-	-	-	-	0.0192%	0.2963%	1.9041%	5.5656%	8.4302%	0.7871%
Hispanic <sup>d</sup>	0.4084%	-	-	-	0.0006%	0.0342%	0.5452%	3.1990%	8.4377%	9.8517%	1.1356%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.18  
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	31,763	12,887	7,717	5,264	2,760	1,529	53,464	62,024	2,098	64,122
	Males	19,112	7,354	4,149	2,604	1,427	813	31,117	35,526	1,199	36,725
	Females	12,651	5,533	3,568	2,660	1,333	716	22,347	26,498	899	27,397
White <sup>b</sup>	Both Sexes	24,366	9,461	5,842	3,972	2,127	1,286	40,474	47,124	1,708	48,832
	Males	14,951	5,420	3,072	1,994	1,129	694	23,834	27,305	922	28,227
	Females	9,415	4,041	2,770	1,978	998	592	16,640	19,819	786	20,605
Black <sup>b</sup>	Both Sexes	4,497	1,951	987	706	393	107	7,561	8,665	270	8,935
	Males	2,436	1,017	606	320	212	69	4,095	4,671	165	4,836
	Females	2,061	934	381	386	181	38	3,466	3,994	105	4,099
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,743	1,313	811	+	+	+	5,002	+	+	+
	Males	1,549	758	454	+	+	+	2,836	+	+	+
	Females	1,194	555	357	+	+	+	2,166	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,790	1,519	853	+	+	+	6,293	+	+	+
	Males	2,048	754	380	+	+	+	3,235	+	+	+
	Females	1,742	765	473	+	+	+	3,058	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0198%	-	-	0.0005%	0.0031%	0.0103%	0.0263%	0.0558%	0.1130%	0.1469%	0.0207%
	Males	0.0228%	-	-	0.0004%	0.0033%	0.0111%	0.0322%	0.0757%	0.1525%	0.1989%	0.0271%
	Females	0.0169%	-	-	0.0005%	0.0030%	0.0095%	0.0206%	0.0378%	0.0826%	0.1180%	0.0158%
White <sup>c</sup>	Both Sexes	0.0165%	-	-	0.0004%	0.0029%	0.0084%	0.0211%	0.0454%	0.0873%	0.1118%	0.0163%
	Males	0.0190%	-	-	0.0004%	0.0030%	0.0094%	0.0263%	0.0624%	0.1184%	0.1484%	0.0214%
	Females	0.0139%	-	-	0.0005%	0.0027%	0.0073%	0.0161%	0.0296%	0.0627%	0.0917%	0.0123%
Black <sup>c</sup>	Both Sexes	0.0187%	-	-	0.0007%	0.0032%	0.0147%	0.0370%	0.0727%	0.1306%	0.1688%	0.0256%
	Males	0.0208%	-	-	-	0.0043%	0.0161%	0.0436%	0.0960%	0.1750%	0.2371%	0.0332%
	Females	0.0167%	-	-	-	0.0023%	0.0136%	0.0314%	0.0546%	0.1012%	0.1356%	0.0205%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0422%	-	-	-	0.0043%	0.0177%	0.0484%	0.1104%	0.2666%	0.4071%	0.0463%
	Males	0.0492%	-	-	-	0.0039%	0.0179%	0.0602%	0.1512%	0.3654%	0.5381%	0.0609%
	Females	0.0355%	-	-	-	0.0046%	0.0176%	0.0381%	0.0759%	0.1942%	0.3208%	0.0354%
Hispanic <sup>d</sup>	Both Sexes	0.0142%	-	-	0.0007%	0.0042%	0.0132%	0.0331%	0.0705%	0.1566%	0.2043%	0.0279%
	Males	0.0141%	-	-	0.0006%	0.0036%	0.0128%	0.0369%	0.0957%	0.1855%	0.2192%	0.0323%
	Females	0.0143%	-	-	0.0009%	0.0049%	0.0135%	0.0294%	0.0493%	0.1355%	0.1956%	0.0244%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 25.14  
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	39,842	36,207	31,222	28,324	20,672	14,328	113,356	172,591	17,674	190,265
White <sup>b</sup>	37,597	34,027	29,646	27,391	19,958	13,852	107,184	164,350	16,986	181,336
Black <sup>b</sup>	903	945	596	347	345	220	2,500	3,412	152	3,564
Asian/Pacific Islander <sup>c</sup>	672	540	482	+	+	+	1,756	+	+	+
Hispanic <sup>d</sup>	4,384	3,247	2,515	+	+	+	10,534	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0690%	0.0021%	0.0061%	0.0613%	0.1397%	0.1538%	0.0844%	0.0355%	0.0156%	0.0092%	0.0675%	
White <sup>c</sup>	0.0836%	0.0021%	0.0070%	0.0748%	0.1733%	0.1850%	0.0991%	0.0419%	0.0185%	0.0102%	0.0817%	
Black <sup>c</sup>	0.0140%	-	0.0024%	0.0099%	0.0260%	0.0360%	0.0231%	0.0051%	-	-	0.0143%	
Asian/Pacific Islander <sup>c</sup>	0.0242%	0.0038%	0.0046%	0.0218%	0.0410%	0.0492%	0.0349%	0.0087%	-	-	0.0228%	
Hispanic <sup>d</sup>	0.0458%	0.0026%	0.0063%	0.0548%	0.0989%	0.0935%	0.0448%	0.0211%	0.0076%	-	0.0448%	

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 26.18  
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	120,479	79,343	57,374	42,937	33,340	28,843	266,724	367,333	43,071	410,404
	Males	26,876	17,561	13,313	8,966	7,186	6,270	59,608	81,388	10,006	91,394
	Females	93,603	61,782	44,061	33,971	26,154	22,573	207,116	285,945	33,065	319,010
White <sup>b</sup>	Both Sexes	105,500	69,731	50,969	38,504	30,106	25,978	234,804	325,378	38,993	364,371
	Males	24,054	15,955	12,190	8,278	6,489	5,792	53,919	73,885	8,467	82,352
	Females	81,446	53,776	38,779	30,226	23,617	20,186	180,885	251,493	30,526	282,019
Black <sup>b</sup>	Both Sexes	7,374	4,510	3,036	1,964	1,512	1,349	15,346	19,969	1,776	21,745
	Males	1,429	664	575	313	330	145	2,746	3,480	195	3,675
	Females	5,945	3,846	2,461	1,651	1,182	1,204	12,600	16,489	1,581	18,070
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	5,407	3,474	2,446	+	+	+	11,738	+	+	+
	Males	996	652	396	+	+	+	2,112	+	+	+
	Females	4,411	2,822	2,050	+	+	+	9,626	+	+	+
Hispanic <sup>d</sup>	Both Sexes	10,871	6,673	4,626	+	+	+	22,903	+	+	+
	Males	1,891	1,135	760	+	+	+	3,924	+	+	+
	Females	8,980	5,538	3,866	+	+	+	18,979	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0831%	-	0.0039%	0.0334%	0.0974%	0.1401%	0.1605%	0.1592%	0.1384%	0.0973%	0.0821%
	Males	0.0368%	-	0.0015%	0.0092%	0.0329%	0.0538%	0.0769%	0.0962%	0.0934%	0.0689%	0.0380%
	Females	0.1285%	-	0.0063%	0.0593%	0.1637%	0.2262%	0.2402%	0.2160%	0.1732%	0.1130%	0.1248%
White <sup>c</sup>	Both Sexes	0.0894%	-	0.0041%	0.0365%	0.1084%	0.1524%	0.1681%	0.1623%	0.1394%	0.0970%	0.0873%
	Males	0.0409%	-	0.0016%	0.0097%	0.0369%	0.0597%	0.0840%	0.1002%	0.0982%	0.0712%	0.0411%
	Females	0.1381%	-	0.0068%	0.0663%	0.1850%	0.2480%	0.2508%	0.2199%	0.1719%	0.1111%	0.1338%
Black <sup>c</sup>	Both Sexes	0.0384%	-	0.0015%	0.0119%	0.0424%	0.0637%	0.0890%	0.1121%	0.0899%	0.0592%	0.0434%
	Males	0.0147%	-	-	0.0036%	0.0127%	0.0204%	0.0337%	0.0610%	0.0600%	0.0442%	0.0189%
	Females	0.0599%	-	0.0020%	0.0199%	0.0684%	0.1017%	0.1349%	0.1518%	0.1098%	0.0664%	0.0637%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0842%	-	0.0043%	0.0334%	0.0833%	0.1299%	0.1659%	0.1695%	0.1576%	0.1206%	0.0817%
	Males	0.0317%	-	-	0.0112%	0.0264%	0.0446%	0.0618%	0.0903%	0.0770%	0.0638%	0.0327%
	Females	0.1335%	-	0.0079%	0.0556%	0.1373%	0.2082%	0.2573%	0.2367%	0.2166%	0.1581%	0.1251%
Hispanic <sup>d</sup>	Both Sexes	0.0521%	-	0.0036%	0.0260%	0.0676%	0.1077%	0.1399%	0.1477%	0.1348%	0.1063%	0.0688%
	Males	0.0170%	-	0.0013%	0.0052%	0.0198%	0.0284%	0.0549%	0.0762%	0.0680%	0.0583%	0.0260%
	Females	0.0891%	-	0.0060%	0.0511%	0.1230%	0.1918%	0.2206%	0.2084%	0.1834%	0.1341%	0.1110%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 27.18  
Cancer of the Urinary Bladder (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	188,227	127,800	86,030	53,796	32,634	18,084	414,603	508,818	18,678	527,496
	Males	140,439	95,158	63,819	39,794	23,482	12,677	308,631	376,871	12,094	388,965
	Females	47,788	32,642	22,211	14,002	9,152	5,407	105,972	131,947	6,584	138,531
White <sup>b</sup>	Both Sexes	174,608	119,935	81,097	51,448	31,099	17,298	387,559	477,550	14,627	492,177
	Males	131,060	89,679	60,169	38,026	22,390	12,187	289,662	354,884	9,316	364,200
	Females	43,548	30,256	20,928	13,422	8,709	5,111	97,897	122,666	5,311	127,977
Black <sup>b</sup>	Both Sexes	8,208	4,408	3,180	1,522	933	422	16,177	18,748	305	19,053
	Males	5,275	2,854	2,322	1,165	641	256	10,719	12,575	245	12,820
	Females	2,933	1,554	858	357	292	166	5,458	6,173	60	6,233
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,828	1,604	1,000	+	+	+	5,569	+	+	+
	Males	2,136	1,179	750	+	+	+	4,162	+	+	+
	Females	692	425	250	+	+	+	1,407	+	+	+
Hispanic <sup>d</sup>	Both Sexes	5,959	3,537	2,232	+	+	+	12,129	+	+	+
	Males	4,439	2,663	1,679	+	+	+	9,067	+	+	+
	Females	1,520	874	553	+	+	+	3,062	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1146%	0.0001%	0.0003%	0.0019%	0.0069%	0.0294%	0.1111%	0.3557%	0.7480%	0.9882%	0.1214%
	Males	0.1735%	-	0.0004%	0.0024%	0.0100%	0.0445%	0.1752%	0.5813%	1.3098%	1.9667%	0.2142%
	Females	0.0567%	-	0.0002%	0.0015%	0.0039%	0.0142%	0.0500%	0.1521%	0.3145%	0.4453%	0.0529%
White <sup>c</sup>	Both Sexes	0.1365%	0.0001%	0.0003%	0.0024%	0.0083%	0.0333%	0.1268%	0.4084%	0.8532%	1.0777%	0.1370%
	Males	0.2057%	-	0.0005%	0.0029%	0.0115%	0.0497%	0.1971%	0.6584%	1.4838%	2.1806%	0.2407%
	Females	0.0672%	-	-	0.0018%	0.0048%	0.0165%	0.0576%	0.1767%	0.3552%	0.4729%	0.0594%
Black <sup>c</sup>	Both Sexes	0.0403%	-	-	-	0.0036%	0.0167%	0.0636%	0.1773%	0.3302%	0.4624%	0.0584%
	Males	0.0571%	-	-	-	0.0058%	0.0273%	0.1042%	0.2996%	0.5451%	0.7724%	0.0975%
	Females	0.0250%	-	-	-	0.0016%	0.0074%	0.0298%	0.0820%	0.1872%	0.3123%	0.0325%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0482%	-	-	0.0007%	0.0020%	0.0131%	0.0404%	0.1272%	0.3125%	0.5751%	0.0539%
	Males	0.0741%	-	-	-	0.0029%	0.0216%	0.0659%	0.2196%	0.5525%	1.0174%	0.0943%
	Females	0.0239%	-	-	-	0.0012%	0.0053%	0.0179%	0.0490%	0.1366%	0.2836%	0.0241%
Hispanic <sup>d</sup>	Both Sexes	0.0268%	-	-	0.0008%	0.0041%	0.0149%	0.0482%	0.1589%	0.3314%	0.4865%	0.0562%
	Males	0.0387%	-	-	0.0007%	0.0052%	0.0211%	0.0735%	0.2692%	0.6008%	0.9210%	0.0991%
	Females	0.0142%	-	-	0.0009%	0.0027%	0.0083%	0.0241%	0.0654%	0.1354%	0.2339%	0.0251%

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 28.12  
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2006  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 <sup>e</sup>	0 to <31 <sup>e</sup>	>=31 <sup>f</sup>	Complete <sup>g</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	60,231	50,458	45,228	38,245	32,402	28,377	163,805	259,480	+	+
	Males	32,416	26,270	23,395	19,937	16,074	13,113	86,544	133,367	+	+
	Females	27,815	24,188	21,833	18,308	16,328	15,264	77,261	126,113	+	+
White <sup>b</sup>	Both Sexes	50,388	42,539	38,148	33,001	28,466	25,200	137,833	221,693	+	+
	Males	27,315	22,085	19,840	17,296	14,280	11,739	73,153	114,499	+	+
	Females	23,073	20,454	18,308	15,705	14,186	13,461	64,680	107,194	+	+
Black <sup>b</sup>	Both Sexes	6,410	5,010	4,628	3,421	2,707	2,234	16,723	24,815	+	+
	Males	3,228	2,620	2,290	1,657	1,235	933	8,463	12,120	+	+
	Females	3,182	2,390	2,338	1,764	1,472	1,301	8,260	12,695	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,112	1,691	1,436	+	+	+	5,520	+	+	+
	Males	1,160	927	709	+	+	+	2,948	+	+	+
	Females	952	764	727	+	+	+	2,572	+	+	+
Hispanic <sup>d</sup>	Both Sexes	10,378	7,526	5,973	+	+	+	24,938	+	+	+
	Males	5,704	3,922	3,231	+	+	+	13,501	+	+	+
	Females	4,674	3,604	2,742	+	+	+	11,437	+	+	+

<sup>a</sup> US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census.

<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f g</sup> (f) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <31 and >=31.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.