

Table 1.23
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity
Both Sexes

	All Races		White		Black	
	Rate ^b 2002-2006	APC ^c 1997-2006	Rate ^b 2002-2006	APC ^c 1997-2006	Rate ^b 2002-2006	APC ^c 1997-2006
All Sites	462.9	-1.0*	470.6	-0.9*	493.6	-1.4*
Prostate ^f	70.6	-1.2	68.7	-2.1*	99.6	-2.7*
Breast	67.1	-1.9*	68.5	-1.1	74.7	-1.8*
Lung and Bronchus	63.1	-1.5*	64.4	-1.5*	67.4	-0.7*
Colon and Rectum	49.1	-2.4*	48.6	-2.6*	59.9	-1.5*
Urinary Bladder	21.0	-0.4	22.9	2.1*	15.6	-0.2
Melanoma of the Skin	19.6	1.6*	22.8	-0.3	15.0	1.5
Non-Hodgkin Lymphoma	19.5	0.1	20.4	0.2	14.8	0.3
Kidney and Renal Pelvis	13.6	2.3*	14.1	2.4*	12.8	-0.1
Corpus and Uterus, NOS ^f	12.5	-1.0*	12.9	-1.3*	12.1	-3.3*
Leukemia	12.2	-0.9*	12.8	-1.0*	11.7	-0.2
Pancreas	11.7	0.5	11.5	0.6*	11.7	1.2
Oral Cavity and Pharynx	10.4	-1.5*	10.5	-1.3*	10.5	-2.7*
Thyroid	9.6	5.9*	10.1	6.2*	9.8	-0.7
Stomach	7.9	-1.7*	7.4	-1.9*	8.2	3.3*
Ovary ^{fh}	7.1	-1.8*	7.1	-0.6	5.8	-1.2
	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2002-2006	APC ^c 1997-2006	Rate ^b 2002-2006	APC ^c 1997-2006	Rate ^b 2002-2006	APC ^c 1997-2006
All Sites	311.1	-0.9*	313.6	-0.3	350.6	-1.0*
Breast	48.9	-0.9*	44.9	0.6	58.2	-1.5*
Colon and Rectum	40.0	-1.9*	42.3	-1.7	47.4	-0.9*
Prostate ^f	39.4	-0.9	40.2	1.4	38.4	-1.4*
Lung and Bronchus	38.9	-0.9*	33.6	-1.3	32.5	-2.4*
Liver & IBD ^g	14.2	0.1	17.5	2.7	16.7	-0.2
Stomach	13.4	-3.2*	11.2	-4.0	13.2	2.0*
Non-Hodgkin Lymphoma	12.9	-0.9	10.8	0.5	11.7	-2.3*
Thyroid	9.4	3.6*	9.9	4.6	11.4	0.0
Urinary Bladder	9.3	0.1	9.7	-0.9	10.6	-0.1
Corpus and Uterus, NOS ^f	9.1	0.6	8.8	1.6	10.2	1.9*
Pancreas	9.0	-0.1	7.4	-	9.7	-0.4
Oral Cavity and Pharynx	7.8	-2.0*	7.1	-4.3	9.5	0.6
Leukemia	7.3	-1.0	7.0	2.1	8.2	5.2*
Kidney and Renal Pelvis	6.9	2.2*	5.9	-2.3	6.5	-4.1*
Ovary ^{fh}	5.4	-0.5	5.8	0.3	6.1	-1.1

- ^a Top 15 cancer sites selected based on 2002-2006 age-adjusted rates for the race/ethnic group.
- ^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^c The APC is the Annual Percent Change over the time interval. Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).
- ^d Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.
- ^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
- ^f The rates for sex-specific cancer sites are calculated using the population for both sexes combined.
- ^g IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- ^h Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- * The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.