

Table 4.25

Cancer of the Female Breast(*In Situ*)Percent Distribution and Counts by Histology among Histologically Confirmed Cases, 2002-2006
Females by Race

Histology ^a	All Races		White		Black		Asian/Pacific Islander		American Indian/ Alaska Native ^b		Hispanic ^c	
	Count	Percent	Count	Percent	Count	Percent	Count	Percent	Count	Percent	Count	Percent
Adenocarcinoma ^d	56,033	95.0%	45,834	94.9%	4,837	95.7%	4,548	94.7%	163	96.4%	4,577	95.6%
Adenocarcinoma <i>in situ</i> , NOS (8140/2)	-	-	-	-	-	-	-	-	-	-	-	-
Cribiform carcinoma <i>in situ</i> (8201/2)	4,737	8.0%	3,816	7.9%	370	7.3%	492	10.2%	-	-	389	8.1%
Ductal carcinoma <i>in situ</i> (8500/2)	20,472	34.7%	16,715	34.6%	1,819	36.0%	1,645	34.3%	72	42.6%	1,648	34.4%
Lobular carcinoma <i>in situ</i> , NOS (8520/2)	6,786	11.5%	5,950	12.3%	408	8.1%	240	5.0%	23	13.6%	496	10.4%
Comedocarcinoma <i>in situ</i> (8501/2)	4,746	8.0%	3,934	8.1%	392	7.8%	372	7.7%	-	-	378	7.9%
Intraductal and lobular carcinoma <i>in situ</i> (8522/2,8523/2,8524/2)	15,358	26.0%	12,336	25.6%	1,405	27.8%	1,459	30.4%	31	18.3%	1,329	27.8%
Intraductal micropapillary (8507/2)	1,734	2.9%	1,412	2.9%	177	3.5%	123	2.6%	-	-	154	3.2%
Other adenocarcinomas ^e	2,185	3.7%	1,657	3.4%	266	5.3%	216	4.5%	-	-	183	3.8%
Other <i>in situ</i> histologies ^f	2,941	5.0%	2,444	5.1%	219	4.3%	253	5.3%	-	-	209	4.4%
Total	58,974	100.0%	48,278	100.0%	5,056	100.0%	4,801	100.0%	169	100.0%	4,786	100.0%

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Percents may not sum to 100 due to rounding.

^a Excludes Kaposi Sarcoma, mesothelioma, lymphomas, leukemias, myelomas, lymphoreticular, and immunoproliferative diseases.

^b Estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Adenocarcinoma includes histologies 8050, 8140-8147, 8160-8162, 8180-8221, 8250-8507, 8514, 8520-8551, 8560, 8570-8574, 8576, 8940-8941.

^e Other adenocarcinomas include 8050, 8141-8147, 8160-8162, 8180-8200, 8202-8221, 8250-8490, 8502-8506, 8514, 8521, 8525-8551, 8560, 8570-8574, 8576, 8940-8941.

^f Other histologies include 8000-8049, 8051-8139, 8148-8159, 8163-8179, 8222-8249, 8508-8513, 8515-8519, 8552-8559, 8561-8569, 8575, 8577-8939, 8942-9989.

- Statistic not shown due to fewer than 16 cases during the time period.