

Table 7.6
Cancer of the Corpus and Uterus, NOS (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>Age-Adjusted Rates, 2002-2006</u>						
Age at Diagnosis/Death:						
All ages	23.3	24.2	20.3	4.1	3.9	7.2
Under 65	14.9	15.5	11.1	1.5	1.4	2.4
65 and over	81.0	84.4	83.6	22.7	21.2	40.3
All ages (IARC world std) ^c	17.1	17.8	14.8	2.6	2.4	4.6
<u>Age-Specific Rates, 2002-2006</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	0.4	0.3	-	-	-	-
25-29	1.4	1.4	1.3	0.1	0.1	-
30-34	3.7	3.7	3.0	0.2	0.2	0.3
35-39	7.2	7.0	5.8	0.4	0.4	0.5
40-44	13.0	13.3	9.1	0.9	0.9	1.2
45-49	23.2	23.6	13.5	1.8	1.7	2.3
50-54	43.7	46.5	24.7	3.7	3.5	4.8
55-59	68.8	72.6	49.8	7.0	6.6	11.1
60-64	83.5	86.9	83.1	11.6	10.4	23.5
65-69	89.3	92.6	99.1	16.4	14.5	34.5
70-74	84.7	88.5	87.0	20.6	18.8	41.1
75-79	80.8	84.5	85.0	23.7	22.2	43.1
80-84	77.3	80.8	69.2	28.1	27.2	44.0
85+	59.9	61.9	56.4	32.9	32.7	42.5

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.