

Table 10.8
Kaposi Sarcoma

Survival Rates^a
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Relative Survival Rates (SEER ^b)																			
1-year	95.7	65.1	59.8	61.5	62.4	63.0	58.9	55.2	61.5	69.3	73.7	73.0	76.7	73.4	79.3	76.5	74.8	81.1	82.5
2-year	89.8	41.7	33.2	32.9	29.6	28.1	29.2	27.1	45.7	59.6	63.9	63.2	64.4	65.6	73.1	69.8	72.2	72.2	
3-year	88.0	32.6	19.2	19.1	14.6	15.9	18.7	20.4	41.7	53.8	60.1	58.5	60.4	59.3	68.3	65.1	66.7		
4-year	85.4	27.9	12.9	13.7	9.9	10.6	14.4	17.9	37.5	50.5	56.1	55.5	55.0	55.7	64.7	64.5			
5-year	85.4	24.0	9.4	9.1	7.6	8.5	12.8	16.7	34.8	47.3	54.5	54.3	53.8	52.9	63.4				
6-year	77.7	20.0	7.7	7.0	6.6	7.9	12.3	15.5	34.1	44.7	51.7	51.8	50.6	51.6					
7-year	76.8	18.3	6.4	6.4	6.2	7.1	11.7	14.5	32.3	43.6	47.7	49.8	49.7						
8-year	76.8	15.8	5.5	5.8	5.7	7.0	11.1	14.0	31.0	43.4	44.4	48.5							
9-year	76.8	14.3	4.9	5.5	5.1	6.9	10.8	13.4	30.0	41.4	41.0								
10-year	76.8	12.4	4.6	5.5	5.0	6.6	10.5	13.1	28.8	41.2									
11-year	72.7	11.0	4.3	5.3	4.8	6.6	9.8	12.7	27.7										
12-year	72.7	10.6	4.1	5.3	4.6	6.4	9.7	12.5											
13-year	72.7	9.8	3.8	5.1	4.3	6.4	9.6												
14-year	72.7	9.6	3.6	4.5	4.3	6.1													
15-year	72.7	8.9	3.5	4.2	4.3														
16-year	72.7	8.6	3.3	4.1															
17-year	72.7	8.3	3.3																
18-year	72.7	7.4																	
19-year	72.7	7.4																	
20-year	72.7	6.8																	

^a Survival rates are relative rates expressed as percents.
^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).