

Table 10.13
Kaposi SarcomaEstimated United States Cancer Prevalence Counts^a on January 1, 2006
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <16 ^e	0 to <31 ^e	>=31 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	6,695	6,311	5,712	1,642	545	231	19,214	21,168	+	+
	Males	6,215	5,970	5,428	1,578	511	115	18,087	19,850	+	+
	Females	480	341	284	64	34	116	1,127	1,318	+	+
White ^b	Both Sexes	4,701	4,681	4,824	1,351	447	172	14,600	16,208	+	+
	Males	4,338	4,457	4,599	1,310	424	79	13,789	15,240	+	+
	Females	363	224	225	41	23	93	811	968	+	+
Black ^b	Both Sexes	1,662	1,254	601	112	52	12	3,541	3,694	+	+
	Males	1,563	1,181	542	112	52	12	3,310	3,463	+	+
	Females	99	73	59	0	0	0	231	231	+	+
Asian/ Pacific Islander ^c	Both Sexes	105	83	48	+	+	+	238	+	+	+
	Males	100	80	48	+	+	+	230	+	+	+
	Females	5	3	0	+	+	+	8	+	+	+
Hispanic ^d	Both Sexes	1,235	1,094	871	+	+	+	3,254	+	+	+
	Males	1,169	1,050	837	+	+	+	3,105	+	+	+
	Females	66	44	34	+	+	+	149	+	+	+

Estimated prevalence percent^a on January 1, 2006, of the SEER 11 population diagnosed in the previous 16 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0074%	-	-	0.0008%	0.0080%	0.0197%	0.0132%	0.0070%	0.0078%	0.0136%	0.0073%
	Males	0.0142%	-	-	0.0014%	0.0156%	0.0389%	0.0265%	0.0136%	0.0149%	0.0259%	0.0143%
	Females	0.0007%	-	-	-	0.0002%	0.0006%	0.0004%	0.0012%	0.0023%	0.0067%	0.0006%
White ^c	Both Sexes	0.0074%	-	-	0.0006%	0.0070%	0.0201%	0.0132%	0.0072%	0.0085%	0.0139%	0.0072%
	Males	0.0141%	-	-	0.0011%	0.0134%	0.0390%	0.0263%	0.0136%	0.0158%	0.0266%	0.0140%
	Females	0.0007%	-	-	-	-	0.0005%	0.0003%	0.0012%	0.0028%	0.0070%	0.0006%
Black ^c	Both Sexes	0.0098%	-	-	0.0019%	0.0187%	0.0272%	0.0160%	0.0085%	0.0045%	0.0111%	0.0103%
	Males	0.0197%	-	-	0.0036%	0.0397%	0.0567%	0.0335%	0.0177%	0.0097%	-	0.0212%
	Females	0.0009%	-	-	-	-	0.0013%	-	-	-	0.0115%	0.0010%
Asian/ Pacific Islander ^c	Both Sexes	0.0018%	-	-	-	0.0014%	0.0033%	0.0045%	0.0023%	0.0026%	0.0044%	0.0017%
	Males	0.0035%	-	-	-	0.0029%	0.0067%	0.0096%	0.0044%	0.0060%	0.0091%	0.0036%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic ^d	Both Sexes	0.0075%	-	-	0.0009%	0.0108%	0.0233%	0.0148%	0.0114%	0.0145%	0.0387%	0.0101%
	Males	0.0139%	-	-	0.0017%	0.0199%	0.0449%	0.0298%	0.0218%	0.0297%	0.0702%	0.0193%
	Females	0.0007%	-	-	-	-	-	-	0.0026%	-	0.0204%	0.0012%

^a US 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER registries and 1/1/2006 US population estimates based on the average of 2005 and 2006 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 31 years for 1975-2006 SEER 9 data; 16 years for 1990-2006 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 31 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <31 and >=31. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.