

Table 15.14
Cancer of the Lung and Bronchus (Invasive)

Survival Rates^a
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Relative Survival Rates (SEER ^b)																			
1-year	36.7	38.7	40.0	40.3	41.2	41.1	40.8	41.0	41.5	41.0	41.7	41.2	42.7	42.3	41.6	42.0	43.1	43.5	44.4
2-year	21.7	22.6	23.1	23.6	23.9	24.7	24.6	24.5	25.4	25.3	25.4	25.8	26.4	26.5	26.2	27.1	27.1	28.0	
3-year	16.7	17.4	17.5	18.0	18.7	19.0	18.9	19.1	19.5	19.7	19.8	20.2	20.2	20.8	20.3	21.4	21.2		
4-year	14.4	14.9	15.0	15.3	16.0	16.2	16.2	16.3	17.0	16.7	17.0	17.4	17.4	18.1	17.3	18.1			
5-year	13.0	13.4	13.5	13.8	14.2	14.4	14.7	14.8	15.1	15.1	15.3	15.1	15.7	16.2	15.6				
6-year	12.0	12.3	12.4	12.7	13.2	13.2	13.6	13.6	13.7	13.8	13.9	13.8	14.4	14.8					
7-year	11.4	11.5	11.5	11.7	12.0	12.2	12.6	12.6	13.0	12.8	12.9	12.9	13.4						
8-year	10.8	10.7	10.8	11.0	11.2	11.3	11.7	11.9	12.2	12.0	12.0	11.9							
9-year	10.2	10.0	10.1	10.1	10.6	10.6	11.1	11.2	11.6	11.2	11.2								
11-year	9.7	9.4	9.6	9.4	10.1	10.0	10.3	10.6	11.0	10.5									
10-year	9.2	9.1	9.0	8.9	9.5	9.4	9.8	10.1	10.5										
12-year	8.9	8.6	8.6	8.3	9.0	9.1	9.2	9.4											
13-year	8.5	8.2	8.2	8.1	8.4	8.6	8.8												
14-year	8.2	7.9	7.8	7.9	7.8	8.4													
15-year	7.8	7.5	7.5	7.6	7.4														
16-year	7.6	7.2	7.2	7.3															
17-year	7.4	6.9	6.8																
18-year	7.0	6.6																	
19-year	6.8	6.2																	
20-year	6.6	5.8																	

^a Survival rates are relative rates expressed as percents.
^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).