

Table 15.15
Cancer of the Lung and Bronchus (Invasive)

Survival Rates^a
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Relative Survival Rates (SEER ^b)																			
1-year	34.7	36.8	37.9	38.1	39.0	39.1	38.2	38.7	39.3	39.1	39.3	38.4	39.6	40.0	38.3	38.4	40.6	40.5	40.8
2-year	20.0	20.8	21.5	21.4	22.1	23.2	22.4	22.1	23.6	23.5	23.5	23.2	23.3	24.3	22.7	23.6	24.2	25.0	
3-year	15.1	15.7	16.0	16.1	17.1	17.6	16.6	16.7	17.6	18.2	18.1	17.8	17.0	18.7	17.7	18.4	18.7		
4-year	12.9	13.4	13.5	13.5	14.2	14.8	14.0	14.0	15.2	15.0	15.3	15.1	14.4	16.1	14.8	15.5			
5-year	11.5	12.0	12.1	12.0	12.6	13.1	12.6	12.7	13.5	13.5	13.8	12.9	13.0	14.6	13.2				
6-year	10.6	10.9	11.1	11.0	11.7	11.8	11.6	11.5	12.2	12.2	12.3	11.4	11.8	13.1					
7-year	10.0	10.1	10.3	10.2	10.6	10.9	10.8	10.6	11.7	11.2	11.3	10.6	10.9						
8-year	9.5	9.3	9.6	9.3	9.7	9.9	9.9	9.9	10.8	10.5	10.6	9.7							
9-year	8.9	8.7	9.1	8.4	9.2	9.2	9.4	9.5	10.2	10.0	9.8								
11-year	8.4	8.2	8.6	7.7	8.9	8.6	8.9	9.0	9.5	9.2									
10-year	8.0	7.9	8.0	7.2	8.2	7.9	8.2	8.7	9.1										
12-year	7.6	7.5	7.5	6.7	7.9	7.7	7.7	7.9											
13-year	7.2	7.1	7.1	6.5	7.2	7.3	7.1												
14-year	7.0	6.8	6.8	6.3	6.6	7.1													
15-year	6.6	6.4	6.5	5.9	6.2														
16-year	6.4	6.1	6.3	5.5															
17-year	6.2	5.7	5.8																
18-year	5.9	5.5																	
19-year	5.7	5.2																	
20-year	5.5	4.9																	

^a

Survival rates are relative rates expressed as percents.

^b

Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).