

Table 35.2
Soft Tissue Sarcomas

Site Distribution of Sarcomas and Other Histologies at Parenchymatous and
Nonparenchymatous sites (Soft Tissues, Heart and Mediastinum, Retroperitoneum and peritoneum, Pleura, Spleen)
Among Histologically Confirmed Cases
2002-2006, All Races, Both Sexes

Primary Site (ICD-O-3 Site Code)	All Sarcomas		Epidermoid Carcinoma		Adenocarcinoma		Other Specified Types	
	Count	Percent	Count	Percent	Count	Percent	Count	Percent
All Nonparenchymatous Sites (C38.0-C38.4, C38.8, C42.2, C48.0-C48.2, C48.8, C49.0-C49.9)	15,276	30.1%	80	0.0%	2,067	0.2%	4,493	1.1%
Soft tissues (Connective, Subcutaneous & Other Soft Tissue) (C49.0-C49.9)	10,192	20.1%	74	0.0%	27	0.0%	1,718	0.4%
Retroperitoneum & Peritoneum (C48.0-C48.2, C48.8)	1,709	3.4%	-	-	2,039	0.2%	777	0.2%
Pleura (C38.4)	3,097	6.1%	-	-	-	-	175	0.0%
Heart & Mediastinum (C38.0-C38.3, C38.8)	250	0.5%	-	-	-	-	670	0.2%
Spleen (C42.2)	28	0.1%	-	-	-	-	1,153	0.3%
Parenchymatous Sites (All Other Primary Sites)	35,530	69.9%	195,485	100.0%	962,663	99.8%	402,654	98.9%
All Sites	50,806	100.0%	195,565	100.0%	964,730	100.0%	407,147	100.0%

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Percents may not sum to 100 due to rounding.

- Statistic not shown due to fewer than 16 cases during the time period.