

Table 34.1
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 1999-2006
Both Sexes by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	1,720,790	65.6	0.0	196,969	58.6	0.1	124,755	64.2	0.2	7,915	57.3	0.6	184,205	64.4	0.1
Oral Cavity & Pharynx	37,215	66.2	0.3	4,408	46.2	0.9	3,371	66.3	1.0	213	59.4	4.0	3,247	61.5	1.0
Esophagus	16,045	19.4	0.4	2,155	13.1	0.9	890	17.6	1.6	95	12.9	3.9	1,291	16.4	1.3
Stomach	24,944	25.1	0.3	4,352	27.6	0.8	5,117	35.3	0.8	276	22.2	2.9	5,659	25.6	0.7
Colon & Rectum	174,839	64.3	0.1	21,844	56.3	0.4	15,478	68.0	0.4	1,006	62.4	1.8	17,711	62.0	0.4
Liver & Intrahepatic Bile Duct	19,971	15.2	0.3	3,274	11.7	0.8	5,645	20.5	0.7	255	16.2	2.9	5,072	15.6	0.7
Pancreas	39,894	5.5	0.1	5,435	5.5	0.4	3,203	7.1	0.6	224	5.8	1.8	4,523	6.1	0.4
Larynx	12,972	67.5	0.5	2,304	55.4	1.2	534	70.0	2.3	47	62.8	8.4	1,228	65.5	1.6
Lung & Bronchus	218,174	17.2	0.1	27,054	14.0	0.3	14,074	18.5	0.4	932	12.3	1.3	13,509	15.2	0.4
Melanoma of the Skin	79,114	89.0	0.1	386	73.3	2.6	551	76.9	2.1	92	78.1	4.8	2,665	84.0	0.8
Breast	261,874	88.3	0.1	28,514	77.5	0.3	21,368	90.3	0.2	1,132	84.6	1.3	27,579	85.3	0.3
Cervix uteri	17,420	73.1	0.4	2,945	64.3	1.0	1,983	73.6	1.1	127	69.5	4.6	5,306	75.0	0.7
Corpus & Uterus, NOS	50,031	83.1	0.2	4,585	62.4	0.8	3,952	84.9	0.6	245	82.5	2.7	5,362	81.1	0.6
Ovary ^d	28,450	43.1	0.3	2,364	36.7	1.2	2,270	55.9	1.2	162	43.9	4.7	3,446	47.1	1.0
Prostate	277,653	93.5	0.1	43,633	91.2	0.2	16,250	93.8	0.2	801	89.7	1.3	28,028	92.3	0.2
Testis	13,666	95.9	0.2	414	89.0	1.6	480	93.9	1.2	99	96.9	1.7	2,553	92.6	0.6
Urinary Bladder	75,198	79.5	0.2	3,983	65.7	0.9	3,028	79.6	0.9	156	69.8	4.3	4,429	75.6	0.8
Kidney & Renal Pelvis	46,710	69.1	0.2	5,370	70.3	0.7	2,532	70.9	1.0	406	67.0	2.7	6,465	69.3	0.7
Brain & Nervous System	27,218	33.7	0.3	1,942	39.8	1.2	1,502	40.4	1.5	111	44.5	5.8	4,027	45.6	0.9
Thyroid	36,920	96.2	0.1	2,637	95.0	0.5	4,409	95.9	0.3	201	95.6	1.6	6,190	96.0	0.3
Hodgkin Lymphoma	12,005	86.0	0.4	1,485	82.8	1.1	625	84.5	1.7	34	86.1	6.7	2,022	83.8	0.9
Non-Hodgkin Lymphoma	73,654	67.3	0.2	6,370	61.6	0.7	5,231	65.0	0.7	286	61.5	3.2	8,973	64.1	0.6
Myeloma	19,134	41.0	0.5	4,369	44.1	1.0	1,280	44.2	1.8	126	33.7	5.3	2,671	42.3	1.3
Leukemia	47,509	54.7	0.3	4,070	49.5	0.9	3,035	51.3	1.0	224	58.9	3.8	7,002	56.5	0.7
Mesothelioma	4,041	7.3	0.5	198	12.9	3.1	134	15.7	3.8	-	-	-	456	10.7	1.9

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006.

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 1999-2006 with follow-up through 2007.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.2
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 1999-2006
Males by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	886,212	65.6	0.1	105,948	60.9	0.2	60,322	59.8	0.2	3,750	53.5	0.9	91,246	63.3	0.2
Oral Cavity & Pharynx	25,845	65.9	0.3	3,084	42.1	1.0	2,123	64.0	1.2	145	55.1	4.9	2,198	59.2	1.2
Esophagus	12,434	19.3	0.4	1,513	11.7	1.0	678	16.3	1.8	74	10.6	4.2	1,026	16.1	1.4
Stomach	15,354	23.5	0.4	2,398	24.7	1.0	2,932	34.9	1.0	170	20.6	3.5	3,183	24.9	0.9
Colon & Rectum	88,982	64.9	0.2	10,076	55.7	0.6	8,084	67.5	0.6	487	59.0	2.7	9,435	61.6	0.6
Liver & Intrahepatic Bile Duct	14,011	15.5	0.4	2,381	10.9	0.9	3,917	21.1	0.8	179	17.1	3.3	3,574	15.1	0.8
Pancreas	19,623	5.4	0.2	2,459	4.5	0.5	1,573	6.4	0.8	102	5.4	2.4	2,118	5.0	0.6
Larynx	10,365	67.9	0.5	1,849	55.1	1.4	469	70.8	2.4	32	52.5	10.8	1,047	63.8	1.7
Lung & Bronchus	116,067	15.0	0.1	15,669	12.5	0.3	8,449	16.0	0.5	503	12.0	1.7	7,629	13.6	0.5
Melanoma of the Skin	43,858	86.6	0.2	165	67.0	4.1	267	72.6	3.3	46	67.8	7.5	1,081	77.3	1.5
Prostate	277,653	93.5	0.1	43,633	91.2	0.2	16,250	93.8	0.2	801	89.7	1.3	28,028	92.3	0.2
Testis	13,666	95.9	0.2	414	89.0	1.6	480	93.9	1.2	99	96.9	1.7	2,553	92.6	0.6
Urinary Bladder	56,583	80.9	0.2	2,518	70.0	1.1	2,296	81.3	1.0	119	73.3	4.8	3,258	78.0	0.9
Kidney & Renal Pelvis	28,693	68.8	0.3	3,200	69.3	0.9	1,573	71.1	1.3	231	62.4	3.7	3,835	67.6	0.9
Brain & Nervous System	15,197	32.1	0.4	1,021	35.0	1.7	812	37.0	1.9	54	45.4	8.3	2,193	44.2	1.2
Thyroid	8,792	93.3	0.3	494	92.2	1.4	859	93.3	1.0	36	88.5	5.4	1,123	92.5	0.9
Hodgkin Lymphoma	6,504	84.9	0.5	792	78.5	1.7	347	83.5	2.3	-	-	-	1,163	83.0	1.2
Non-Hodgkin Lymphoma	39,292	66.2	0.3	3,480	57.7	0.9	2,792	63.1	1.0	150	60.4	4.5	4,874	61.7	0.8
Myeloma	10,505	42.6	0.6	2,114	44.4	1.4	650	46.1	2.6	55	40.0	8.4	1,390	45.6	1.8
Leukemia	27,363	55.1	0.3	2,197	49.4	1.2	1,717	51.1	1.3	123	60.4	4.9	3,939	56.8	0.9
Mesothelioma	3,136	5.6	0.5	138	11.2	3.8	101	14.4	4.4	-	-	-	332	9.7	2.1

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006.

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 1999-2006 with follow-up through 2007.

^b Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.3
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 1999-2006
Females by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	834,578	65.6	0.1	91,021	56.0	0.2	64,433	68.2	0.2	4,165	60.6	0.9	92,959	65.5	0.2
Oral Cavity & Pharynx	11,370	66.7	0.5	1,324	55.6	1.5	1,248	70.2	1.5	68	70.6	6.2	1,049	66.2	1.7
Esophagus	3,611	19.9	0.8	642	16.7	1.8	212	21.2	3.3	-	-	-	265	17.8	2.9
Stomach	9,590	27.9	0.5	1,954	31.2	1.2	2,185	35.7	1.2	106	24.6	4.9	2,476	26.6	1.1
Colon & Rectum	85,857	63.7	0.2	11,768	56.7	0.5	7,394	68.6	0.6	519	65.5	2.4	8,276	62.4	0.6
Liver & Intrahepatic Bile Duct	5,960	14.7	0.6	893	13.5	1.6	1,728	19.2	1.2	76	16.6	4.9	1,498	16.6	1.2
Pancreas	20,271	5.6	0.2	2,976	6.3	0.5	1,630	7.8	0.8	122	6.1	2.7	2,405	7.1	0.7
Larynx	2,607	65.8	1.1	455	56.6	2.8	65	65.2	7.0	-	-	-	181	74.9	3.5
Lung & Bronchus	102,107	19.7	0.1	11,385	16.1	0.4	5,625	22.1	0.7	429	12.5	2.0	5,880	17.2	0.6
Melanoma of the Skin	35,256	92.0	0.2	221	78.1	3.3	284	80.7	2.7	46	90.6	4.5	1,584	88.5	0.9
Breast	260,127	88.3	0.1	28,249	77.5	0.3	21,271	90.3	0.2	1,129	84.6	1.3	27,446	85.3	0.3
Cervix uteri	17,420	73.1	0.4	2,945	64.3	1.0	1,983	73.6	1.1	127	69.5	4.6	5,306	75.0	0.7
Corpus & Uterus, NOS	50,031	83.1	0.2	4,585	62.4	0.8	3,952	84.9	0.6	245	82.5	2.7	5,362	81.1	0.6
Ovary ^d	28,450	43.1	0.3	2,364	36.7	1.2	2,270	55.9	1.2	162	43.9	4.7	3,446	47.1	1.0
Urinary Bladder	18,615	75.2	0.4	1,465	58.2	1.5	732	74.4	1.9	37	59.2	8.6	1,171	68.8	1.6
Kidney & Renal Pelvis	18,017	69.7	0.4	2,170	71.7	1.1	959	70.6	1.7	175	73.6	3.6	2,630	71.8	1.0
Brain & Nervous System	12,021	35.6	0.5	921	45.2	1.8	690	44.4	2.2	57	44.6	8.0	1,834	47.2	1.3
Thyroid	28,128	97.1	0.1	2,143	95.6	0.5	3,550	96.5	0.3	165	97.1	1.4	5,067	96.7	0.3
Hodgkin Lymphoma	5,501	87.2	0.5	693	87.6	1.4	278	85.6	2.5	-	-	-	859	84.9	1.4
Non-Hodgkin Lymphoma	34,362	68.6	0.3	2,890	66.2	1.0	2,439	67.0	1.1	136	62.7	4.7	4,099	66.9	0.8
Myeloma	8,629	39.1	0.7	2,255	43.8	1.4	630	42.3	2.5	71	28.8	6.7	1,281	38.8	1.8
Leukemia	20,146	54.2	0.4	1,873	49.7	1.4	1,318	51.5	1.6	101	56.6	6.2	3,063	56.1	1.0
Mesothelioma	905	13.1	1.4	60	15.7	5.6	33	18.5	7.4	-	-	-	124	12.7	4.2

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006.

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 1999-2006 with follow-up through 2007.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.4

Cancer of the Female Breast

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 1999-2006

Race/Ethnicity	All Ages				Ages <50				Ages 50+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	312,972	100.0	87.5	0.1	78,736	100.0	88.3	0.1	234,236	100.0	87.2	0.1
White	260,127	83.1	88.3	0.1	61,027	77.5	89.7	0.1	199,100	85.0	87.9	0.1
White Non-Hispanic	233,660	74.7	88.6	0.1	50,981	64.7	90.6	0.2	182,679	78.0	88.1	0.1
White Hispanic	26,467	8.5	85.4	0.3	10,046	12.8	84.9	0.4	16,421	7.0	85.7	0.3
Black	28,249	9.0	77.5	0.3	9,318	11.8	77.7	0.5	18,931	8.1	77.5	0.4
American Indian/Alaska Native ^c	1,129	0.4	84.6	1.3	391	0.5	82.9	2.3	738	0.3	85.4	1.6
Asian	19,644	6.3	90.7	0.3	6,815	8.7	91.1	0.4	12,829	5.5	90.5	0.3
Asian Indian, Pakistani	1,396	0.4	89.3	1.0	544	0.7	90.7	1.6	852	0.4	88.3	1.4
Chinese	3,952	1.3	90.3	0.6	1,499	1.9	91.8	0.9	2,453	1.0	89.4	0.8
Filipino	5,475	1.7	88.6	0.5	1,789	2.3	88.6	0.9	3,686	1.6	88.5	0.6
Japanese	3,629	1.2	93.0	0.5	731	0.9	93.3	1.1	2,898	1.2	93.0	0.5
Korean	1,341	0.4	90.8	1.0	611	0.8	90.2	1.5	730	0.3	91.3	1.3
Vietnamese	1,126	0.4	90.3	1.1	507	0.6	89.8	1.6	619	0.3	90.8	1.4
Other Asian ^d	2,725	0.9	93.3	0.6	1,134	1.4	94.3	0.8	1,591	0.7	92.5	0.8
Pacific Islander	1,627	0.5	85.8	1.0	517	0.7	83.6	1.9	1,110	0.5	86.9	1.2
Hawaiian	1,143	0.4	87.9	1.1	341	0.4	88.2	2.1	802	0.3	87.9	1.3
Other Pacific Islander ^e	484	0.2	80.6	2.1	176	0.2	74.4	3.8	308	0.1	84.2	2.5
Hispanic ^f	27,446	8.8	85.3	0.3	10,399	13.2	84.8	0.4	17,047	7.3	85.6	0.3

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006.

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 1999-2006 with follow-up through 2007.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.5

Cancer of the Colon and Rectum

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 1999-2006

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	214,789	100.0	63.9	0.1	108,501	100.0	64.4	0.2	106,288	100.0	63.5	0.2
White	174,839	81.4	64.3	0.1	88,982	82.0	64.9	0.2	85,857	80.8	63.7	0.2
White Non-Hispanic	157,755	73.4	64.6	0.1	79,837	73.6	65.3	0.2	77,918	73.3	63.8	0.2
White Hispanic	17,084	8.0	61.8	0.4	9,145	8.4	61.3	0.6	7,939	7.5	62.4	0.7
Black	21,844	10.2	56.3	0.4	10,076	9.3	55.7	0.6	11,768	11.1	56.7	0.5
American Indian/Alaska Native ^c	1,006	0.5	62.4	1.8	487	0.4	59.0	2.7	519	0.5	65.5	2.4
Asian	14,607	6.8	68.3	0.5	7,580	7.0	67.9	0.6	7,027	6.6	68.9	0.6
Asian Indian, Pakistani	587	0.3	70.3	2.3	336	0.3	72.2	3.0	251	0.2	67.3	3.8
Chinese	3,633	1.7	69.4	0.9	1,855	1.7	69.4	1.2	1,778	1.7	69.4	1.3
Filipino	3,034	1.4	65.3	1.0	1,643	1.5	64.3	1.4	1,391	1.3	66.6	1.5
Japanese	3,453	1.6	69.2	0.9	1,729	1.6	69.2	1.3	1,724	1.6	69.2	1.3
Korean	1,316	0.6	66.8	1.5	682	0.6	64.9	2.2	634	0.6	68.9	2.2
Vietnamese	1,018	0.5	65.1	1.8	534	0.5	63.5	2.5	484	0.5	67.1	2.5
Other Asian ^d	1,566	0.7	72.5	1.3	801	0.7	72.5	1.9	765	0.7	72.6	1.9
Pacific Islander	871	0.4	62.4	1.9	504	0.5	61.5	2.5	367	0.3	63.6	2.9
Hawaiian	631	0.3	62.4	2.3	375	0.3	61.2	3.0	256	0.2	64.1	3.5
Other Pacific Islander ^e	240	0.1	62.4	3.6	129	0.1	62.3	4.7	111	0.1	62.4	5.6
Hispanic ^f	17,711	8.2	62.0	0.4	9,435	8.7	61.6	0.6	8,276	7.8	62.4	0.6

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006.

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 1999-2006 with follow-up through 2007.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.6

Cancer of the Liver and Intrahepatic Bile Duct

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 1999-2006

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	29,342	100.0	15.9	0.3	20,621	100.0	16.1	0.3	8,721	100.0	15.6	0.5
White	19,971	68.1	15.2	0.3	14,011	67.9	15.5	0.4	5,960	68.3	14.7	0.6
White Non-Hispanic	15,041	51.3	15.2	0.4	10,523	51.0	15.5	0.5	4,518	51.8	14.3	0.6
White Hispanic	4,930	16.8	15.4	0.7	3,488	16.9	15.1	0.8	1,442	16.5	16.0	1.2
Black	3,274	11.2	11.7	0.8	2,381	11.5	10.9	0.9	893	10.2	13.5	1.6
American Indian/Alaska Native ^c	255	0.9	16.2	2.9	179	0.9	17.1	3.3	76	0.9	16.6	4.9
Asian	5,378	18.3	20.8	0.7	3,718	18.0	21.5	0.9	1,660	19.0	19.3	1.2
Asian Indian, Pakistani	147	0.5	23.2	4.3	105	0.5	23.6	5.4	42	0.5	21.7	6.8
Chinese	1,394	4.8	23.0	1.4	983	4.8	24.5	1.7	411	4.7	19.3	2.4
Filipino	792	2.7	17.4	1.8	564	2.7	18.7	2.2	228	2.6	15.4	3.1
Japanese	583	2.0	18.1	2.1	270	1.3	21.4	3.2	313	3.6	15.8	2.7
Korean	755	2.6	22.4	2.0	500	2.4	20.1	2.3	255	2.9	27.5	3.5
Vietnamese	1,050	3.6	20.6	1.7	808	3.9	20.9	1.9	242	2.8	19.3	3.4
Other Asian ^d	657	2.2	20.5	1.9	488	2.4	21.3	2.3	169	1.9	18.1	3.7
Pacific Islander	267	0.9	15.0	2.7	199	1.0	13.2	3.0	68	0.8	20.4	6.0
Hawaiian	149	0.5	13.0	3.4	106	0.5	11.9	3.8	43	0.5	16.9	6.4
Other Pacific Islander ^e	118	0.4	18.7	4.4	93	0.5	16.5	4.3	25	0.3	25.4	12.4
Hispanic ^f	5,072	17.3	15.6	0.7	3,574	17.3	15.1	0.8	1,498	17.2	16.6	1.2

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006.

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 1999-2006 with follow-up through 2007.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.7

Cancer of the Lung and Bronchus

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 1999-2006

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	261,034	100.0	17.0	0.1	141,130	100.0	14.8	0.1	119,904	100.0	19.5	0.1
White	218,174	83.6	17.2	0.1	116,067	82.2	15.0	0.1	102,107	85.2	19.7	0.1
White Non-Hispanic	205,127	78.6	17.3	0.1	108,716	77.0	15.1	0.1	96,411	80.4	19.8	0.2
White Hispanic	13,047	5.0	15.2	0.4	7,351	5.2	13.5	0.5	5,696	4.8	17.3	0.6
Black	27,054	10.4	14.0	0.3	15,669	11.1	12.5	0.3	11,385	9.5	16.1	0.4
American Indian/Alaska Native ^c	932	0.4	12.3	1.3	503	0.4	12.0	1.7	429	0.4	12.5	2.0
Asian	12,801	4.9	18.7	0.4	7,706	5.5	16.2	0.5	5,095	4.2	22.5	0.7
Asian Indian, Pakistani	458	0.2	23.3	2.4	311	0.2	22.9	2.8	147	0.1	24.2	4.6
Chinese	3,360	1.3	18.7	0.9	1,965	1.4	15.9	1.0	1,395	1.2	22.6	1.4
Filipino	3,388	1.3	18.8	0.8	2,173	1.5	15.9	1.0	1,215	1.0	23.7	1.5
Japanese	2,155	0.8	16.7	1.0	1,181	0.8	13.5	1.2	974	0.8	20.6	1.6
Korean	987	0.4	16.1	1.5	571	0.4	14.1	1.8	416	0.3	18.7	2.5
Vietnamese	1,336	0.5	19.7	1.4	871	0.6	18.7	1.6	465	0.4	21.6	2.4
Other Asian ^d	1,117	0.4	22.1	1.6	634	0.4	18.4	2.1	483	0.4	26.8	2.6
Pacific Islander	1,273	0.5	16.3	1.2	743	0.5	14.6	1.6	530	0.4	18.4	2.0
Hawaiian	930	0.4	16.3	1.4	525	0.4	14.4	1.9	405	0.3	18.5	2.2
Other Pacific Islander ^e	343	0.1	15.6	2.8	218	0.2	13.6	3.9	125	0.1	17.6	4.5
Hispanic ^f	13,509	5.2	15.2	0.4	7,629	5.4	13.6	0.5	5,880	4.9	17.2	0.6

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006.

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 1999-2006 with follow-up through 2007.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.8

Cancer of the Prostate

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 17 Registries, 1999-2006

Race/Ethnicity	All Ages				Ages <65				Ages 65+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	348,441	100.0	93.3	0.1	134,545	100.0	96.5	0.1	213,896	100.0	91.2	0.1
White	277,653	79.7	93.5	0.1	105,036	78.1	96.8	0.1	172,617	80.7	91.4	0.1
White Non-Hispanic	250,962	72.0	93.6	0.1	94,883	70.5	97.0	0.1	156,079	73.0	91.5	0.1
White Hispanic	26,691	7.7	92.3	0.2	10,153	7.5	94.8	0.3	16,538	7.7	90.7	0.3
Black	43,633	12.5	91.2	0.2	21,377	15.9	94.8	0.2	22,256	10.4	87.7	0.3
American Indian/Alaska Native ^c	801	0.2	89.7	1.3	342	0.3	92.1	1.8	459	0.2	87.9	1.8
Asian	15,223	4.4	94.2	0.2	4,243	3.2	97.1	0.3	10,980	5.1	93.0	0.3
Asian Indian, Pakistani	1,200	0.3	95.2	0.8	485	0.4	98.5	0.7	715	0.3	92.8	1.2
Chinese	3,422	1.0	94.8	0.5	743	0.6	97.6	0.7	2,679	1.3	94.0	0.6
Filipino	4,325	1.2	93.2	0.5	1,379	1.0	96.0	0.6	2,946	1.4	91.9	0.6
Japanese	3,248	0.9	93.7	0.5	650	0.5	97.0	0.8	2,598	1.2	92.9	0.6
Korean	653	0.2	93.7	1.2	153	0.1	97.0	1.5	500	0.2	92.5	1.6
Vietnamese	752	0.2	93.3	1.1	252	0.2	95.9	1.6	500	0.2	92.0	1.4
Other Asian ^d	1,623	0.5	96.3	0.6	581	0.4	98.4	0.6	1,042	0.5	95.1	0.8
Pacific Islander	1,027	0.3	88.5	1.2	352	0.3	91.9	1.6	675	0.3	86.5	1.7
Hawaiian	662	0.2	89.7	1.4	219	0.2	93.9	1.7	443	0.2	87.0	2.0
Other Pacific Islander ^e	365	0.1	86.4	2.2	133	0.1	87.8	3.4	232	0.1	85.5	2.9
Hispanic ^f	28,028	8.0	92.3	0.2	10,621	7.9	94.9	0.3	17,407	8.1	90.8	0.3

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006.

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 1999-2006 with follow-up through 2007.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.