

Table 1.21  
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2007<sup>a</sup>  
Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			32-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 32 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 32 Years <sup>d</sup>
All Sites	Both Sexes	4,128,739	4,208,495	4,587,516	11,196,431	11,396,724
	Male	2,110,305	2,140,911	2,323,221	5,248,197	5,310,771
	Female	2,018,434	2,067,584	2,264,295	5,948,234	6,085,953
Oral Cavity & Pharynx	Both Sexes	91,178	103,795	107,311	235,609	258,515
	Male	62,485	70,353	72,387	153,118	166,208
	Female	28,693	33,442	34,924	82,491	92,307
Esophagus	Both Sexes	18,697	22,371	22,384	28,549	33,324
	Male	14,341	17,140	17,153	21,695	25,162
	Female	4,356	5,231	5,231	6,854	8,162
Stomach	Both Sexes	32,686	39,128	39,293	63,730	72,980
	Male	19,323	23,318	23,348	36,215	41,618
	Female	13,363	15,810	15,945	27,515	31,362
Colon & Rectum	Both Sexes	404,997	468,117	476,559	1,076,622	1,199,140
	Male	202,765	234,821	238,838	527,903	584,590
	Female	202,232	233,296	237,721	548,719	614,550
Liver & Intrahepatic Bile Duct	Both Sexes	20,688	23,505	23,517	27,353	30,585
	Male	14,728	16,576	16,588	18,577	20,660
	Female	5,960	6,929	6,929	8,776	9,925
Pancreas	Both Sexes	24,579	29,347	29,358	32,616	38,193
	Male	12,128	14,558	14,569	15,897	18,650
	Female	12,451	14,789	14,789	16,719	19,543
Larynx	Both Sexes	30,163	35,886	36,142	87,367	97,387
	Male	23,925	28,340	28,574	69,746	77,350
	Female	6,238	7,546	7,568	17,621	20,037
Lung & Bronchus	Both Sexes	213,542	267,851	273,542	360,606	436,310
	Male	100,425	126,335	128,613	167,066	201,817
	Female	113,117	141,516	144,929	193,540	234,493
Melanoma of the Skin	Both Sexes	242,029	271,164	281,289	751,279	806,659
	Male	127,243	145,557	152,065	371,049	402,671
	Female	114,786	125,607	129,224	380,230	403,988
Breast	Female	796,935	857,596	904,022	2,509,315	2,661,082
Cervix	Female	38,618	40,422	40,532	195,066	200,672
Corpus & Uterus, NOS	Female	148,471	167,653	167,699	527,264	573,628
Ovary <sup>f</sup>	Female	53,433	62,494	62,552	156,901	176,514

<sup>a</sup> U.S. 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER 9 registries and 1/1/2007 U.S. population estimates based on the average of 2006 and 2007 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 32 years (1975-2006)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2002-2006)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2002; Melanoma in 2003.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 32-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2007.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 32-year limited duration prevalence. The 2002 breast cancer is counted for the breast 5-year and 32-year limited duration prevalence.

In method (e) the 2002 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2003 melanoma is counted for 5-year limited duration prevalence for melanoma.

<sup>f</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.21 - continued  
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2007<sup>a</sup>  
 Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			32-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 32 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 32 Years <sup>d</sup>
Prostate	Male	938,613	1,013,875	1,013,911	2,273,570	2,425,802
Testis	Male	39,905	40,672	41,227	179,020	181,360
Urinary Bladder	Both Sexes	189,135	233,199	237,692	518,286	598,692
	Male	141,812	175,570	179,094	384,498	443,555
	Female	47,323	57,629	58,598	133,788	155,137
Kidney & Renal Pelvis	Both Sexes	118,493	143,522	144,764	270,625	314,100
	Male	70,932	86,672	87,600	158,850	185,673
	Female	47,561	56,850	57,164	111,775	128,427
Brain & Nervous System	Both Sexes	40,293	42,551	42,885	112,726	116,105
	Male	22,134	23,505	23,659	60,362	62,268
	Female	18,159	19,046	19,226	52,364	53,837
Thyroid	Both Sexes	129,436	141,554	141,860	392,263	415,126
	Male	28,915	33,014	33,071	86,537	93,189
	Female	100,521	108,540	108,789	305,726	321,937
Hodgkin Lymphoma	Both Sexes	36,160	38,130	38,141	149,892	153,535
	Male	18,823	19,773	19,784	77,318	79,103
	Female	17,337	18,357	18,357	72,574	74,432
Non-Hodgkin Lymphoma	Both Sexes	182,913	212,380	213,902	427,401	474,880
	Male	95,299	111,717	112,437	221,132	245,677
	Female	87,614	100,663	101,465	206,269	229,203
Myeloma	Both Sexes	40,775	47,047	47,082	61,360	69,598
	Male	21,757	25,411	25,446	33,418	38,267
	Female	19,018	21,636	21,636	27,942	31,331
Leukemia	Both Sexes	99,568	112,939	113,079	238,435	259,889
	Male	57,160	65,055	65,125	134,732	147,037
	Female	42,408	47,884	47,954	103,703	112,852
Acute Lymphocytic Leukemia	Both Sexes	15,056	15,397	15,397	57,526	58,016
	Male	8,617	8,718	8,718	31,667	31,805
	Female	6,439	6,679	6,679	25,859	26,211
Childhood (Ages 0-19)	Both Sexes	59,446	59,545	59,919	269,403	269,864
	Male	32,002	32,057	32,201	138,408	138,607
	Female	27,444	27,488	27,718	130,995	131,257
Kaposi Sarcoma	Both Sexes	6,329	6,781	6,781	21,513	22,595
	Male	5,909	6,235	6,235	20,237	21,145
	Female	420	546	546	1,276	1,450
Mesothelioma	Both Sexes	2,943	3,641	3,641	4,427	5,271
	Male	2,066	2,537	2,537	2,655	3,237
	Female	877	1,104	1,104	1,772	2,034

<sup>a</sup> U.S. 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER 9 registries and 1/1/2007 U.S. population estimates based on the average of 2006 and 2007 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 32 years (1975-2006)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2002-2006)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2002; Melanoma in 2003.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 32-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2007.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 32-year limited duration prevalence. The 2002 breast cancer is counted for the breast 5-year and 32-year limited duration prevalence.

In method (e) the 2002 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2003 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.22  
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2007<sup>a</sup>  
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	5,353,054	17,384	39,980	78,051	145,588	336,123	749,038	1,258,920	2,727,970
Females	6,360,682	15,217	33,503	84,509	213,851	604,412	1,155,959	1,395,096	2,858,136
Oral Cavity & Pharynx									
Males	161,112	34	462	1,460	3,306	14,446	38,574	42,305	60,526
Females	88,254	112	478	1,499	3,730	8,328	16,027	18,737	39,343
Esophagus									
Males	21,782	0	0	16	110	1,155	4,158	7,094	9,250
Females	6,947	0	0	12	71	215	906	1,589	4,154
Stomach									
Males	37,297	0	5	29	621	2,421	5,852	9,147	19,223
Females	28,342	5	11	106	575	2,139	3,970	4,882	16,654
Colon & Rectum									
Males	540,636	11	46	1,079	5,377	23,458	70,527	118,887	321,251
Females	571,857	0	109	1,421	4,924	21,129	59,895	98,155	386,224
Liver & Intrahep									
Males	18,644	408	525	363	435	1,451	6,731	4,464	4,268
Females	9,109	411	346	356	427	842	1,852	1,835	3,041
Pancreas									
Males	16,057	0	23	93	227	1,283	3,502	4,577	6,353
Females	16,936	0	53	137	418	1,208	3,029	4,061	8,031
Larynx									
Males	72,246	0	0	24	215	2,470	10,579	19,435	39,524
Females	18,192	0	0	57	77	1,144	3,131	4,429	9,355
Lung & Bronchus									
Males	172,739	34	75	375	1,109	6,021	23,958	50,028	91,141
Females	197,878	11	55	332	1,269	8,266	26,154	53,184	108,608
Melanoma of the Skin									
Males	385,054	94	750	5,271	16,607	46,961	86,015	93,408	135,947
Females	408,229	72	991	11,270	32,943	70,027	96,409	80,585	115,933

<sup>a</sup> U.S. 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2007 U.S. population estimates based on the average of 2006 and 2007 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.22 - continued  
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2007<sup>a</sup>  
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	13,326	0	0	29	93	497	2,014	3,353	7,340
Females	2,591,855	0	81	2,530	35,087	208,053	505,550	645,055	1,195,498
Cervix									
Females	247,180	11	81	2,348	17,537	43,343	58,279	48,897	76,684
Corpus & Uterus, NOS									
Females	575,108	0	67	452	5,116	23,123	78,436	129,723	338,192
Ovary <sup>d</sup>									
Females	177,162	84	949	3,571	6,786	20,117	37,716	40,600	67,339
Prostate									
Males	2,276,112	50	69	80	387	17,453	197,739	585,651	1,474,683
Urinary Bladder									
Males	395,480	11	131	702	2,434	12,104	42,058	87,203	250,837
Females	139,756	33	46	342	1,153	4,415	13,116	27,208	93,443
Kidney & Renal Pelvis									
Males	164,839	1,413	2,428	2,472	4,088	14,777	33,714	43,718	62,228
Females	116,651	1,441	2,633	2,809	3,988	10,792	19,272	25,537	50,178
Hodgkin Lymphoma									
Males	84,583	162	2,193	9,019	16,237	22,676	18,661	9,862	5,771
Females	79,690	107	1,397	10,426	16,314	21,812	16,409	7,293	5,934
Non-Hodgkin Lymphoma									
Males	226,855	689	3,434	7,073	12,028	27,188	44,816	51,557	80,070
Females	211,470	393	1,596	4,130	7,622	18,650	37,239	45,364	96,476
Myeloma									
Males	33,522	0	5	40	488	2,380	6,786	9,878	13,944
Females	28,120	0	0	23	215	1,671	5,234	7,536	13,441
Leukemia									
Males	137,398	6,452	11,601	11,047	9,522	11,671	18,275	24,790	44,041
Females	106,874	5,226	9,722	9,893	8,060	8,514	12,311	15,613	37,535
Acute Lymphocytic Leuk									
Males	33,394	5,340	10,132	8,470	5,396	2,320	905	538	291
Females	27,389	4,366	8,155	7,282	4,309	1,895	706	455	220

<sup>a</sup> U.S. 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2007 U.S. population estimates based on the average of 2006 and 2007 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.21  
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	4,128,739	2,849,626	1,881,402	1,133,219	660,646	414,727	9,393,312	11,196,431	517,305	11,713,736
	Males	2,110,305	1,414,018	900,238	446,691	221,063	123,842	4,652,707	5,248,197	104,857	5,353,054
	Females	2,018,434	1,435,608	981,164	686,528	439,583	290,885	4,740,605	5,948,234	412,448	6,360,682
White <sup>b</sup>	Both Sexes	3,549,667	2,501,487	1,669,462	1,028,390	601,419	379,225	8,203,121	9,846,464	456,479	10,302,943
	Males	1,806,202	1,232,561	795,374	408,847	203,179	114,733	4,042,339	4,590,344	97,851	4,688,195
	Females	1,743,465	1,268,926	874,088	619,543	398,240	264,492	4,160,782	5,256,120	358,628	5,614,748
Black <sup>b</sup>	Both Sexes	393,293	236,404	145,265	69,698	38,611	22,817	809,114	913,188	29,179	942,367
	Males	210,554	129,547	75,909	25,938	11,280	5,418	429,854	460,141	2,858	462,999
	Females	182,739	106,857	69,356	43,760	27,331	17,399	379,260	453,047	26,321	479,368
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	111,468	66,868	39,327	+	+	+	227,997	+	+	+
	Males	48,693	27,237	14,994	+	+	+	94,285	+	+	+
	Females	62,775	39,631	24,333	+	+	+	133,712	+	+	+
Hispanic <sup>d</sup>	Both Sexes	242,034	146,222	88,702	+	+	+	501,611	+	+	+
	Males	116,603	68,482	38,827	+	+	+	233,031	+	+	+
	Females	125,431	77,740	49,875	+	+	+	268,580	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	2.6612%	0.0795%	0.1575%	0.2794%	0.6764%	1.5898%	3.7412%	8.2074%	13.6274%	14.3849%	2.7186%
	Males	2.6537%	0.0843%	0.1675%	0.2592%	0.5230%	1.0880%	3.0641%	9.0062%	18.2916%	21.0395%	3.1045%
	Females	2.6687%	0.0744%	0.1470%	0.3012%	0.8343%	2.0922%	4.3879%	7.4843%	9.9995%	10.6734%	2.4951%
White <sup>c</sup>	Both Sexes	2.9016%	0.0854%	0.1709%	0.3117%	0.7504%	1.6847%	3.8710%	8.5468%	14.3057%	14.8471%	2.8460%
	Males	2.8623%	0.0909%	0.1820%	0.2906%	0.5919%	1.1674%	3.1413%	9.1816%	18.8168%	21.3703%	3.1973%
	Females	2.9410%	0.0796%	0.1593%	0.3351%	0.9205%	2.2202%	4.5910%	7.9563%	10.7117%	11.2426%	2.6522%
Black <sup>c</sup>	Both Sexes	1.9389%	0.0563%	0.1094%	0.1700%	0.4733%	1.2738%	3.6502%	8.5816%	13.5567%	12.4189%	2.5670%
	Males	2.1738%	0.0552%	0.1155%	0.1462%	0.3244%	0.8913%	3.6847%	11.8095%	21.6903%	21.1353%	3.5022%
	Females	1.7257%	0.0576%	0.1031%	0.1929%	0.6031%	1.6098%	3.6215%	6.0800%	8.1048%	8.1833%	1.9851%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1.7429%	0.0639%	0.1214%	0.1783%	0.4091%	1.1723%	2.7145%	5.0524%	8.4354%	10.2622%	1.8025%
	Males	1.5477%	0.0703%	0.1314%	0.1594%	0.2530%	0.6160%	1.6415%	4.7249%	10.7800%	14.9286%	1.8500%
	Females	1.9264%	0.0571%	0.1108%	0.1975%	0.5573%	1.6852%	3.6549%	5.3311%	6.7103%	7.2408%	1.8227%
Hispanic <sup>d</sup>	Both Sexes	1.0760%	0.0709%	0.1418%	0.1980%	0.4572%	1.1046%	2.5135%	5.4879%	9.3151%	9.7153%	1.8600%
	Males	0.9671%	0.0765%	0.1502%	0.1882%	0.3375%	0.6500%	1.7790%	5.8345%	12.9875%	14.6503%	2.1041%
	Females	1.1911%	0.0650%	0.1331%	0.2096%	0.5964%	1.5917%	3.2150%	5.1912%	6.6245%	6.8413%	1.7519%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 3.21  
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	40,293	23,264	18,341	14,120	8,689	6,051	88,389	112,726	13,603	126,329
	Males	22,134	12,039	10,128	7,344	4,690	3,037	47,864	60,362	6,260	66,622
	Females	18,159	11,225	8,213	6,776	3,999	3,014	40,525	52,364	7,343	59,707
White <sup>b</sup>	Both Sexes	35,646	20,578	16,082	12,500	7,789	5,385	77,951	99,755	6,327	106,082
	Males	19,983	10,701	8,914	6,431	4,178	2,711	42,656	53,819	3,494 <sup>i</sup>	57,313
	Females	15,663	9,877	7,168	6,069	3,611	2,674	35,295	45,936	2,833 <sup>i</sup>	48,769
Black <sup>b</sup>	Both Sexes	2,955	1,769	1,440	1,139	630	496	6,758	8,597	+	+
	Males	1,318	936	743	630	369	233	3,358	4,304	+	+
	Females	1,637	833	697	509	261	263	3,400	4,293	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1,061	562	481	+	+	+	2,232	+	+	+
	Males	525	279	254	+	+	+	1,127	+	+	+
	Females	536	283	227	+	+	+	1,105	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,037	2,360	1,451	+	+	+	8,389	+	+	+
	Males	2,245	1,248	720	+	+	+	4,499	+	+	+
	Females	1,792	1,112	731	+	+	+	3,890	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted <sup>f</sup>
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0254%	0.0137%	0.0285%	0.0243%	0.0247%	0.0305%	0.0310%	0.0299%	0.0252%	0.0133%	0.0253%
	Males	0.0277%	0.0141%	0.0310%	0.0251%	0.0270%	0.0331%	0.0353%	0.0332%	0.0281%	0.0139%	0.0275%
	Females	0.0231%	0.0132%	0.0259%	0.0234%	0.0222%	0.0278%	0.0268%	0.0270%	0.0230%	0.0129%	0.0232%
White <sup>c</sup>	Both Sexes	0.0285%	0.0149%	0.0319%	0.0267%	0.0284%	0.0353%	0.0346%	0.0330%	0.0280%	0.0137%	0.0284%
	Males	0.0313%	0.0155%	0.0350%	0.0275%	0.0312%	0.0382%	0.0395%	0.0371%	0.0315%	0.0152%	0.0309%
	Females	0.0257%	0.0141%	0.0285%	0.0259%	0.0255%	0.0323%	0.0299%	0.0293%	0.0251%	0.0128%	0.0258%
Black <sup>c</sup>	Both Sexes	0.0159%	0.0102%	0.0188%	0.0174%	0.0145%	0.0127%	0.0195%	0.0220%	0.0186%	0.0141%	0.0160%
	Males	0.0160%	0.0092%	0.0197%	0.0182%	0.0160%	0.0107%	0.0233%	0.0204%	0.0159%	-	0.0157%
	Females	0.0158%	0.0113%	0.0179%	0.0167%	0.0132%	0.0145%	0.0163%	0.0233%	0.0203%	0.0193%	0.0160%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0145%	0.0095%	0.0176%	0.0150%	0.0128%	0.0166%	0.0173%	0.0151%	0.0128%	0.0075%	0.0145%
	Males	0.0150%	0.0106%	0.0179%	0.0145%	0.0127%	0.0192%	0.0165%	0.0151%	0.0137%	0.0087%	0.0150%
	Females	0.0140%	0.0083%	0.0173%	0.0155%	0.0130%	0.0141%	0.0180%	0.0151%	0.0121%	0.0067%	0.0140%
Hispanic <sup>d</sup>	Both Sexes	0.0178%	0.0103%	0.0200%	0.0135%	0.0164%	0.0264%	0.0251%	0.0259%	0.0220%	0.0102%	0.0191%
	Males	0.0185%	0.0106%	0.0211%	0.0132%	0.0178%	0.0274%	0.0254%	0.0291%	0.0255%	0.0117%	0.0202%
	Females	0.0171%	0.0099%	0.0190%	0.0139%	0.0148%	0.0253%	0.0248%	0.0232%	0.0193%	0.0093%	0.0181%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 4.25  
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	802,387	665,429	448,662	305,856	170,696	97,699	2,053,712	2,522,416	82,765	2,605,181
	Males	5,452	3,400	1,911	1,058	740	395	11,283	13,101	225	13,326
	Females	796,935	662,029	446,751	304,798	169,956	97,304	2,042,429	2,509,315	82,540	2,591,855
White <sup>b</sup>	Both Sexes	692,879	589,290	401,110	278,139	155,068	89,158	1,807,186	2,234,644	73,464	2,308,108
	Males	4,658	2,914	1,706	946	675	370	9,735	11,414	+ <sup>i</sup>	11,410
	Females	688,221	586,376	399,404	277,193	154,393	88,788	1,797,451	2,223,230	73,468	2,296,698
Black <sup>b</sup>	Both Sexes	75,195	50,129	32,942	19,351	11,205	6,099	167,447	196,730	+	+
	Males	630	353	133	108	50	25	1,176	1,299	+	+
	Females	74,565	49,776	32,809	19,243	11,155	6,074	166,271	195,431	5,845	201,276
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	26,068	18,875	10,771	+	+	+	58,789	+	+	+
	Males	126	45	52	+	+	+	227	+	+	+
	Females	25,942	18,830	10,719	+	+	+	58,562	+	+	+
Hispanic <sup>d</sup>	Both Sexes	46,542	31,567	19,047	+	+	+	102,828	+	+	+
	Males	242	102	33	+	+	+	392	+	+	+
	Females	46,300	31,465	19,014	+	+	+	102,436	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.5838%	-	0.0001%	0.0060%	0.0835%	0.4303%	1.1069%	1.9143%	2.4772%	2.8655%	0.5825%
	Males	0.0065%	-	-	-	0.0004%	0.0023%	0.0089%	0.0238%	0.0425%	0.0554%	0.0075%
	Females	1.1513%	-	0.0003%	0.0122%	0.1690%	0.8587%	2.1556%	3.6255%	4.3709%	4.4327%	1.0660%
White <sup>c</sup>	Both Sexes	0.6383%	-	0.0001%	0.0054%	0.0828%	0.4342%	1.1393%	2.0226%	2.6593%	3.0648%	0.6116%
	Males	0.0069%	-	-	-	0.0003%	0.0021%	0.0084%	0.0239%	0.0447%	0.0609%	0.0077%
	Females	1.2709%	-	0.0003%	0.0113%	0.1714%	0.8814%	2.2552%	3.8818%	4.7423%	4.7247%	1.1306%
Black <sup>c</sup>	Both Sexes	0.4046%	-	-	0.0084%	0.0968%	0.4224%	1.0009%	1.6173%	2.0665%	2.3180%	0.5089%
	Males	0.0060%	-	-	-	-	0.0027%	0.0154%	0.0387%	0.0430%	0.0364%	0.0088%
	Females	0.7663%	-	-	0.0162%	0.1809%	0.7912%	1.8199%	2.8407%	3.4228%	3.4268%	0.8759%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.4362%	-	-	0.0067%	0.0766%	0.4117%	0.9833%	1.4185%	1.6038%	1.6166%	0.4363%
	Males	0.0037%	-	-	-	-	0.0015%	0.0065%	0.0119%	0.0275%	0.0226%	0.0043%
	Females	0.8427%	-	-	0.0133%	0.1489%	0.7899%	1.8395%	2.6156%	2.7636%	2.6488%	0.7904%
Hispanic <sup>d</sup>	Both Sexes	0.2226%	-	-	0.0046%	0.0568%	0.2942%	0.7558%	1.2669%	1.5878%	1.6452%	0.3811%
	Males	0.0016%	-	-	-	-	0.0015%	0.0048%	0.0116%	0.0193%	0.0164%	0.0034%
	Females	0.4562%	-	-	0.0099%	0.1224%	0.6079%	1.4730%	2.3412%	2.7371%	2.5938%	0.6979%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 5.15  
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	38,618	37,290	35,406	31,088	23,985	20,517	124,596	195,066	52,114	247,180
White <sup>b</sup>	30,478	29,632	28,974	25,843	19,875	16,823	99,950	158,316	43,651	201,967
Black <sup>b</sup>	5,577	5,091	3,981	3,525	3,010	2,735	16,273	25,096	+	+
Asian/Pacific Islander <sup>c</sup>	1,979	1,737	1,621	+	+	+	5,824	+	+	+
Hispanic <sup>d</sup>	7,913	6,838	5,801	+	+	+	22,701	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0808%	-	0.0003%	0.0102%	0.0830%	0.1570%	0.1673%	0.1447%	0.1312%	0.0788%	0.0777%
White <sup>c</sup>	0.0821%	-	0.0004%	0.0108%	0.0944%	0.1676%	0.1643%	0.1340%	0.1171%	0.0625%	0.0787%
Black <sup>c</sup>	0.0712%	-	-	0.0103%	0.0656%	0.1295%	0.1616%	0.1685%	0.1593%	0.1522%	0.0759%
Asian/Pacific Islander <sup>c</sup>	0.0784%	-	-	0.0033%	0.0349%	0.1120%	0.1799%	0.1870%	0.1943%	0.1552%	0.0732%
Hispanic <sup>d</sup>	0.0973%	-	-	0.0108%	0.0863%	0.2379%	0.2981%	0.3072%	0.2807%	0.1758%	0.1296%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 6.28  
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	404,997	270,735	166,731	115,238	70,411	38,673	895,585	1,076,622	35,871	1,112,493
	Males	202,765	135,385	81,371	56,362	32,422	15,808	445,736	527,903	12,733	540,636
	Females	202,232	135,350	85,360	58,876	37,989	22,865	449,849	548,719	23,138	571,857
White <sup>b</sup>	Both Sexes	341,352	235,165	145,812	101,665	62,835	34,516	769,126	930,221	31,871	962,092
	Males	173,461	118,712	71,548	50,351	29,025	14,163	386,960	460,739	11,452	472,191
	Females	167,891	116,453	74,264	51,314	33,810	20,353	382,166	469,482	20,419	489,901
Black <sup>b</sup>	Both Sexes	42,942	23,884	14,018	9,625	5,473	2,713	85,238	99,301	2,547	101,848
	Males	18,611	10,360	6,108	4,001	2,227	1,002	37,089	42,489	813	43,302
	Females	24,331	13,524	7,910	5,624	3,246	1,711	48,149	56,812	1,734	58,546
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	14,736	8,438	5,000	+	+	+	29,538	+	+	+
	Males	7,684	4,367	2,608	+	+	+	15,315	+	+	+
	Females	7,052	4,071	2,392	+	+	+	14,223	+	+	+
Hispanic <sup>d</sup>	Both Sexes	23,522	12,310	6,624	+	+	+	44,705	+	+	+
	Males	12,416	6,562	3,415	+	+	+	23,519	+	+	+
	Females	11,106	5,748	3,209	+	+	+	21,186	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.2552%	-	0.0003%	0.0050%	0.0233%	0.0916%	0.3014%	0.7335%	1.4720%	2.1229%	0.2630%
	Males	0.2587%	-	0.0002%	0.0045%	0.0241%	0.0961%	0.3361%	0.8653%	1.7801%	2.3888%	0.3052%
	Females	0.2518%	-	0.0004%	0.0055%	0.0225%	0.0871%	0.2682%	0.6142%	1.2324%	1.9745%	0.2300%
White <sup>c</sup>	Both Sexes	0.2670%	-	0.0003%	0.0053%	0.0233%	0.0884%	0.2874%	0.7177%	1.4779%	2.1629%	0.2614%
	Males	0.2702%	-	0.0003%	0.0047%	0.0245%	0.0915%	0.3257%	0.8555%	1.7819%	2.4334%	0.3043%
	Females	0.2638%	-	0.0004%	0.0059%	0.0220%	0.0852%	0.2496%	0.5896%	1.2357%	2.0134%	0.2271%
Black <sup>c</sup>	Both Sexes	0.2031%	-	-	0.0035%	0.0249%	0.1011%	0.3729%	0.8885%	1.5260%	1.8955%	0.2803%
	Males	0.1900%	-	-	0.0035%	0.0226%	0.1070%	0.3893%	0.9589%	1.7513%	1.9471%	0.3033%
	Females	0.2149%	-	-	0.0034%	0.0269%	0.0959%	0.3592%	0.8340%	1.3750%	1.8704%	0.2644%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.2348%	-	-	0.0037%	0.0215%	0.0963%	0.3113%	0.6905%	1.3649%	1.9298%	0.2485%
	Males	0.2502%	-	-	0.0042%	0.0228%	0.1088%	0.3371%	0.8236%	1.7300%	2.2758%	0.2968%
	Females	0.2204%	-	-	0.0032%	0.0203%	0.0847%	0.2887%	0.5773%	1.0963%	1.7057%	0.2120%
Hispanic <sup>d</sup>	Both Sexes	0.0954%	-	-	0.0037%	0.0176%	0.0659%	0.2194%	0.5625%	1.0195%	1.3458%	0.1851%
	Males	0.0968%	-	-	0.0037%	0.0173%	0.0655%	0.2435%	0.6708%	1.3310%	1.5392%	0.2209%
	Females	0.0940%	-	-	0.0037%	0.0179%	0.0664%	0.1963%	0.4699%	0.7912%	1.2331%	0.1582%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 7.15  
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	148,471	117,687	88,626	64,993	48,948	40,742	383,228	527,264	47,844	575,108
White <sup>b</sup>	130,470	106,974	81,701	60,668	46,361	38,650	345,716	481,851	41,762	523,613
Black <sup>b</sup>	10,536	5,533	3,731	2,507	1,418	955	20,861	25,054	1,121	26,175
Asian/Pacific Islander <sup>c</sup>	4,830	3,163	2,012	+	+	+	10,533	+	+	+
Hispanic <sup>d</sup>	8,999	5,565	3,626	+	+	+	19,390	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	0.2096%	-	0.0002%	0.0028%	0.0261%	0.1006%	0.3409%	0.7675%	0.9240%	0.8180%	0.1952%
White <sup>c</sup>	0.2382%	-	-	0.0025%	0.0257%	0.1008%	0.3612%	0.8429%	1.0362%	0.9063%	0.2126%
Black <sup>c</sup>	0.0948%	-	-	0.0026%	0.0223%	0.0604%	0.1645%	0.4575%	0.5918%	0.4106%	0.1135%
Asian/Pacific Islander <sup>c</sup>	0.1505%	-	-	0.0048%	0.0282%	0.1281%	0.3597%	0.5055%	0.4628%	0.3601%	0.1401%
Hispanic <sup>d</sup>	0.0856%	-	-	0.0037%	0.0344%	0.0968%	0.2576%	0.5098%	0.5101%	0.4003%	0.1303%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 8.21  
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	18,697	5,830	2,497	956	421	108	27,620	28,549	180	28,729
	Males	14,341	4,441	1,879	653	284	84	21,077	21,695	87	21,782
	Females	4,356	1,389	618	303	137	24	6,543	6,854	93	6,947
White <sup>b</sup>	Both Sexes	16,415	5,127	2,231	780	327	96	24,244	25,000	146	25,146
	Males	12,832	4,001	1,681	539	231	84	18,852	19,380	81	19,461
	Females	3,583	1,126	550	241	96	12	5,392	5,620	65	5,685
Black <sup>b</sup>	Both Sexes	1,770	519	209	138	89	13	2,600	2,753	48	2,801
	Males	1,115	304	152	89	52	0	1,635	1,712	14	1,726
	Females	655	215	57	49	37	13	965	1,041	34	1,075
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	351	109	47	+	+	+	520	+	+	+
	Males	268	77	36	+	+	+	390	+	+	+
	Females	83	32	11	+	+	+	130	+	+	+
Hispanic <sup>d</sup>	Both Sexes	608	168	102	+	+	+	920	+	+	+
	Males	487	142	76	+	+	+	747	+	+	+
	Females	121	26	26	+	+	+	173	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0075%	-	-	-	0.0004%	0.0026%	0.0108%	0.0299%	0.0447%	0.0353%	0.0077%
	Males	0.0114%	-	-	-	0.0005%	0.0043%	0.0182%	0.0508%	0.0739%	0.0622%	0.0129%
	Females	0.0036%	-	-	-	0.0002%	0.0008%	0.0038%	0.0109%	0.0220%	0.0203%	0.0034%
White <sup>c</sup>	Both Sexes	0.0083%	-	-	-	0.0005%	0.0029%	0.0114%	0.0320%	0.0480%	0.0359%	0.0082%
	Males	0.0129%	-	-	-	0.0007%	0.0049%	0.0195%	0.0555%	0.0794%	0.0636%	0.0139%
	Females	0.0037%	-	-	-	0.0002%	0.0009%	0.0035%	0.0101%	0.0230%	0.0207%	0.0034%
Black <sup>c</sup>	Both Sexes	0.0057%	-	-	-	-	0.0018%	0.0131%	0.0344%	0.0406%	0.0240%	0.0075%
	Males	0.0073%	-	-	-	-	0.0024%	0.0183%	0.0494%	0.0628%	0.0331%	0.0109%
	Females	0.0042%	-	-	-	-	0.0013%	0.0088%	0.0227%	0.0256%	0.0195%	0.0050%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0042%	-	-	-	-	0.0014%	0.0048%	0.0135%	0.0265%	0.0355%	0.0045%
	Males	0.0066%	-	-	-	-	0.0027%	0.0090%	0.0208%	0.0460%	0.0658%	0.0078%
	Females	0.0021%	-	-	-	-	-	0.0073%	0.0121%	0.0159%	0.0159%	0.0020%
Hispanic <sup>d</sup>	Both Sexes	0.0020%	-	-	-	0.0005%	0.0013%	0.0038%	0.0139%	0.0245%	0.0180%	0.0038%
	Males	0.0030%	-	-	-	0.0007%	0.0021%	0.0062%	0.0252%	0.0462%	0.0332%	0.0068%
	Females	0.0008%	-	-	-	-	-	0.0015%	0.0042%	0.0087%	0.0091%	0.0014%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 9.21  
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	36,160	30,045	26,050	23,411	18,079	12,529	102,168	149,892	14,381	164,273
	Males	18,823	15,470	13,400	11,828	9,506	6,503	52,938	77,318	7,265	84,583
	Females	17,337	14,575	12,650	11,583	8,573	6,026	49,230	72,574	7,116	79,690
White <sup>b</sup>	Both Sexes	30,654	26,172	23,213	21,619	16,737	11,827	89,094	133,704	16,051	149,755
	Males	16,066	13,446	12,011	11,018	8,640	6,053	46,375	68,904	8,284	77,188
	Females	14,588	12,726	11,202	10,601	8,097	5,774	42,719	64,800	7,767	72,567
Black <sup>b</sup>	Both Sexes	4,130	3,024	2,342	1,419	1,076	586	10,212	12,638	1,034	13,672
	Males	1,991	1,558	1,156	623	727	371	5,024	6,497	710	7,207
	Females	2,139	1,466	1,186	796	349	215	5,188	6,141	324	6,465
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	799	503	285	+	+	+	1,673	+	+	+
	Males	404	278	116	+	+	+	842	+	+	+
	Females	395	225	169	+	+	+	831	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,193	2,504	1,620	+	+	+	7,908	+	+	+
	Males	1,801	1,426	915	+	+	+	4,424	+	+	+
	Females	1,392	1,078	705	+	+	+	3,484	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0282%	0.0006%	0.0093%	0.0389%	0.0518%	0.0421%	0.0295%	0.0253%	0.0253%	0.0165%	0.0281%
	Males	0.0297%	0.0008%	0.0112%	0.0349%	0.0504%	0.0458%	0.0349%	0.0311%	0.0287%	0.0199%	0.0295%
	Females	0.0267%	0.0005%	0.0073%	0.0433%	0.0532%	0.0384%	0.0243%	0.0201%	0.0227%	0.0146%	0.0268%
White <sup>c</sup>	Both Sexes	0.0315%	0.0007%	0.0101%	0.0439%	0.0594%	0.0471%	0.0325%	0.0282%	0.0282%	0.0188%	0.0316%
	Males	0.0331%	0.0008%	0.0120%	0.0389%	0.0577%	0.0504%	0.0383%	0.0336%	0.0324%	0.0227%	0.0329%
	Females	0.0299%	0.0006%	0.0081%	0.0493%	0.0611%	0.0436%	0.0267%	0.0231%	0.0248%	0.0166%	0.0305%
Black <sup>c</sup>	Both Sexes	0.0247%	-	0.0082%	0.0295%	0.0463%	0.0414%	0.0298%	0.0228%	0.0226%	0.0108%	0.0252%
	Males	0.0256%	-	0.0092%	0.0250%	0.0426%	0.0488%	0.0357%	0.0318%	0.0233%	0.0165%	0.0269%
	Females	0.0238%	-	0.0072%	0.0339%	0.0495%	0.0349%	0.0249%	0.0158%	0.0221%	0.0080%	0.0239%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0107%	-	0.0056%	0.0195%	0.0178%	0.0122%	0.0097%	0.0076%	0.0090%	-	0.0103%
	Males	0.0111%	-	0.0081%	0.0193%	0.0161%	0.0125%	0.0099%	0.0119%	0.0088%	-	0.0107%
	Females	0.0103%	-	0.0030%	0.0196%	0.0193%	0.0120%	0.0096%	0.0040%	0.0091%	-	0.0098%
Hispanic <sup>d</sup>	Both Sexes	0.0165%	0.0008%	0.0090%	0.0228%	0.0256%	0.0241%	0.0198%	0.0254%	0.0255%	0.0169%	0.0181%
	Males	0.0181%	0.0010%	0.0111%	0.0221%	0.0254%	0.0276%	0.0255%	0.0265%	0.0377%	0.0257%	0.0207%
	Females	0.0148%	-	0.0067%	0.0235%	0.0258%	0.0203%	0.0144%	0.0245%	0.0166%	0.0117%	0.0159%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 10.15  
Kaposi SarcomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	6,329	5,120	7,221	1,941	616	206	19,682	21,513	+	+
	Males	5,909	4,771	6,982	1,798	582	136	18,560	20,237	+	+
	Females	420	349	239	143	34	70	1,122	1,276	+	+
White <sup>b</sup>	Both Sexes	4,583	3,682	5,965	1,604	508	170	15,072	16,594	+	+
	Males	4,259	3,433	5,797	1,494	485	100	14,250	15,628	+	+
	Females	324	249	168	110	23	70	822	966	+	+
Black <sup>b</sup>	Both Sexes	1,483	1,113	863	155	63	12	3,537	3,689	+	+
	Males	1,404	1,047	803	145	63	12	3,322	3,474	+	+
	Females	79	66	60	10	0	0	215	215	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	100	68	76	+	+	+	247	+	+	+
	Males	95	68	76	+	+	+	242	+	+	+
	Females	5	0	0	+	+	+	5	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,142	945	1,123	+	+	+	3,318	+	+	+
	Males	1,093	886	1,103	+	+	+	3,170	+	+	+
	Females	49	59	20	+	+	+	148	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0073%	-	-	0.0008%	0.0075%	0.0193%	0.0140%	0.0069%	0.0077%	0.0130%	0.0072%
	Males	0.0141%	-	-	0.0014%	0.0147%	0.0379%	0.0281%	0.0137%	0.0152%	0.0256%	0.0141%
	Females	0.0006%	-	-	-	-	0.0006%	0.0005%	0.0008%	0.0018%	0.0059%	0.0006%
White <sup>c</sup>	Both Sexes	0.0074%	-	-	0.0006%	0.0068%	0.0195%	0.0142%	0.0072%	0.0084%	0.0130%	0.0071%
	Males	0.0141%	-	-	0.0010%	0.0130%	0.0378%	0.0282%	0.0141%	0.0161%	0.0260%	0.0139%
	Females	0.0006%	-	-	-	-	0.0006%	0.0003%	0.0008%	0.0022%	0.0059%	0.0005%
Black <sup>c</sup>	Both Sexes	0.0097%	-	-	0.0026%	0.0161%	0.0275%	0.0175%	0.0070%	0.0045%	0.0105%	0.0101%
	Males	0.0195%	-	-	0.0047%	0.0347%	0.0576%	0.0364%	0.0153%	0.0096%	-	0.0208%
	Females	0.0008%	-	-	-	-	-	0.0017%	-	-	0.0107%	0.0009%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0018%	-	-	-	0.0014%	0.0036%	0.0043%	0.0019%	0.0033%	0.0055%	0.0018%
	Males	0.0037%	-	-	-	0.0029%	0.0072%	0.0092%	0.0042%	0.0078%	0.0122%	0.0038%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic <sup>d</sup>	Both Sexes	0.0073%	-	-	0.0011%	0.0095%	0.0236%	0.0152%	0.0103%	0.0140%	0.0318%	0.0096%
	Males	0.0136%	-	-	0.0019%	0.0175%	0.0450%	0.0305%	0.0206%	0.0287%	0.0632%	0.0187%
	Females	0.0006%	-	-	-	-	-	-	-	-	0.0136%	0.0010%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 11.21  
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	118,493	64,286	40,411	24,112	13,710	7,600	234,402	270,625	10,865	281,490
	Males	70,932	37,501	23,644	13,686	8,044	3,972	138,797	158,850	5,989	164,839
	Females	47,561	26,785	16,767	10,426	5,666	3,628	95,605	111,775	4,876	116,651
White <sup>b</sup>	Both Sexes	101,449	55,278	34,902	21,540	12,276	6,674	201,547	234,006	12,430	246,436
	Males	60,840	32,703	20,849	12,378	7,347	3,589	120,399	138,699	5,639	144,338
	Females	40,609	22,575	14,053	9,162	4,929	3,085	81,148	95,307	6,791	102,098
Black <sup>b</sup>	Both Sexes	11,896	6,735	4,232	2,038	1,170	772	23,848	26,947	1,928	28,875
	Males	7,010	3,421	2,082	1,028	526	314	13,064	14,444	653	15,097
	Females	4,886	3,314	2,150	1,010	644	458	10,784	12,503	1,275	13,778
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,648	1,260	648	+	+	+	4,697	+	+	+
	Males	1,640	762	408	+	+	+	2,881	+	+	+
	Females	1,008	498	240	+	+	+	1,816	+	+	+
Hispanic <sup>d</sup>	Both Sexes	9,278	4,670	2,396	+	+	+	17,119	+	+	+
	Males	5,458	2,684	1,299	+	+	+	9,831	+	+	+
	Females	3,820	1,986	1,097	+	+	+	7,288	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0662%	0.0063%	0.0087%	0.0027%	0.0107%	0.0421%	0.1127%	0.2256%	0.3102%	0.2689%	0.0669%
	Males	0.0802%	0.0061%	0.0076%	0.0027%	0.0112%	0.0511%	0.1498%	0.3067%	0.4132%	0.3819%	0.0880%
	Females	0.0524%	0.0066%	0.0099%	0.0028%	0.0103%	0.0330%	0.0772%	0.1522%	0.2301%	0.2059%	0.0495%
White <sup>c</sup>	Both Sexes	0.0723%	0.0067%	0.0092%	0.0027%	0.0116%	0.0441%	0.1166%	0.2391%	0.3272%	0.2832%	0.0704%
	Males	0.0875%	0.0065%	0.0083%	0.0026%	0.0119%	0.0528%	0.1535%	0.3221%	0.4369%	0.4068%	0.0924%
	Females	0.0572%	0.0068%	0.0102%	0.0028%	0.0112%	0.0350%	0.0802%	0.1618%	0.2397%	0.2149%	0.0519%
Black <sup>c</sup>	Both Sexes	0.0553%	0.0073%	0.0104%	0.0038%	0.0112%	0.0449%	0.1309%	0.2245%	0.3063%	0.2456%	0.0689%
	Males	0.0653%	0.0065%	0.0074%	0.0038%	0.0114%	0.0544%	0.1825%	0.3187%	0.3875%	0.3199%	0.0898%
	Females	0.0462%	0.0081%	0.0134%	0.0037%	0.0111%	0.0366%	0.0880%	0.1516%	0.2518%	0.2095%	0.0536%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0363%	0.0028%	0.0030%	0.0007%	0.0047%	0.0235%	0.0635%	0.1206%	0.1844%	0.1624%	0.0373%
	Males	0.0459%	0.0024%	0.0028%	-	0.0051%	0.0321%	0.0881%	0.1697%	0.2508%	0.2313%	0.0512%
	Females	0.0272%	0.0032%	0.0033%	-	0.0044%	0.0155%	0.0419%	0.0788%	0.1356%	0.1177%	0.0263%
Hispanic <sup>d</sup>	Both Sexes	0.0369%	0.0050%	0.0066%	0.0020%	0.0095%	0.0347%	0.1010%	0.2138%	0.3092%	0.2853%	0.0636%
	Males	0.0410%	0.0051%	0.0049%	0.0019%	0.0087%	0.0395%	0.1258%	0.2869%	0.4241%	0.3553%	0.0813%
	Females	0.0324%	0.0049%	0.0084%	0.0020%	0.0104%	0.0296%	0.0773%	0.1512%	0.2251%	0.2445%	0.0496%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 12.21  
Cancer of the Larynx (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	30,163	21,068	14,575	10,259	6,493	3,967	70,734	87,367	3,071	90,438
	Males	23,925	16,934	11,607	8,122	5,238	3,203	56,324	69,746	2,500	72,246
	Females	6,238	4,134	2,968	2,137	1,255	764	14,410	17,621	571	18,192
White <sup>b</sup>	Both Sexes	25,165	18,066	12,577	8,770	5,795	3,578	60,051	74,683	2,464	77,147
	Males	19,911	14,568	10,075	7,021	4,747	2,887	47,848	59,826	1,998	61,824
	Females	5,254	3,498	2,502	1,749	1,048	691	12,203	14,857	466	15,323
Black <sup>b</sup>	Both Sexes	4,238	2,454	1,679	1,238	597	283	8,940	10,596	211	10,807
	Males	3,353	1,884	1,290	932	418	226	6,988	8,200	180	8,380
	Females	885	570	389	306	179	57	1,952	2,396	31	2,427
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	494	329	204	+	+	+	1,095	+	+	+
	Males	439	285	178	+	+	+	959	+	+	+
	Females	55	44	26	+	+	+	136	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,488	961	611	+	+	+	3,291	+	+	+
	Males	1,276	794	512	+	+	+	2,776	+	+	+
	Females	212	167	99	+	+	+	515	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0188%	-	-	0.0002%	0.0008%	0.0062%	0.0276%	0.0755%	0.1143%	0.0863%	0.0193%
	Males	0.0306%	-	-	0.0002%	0.0010%	0.0088%	0.0447%	0.1326%	0.2104%	0.1974%	0.0352%
	Females	0.0072%	-	-	0.0003%	0.0005%	0.0036%	0.0112%	0.0238%	0.0396%	0.0243%	0.0068%
White <sup>c</sup>	Both Sexes	0.0203%	-	-	0.0003%	0.0008%	0.0067%	0.0283%	0.0770%	0.1197%	0.0880%	0.0200%
	Males	0.0327%	-	-	-	0.0009%	0.0092%	0.0451%	0.1337%	0.2158%	0.2010%	0.0358%
	Females	0.0080%	-	-	0.0004%	0.0007%	0.0040%	0.0117%	0.0244%	0.0432%	0.0256%	0.0073%
Black <sup>c</sup>	Both Sexes	0.0204%	-	-	-	0.0011%	0.0075%	0.0442%	0.1205%	0.1469%	0.1000%	0.0271%
	Males	0.0339%	-	-	-	0.0018%	0.0105%	0.0752%	0.2255%	0.2928%	0.2462%	0.0521%
	Females	0.0081%	-	-	-	-	0.0049%	0.0184%	0.0391%	0.0492%	0.0289%	0.0097%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0086%	-	-	-	-	0.0026%	0.0091%	0.0296%	0.0586%	0.0632%	0.0092%
	Males	0.0156%	-	-	-	-	0.0049%	0.0167%	0.0561%	0.1224%	0.1420%	0.0187%
	Females	0.0020%	-	-	-	-	-	0.0024%	0.0071%	0.0118%	0.0122%	0.0020%
Hispanic <sup>d</sup>	Both Sexes	0.0068%	-	-	-	0.0008%	0.0034%	0.0138%	0.0523%	0.0841%	0.0669%	0.0134%
	Males	0.0111%	-	-	-	0.0007%	0.0040%	0.0231%	0.1045%	0.1694%	0.1513%	0.0263%
	Females	0.0022%	-	-	-	0.0009%	0.0028%	0.0049%	0.0075%	0.0216%	0.0177%	0.0036%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.29  
LeukemiaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	99,568	56,704	36,263	22,054	13,763	8,044	202,425	238,435	5,837	244,272
	Males	57,160	32,439	20,489	12,144	7,615	3,875	115,334	134,732	2,666	137,398
	Females	42,408	24,265	15,774	9,910	6,148	4,169	87,091	103,703	3,171	106,874
White <sup>b</sup>	Both Sexes	87,445	50,650	32,627	19,931	12,643	7,410	179,727	212,560	4,933	217,493
	Males	50,580	29,024	18,501	11,179	7,018	3,603	102,972	120,851	1,919	122,770
	Females	36,865	21,626	14,126	8,752	5,625	3,807	76,755	91,709	3,014	94,723
Black <sup>b</sup>	Both Sexes	7,126	3,438	1,954	1,176	566	362	13,027	14,752	672	15,424
	Males	3,824	1,724	997	515	324	126	6,763	7,564	313	7,877
	Females	3,302	1,714	957	661	242	236	6,264	7,188	359	7,547
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,359	1,280	950	+	+	+	4,870	+	+	+
	Males	1,262	755	510	+	+	+	2,685	+	+	+
	Females	1,097	525	440	+	+	+	2,185	+	+	+
Hispanic <sup>d</sup>	Both Sexes	8,614	4,762	3,457	+	+	+	17,809	+	+	+
	Males	4,998	2,595	2,023	+	+	+	10,181	+	+	+
	Females	3,616	2,167	1,434	+	+	+	7,628	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0587%	0.0298%	0.0504%	0.0226%	0.0215%	0.0323%	0.0630%	0.1311%	0.2141%	0.2369%	0.0600%
	Males	0.0680%	0.0323%	0.0546%	0.0245%	0.0245%	0.0383%	0.0783%	0.1716%	0.2861%	0.3203%	0.0743%
	Females	0.0496%	0.0272%	0.0459%	0.0204%	0.0185%	0.0262%	0.0483%	0.0945%	0.1581%	0.1903%	0.0481%
White <sup>c</sup>	Both Sexes	0.0659%	0.0326%	0.0563%	0.0252%	0.0229%	0.0338%	0.0674%	0.1447%	0.2376%	0.2604%	0.0657%
	Males	0.0762%	0.0356%	0.0606%	0.0275%	0.0258%	0.0403%	0.0830%	0.1888%	0.3160%	0.3545%	0.0813%
	Females	0.0557%	0.0294%	0.0518%	0.0226%	0.0199%	0.0270%	0.0519%	0.1037%	0.1751%	0.2085%	0.0525%
Black <sup>c</sup>	Both Sexes	0.0314%	0.0141%	0.0230%	0.0149%	0.0155%	0.0247%	0.0443%	0.0773%	0.1480%	0.1263%	0.0369%
	Males	0.0354%	0.0147%	0.0248%	0.0144%	0.0168%	0.0303%	0.0569%	0.1012%	0.1827%	0.1748%	0.0451%
	Females	0.0279%	0.0135%	0.0211%	0.0154%	0.0143%	0.0198%	0.0339%	0.0588%	0.1247%	0.1027%	0.0310%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0329%	0.0291%	0.0444%	0.0155%	0.0186%	0.0260%	0.0365%	0.0533%	0.0701%	0.0705%	0.0340%
	Males	0.0373%	0.0311%	0.0497%	0.0172%	0.0230%	0.0275%	0.0427%	0.0587%	0.0940%	0.0956%	0.0397%
	Females	0.0287%	0.0270%	0.0388%	0.0138%	0.0144%	0.0245%	0.0310%	0.0487%	0.0526%	0.0542%	0.0292%
Hispanic <sup>d</sup>	Both Sexes	0.0380%	0.0297%	0.0584%	0.0249%	0.0218%	0.0263%	0.0459%	0.0763%	0.0967%	0.1077%	0.0437%
	Males	0.0421%	0.0323%	0.0639%	0.0288%	0.0248%	0.0309%	0.0540%	0.0864%	0.1188%	0.1366%	0.0506%
	Females	0.0338%	0.0271%	0.0527%	0.0203%	0.0184%	0.0214%	0.0382%	0.0677%	0.0805%	0.0908%	0.0375%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 13.30  
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	15,056	11,555	9,860	8,516	6,647	4,876	40,117	57,526	3,257	60,783
	Males	8,617	6,300	5,530	4,660	3,644	2,408	22,431	31,667	1,727	33,394
	Females	6,439	5,255	4,330	3,856	3,003	2,468	17,686	25,859	1,530	27,389
White <sup>b</sup>	Both Sexes	12,844	10,163	8,537	7,495	6,030	4,505	34,815	50,529	+	+
	Males	7,390	5,544	4,856	4,178	3,325	2,262	19,610	28,027	+	+
	Females	5,454	4,619	3,681	3,317	2,705	2,243	15,205	22,502	+	+
Black <sup>b</sup>	Both Sexes	1,088	745	655	549	297	178	2,692	3,539	+	+
	Males	639	368	306	241	173	57	1,405	1,814	+	+
	Females	449	377	349	308	124	121	1,287	1,725	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	770	442	433	+	+	+	1,778	+	+	+
	Males	416	269	238	+	+	+	989	+	+	+
	Females	354	173	195	+	+	+	789	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,908	2,479	2,096	+	+	+	9,069	+	+	+
	Males	2,265	1,321	1,266	+	+	+	5,170	+	+	+
	Females	1,643	1,158	830	+	+	+	3,899	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0133%	0.0252%	0.0434%	0.0133%	0.0051%	0.0032%	0.0028%	0.0027%	0.0017%	0.0012%	0.0136%
	Males	0.0152%	0.0271%	0.0476%	0.0152%	0.0064%	0.0035%	0.0032%	0.0030%	0.0017%	0.0018%	0.0151%
	Females	0.0115%	0.0232%	0.0391%	0.0112%	0.0037%	0.0029%	0.0025%	0.0025%	0.0017%	0.0009%	0.0121%
White <sup>c</sup>	Both Sexes	0.0147%	0.0279%	0.0493%	0.0155%	0.0056%	0.0034%	0.0032%	0.0029%	0.0016%	0.0013%	0.0153%
	Males	0.0167%	0.0303%	0.0536%	0.0178%	0.0072%	0.0036%	0.0036%	0.0034%	0.0017%	0.0020%	0.0170%
	Females	0.0127%	0.0254%	0.0447%	0.0130%	0.0038%	0.0032%	0.0028%	0.0024%	0.0016%	0.0009%	0.0136%
Black <sup>c</sup>	Both Sexes	0.0063%	0.0106%	0.0170%	0.0068%	0.0022%	0.0015%	-	-	-	-	0.0057%
	Males	0.0070%	0.0116%	0.0183%	0.0067%	0.0015%	0.0021%	-	-	-	-	0.0060%
	Females	0.0058%	0.0095%	0.0157%	0.0068%	0.0027%	-	-	-	-	-	0.0054%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0109%	0.0240%	0.0375%	0.0071%	0.0048%	0.0038%	0.0019%	0.0030%	0.0029%	-	0.0119%
	Males	0.0126%	0.0241%	0.0428%	0.0092%	0.0060%	0.0041%	0.0027%	-	-	-	0.0131%
	Females	0.0092%	0.0239%	0.0319%	0.0050%	0.0037%	0.0035%	-	0.0040%	-	-	0.0106%
Hispanic <sup>d</sup>	Both Sexes	0.0192%	0.0256%	0.0520%	0.0157%	0.0063%	0.0048%	0.0057%	0.0038%	0.0024%	-	0.0162%
	Males	0.0213%	0.0271%	0.0577%	0.0187%	0.0072%	0.0057%	0.0055%	0.0035%	-	-	0.0178%
	Females	0.0170%	0.0240%	0.0461%	0.0122%	0.0054%	0.0037%	0.0058%	0.0040%	-	-	0.0144%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 14.21  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	20,688	3,774	1,321	792	558	198	26,148	27,353	400	27,753
	Males	14,728	2,391	765	350	244	65	18,083	18,577	67	18,644
	Females	5,960	1,383	556	442	314	133	8,065	8,776	333	9,109
White <sup>b</sup>	Both Sexes	14,933	2,777	993	626	446	131	18,978	19,928	429	20,357
	Males	10,862	1,720	562	251	189	35	13,292	13,653	174 <sup>i</sup>	13,827
	Females	4,071	1,057	431	375	257	96	5,686	6,275	255 <sup>i</sup>	6,530
Black <sup>b</sup>	Both Sexes	2,856	286	90	94	61	45	3,279	3,432	+	+
	Males	1,869	172	42	48	26	22	2,105	2,178	+	+
	Females	987	114	48	46	35	23	1,174	1,254	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,525	610	155	+	+	+	3,331	+	+	+
	Males	1,746	423	116	+	+	+	2,313	+	+	+
	Females	779	187	39	+	+	+	1,018	+	+	+
Hispanic <sup>d</sup>	Both Sexes	2,846	453	230	+	+	+	3,555	+	+	+
	Males	2,024	271	154	+	+	+	2,470	+	+	+
	Females	822	182	76	+	+	+	1,085	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted <sup>f</sup>
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0103%	0.0022%	0.0015%	0.0007%	0.0016%	0.0056%	0.0244%	0.0341%	0.0426%	0.0247%	0.0102%
	Males	0.0144%	0.0024%	0.0017%	0.0007%	0.0020%	0.0080%	0.0400%	0.0501%	0.0594%	0.0381%	0.0150%
	Females	0.0062%	0.0020%	0.0012%	0.0007%	0.0011%	0.0031%	0.0095%	0.0196%	0.0295%	0.0173%	0.0060%
White <sup>c</sup>	Both Sexes	0.0081%	0.0023%	0.0016%	0.0006%	0.0012%	0.0040%	0.0203%	0.0244%	0.0280%	0.0175%	0.0077%
	Males	0.0114%	0.0025%	0.0019%	0.0007%	0.0014%	0.0055%	0.0334%	0.0361%	0.0407%	0.0253%	0.0113%
	Females	0.0047%	0.0021%	0.0012%	0.0006%	0.0009%	0.0025%	0.0074%	0.0136%	0.0179%	0.0132%	0.0044%
Black <sup>c</sup>	Both Sexes	0.0079%	-	-	0.0014%	0.0013%	0.0045%	0.0285%	0.0322%	0.0310%	0.0207%	0.0092%
	Males	0.0107%	-	-	-	-	0.0048%	0.0471%	0.0514%	0.0363%	0.0365%	0.0135%
	Females	0.0052%	-	-	0.0020%	-	0.0043%	0.0130%	0.0173%	0.0274%	0.0130%	0.0059%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0259%	0.0028%	0.0011%	-	0.0036%	0.0158%	0.0454%	0.0979%	0.1400%	0.0794%	0.0267%
	Males	0.0368%	0.0028%	-	-	0.0054%	0.0265%	0.0759%	0.1467%	0.1956%	0.1286%	0.0404%
	Females	0.0158%	0.0029%	-	-	0.0020%	0.0060%	0.0186%	0.0564%	0.0991%	0.0476%	0.0155%
Hispanic <sup>d</sup>	Both Sexes	0.0079%	0.0021%	0.0017%	0.0006%	0.0010%	0.0058%	0.0297%	0.0492%	0.0550%	0.0338%	0.0130%
	Males	0.0107%	0.0026%	0.0021%	-	0.0015%	0.0090%	0.0490%	0.0723%	0.0721%	0.0370%	0.0187%
	Females	0.0050%	0.0015%	0.0012%	-	-	0.0025%	0.0113%	0.0294%	0.0425%	0.0319%	0.0079%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 15.27  
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	213,542	69,437	38,895	20,533	10,999	5,743	331,521	360,606	10,011	370,617
	Males	100,425	30,787	17,826	9,540	4,966	2,750	153,231	167,066	5,673	172,739
	Females	113,117	38,650	21,069	10,993	6,033	2,993	178,290	193,540	4,338	197,878
White <sup>b</sup>	Both Sexes	183,440	61,075	34,181	18,319	9,701	5,177	287,364	313,121	4,676	317,797
	Males	85,221	26,665	15,348	8,367	4,347	2,440	130,986	143,006	2,240	145,246
	Females	98,219	34,410	18,833	9,952	5,354	2,737	156,378	170,115	2,436	172,551
Black <sup>b</sup>	Both Sexes	22,255	6,098	3,559	1,711	1,022	436	32,646	35,256	429	35,685
	Males	11,033	2,890	1,817	927	466	230	16,061	17,500	228	17,728
	Females	11,222	3,208	1,742	784	556	206	16,585	17,756	201	17,957
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	5,962	1,642	760	+	+	+	8,553	+	+	+
	Males	3,109	854	428	+	+	+	4,504	+	+	+
	Females	2,853	788	332	+	+	+	4,049	+	+	+
Hispanic <sup>d</sup>	Both Sexes	6,506	1,898	1,016	+	+	+	9,674	+	+	+
	Males	3,341	929	508	+	+	+	4,879	+	+	+
	Females	3,165	969	508	+	+	+	4,795	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0917%	0.0001%	0.0003%	0.0015%	0.0051%	0.0256%	0.1080%	0.3420%	0.6224%	0.5033%	0.0958%
	Males	0.0867%	-	0.0003%	0.0014%	0.0049%	0.0225%	0.1067%	0.3527%	0.6718%	0.5928%	0.1017%
	Females	0.0967%	-	0.0002%	0.0015%	0.0053%	0.0287%	0.1093%	0.3323%	0.5839%	0.4534%	0.0918%
White <sup>c</sup>	Both Sexes	0.0983%	0.0001%	0.0003%	0.0017%	0.0053%	0.0252%	0.1053%	0.3507%	0.6452%	0.5172%	0.0980%
	Males	0.0901%	-	0.0003%	0.0016%	0.0049%	0.0212%	0.1017%	0.3476%	0.6773%	0.5955%	0.1010%
	Females	0.1064%	-	-	0.0017%	0.0058%	0.0293%	0.1089%	0.3536%	0.6196%	0.4738%	0.0964%
Black <sup>c</sup>	Both Sexes	0.0781%	-	-	0.0009%	0.0055%	0.0349%	0.1631%	0.3971%	0.6395%	0.4228%	0.1054%
	Males	0.0808%	-	-	-	0.0059%	0.0339%	0.1773%	0.4842%	0.7644%	0.5531%	0.1244%
	Females	0.0757%	-	-	-	0.0051%	0.0358%	0.1513%	0.3296%	0.5558%	0.3595%	0.0924%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0695%	-	-	-	0.0039%	0.0204%	0.0804%	0.2483%	0.4793%	0.4449%	0.0741%
	Males	0.0750%	-	-	-	0.0042%	0.0213%	0.0836%	0.3000%	0.5872%	0.5900%	0.0897%
	Females	0.0642%	-	-	-	0.0037%	0.0196%	0.0776%	0.2043%	0.4000%	0.3510%	0.0625%
Hispanic <sup>d</sup>	Both Sexes	0.0211%	-	-	0.0009%	0.0031%	0.0109%	0.0430%	0.1323%	0.2727%	0.2656%	0.0419%
	Males	0.0205%	-	-	0.0008%	0.0033%	0.0108%	0.0452%	0.1357%	0.3320%	0.3529%	0.0488%
	Females	0.0217%	-	-	0.0010%	0.0027%	0.0110%	0.0409%	0.1293%	0.2293%	0.2148%	0.0372%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 16.21  
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	242,029	177,259	124,403	90,820	63,982	42,033	583,389	751,279	42,004	793,283
	Males	127,243	90,751	62,112	42,063	27,714	16,942	299,272	371,049	14,005	385,054
	Females	114,786	86,508	62,291	48,757	36,268	25,091	284,117	380,230	27,999	408,229
White <sup>b</sup>	Both Sexes	233,368	171,359	119,209	87,317	60,938	40,688	562,233	722,928	39,698	762,626
	Males	123,001	87,802	59,552	40,846	26,393	16,310	288,984	357,863	13,356	371,219
	Females	110,367	83,557	59,657	46,471	34,545	24,378	273,249	365,065	26,342	391,407
Black <sup>b</sup>	Both Sexes	852	682	395	359	140	60	2,084	2,574	114	2,688
	Males	313	321	212	124	32	11	925	1,038	45	1,083
	Females	539	361	183	235	108	49	1,159	1,536	69	1,605
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	645	426	225	+	+	+	1,345	+	+	+
	Males	313	171	103	+	+	+	595	+	+	+
	Females	332	255	122	+	+	+	750	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,076	3,066	1,958	+	+	+	9,786	+	+	+
	Males	1,647	1,139	708	+	+	+	3,723	+	+	+
	Females	2,429	1,927	1,250	+	+	+	6,063	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1523%	0.0004%	0.0034%	0.0307%	0.0906%	0.1749%	0.2766%	0.4034%	0.5014%	0.5019%	0.1515%
	Males	0.1603%	0.0004%	0.0031%	0.0198%	0.0628%	0.1510%	0.2952%	0.5112%	0.7141%	0.7798%	0.1739%
	Females	0.1444%	0.0004%	0.0038%	0.0426%	0.1192%	0.1988%	0.2589%	0.3059%	0.3359%	0.3469%	0.1371%
White <sup>c</sup>	Both Sexes	0.1923%	0.0004%	0.0043%	0.0392%	0.1161%	0.2188%	0.3424%	0.4873%	0.6097%	0.5853%	0.1858%
	Males	0.2009%	0.0005%	0.0038%	0.0246%	0.0781%	0.1859%	0.3605%	0.6090%	0.8580%	0.9198%	0.2097%
	Females	0.1836%	0.0004%	0.0048%	0.0553%	0.1569%	0.2529%	0.3245%	0.3741%	0.4119%	0.4006%	0.1714%
Black <sup>c</sup>	Both Sexes	0.0048%	-	-	0.0014%	0.0037%	0.0048%	0.0065%	0.0178%	0.0265%	0.0295%	0.0061%
	Males	0.0044%	-	-	-	0.0015%	0.0042%	0.0053%	0.0212%	0.0424%	0.0397%	0.0071%
	Females	0.0051%	-	-	0.0023%	0.0056%	0.0053%	0.0075%	0.0152%	0.0158%	0.0245%	0.0057%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0098%	-	0.0011%	0.0024%	0.0068%	0.0115%	0.0146%	0.0266%	0.0251%	0.0414%	0.0098%
	Males	0.0092%	-	0.0018%	0.0027%	0.0043%	0.0086%	0.0144%	0.0297%	0.0353%	0.0447%	0.0100%
	Females	0.0103%	-	-	0.0021%	0.0091%	0.0141%	0.0149%	0.0240%	0.0176%	0.0393%	0.0097%
Hispanic <sup>d</sup>	Both Sexes	0.0207%	-	0.0008%	0.0045%	0.0166%	0.0363%	0.0571%	0.0787%	0.1057%	0.1379%	0.0319%
	Males	0.0152%	-	0.0009%	0.0026%	0.0076%	0.0236%	0.0424%	0.0694%	0.1222%	0.2027%	0.0293%
	Females	0.0265%	-	-	0.0068%	0.0270%	0.0499%	0.0711%	0.0866%	0.0937%	0.1001%	0.0360%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 17.13  
MesotheliomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	2,943	596	320	238	150	159	3,965	4,427	+	+
	Males	2,066	286	105	58	80	60	2,489	2,655	+	+
	Females	877	310	215	180	70	99	1,476	1,772	+	+
White <sup>b</sup>	Both Sexes	2,749	485	283	235	127	136	3,624	4,037	+	+
	Males	1,934	231	92	55	57	47	2,290	2,417	+	+
	Females	815	254	191	180	70	89	1,334	1,620	+	+
Black <sup>b</sup>	Both Sexes	97	69	26	0	22	24	191	236	+	+
	Males	61	34	13	0	22	13	107	142	+	+
	Females	36	35	13	0	0	11	84	94	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	47	12	5	+	+	+	64	+	+	+
	Males	35	9	5	+	+	+	49	+	+	+
	Females	12	3	0	+	+	+	15	+	+	+
Hispanic <sup>d</sup>	Both Sexes	261	62	10	+	+	+	338	+	+	+
	Males	153	46	0	+	+	+	199	+	+	+
	Females	108	16	10	+	+	+	139	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0012%	-	-	-	0.0002%	0.0006%	0.0016%	0.0038%	0.0063%	0.0066%	0.0012%
	Males	0.0015%	-	-	-	-	0.0004%	0.0019%	0.0057%	0.0106%	0.0124%	0.0018%
	Females	0.0008%	-	-	-	0.0003%	0.0008%	0.0013%	0.0020%	0.0030%	0.0034%	0.0008%
White <sup>c</sup>	Both Sexes	0.0013%	-	-	-	0.0002%	0.0006%	0.0019%	0.0041%	0.0073%	0.0070%	0.0013%
	Males	0.0017%	-	-	-	-	0.0004%	0.0021%	0.0060%	0.0119%	0.0127%	0.0019%
	Females	0.0010%	-	-	-	0.0003%	0.0008%	0.0016%	0.0023%	0.0036%	0.0039%	0.0009%
Black <sup>c</sup>	Both Sexes	0.0005%	-	-	-	-	-	-	0.0028%	0.0032%	0.0054%	0.0007%
	Males	0.0008%	-	-	-	-	-	-	0.0041%	-	-	0.0012%
	Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0004%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0005%	-	-	-	-	-	-	0.0016%	-	0.0041%	0.0005%
	Males	0.0008%	-	-	-	-	-	-	0.0036%	-	0.0087%	0.0010%
	Females	0.0002%	-	-	-	-	-	-	-	-	-	0.0002%
Hispanic <sup>d</sup>	Both Sexes	0.0008%	-	-	-	-	0.0007%	0.0020%	0.0036%	0.0078%	0.0095%	0.0014%
	Males	0.0009%	-	-	-	-	-	0.0027%	0.0067%	0.0092%	0.0117%	0.0018%
	Females	0.0006%	-	-	-	-	-	0.0014%	-	0.0068%	0.0082%	0.0011%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 18.21  
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	40,775	12,641	4,956	2,062	611	277	59,531	61,360	282	61,642	
	Males	21,757	7,215	2,742	1,190	394	120	32,395	33,418	104	33,522	
	Females	19,018	5,426	2,214	872	217	157	27,136	27,942	178	28,120	
White <sup>b</sup>	Both Sexes	31,926	9,905	3,851	1,515	437	217	46,575	47,863	97	47,960	
	Males	17,731	5,904	2,177	903	285	95	26,365	27,095	0 <sup>i</sup>	27,095	
	Females	14,195	4,001	1,674	612	152	122	20,210	20,768	97 <sup>i</sup>	20,865	
Black <sup>b</sup>	Both Sexes	7,389	2,183	919	469	136	51	10,718	11,172	+	+	
	Males	3,325	1,016	491	261	86	25	4,938	5,204	+	+	
	Females	4,064	1,167	428	208	50	26	5,780	5,968	+	+	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	841	308	126	+	+	+	1,295	+	+	+	
	Males	420	154	64	+	+	+	651	+	+	+	
	Females	421	154	62	+	+	+	644	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	2,737	939	275	+	+	+	4,022	+	+	+	
	Males	1,439	524	140	+	+	+	2,152	+	+	+	
	Females	1,298	415	135	+	+	+	1,870	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Race	Sex	Age Specific (Crude)								Age-Adjusted <sup>f</sup>	
			All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+
All Races <sup>c</sup>	Both Sexes	0.0178%	-	-	0.0001%	0.0020%	0.0086%	0.0288%	0.0650%	0.0962%	0.0866%	0.0181%
	Males	0.0194%	-	-	0.0002%	0.0027%	0.0100%	0.0330%	0.0774%	0.1198%	0.1044%	0.0218%
	Females	0.0163%	-	-	-	0.0012%	0.0071%	0.0248%	0.0539%	0.0777%	0.0766%	0.0151%
White <sup>c</sup>	Both Sexes	0.0174%	-	-	-	0.0018%	0.0080%	0.0265%	0.0609%	0.0915%	0.0804%	0.0170%
	Males	0.0196%	-	-	-	0.0026%	0.0099%	0.0311%	0.0741%	0.1172%	0.0998%	0.0210%
	Females	0.0153%	-	-	-	0.0010%	0.0059%	0.0220%	0.0486%	0.0709%	0.0696%	0.0136%
Black <sup>c</sup>	Both Sexes	0.0270%	-	-	-	0.0043%	0.0170%	0.0594%	0.1278%	0.1793%	0.1722%	0.0356%
	Males	0.0268%	-	-	-	0.0053%	0.0164%	0.0645%	0.1502%	0.2098%	0.1753%	0.0395%
	Females	0.0273%	-	-	-	0.0035%	0.0176%	0.0552%	0.1105%	0.1589%	0.1707%	0.0327%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0105%	-	-	-	0.0006%	0.0038%	0.0156%	0.0393%	0.0572%	0.0674%	0.0109%
	Males	0.0107%	-	-	-	-	0.0042%	0.0177%	0.0417%	0.0643%	0.0830%	0.0124%
	Females	0.0102%	-	-	-	-	0.0035%	0.0138%	0.0373%	0.0521%	0.0573%	0.0098%
Hispanic <sup>d</sup>	Both Sexes	0.0089%	-	-	-	0.0024%	0.0076%	0.0237%	0.0589%	0.0918%	0.0708%	0.0163%
	Males	0.0091%	-	-	-	0.0033%	0.0082%	0.0264%	0.0690%	0.1013%	0.0892%	0.0189%
	Females	0.0087%	-	-	-	0.0014%	0.0069%	0.0211%	0.0502%	0.0848%	0.0601%	0.0144%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 19.25  
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	182,913	110,861	63,509	35,794	20,435	10,965	373,820	427,401	10,924	438,325	
	Males	95,299	57,035	32,947	18,470	10,552	5,417	193,838	221,132	5,723	226,855	
	Females	87,614	53,826	30,562	17,324	9,883	5,548	179,982	206,269	5,201	211,470	
White <sup>b</sup>	Both Sexes	161,338	98,813	56,443	32,481	18,623	10,069	331,476	380,479	11,206	391,685	
	Males	84,198	50,949	29,300	16,811	9,634	4,977	172,148	197,183	6,253	203,436	
	Females	77,140	47,864	27,143	15,670	8,989	5,092	159,328	183,296	4,953	188,249	
Black <sup>b</sup>	Both Sexes	14,184	7,525	4,395	2,174	1,167	613	27,218	30,201	848	31,049	
	Males	7,216	3,766	2,221	1,066	580	269	13,735	15,187	372	15,559	
	Females	6,968	3,759	2,174	1,108	587	344	13,483	15,014	476	15,490	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	4,512	2,708	1,515	+	+	+	9,114	+	+	+	
	Males	2,334	1,432	761	+	+	+	4,732	+	+	+	
	Females	2,178	1,276	754	+	+	+	4,382	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	11,455	6,593	3,495	+	+	+	22,381	+	+	+	
	Males	5,943	3,499	1,841	+	+	+	11,662	+	+	+	
	Females	5,512	3,094	1,654	+	+	+	10,719	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.1078%	0.0023%	0.0110%	0.0209%	0.0371%	0.0814%	0.1652%	0.3072%	0.4837%	0.5123%	0.1092%
	Males	0.1134%	0.0029%	0.0145%	0.0253%	0.0431%	0.0969%	0.1856%	0.3453%	0.5413%	0.5781%	0.1244%
	Females	0.1023%	0.0017%	0.0074%	0.0161%	0.0309%	0.0658%	0.1456%	0.2728%	0.4390%	0.4756%	0.0961%
White <sup>c</sup>	Both Sexes	0.1197%	0.0024%	0.0111%	0.0219%	0.0388%	0.0853%	0.1751%	0.3304%	0.5290%	0.5480%	0.1169%
	Males	0.1251%	0.0031%	0.0146%	0.0268%	0.0453%	0.1014%	0.1965%	0.3680%	0.5907%	0.6147%	0.1326%
	Females	0.1142%	0.0017%	0.0074%	0.0166%	0.0318%	0.0687%	0.1541%	0.2955%	0.4798%	0.5112%	0.1030%
Black <sup>c</sup>	Both Sexes	0.0655%	0.0019%	0.0102%	0.0196%	0.0360%	0.0784%	0.1409%	0.2251%	0.2763%	0.2352%	0.0779%
	Males	0.0703%	0.0022%	0.0142%	0.0224%	0.0449%	0.0942%	0.1595%	0.2670%	0.2746%	0.2769%	0.0890%
	Females	0.0610%	0.0017%	0.0060%	0.0168%	0.0283%	0.0644%	0.1255%	0.1927%	0.2775%	0.2149%	0.0690%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0686%	0.0017%	0.0108%	0.0156%	0.0244%	0.0533%	0.1043%	0.1810%	0.3033%	0.3623%	0.0705%
	Males	0.0726%	0.0021%	0.0134%	0.0204%	0.0251%	0.0627%	0.1135%	0.2037%	0.3484%	0.4261%	0.0807%
	Females	0.0650%	-	0.0079%	0.0108%	0.0238%	0.0445%	0.0963%	0.1617%	0.2701%	0.3209%	0.0623%
Hispanic <sup>d</sup>	Both Sexes	0.0488%	0.0018%	0.0087%	0.0145%	0.0275%	0.0586%	0.1182%	0.2120%	0.3666%	0.3778%	0.0797%
	Males	0.0492%	0.0020%	0.0117%	0.0181%	0.0325%	0.0658%	0.1316%	0.2061%	0.3821%	0.4041%	0.0855%
	Females	0.0482%	0.0015%	0.0056%	0.0101%	0.0216%	0.0510%	0.1053%	0.2170%	0.3553%	0.3624%	0.0742%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 20.27  
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	91,178	55,462	36,425	23,483	15,715	10,014	194,136	235,609	13,757	249,366
	Males	62,485	36,207	23,169	14,155	9,229	5,863	128,665	153,118	7,994	161,112
	Females	28,693	19,255	13,256	9,328	6,486	4,151	65,471	82,491	5,763	88,254
White <sup>b</sup>	Both Sexes	78,100	48,449	31,721	20,976	14,013	8,823	168,105	205,072	11,682	216,754
	Males	54,180	31,962	20,506	12,855	8,448	5,348	112,812	135,130	6,846	141,976
	Females	23,920	16,487	11,215	8,121	5,565	3,475	55,293	69,942	4,836	74,778
Black <sup>b</sup>	Both Sexes	7,594	3,986	2,704	1,340	1,057	672	14,962	17,544	705	18,249
	Males	4,883	2,369	1,468	667	455	233	9,033	10,157	279	10,436
	Females	2,711	1,617	1,236	673	602	439	5,929	7,387	426	7,813
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,410	2,033	1,285	+	+	+	7,068	+	+	+
	Males	2,130	1,232	752	+	+	+	4,324	+	+	+
	Females	1,280	801	533	+	+	+	2,744	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,180	2,382	1,565	+	+	+	8,517	+	+	+
	Males	2,682	1,475	872	+	+	+	5,218	+	+	+
	Females	1,498	907	693	+	+	+	3,299	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0560%	0.0004%	0.0020%	0.0059%	0.0147%	0.0431%	0.1132%	0.1848%	0.2205%	0.2027%	0.0556%
	Males	0.0748%	0.0002%	0.0019%	0.0055%	0.0144%	0.0563%	0.1671%	0.2806%	0.3238%	0.2993%	0.0797%
	Females	0.0376%	0.0005%	0.0021%	0.0063%	0.0150%	0.0299%	0.0617%	0.0980%	0.1402%	0.1488%	0.0353%
White <sup>c</sup>	Both Sexes	0.0594%	0.0004%	0.0019%	0.0055%	0.0132%	0.0407%	0.1145%	0.1941%	0.2330%	0.2132%	0.0569%
	Males	0.0801%	0.0003%	0.0015%	0.0049%	0.0130%	0.0542%	0.1709%	0.2935%	0.3433%	0.3211%	0.0823%
	Females	0.0387%	0.0005%	0.0023%	0.0063%	0.0134%	0.0268%	0.0588%	0.1016%	0.1450%	0.1537%	0.0350%
Black <sup>c</sup>	Both Sexes	0.0370%	-	0.0029%	0.0053%	0.0128%	0.0381%	0.0985%	0.1488%	0.1672%	0.1164%	0.0444%
	Males	0.0479%	-	0.0044%	0.0066%	0.0110%	0.0455%	0.1452%	0.2356%	0.2555%	0.1377%	0.0629%
	Females	0.0270%	-	0.0014%	0.0040%	0.0143%	0.0316%	0.0597%	0.0815%	0.1080%	0.1061%	0.0303%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0513%	-	0.0012%	0.0085%	0.0218%	0.0587%	0.1097%	0.1392%	0.1669%	0.1539%	0.0509%
	Males	0.0646%	-	0.0016%	0.0083%	0.0225%	0.0750%	0.1491%	0.2096%	0.2225%	0.2109%	0.0682%
	Females	0.0388%	-	-	0.0086%	0.0211%	0.0437%	0.0752%	0.0793%	0.1259%	0.1169%	0.0366%
Hispanic <sup>d</sup>	Both Sexes	0.0183%	0.0003%	0.0010%	0.0029%	0.0076%	0.0215%	0.0562%	0.1000%	0.1270%	0.1316%	0.0306%
	Males	0.0219%	-	0.0009%	0.0022%	0.0072%	0.0257%	0.0737%	0.1477%	0.1938%	0.1730%	0.0419%
	Females	0.0145%	-	0.0012%	0.0038%	0.0080%	0.0169%	0.0395%	0.0592%	0.0781%	0.1075%	0.0215%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 21.15  
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	53,433	31,181	22,816	18,814	15,719	11,447	115,526	156,901	20,261	177,162
White <sup>b</sup>	46,925	27,547	20,126	17,210	14,082	10,279	102,101	139,282	17,539	156,821
Black <sup>b</sup>	3,713	1,948	1,573	985	1,049	785	7,541	10,338	1,734	12,072
Asian/Pacific Islander <sup>c</sup>	2,165	1,209	799	+	+	+	4,413	+	+	+
Hispanic <sup>d</sup>	3,993	2,093	1,483	+	+	+	8,061	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0677%	0.0004%	0.0056%	0.0171%	0.0268%	0.0629%	0.1317%	0.1953%	0.2091%	0.1691%	0.0636%	
White <sup>c</sup>	0.0741%	0.0004%	0.0059%	0.0177%	0.0261%	0.0658%	0.1398%	0.2094%	0.2295%	0.1811%	0.0675%	
Black <sup>c</sup>	0.0357%	-	0.0035%	0.0105%	0.0256%	0.0399%	0.0747%	0.1188%	0.1202%	0.0979%	0.0394%	
Asian/Pacific Islander <sup>c</sup>	0.0608%	-	0.0052%	0.0213%	0.0302%	0.0652%	0.1314%	0.1610%	0.1442%	0.1184%	0.0568%	
Hispanic <sup>d</sup>	0.0356%	-	0.0065%	0.0175%	0.0222%	0.0449%	0.0990%	0.1389%	0.1753%	0.1408%	0.0498%	

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.  
<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census.  
<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.  
<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.  
<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.  
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
 + Not available.

Table 22.21  
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	24,579	3,945	1,995	1,118	532	344	31,057	32,616	377	32,993
	Males	12,128	1,973	851	536	177	159	15,239	15,897	160	16,057
	Females	12,451	1,972	1,144	582	355	185	15,818	16,719	217	16,936
White <sup>b</sup>	Both Sexes	21,067	3,218	1,570	866	400	293	26,324	27,499	312	27,811
	Males	10,738	1,760	660	461	132	140	13,424	13,947	146	14,093
	Females	10,329	1,458	910	405	268	153	12,900	13,552	166	13,718
Black <sup>b</sup>	Both Sexes	2,376	477	260	205	95	42	3,160	3,473	148	3,621
	Males	883	134	94	64	38	10	1,124	1,240	51	1,291
	Females	1,493	343	166	141	57	32	2,036	2,233	97	2,330
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	815	152	57	+	+	+	1,040	+	+	+
	Males	363	54	26	+	+	+	451	+	+	+
	Females	452	98	31	+	+	+	589	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,446	245	203	+	+	+	1,922	+	+	+
	Males	672	94	55	+	+	+	840	+	+	+
	Females	774	151	148	+	+	+	1,082	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0094%	-	0.0001%	0.0006%	0.0016%	0.0051%	0.0155%	0.0307%	0.0470%	0.0488%	0.0095%
	Males	0.0091%	-	-	0.0004%	0.0010%	0.0052%	0.0166%	0.0348%	0.0507%	0.0530%	0.0102%
	Females	0.0096%	-	0.0002%	0.0007%	0.0021%	0.0051%	0.0144%	0.0270%	0.0441%	0.0464%	0.0089%
White <sup>c</sup>	Both Sexes	0.0097%	-	0.0001%	0.0006%	0.0015%	0.0048%	0.0161%	0.0305%	0.0471%	0.0479%	0.0095%
	Males	0.0098%	-	-	0.0005%	0.0010%	0.0052%	0.0176%	0.0350%	0.0528%	0.0535%	0.0105%
	Females	0.0097%	-	0.0002%	0.0006%	0.0021%	0.0044%	0.0147%	0.0263%	0.0425%	0.0448%	0.0086%
Black <sup>c</sup>	Both Sexes	0.0077%	-	-	-	0.0017%	0.0054%	0.0165%	0.0370%	0.0478%	0.0436%	0.0101%
	Males	0.0063%	-	-	-	-	0.0052%	0.0162%	0.0369%	0.0383%	0.0331%	0.0090%
	Females	0.0089%	-	-	-	0.0022%	0.0055%	0.0167%	0.0370%	0.0541%	0.0487%	0.0107%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0083%	-	-	-	0.0015%	0.0061%	0.0105%	0.0257%	0.0412%	0.0578%	0.0087%
	Males	0.0074%	-	-	-	-	0.0047%	0.0095%	0.0325%	0.0384%	0.0570%	0.0085%
	Females	0.0092%	-	-	-	0.0020%	0.0074%	0.0115%	0.0199%	0.0433%	0.0584%	0.0088%
Hispanic <sup>d</sup>	Both Sexes	0.0043%	-	-	0.0005%	0.0012%	0.0043%	0.0113%	0.0236%	0.0387%	0.0437%	0.0076%
	Males	0.0036%	-	-	-	0.0008%	0.0036%	0.0100%	0.0260%	0.0405%	0.0375%	0.0075%
	Females	0.0050%	-	-	0.0007%	0.0018%	0.0051%	0.0125%	0.0215%	0.0374%	0.0472%	0.0079%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 23.15  
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	938,613	715,069	438,415	143,491	27,982	8,819	2,182,244	2,273,570	2,542	2,276,112
White <sup>b</sup>	772,875	604,521	378,613	130,377	25,520	8,083	1,837,937	1,921,072	2,819	1,923,891
Black <sup>b</sup>	123,952	87,191	48,959	10,618	2,158	632	266,714	273,584	229	273,813
Asian/Pacific Islander <sup>c</sup>	17,867	11,359	5,396	+	+	+	35,457	+	+	+
Hispanic <sup>d</sup>	48,964	32,063	16,287	+	+	+	99,902	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	1.2331%	0.0002%	0.0002%	-	0.0014%	0.0699%	0.9490%	4.6574%	10.9096%	12.3145%	1.5176%
White <sup>c</sup>	1.2897%	-	0.0002%	-	0.0011%	0.0631%	0.9102%	4.6011%	10.9849%	12.1470%	1.5067%
Black <sup>c</sup>	1.3375%	-	-	-	0.0031%	0.1660%	1.8572%	8.0866%	16.2151%	15.7412%	2.3194%
Asian/Pacific Islander <sup>c</sup>	0.6262%	-	-	-	-	0.0166%	0.3086%	1.9409%	5.6545%	8.5061%	0.8001%
Hispanic <sup>d</sup>	0.4163%	-	-	-	0.0006%	0.0355%	0.5303%	3.1651%	8.3342%	9.5099%	1.1150%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.21  
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	32,686	13,495	7,594	5,188	2,989	1,434	55,990	63,730	1,909	65,639
	Males	19,323	7,719	4,217	2,374	1,624	735	32,190	36,215	1,082	37,297
	Females	13,363	5,776	3,377	2,814	1,365	699	23,800	27,515	827	28,342
White <sup>b</sup>	Both Sexes	24,914	9,954	5,713	3,812	2,311	1,195	42,136	48,171	1,549	49,720
	Males	15,175	5,732	3,106	1,772	1,290	649	24,651	27,901	839	28,740
	Females	9,739	4,222	2,607	2,040	1,021	546	17,485	20,270	710	20,980
Black <sup>b</sup>	Both Sexes	4,890	2,000	1,012	734	406	101	8,231	9,182	248	9,430
	Males	2,458	1,052	605	317	210	40	4,271	4,707	143	4,850
	Females	2,432	948	407	417	196	61	3,960	4,475	105	4,580
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,776	1,350	824	+	+	+	5,216	+	+	+
	Males	1,586	757	479	+	+	+	2,957	+	+	+
	Females	1,190	593	345	+	+	+	2,259	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,135	1,594	883	+	+	+	6,870	+	+	+
	Males	2,226	856	394	+	+	+	3,580	+	+	+
	Females	1,909	738	489	+	+	+	3,290	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0202%	-	-	0.0003%	0.0031%	0.0108%	0.0262%	0.0562%	0.1131%	0.1463%	0.0208%
	Males	0.0232%	-	-	0.0002%	0.0032%	0.0115%	0.0317%	0.0770%	0.1518%	0.1982%	0.0271%
	Females	0.0172%	-	-	0.0004%	0.0030%	0.0101%	0.0210%	0.0373%	0.0831%	0.1174%	0.0159%
White <sup>c</sup>	Both Sexes	0.0168%	-	-	0.0003%	0.0028%	0.0089%	0.0211%	0.0467%	0.0883%	0.1104%	0.0165%
	Males	0.0196%	-	-	0.0002%	0.0030%	0.0094%	0.0264%	0.0651%	0.1183%	0.1492%	0.0216%
	Females	0.0141%	-	-	0.0005%	0.0026%	0.0084%	0.0159%	0.0296%	0.0643%	0.0889%	0.0124%
Black <sup>c</sup>	Both Sexes	0.0194%	-	-	-	0.0038%	0.0163%	0.0377%	0.0754%	0.1211%	0.1793%	0.0261%
	Males	0.0205%	-	-	-	0.0040%	0.0190%	0.0396%	0.0986%	0.1570%	0.2191%	0.0319%
	Females	0.0185%	-	-	-	0.0036%	0.0140%	0.0361%	0.0574%	0.0970%	0.1599%	0.0222%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0418%	-	-	-	0.0039%	0.0166%	0.0478%	0.1040%	0.2636%	0.3947%	0.0448%
	Males	0.0490%	-	-	-	0.0032%	0.0173%	0.0582%	0.1420%	0.3679%	0.5243%	0.0595%
	Females	0.0351%	-	-	-	0.0046%	0.0159%	0.0387%	0.0716%	0.1868%	0.3109%	0.0340%
Hispanic <sup>d</sup>	Both Sexes	0.0149%	-	-	0.0006%	0.0041%	0.0143%	0.0340%	0.0758%	0.1537%	0.2018%	0.0282%
	Males	0.0151%	-	-	0.0006%	0.0041%	0.0138%	0.0373%	0.1042%	0.1845%	0.2150%	0.0329%
	Females	0.0147%	-	-	-	0.0040%	0.0148%	0.0308%	0.0516%	0.1312%	0.1942%	0.0244%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 25.15  
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	39,905	36,907	31,956	28,879	21,830	15,502	120,716	179,020	16,949	195,969
White <sup>b</sup>	37,551	34,584	30,434	27,738	21,118	15,009	113,978	170,262	16,283	186,545
Black <sup>b</sup>	944	1,034	616	378	324	226	2,745	3,637	155	3,792
Asian/Pacific Islander <sup>c</sup>	648	604	468	+	+	+	1,891	+	+	+
Hispanic <sup>d</sup>	4,490	3,498	2,489	+	+	+	11,398	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0694%	0.0018%	0.0056%	0.0638%	0.1426%	0.1540%	0.0857%	0.0349%	0.0148%	0.0093%	0.0682%	
White <sup>c</sup>	0.0841%	0.0017%	0.0063%	0.0777%	0.1771%	0.1858%	0.1005%	0.0414%	0.0171%	0.0104%	0.0826%	
Black <sup>c</sup>	0.0146%	-	0.0022%	0.0127%	0.0263%	0.0367%	0.0222%	0.0041%	-	-	0.0149%	
Asian/Pacific Islander <sup>c</sup>	0.0240%	0.0034%	0.0053%	0.0217%	0.0397%	0.0488%	0.0365%	0.0083%	0.0049%	-	0.0228%	
Hispanic <sup>d</sup>	0.0458%	0.0022%	0.0054%	0.0586%	0.1013%	0.0902%	0.0436%	0.0199%	-	-	0.0446%	

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 26.21  
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	129,436	84,397	60,842	43,865	34,734	28,796	293,782	392,263	41,993	434,256
	Males	28,915	18,810	13,638	9,320	7,506	5,976	65,378	86,537	9,693	96,230
	Females	100,521	65,587	47,204	34,545	27,228	22,820	228,404	305,726	32,300	338,026
White <sup>b</sup>	Both Sexes	113,306	74,195	53,885	39,494	31,213	25,971	258,618	347,261	37,998	385,259
	Males	25,872	17,067	12,412	8,663	6,765	5,476	59,105	78,450	8,184	86,634
	Females	87,434	57,128	41,473	30,831	24,448	20,495	199,513	268,811	29,814	298,625
Black <sup>b</sup>	Both Sexes	7,967	4,681	3,266	1,980	1,621	1,348	16,820	21,343	1,766	23,109
	Males	1,505	766	606	340	306	180	3,014	3,761	193	3,954
	Females	6,462	3,915	2,660	1,640	1,315	1,168	13,806	17,582	1,573	19,155
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	5,842	3,838	2,659	+	+	+	13,108	+	+	+
	Males	1,067	714	445	+	+	+	2,347	+	+	+
	Females	4,775	3,124	2,214	+	+	+	10,761	+	+	+
Hispanic <sup>d</sup>	Both Sexes	11,622	7,347	5,057	+	+	+	25,316	+	+	+
	Males	1,999	1,238	843	+	+	+	4,275	+	+	+
	Females	9,623	6,109	4,214	+	+	+	21,041	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0882%	0.0001%	0.0040%	0.0343%	0.1005%	0.1496%	0.1698%	0.1686%	0.1495%	0.1008%	0.0866%
	Males	0.0389%	-	0.0019%	0.0095%	0.0331%	0.0579%	0.0797%	0.0998%	0.1008%	0.0679%	0.0397%
	Females	0.1366%	-	0.0062%	0.0611%	0.1699%	0.2413%	0.2559%	0.2308%	0.1874%	0.1192%	0.1321%
White <sup>c</sup>	Both Sexes	0.0950%	-	0.0043%	0.0379%	0.1113%	0.1633%	0.1780%	0.1720%	0.1517%	0.0994%	0.0921%
	Males	0.0431%	-	0.0020%	0.0100%	0.0368%	0.0644%	0.0869%	0.1043%	0.1074%	0.0679%	0.0429%
	Females	0.1469%	-	0.0067%	0.0688%	0.1913%	0.2657%	0.2679%	0.2349%	0.1869%	0.1167%	0.1417%
Black <sup>c</sup>	Both Sexes	0.0409%	-	0.0015%	0.0123%	0.0433%	0.0670%	0.0918%	0.1191%	0.0994%	0.0726%	0.0459%
	Males	0.0157%	-	0.0012%	0.0032%	0.0132%	0.0233%	0.0352%	0.0603%	0.0585%	0.0629%	0.0200%
	Females	0.0638%	-	0.0017%	0.0211%	0.0696%	0.1053%	0.1389%	0.1647%	0.1268%	0.0773%	0.0672%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0896%	-	0.0039%	0.0334%	0.0900%	0.1373%	0.1772%	0.1790%	0.1649%	0.1246%	0.0863%
	Males	0.0336%	-	-	0.0119%	0.0282%	0.0459%	0.0660%	0.0940%	0.0802%	0.0684%	0.0342%
	Females	0.1422%	-	0.0070%	0.0552%	0.1486%	0.2215%	0.2747%	0.2514%	0.2272%	0.1609%	0.1324%
Hispanic <sup>d</sup>	Both Sexes	0.0549%	-	0.0035%	0.0272%	0.0695%	0.1110%	0.1502%	0.1512%	0.1512%	0.1010%	0.0718%
	Males	0.0179%	-	0.0013%	0.0058%	0.0193%	0.0299%	0.0575%	0.0755%	0.0788%	0.0510%	0.0267%
	Females	0.0940%	-	0.0058%	0.0527%	0.1279%	0.1978%	0.2387%	0.2160%	0.2042%	0.1301%	0.1164%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 27.21  
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	189,135	130,190	87,503	54,199	33,694	19,021	431,275	518,286	16,950	535,236
	Males	141,812	96,879	64,780	40,296	24,190	13,450	321,574	384,498	10,982	395,480
	Females	47,323	33,311	22,723	13,903	9,504	5,571	109,701	133,788	5,968	139,756
White <sup>b</sup>	Both Sexes	174,734	122,149	82,491	51,862	31,990	18,207	402,630	485,686	13,238	498,924
	Males	131,954	91,226	61,074	38,560	22,936	12,944	301,499	361,591	8,443	370,034
	Females	42,780	30,923	21,417	13,302	9,054	5,263	101,131	124,095	4,795	128,890
Black <sup>b</sup>	Both Sexes	8,035	4,671	3,048	1,513	1,045	433	16,500	18,900	271	19,171
	Males	5,071	3,050	2,256	1,121	761	255	10,882	12,619	216	12,835
	Females	2,964	1,621	792	392	284	178	5,618	6,281	55	6,336
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,016	1,663	1,086	+	+	+	6,030	+	+	+
	Males	2,333	1,215	819	+	+	+	4,559	+	+	+
	Females	683	448	267	+	+	+	1,471	+	+	+
Hispanic <sup>d</sup>	Both Sexes	6,110	3,562	2,348	+	+	+	12,666	+	+	+
	Males	4,588	2,706	1,735	+	+	+	9,507	+	+	+
	Females	1,522	856	613	+	+	+	3,159	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1153%	0.0001%	0.0003%	0.0017%	0.0070%	0.0283%	0.1093%	0.3464%	0.7456%	0.9890%	0.1203%
	Males	0.1747%	-	0.0005%	0.0022%	0.0097%	0.0425%	0.1718%	0.5640%	1.2994%	1.9798%	0.2122%
	Females	0.0569%	-	0.0002%	0.0012%	0.0042%	0.0140%	0.0496%	0.1494%	0.3149%	0.4363%	0.0525%
White <sup>c</sup>	Both Sexes	0.1370%	-	0.0003%	0.0021%	0.0084%	0.0319%	0.1241%	0.3959%	0.8512%	1.0809%	0.1357%
	Males	0.2067%	-	0.0005%	0.0026%	0.0115%	0.0472%	0.1920%	0.6358%	1.4738%	2.1941%	0.2382%
	Females	0.0673%	-	-	0.0015%	0.0051%	0.0161%	0.0572%	0.1727%	0.3552%	0.4658%	0.0589%
Black <sup>c</sup>	Both Sexes	0.0407%	-	-	-	0.0033%	0.0166%	0.0610%	0.1766%	0.3332%	0.4553%	0.0579%
	Males	0.0572%	-	-	-	0.0046%	0.0247%	0.1020%	0.2985%	0.5302%	0.7745%	0.0954%
	Females	0.0257%	-	-	-	0.0022%	0.0095%	0.0270%	0.0821%	0.2011%	0.3002%	0.0329%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0498%	-	-	0.0007%	0.0025%	0.0127%	0.0427%	0.1270%	0.3143%	0.5687%	0.0541%
	Males	0.0775%	-	-	-	0.0032%	0.0210%	0.0710%	0.2194%	0.5609%	1.0399%	0.0961%
	Females	0.0238%	-	-	-	0.0019%	0.0050%	0.0180%	0.0483%	0.1329%	0.2635%	0.0232%
Hispanic <sup>d</sup>	Both Sexes	0.0268%	-	-	0.0007%	0.0037%	0.0137%	0.0482%	0.1548%	0.3327%	0.4518%	0.0546%
	Males	0.0389%	-	-	0.0007%	0.0045%	0.0188%	0.0731%	0.2649%	0.5917%	0.8790%	0.0965%
	Females	0.0141%	-	-	0.0007%	0.0027%	0.0083%	0.0245%	0.0606%	0.1429%	0.2030%	0.0242%

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 28.12  
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2007  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 <sup>e</sup>	0 to <32 <sup>e</sup>	>=32 <sup>f</sup>	Complete <sup>g</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	59,446	51,673	46,411	39,623	34,031	28,556	174,233	269,403	+	+
	Males	32,002	26,696	24,162	20,563	17,099	13,472	91,836	138,408	+	+
	Females	27,444	24,977	22,249	19,060	16,932	15,084	82,397	130,995	+	+
White <sup>b</sup>	Both Sexes	49,454	43,811	38,884	33,972	29,965	25,297	146,398	229,853	+	+
	Males	26,852	22,546	20,340	17,716	15,170	12,095	77,505	118,610	+	+
	Females	22,602	21,265	18,544	16,256	14,795	13,202	68,893	111,243	+	+
Black <sup>b</sup>	Both Sexes	6,466	5,019	4,741	3,695	2,687	2,332	17,784	25,775	+	+
	Males	3,305	2,601	2,354	1,820	1,250	921	9,017	12,643	+	+
	Females	3,161	2,418	2,387	1,875	1,437	1,411	8,767	13,132	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,130	1,725	1,609	+	+	+	5,985	+	+	+
	Males	1,146	952	817	+	+	+	3,184	+	+	+
	Females	984	773	792	+	+	+	2,801	+	+	+
Hispanic <sup>d</sup>	Both Sexes	10,268	7,723	6,255	+	+	+	26,354	+	+	+
	Males	5,698	4,077	3,403	+	+	+	14,393	+	+	+
	Females	4,570	3,646	2,852	+	+	+	11,961	+	+	+

<sup>a</sup> US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census.

<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f g</sup> (f) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <32 and >=32.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 29.7  
U.S. Childhood Cancer Survivors at January 1, 2007  
Number of People Previously Diagnosed with Cancer as Children (Ages 0-19 Years) in the United States and Alive January 1, 2007  
By International Classification of Childhood Cancer (ICCC)<sup>a</sup> Selected Group and Subgroup, Age Group, and Sex

Site	Sex	Complete Prevalence Counts by Age at Prevalence							All Ages		
		0-19	20-29	30-39	40-49	50-59	60+	All Ages	Diagnosed between 1975-2006	Diagnosed before 1975	% <sup>b</sup>
									Count	Count	
All Sites	Both	106,727	87,033	58,533	51,221	28,175	12,258	343,947	269,267	74,680	22
	Male	57,801	44,616	28,369	22,188	12,365	6,824	172,163	138,567	33,596	20
	Female <sup>c</sup>	48,926	42,417	30,164	29,033	15,810	5,434	171,784	130,700	41,084	24
I Leukemia	Both	33,113	17,870	9,164	3,369	867	61	64,444	60,174	4,270	7
	Male	18,119	9,553	4,614	1,725	400	28	34,439	32,468	1,971	6
	Female	14,994	8,317	4,550	1,644	467	33	30,005	27,706	2,299	8
Acute lymphocytic leukemia <sup>d</sup>	Both	28,060	14,859	8,018	2,420	119	0	53,476	50,221	3,255	6
	Male	15,511	7,954	4,262	1,320	80	0	29,127	27,390	1,737	6
	Female	12,549	6,905	3,756	1,100	39	0	24,349	22,831	1,518	6
I(b) Acute myeloid leukemia	Both	3,689	1,883	995	462	117	0	7,146	6,677	469	7
	Male	1,864	955	399	259	56	0	3,533	3,291	242	7
	Female	1,825	928	596	203	61	0	3,613	3,386	227	6
II(a) Hodgkin lymphoma	Both	3,882	9,149	7,899	7,342	3,560	566	32,398	26,669	5,729	18
	Male	2,373	4,327	4,005	3,515	1,819	305	16,344	13,261	3,083	19
	Female	1,509	4,822	3,894	3,827	1,741	261	16,054	13,408	2,646	16
II(b) Non-Hodgkin lymphoma (except Burkitt lymphoma)	Both	6,010	6,179	3,457	2,341	1,408	760	20,155	16,703	3,452	17
	Male	4,045	4,112	2,407	1,361	713	464	13,102	11,226	1,876	14
	Female	1,965	2,067	1,050	980	695	296	7,053	5,477	1,576	22
III CNS and misc intracranial & intraspinial neoplasms	Both	19,496	13,338	9,023	7,857	3,428	837	53,979	42,532	11,447	21
	Male	10,921	7,426	4,790	3,656	1,557	366	28,716	23,430	5,286	18
	Female	8,575	5,912	4,233	4,201	1,871	471	25,263	19,102	6,161	24
IV Neuroblastoma & other peripheral nervous cell tumor	Both	9,176	3,352	1,725	1,414	1,197	451	17,315	13,547	3,768	22
	Male	4,685	1,576	702	528	352	128	7,971	6,744	1,227	15
	Female	4,491	1,776	1,023	886	845	323	9,344	6,803	2,541	27
VI Renal tumors	Both	7,823	4,531	3,333	3,707	1,636	397	21,427	14,221	7,206	34
	Male	3,803	2,027	1,600	1,987	937	236	10,590	6,637	3,953	37
	Female	4,020	2,504	1,733	1,720	699	161	10,837	7,584	3,253	30
VIII Malignant bone tumors	Both	3,329	4,164	2,754	2,497	1,852	486	15,082	11,447	3,635	24
	Male	1,751	2,392	1,483	1,119	856	291	7,892	6,213	1,679	21
	Female	1,578	1,772	1,271	1,378	996	195	7,190	5,234	1,956	27
VIII(a) Osteosarcoma	Both	1,612	2,261	1,554	1,179	1,081	629	8,316	6,054	2,262	27
	Male	753	1,314	852	499	460	370	4,248	3,194	1,054	25
	Female	859	947	702	680	621	259	4,068	2,860	1,208	30
VIII(c) Ewing tumor & related sarcomas of bone	Both	1,258	1,285	819	518	389	38	4,307	3,654	653	15
	Male	734	706	362	269	130	11	2,212	1,972	240	11
	Female	524	579	457	249	259	27	2,095	1,682	413	20
IX Soft tissue & other extraosseous sarcomas	Both	6,956	6,195	4,023	4,738	3,579	3,590	29,081	18,706	10,375	36
	Male	3,842	3,360	1,984	2,179	1,627	2,041	15,033	9,945	5,088	34
	Female	3,114	2,835	2,039	2,559	1,952	1,549	14,048	8,761	5,287	38
X Germ cell & trophoblastic tumors & neoplasms of gonads	Both	4,733	7,196	5,795	6,423	5,746	5,401	35,294	20,932	14,362	41
	Male	2,493	4,497	3,361	3,305	2,892	2,366	18,914	12,183	6,731	36
	Female	2,240	2,699	2,434	3,118	2,854	3,035	16,380	8,749	7,631	47

<sup>a</sup> International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.

<sup>b</sup> Percent diagnosed before 1975 of all ages complete prevalence counts.

<sup>c</sup> All sites females excludes breast cancer.

<sup>d</sup> SEER site recode Acute Lymphocytic Leukemia was used instead of the ICCC classification lymphocytic leukemia (Ia) because of statistical modeling requirements.

For more information regarding the modeling of the prevalence estimates, see Mariotto et al. Long-Term Survivors of Childhood Cancers in the United States. Cancer Epidemiol Biomarkers Prev 2009, page 1034.