

Table 2.10
All Cancer Sites (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Survival Time																				
1-year	64.6	67.6	70.7	73.8	76.4	77.6	76.7	76.2	76.0	76.9	77.6	77.4	78.7	79.2	79.7	79.8	79.5	80.2	80.4	81.2
2-year	54.0	57.1	60.9	64.7	68.1	70.0	69.3	68.3	68.5	70.0	70.7	70.9	72.0	73.0	73.6	73.9	73.5	74.3	74.3	
3-year	48.7	51.8	56.0	60.3	64.2	66.5	65.7	64.9	65.2	66.7	67.6	67.7	69.0	70.0	71.0	71.4	70.7	71.6		
4-year	45.4	48.3	52.7	57.4	61.5	64.3	63.5	62.5	63.0	64.5	65.6	65.8	67.3	68.2	69.2	69.8	69.1			
5-year	42.9	45.7	50.4	55.4	59.5	62.5	61.7	61.0	61.5	63.2	64.1	64.4	66.0	67.0	68.2	68.8				
6-year	41.0	43.8	48.6	53.7	57.8	61.0	60.6	59.7	60.3	62.1	62.9	63.4	65.1	66.1	67.4					
7-year	39.5	42.2	46.9	52.0	56.3	59.6	59.2	58.6	59.3	61.2	62.3	62.6	64.3	65.3						
8-year	38.1	40.9	45.6	50.8	55.0	58.3	58.3	57.6	58.6	60.5	61.5	61.9	63.5							
9-year	37.0	39.9	44.5	49.7	54.0	57.3	57.2	56.7	57.8	59.9	60.8	61.2								
10-year	36.0	39.0	43.5	48.6	53.1	56.2	56.3	56.0	57.1	59.2	60.4									
11-year	35.1	38.2	42.7	47.6	52.1	55.4	55.5	55.2	56.4	58.5										
12-year	34.4	37.5	41.8	46.9	51.4	54.6	54.7	54.4	55.6											
13-year	33.9	36.9	41.0	46.3	50.6	53.8	53.9	53.9												
14-year	33.4	36.4	40.4	45.6	49.9	53.0	53.3													
15-year	32.9	36.0	39.8	44.9	48.9	52.2														
16-year	32.4	35.5	39.2	44.2	48.1															
17-year	32.1	35.1	38.9	43.7																
18-year	31.8	34.9	38.5																	
19-year	31.7	34.7																		
20-year	31.6	34.6																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).