

Table 2.11
All Cancer Sites (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Survival Time																				
1-year	75.4	75.7	77.5	77.9	78.5	78.4	78.2	78.2	78.6	78.6	79.2	79.9	80.0	80.2	80.6	81.0	80.7	80.9	81.2	81.4
2-year	66.8	67.0	69.4	70.3	70.7	71.0	70.9	71.2	71.5	71.8	72.4	73.3	73.8	73.8	74.7	75.1	74.6	74.8	75.1	
3-year	62.2	62.2	65.2	66.1	66.5	67.1	67.0	67.5	67.9	68.3	68.9	70.0	70.5	70.7	71.2	71.8	71.4	71.7		
4-year	59.1	59.1	62.5	63.6	64.1	64.5	64.5	65.0	65.6	66.1	66.7	67.9	68.5	68.7	69.3	69.7	69.3			
5-year	57.0	56.8	60.5	62.0	62.3	62.7	62.8	63.4	64.0	64.5	65.2	66.3	66.9	67.4	67.8	68.2				
6-year	55.3	55.1	59.0	60.5	61.0	61.5	61.6	62.1	62.9	63.3	64.0	65.2	65.8	66.4	66.9					
7-year	54.0	53.7	57.7	59.4	59.9	60.3	60.5	61.1	61.9	62.7	63.2	64.4	65.0	65.5						
8-year	53.0	52.6	56.8	58.5	59.0	59.7	59.5	60.6	61.3	61.9	62.6	63.8	64.3							
9-year	52.1	51.7	56.0	57.7	58.3	59.2	58.9	60.0	60.7	61.2	61.9	63.1								
10-year	51.4	50.8	55.3	57.0	57.8	58.7	58.4	59.4	60.3	60.7	61.4									
11-year	50.7	50.1	54.7	56.5	57.5	58.2	57.9	58.8	59.7	60.5										
12-year	50.2	49.6	54.2	56.1	57.0	57.8	57.6	58.4	59.4											
13-year	49.9	49.2	53.8	55.7	56.6	57.4	57.4	58.2												
14-year	49.5	48.8	53.5	55.3	56.3	57.3	57.1													
15-year	49.1	48.5	53.2	55.1	56.0	57.0														
16-year	48.9	48.1	53.0	54.9	55.7															
17-year	48.7	47.8	52.8	54.7																
18-year	48.6	47.7	52.8																	
19-year	48.4	47.5																		
20-year	48.3	47.4																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).