

Table 4.14
Cancer of the Female Breast (Invasive)

5-Year Relative and Period Survival (Percent) by Race, Diagnosis Year, Stage and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1960-1963 ^a	-	-	-	63	-	-	46	-	-
1970-1973 ^a	-	-	-	68	-	-	51	-	-
1975-1977 ^b	75.3	76.7	74.8	76.1	77.9	75.4	62.4	62.4	62.3
1978-1980 ^b	74.8	75.6	74.6	75.5	76.2	75.3	63.9	67.2	62.0
1981-1983 ^b	76.6	77.5	76.4	77.6	78.7	77.3	64.1	65.4	63.4
1984-1986 ^b	79.4	77.9	79.9	80.5	79.7	80.8	65.3	63.9	66.1
1987-1989 ^b	84.3	81.5	85.3	85.4	82.9	86.2	71.3	69.6	72.3
1990-1992 ^b	85.4	83.4	86.2	86.7	85.2	87.3	71.7	70.2	72.6
1993-1995 ^b	86.7	84.3	87.5	88.0	86.3	88.6	72.8	70.2	74.4
1996-1998 ^b	88.5	86.6	89.3	89.7	88.3	90.2	76.4	74.0	77.9
1999-2006 ^b	89.9 ^e	88.7 ^e	90.4 ^e	91.2 ^e	90.4 ^e	91.6 ^e	78.4 ^e	77.2 ^e	79.0 ^e
5-Year Period Survival (Percent) ^{cd}									
Year of Diagnosis:									
1999-2006	89.2	88.2	89.5	90.5	89.9	90.7	77.2	76.2	77.8
Stage Distribution (%) 1999-2006 ^c									
All Stages									
Number of cases	314,500	79,068	235,432	261,313	61,267	200,046	28,425	9,372	19,053
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	60	53	63	61	54	64	51	46	53
Regional	33	41	30	32	41	29	39	45	35
Distant	5	4	5	5	4	5	8	7	9
Unstaged	2	2	2	2	2	2	3	2	3
5-Year Relative Survival (Percent), 1999-2006 ^c									
Age at Diagnosis:									
Ages <45	86.3	-	-	87.7	-	-	75.4	-	-
Ages 45-54	89.5	-	-	90.9	-	-	78.4	-	-
Ages 55-64	90.0	-	-	91.0	-	-	78.8	-	-
Ages 65-74	90.3	-	-	91.2	-	-	80.7	-	-
Ages 75+	88.2	-	-	89.5	-	-	72.5	-	-
Ages <65	89.0	-	-	90.3	-	-	77.7	-	-
Ages 65+	89.4	-	-	90.4	-	-	77.4	-	-
Stage:									
All Stages	89.0	87.9	89.4	90.2	89.4	90.5	77.5	76.7	78.0
Localized	98.0	96.1	98.7	98.6	96.6	99.3	93.0	91.6	93.7
Regional	83.6	83.4	83.7	84.9	85.2	84.9	72.2	72.3	72.0
Distant	23.4	29.3	21.6	24.8	33.5	22.6	15.2	14.5	15.4
Unstaged	57.9	73.8	52.9	58.5	77.6	53.1	47.6	52.3	45.2
<i>In Situ</i> ^h	100.0	100.0	100.0	100.0	100.0	100.0	100.0	99.5	100.0

^a Based on End Results data from a series of hospital registries and one population-based registry.

^b Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Based on follow-up of patients into 2007.

^c Based on the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006. Based on follow-up of patients into 2007.

^d Period survival provides a more up-to-date estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2004-2006: 0-1 year survival), (2003-2005: 1-2 year survival), (2002-2004: 2-3 year survival), (2001-2003: 3-4 year survival), (2000-2002: 4-5 years survival).

^e The difference between 1975-1977 and 1999-2006 is statistically significant ($p < .05$).

^f The standard error is between 5 and 10 percentage points.

^g The standard error is greater than 10 percentage points.

^h *In situ* cases are not included in the All Stages group.

- Statistic could not be calculated due to fewer than 25 cases during the time period.