

Table 5.7
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>Age-Adjusted Rates, 2003-2007</u>						
Age at Diagnosis/Death:						
All ages	8.1	7.9	10.1	2.4	2.2	4.4
Under 65	7.5	7.5	8.4	1.9	1.7	3.0
65 and over	12.2	10.9	21.6	6.3	5.4	14.3
All ages (IARC world std) ^c	6.4	6.3	7.8	1.8	1.6	3.1
<u>Age-Specific Rates, 2003-2007</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	0.1	-	-	-	-	-
20-24	1.6	1.6	2.0	0.1	0.1	-
25-29	5.3	5.6	5.3	0.5	0.5	1.0
30-34	10.7	11.7	9.6	1.4	1.3	2.1
35-39	13.5	14.5	11.3	2.4	2.3	3.3
40-44	15.1	15.0	16.7	3.4	3.1	5.4
45-49	14.0	13.5	15.7	4.0	3.7	6.4
50-54	13.3	12.6	16.7	4.5	4.2	7.1
55-59	13.2	12.3	18.4	5.2	4.8	8.1
60-64	13.5	11.9	22.0	5.3	4.8	10.0
65-69	14.1	12.9	22.7	5.8	5.1	12.0
70-74	13.0	11.9	20.0	5.8	5.0	12.4
75-79	11.5	10.0	22.0	6.0	5.1	14.7
80-84	10.8	9.4	22.4	7.3	6.5	15.9
85+	9.3	8.1	21.0	7.6	6.4	20.9

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.