

Table 1.10

Age Distribution (%) of Incidence Cases by Site, 2004-2008

All Races, Both Sexes

| Site | Age at Diagnosis | | | | | | | | All Ages | Cases |
|------------------------------------|------------------|-------|-------|-------|-------|-------|-------|------|----------|-----------|
| | <20 | 20-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75-84 | 85+ | | |
| All Sites | 1.1 | 2.7 | 5.6 | 14.1 | 22.7 | 24.7 | 21.4 | 7.8 | 100.0% | 1,751,864 |
| Oral Cavity & Pharynx: | 0.6 | 2.3 | 6.3 | 20.5 | 27.5 | 21.2 | 15.6 | 6.0 | 100.0% | 40,659 |
| Lip | 0.1 | 1.4 | 6.1 | 14.4 | 17.5 | 23.2 | 25.8 | 11.6 | 100.0% | 2,665 |
| Tongue | 0.2 | 2.0 | 6.0 | 20.8 | 31.0 | 21.9 | 13.5 | 4.6 | 100.0% | 11,650 |
| Salivary gland | 2.0 | 6.7 | 7.6 | 15.1 | 18.7 | 18.8 | 20.6 | 10.5 | 100.0% | 4,865 |
| Floor of mouth | 0.1 | 0.2 | 4.0 | 21.1 | 30.2 | 24.5 | 15.7 | 4.2 | 100.0% | 2,245 |
| Gum & other oral cavity | 0.8 | 2.1 | 4.8 | 14.4 | 22.2 | 22.9 | 22.0 | 10.9 | 100.0% | 5,589 |
| Nasopharynx | 3.4 | 6.4 | 13.9 | 24.8 | 23.7 | 16.1 | 8.7 | 2.9 | 100.0% | 2,554 |
| Tonsil | 0.0 | 0.5 | 7.4 | 32.2 | 34.9 | 16.5 | 7.1 | 1.4 | 100.0% | 6,393 |
| Oropharynx | 0.0 | 0.2 | 4.0 | 20.5 | 32.6 | 23.3 | 14.4 | 5.0 | 100.0% | 1,431 |
| Hypopharynx | 0.0 | 0.2 | 1.7 | 16.8 | 29.9 | 28.5 | 18.7 | 4.2 | 100.0% | 2,498 |
| Other oral cavity & pharynx | 0.4 | 0.9 | 3.1 | 15.6 | 30.3 | 24.7 | 18.3 | 6.6 | 100.0% | 769 |
| Digestive System: | 0.2 | 1.0 | 3.7 | 12.9 | 20.8 | 24.3 | 25.6 | 11.4 | 100.0% | 326,516 |
| Esophagus | 0.0 | 0.3 | 2.2 | 11.9 | 24.8 | 27.6 | 25.0 | 8.1 | 100.0% | 16,975 |
| Stomach | 0.1 | 1.6 | 4.7 | 12.1 | 18.5 | 24.2 | 26.6 | 12.2 | 100.0% | 28,959 |
| Small intestine | 0.1 | 1.5 | 5.8 | 15.0 | 23.3 | 24.5 | 21.8 | 8.0 | 100.0% | 7,401 |
| Colon & Rectum: | 0.1 | 1.1 | 3.9 | 12.8 | 19.6 | 24.1 | 26.2 | 12.2 | 100.0% | 176,815 |
| Colon | 0.1 | 0.9 | 3.3 | 10.8 | 18.2 | 24.5 | 28.4 | 13.9 | 100.0% | 126,629 |
| Rectum | 0.0 | 1.5 | 5.6 | 17.9 | 23.0 | 23.3 | 20.6 | 8.1 | 100.0% | 50,186 |
| Colon & Rectum (Male) | 0.1 | 1.1 | 4.0 | 13.7 | 22.0 | 26.1 | 24.4 | 8.6 | 100.0% | 90,265 |
| Colon & Rectum (Female) | 0.1 | 1.1 | 3.8 | 11.8 | 17.0 | 22.1 | 28.0 | 16.0 | 100.0% | 86,550 |
| Anus, anal canal & anorectum | 0.0 | 1.1 | 9.4 | 24.7 | 25.0 | 18.0 | 15.2 | 6.6 | 100.0% | 6,361 |
| Liver & intrahep. bile duct: | 1.1 | 0.9 | 3.2 | 19.2 | 28.4 | 22.5 | 18.9 | 5.9 | 100.0% | 27,949 |
| Liver | 1.2 | 0.9 | 3.2 | 19.8 | 29.1 | 22.2 | 18.3 | 5.4 | 100.0% | 25,644 |
| Intrahepatic bile duct | 0.0 | 1.1 | 3.6 | 12.0 | 20.2 | 25.7 | 25.9 | 11.5 | 100.0% | 2,305 |
| Gallbladder | 0.0 | 0.5 | 2.6 | 8.7 | 17.8 | 25.3 | 30.6 | 14.5 | 100.0% | 4,384 |
| Other biliary | 0.1 | 0.5 | 2.8 | 8.7 | 17.6 | 25.4 | 29.7 | 15.3 | 100.0% | 6,810 |
| Pancreas | 0.0 | 0.4 | 2.2 | 9.7 | 20.3 | 25.4 | 28.6 | 13.3 | 100.0% | 44,804 |
| Retroperitoneum | 9.1 | 4.9 | 9.5 | 15.6 | 20.9 | 19.1 | 15.5 | 5.5 | 100.0% | 1,496 |
| Peritoneum, omentum & mesentery | 0.5 | 1.3 | 3.4 | 10.3 | 24.8 | 30.4 | 23.7 | 5.6 | 100.0% | 2,574 |
| Other digestive system | 0.1 | 1.1 | 3.5 | 10.4 | 18.6 | 22.0 | 29.3 | 15.0 | 100.0% | 1,988 |
| Respiratory System: | 0.1 | 0.4 | 1.7 | 9.2 | 21.3 | 30.8 | 28.3 | 8.2 | 100.0% | 245,440 |
| Nose, nasal cavity & middle ear | 2.2 | 4.9 | 7.8 | 15.6 | 20.6 | 21.4 | 18.7 | 8.8 | 100.0% | 2,702 |
| Larynx | 0.1 | 0.4 | 2.8 | 15.7 | 29.2 | 28.9 | 18.1 | 4.9 | 100.0% | 12,903 |
| Lung & bronchus | 0.0 | 0.2 | 1.6 | 8.8 | 20.9 | 31.1 | 29.0 | 8.3 | 100.0% | 228,955 |
| Lung & bronchus (Male) | 0.0 | 0.2 | 1.5 | 8.7 | 21.9 | 31.8 | 28.6 | 7.4 | 100.0% | 121,024 |
| Lung & bronchus (Female) | 0.0 | 0.3 | 1.7 | 8.9 | 19.8 | 30.4 | 29.6 | 9.4 | 100.0% | 107,931 |
| Pleura | 4.0 | 1.6 | 4.0 | 4.8 | 20.0 | 17.6 | 35.2 | 12.8 | 100.0% | 125 |
| Trachea & other respiratory organs | 17.7 | 21.7 | 9.1 | 13.0 | 12.1 | 11.3 | 9.4 | 5.7 | 100.0% | 755 |
| Bones & joints | 29.0 | 15.4 | 10.5 | 13.0 | 11.4 | 8.3 | 9.1 | 3.5 | 100.0% | 3,457 |
| Soft tissue (including heart) | 9.2 | 9.5 | 10.7 | 15.0 | 16.3 | 15.6 | 16.5 | 7.2 | 100.0% | 12,467 |
| Skin (excl. basal & squamous): | 0.8 | 7.0 | 10.9 | 17.9 | 20.4 | 18.1 | 17.7 | 7.1 | 100.0% | 86,430 |
| Melanoma of the skin | 0.7 | 7.1 | 11.2 | 18.6 | 20.9 | 18.1 | 16.9 | 6.4 | 100.0% | 79,221 |
| Other non-epithelial skin | 1.5 | 6.0 | 7.6 | 10.7 | 14.5 | 17.9 | 27.0 | 14.7 | 100.0% | 7,209 |
| Breast (Female) | 0.0 | 1.9 | 10.2 | 22.6 | 24.4 | 19.7 | 15.5 | 5.6 | 100.0% | 254,683 |
| Breast (Female -in situ) | 0.0 | 0.7 | 11.1 | 28.9 | 26.3 | 18.8 | 11.9 | 2.3 | 100.0% | 63,173 |

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Percents may not sum to 100 due to rounding.

Table 1.10 - continued

Age Distribution (%) of Incidence Cases by Site, 2004-2008

All Races, Both Sexes

| Site | Age at Diagnosis | | | | | | | | All Ages | Cases |
|---------------------------------------|------------------|-------|-------|-------|-------|-------|-------|------|----------|---------|
| | <20 | 20-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75-84 | 85+ | | |
| Female Genital System: | 0.4 | 4.2 | 9.6 | 19.7 | 26.1 | 19.4 | 14.7 | 5.9 | 100.0% | 100,068 |
| Cervix uteri | 0.2 | 14.3 | 25.8 | 23.9 | 16.4 | 10.6 | 6.4 | 2.5 | 100.0% | 15,953 |
| Corpus uteri | 0.0 | 1.6 | 6.1 | 19.3 | 32.1 | 22.2 | 14.1 | 4.5 | 100.0% | 48,508 |
| Uterus, NOS | 0.3 | 2.1 | 5.2 | 15.1 | 21.0 | 19.0 | 20.2 | 17.0 | 100.0% | 1,215 |
| Ovary ^a | 1.2 | 3.5 | 7.3 | 19.1 | 23.1 | 19.7 | 18.2 | 8.0 | 100.0% | 26,550 |
| Vagina | 0.9 | 0.9 | 5.5 | 14.5 | 21.5 | 22.2 | 21.4 | 13.1 | 100.0% | 1,485 |
| Vulva | 0.1 | 2.0 | 7.3 | 15.9 | 18.3 | 17.5 | 24.3 | 14.6 | 100.0% | 4,737 |
| Other female genital system | 0.9 | 6.9 | 8.6 | 14.9 | 24.1 | 20.9 | 16.8 | 6.9 | 100.0% | 1,620 |
| Male Genital System: | 0.2 | 1.9 | 1.7 | 9.3 | 29.6 | 33.9 | 19.1 | 4.3 | 100.0% | 274,701 |
| Prostate | 0.0 | 0.0 | 0.6 | 9.1 | 30.7 | 35.3 | 19.9 | 4.4 | 100.0% | 262,238 |
| Testis | 6.1 | 47.6 | 26.6 | 13.8 | 4.0 | 1.2 | 0.6 | 0.2 | 100.0% | 10,726 |
| Penis | 0.1 | 2.1 | 6.8 | 11.8 | 20.2 | 23.8 | 23.7 | 11.5 | 100.0% | 1,327 |
| Other male genital system | 2.4 | 2.4 | 7.1 | 13.2 | 21.2 | 21.7 | 22.9 | 9.0 | 100.0% | 410 |
| Urinary System: | 0.6 | 0.9 | 3.5 | 10.9 | 20.9 | 26.1 | 26.9 | 10.3 | 100.0% | 136,851 |
| Urinary bladder | 0.1 | 0.4 | 1.7 | 7.4 | 18.0 | 27.2 | 32.0 | 13.2 | 100.0% | 78,104 |
| Kidney & renal pelvis | 1.3 | 1.7 | 6.0 | 16.3 | 25.2 | 24.5 | 19.2 | 5.9 | 100.0% | 55,465 |
| Ureter | 0.0 | 0.1 | 0.9 | 4.2 | 14.2 | 29.0 | 37.1 | 14.6 | 100.0% | 2,167 |
| Other urinary system | 0.2 | 0.5 | 2.2 | 7.9 | 16.6 | 24.6 | 30.4 | 17.6 | 100.0% | 1,115 |
| Eye & Orbit | 13.7 | 3.4 | 7.2 | 14.2 | 20.1 | 19.3 | 16.3 | 5.9 | 100.0% | 3,101 |
| Brain & Nervous System: | 12.9 | 8.9 | 9.6 | 15.2 | 18.6 | 16.3 | 13.9 | 4.5 | 100.0% | 24,633 |
| Brain | 12.3 | 8.8 | 9.4 | 15.2 | 18.8 | 16.7 | 14.2 | 4.6 | 100.0% | 23,107 |
| Cranial nerves & other nervous system | 22.6 | 10.6 | 11.7 | 15.9 | 15.7 | 10.4 | 9.6 | 3.7 | 100.0% | 1,526 |
| Endocrine System: | 3.0 | 15.5 | 20.1 | 23.6 | 18.3 | 11.7 | 6.4 | 1.5 | 100.0% | 45,237 |
| Thyroid | 1.7 | 16.0 | 20.8 | 24.2 | 18.3 | 11.4 | 6.1 | 1.4 | 100.0% | 42,389 |
| Other endocrine & thymus | 21.6 | 7.1 | 9.1 | 15.1 | 18.4 | 15.7 | 10.4 | 2.6 | 100.0% | 2,848 |
| Lymphoma: | 3.0 | 7.3 | 7.9 | 13.5 | 18.5 | 20.4 | 21.1 | 8.2 | 100.0% | 85,449 |
| Hodgkin lymphoma | 12.3 | 31.5 | 15.8 | 12.5 | 9.7 | 8.5 | 7.2 | 2.3 | 100.0% | 10,880 |
| Non-Hodgkin lymphoma | 1.7 | 3.8 | 6.8 | 13.7 | 19.7 | 22.1 | 23.2 | 9.0 | 100.0% | 74,569 |
| Myeloma | 0.0 | 0.5 | 3.3 | 12.0 | 21.6 | 26.5 | 26.4 | 9.8 | 100.0% | 21,504 |
| Leukemia: | 10.8 | 4.8 | 5.3 | 10.3 | 15.7 | 19.7 | 22.5 | 10.8 | 100.0% | 46,877 |
| Lymphocytic: | 16.3 | 3.1 | 3.4 | 9.0 | 16.5 | 20.2 | 21.5 | 10.1 | 100.0% | 23,694 |
| Acute lymphocytic | 60.3 | 10.3 | 5.9 | 6.7 | 6.1 | 5.0 | 4.0 | 1.7 | 100.0% | 6,373 |
| Chronic lymphocytic | 0.1 | 0.2 | 1.5 | 8.8 | 20.3 | 26.6 | 28.9 | 13.6 | 100.0% | 15,761 |
| Other lymphocytic | 0.3 | 2.4 | 11.7 | 20.3 | 20.8 | 17.5 | 18.2 | 8.8 | 100.0% | 1,560 |
| Myeloid & Monocytic: | 5.2 | 6.9 | 7.6 | 12.1 | 15.3 | 19.4 | 23.2 | 10.2 | 100.0% | 20,728 |
| Acute myeloid | 6.1 | 6.6 | 6.6 | 11.4 | 15.1 | 20.0 | 24.2 | 10.1 | 100.0% | 13,240 |
| Chronic myeloid | 2.6 | 7.7 | 9.9 | 13.9 | 15.6 | 19.1 | 21.3 | 10.0 | 100.0% | 5,833 |
| Acute monocytic | 10.4 | 6.7 | 7.9 | 13.4 | 18.1 | 15.1 | 18.9 | 9.4 | 100.0% | 1,057 |
| Other myeloid & monocytic | 3.7 | 7.2 | 8.7 | 8.7 | 12.9 | 19.2 | 25.4 | 14.2 | 100.0% | 598 |
| Other leukemia: | 4.6 | 4.2 | 4.4 | 7.5 | 11.4 | 17.1 | 27.4 | 23.4 | 100.0% | 2,455 |
| Other acute leukemia | 8.0 | 4.5 | 3.8 | 6.2 | 9.7 | 15.9 | 29.0 | 23.0 | 100.0% | 952 |
| Aleukemic, subleukemic & NOS | 2.5 | 4.0 | 4.7 | 8.3 | 12.6 | 17.9 | 26.3 | 23.6 | 100.0% | 1,503 |
| Kaposi Sarcoma | 0.1 | 17.6 | 34.2 | 20.2 | 7.8 | 6.5 | 8.2 | 5.5 | 100.0% | 2,293 |
| Mesothelioma | 0.1 | 0.7 | 2.2 | 6.4 | 16.5 | 26.9 | 35.0 | 12.3 | 100.0% | 3,913 |
| Ill-defined & unspecified | 0.5 | 0.9 | 2.7 | 9.6 | 17.5 | 22.3 | 28.9 | 17.7 | 100.0% | 35,604 |

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).
 Percents may not sum to 100 due to rounding.

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.12

Age Distribution (%) of Deaths by Site, 2004-2008

All Races, Both Sexes

| Site | Age at Death | | | | | | | | All Ages | Deaths |
|------------------------------------|--------------|-------|-------|-------|-------|-------|-------|------|----------|-----------|
| | <20 | 20-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75-84 | 85+ | | |
| All Sites | 0.4 | 0.8 | 2.5 | 9.0 | 18.0 | 24.8 | 29.3 | 15.2 | 100.0% | 2,801,390 |
| Oral Cavity & Pharynx: | 0.1 | 0.8 | 3.2 | 14.5 | 24.2 | 23.8 | 21.8 | 11.5 | 100.0% | 39,405 |
| Lip | 0.0 | 0.6 | 1.9 | 9.1 | 10.7 | 16.0 | 31.0 | 30.7 | 100.0% | 319 |
| Tongue | 0.1 | 1.1 | 3.9 | 15.4 | 24.8 | 23.8 | 20.3 | 10.7 | 100.0% | 9,752 |
| Salivary gland | 0.1 | 0.8 | 3.3 | 9.3 | 16.8 | 21.8 | 28.8 | 19.1 | 100.0% | 3,557 |
| Floor of mouth | 0.0 | 0.5 | 2.8 | 16.6 | 28.7 | 25.3 | 18.0 | 8.0 | 100.0% | 572 |
| Gum & other oral cavity | 0.3 | 0.5 | 2.1 | 9.5 | 18.5 | 22.1 | 26.5 | 20.6 | 100.0% | 5,663 |
| Nasopharynx | 0.7 | 3.2 | 7.1 | 19.1 | 24.5 | 21.8 | 16.5 | 7.0 | 100.0% | 3,207 |
| Tonsil | 0.0 | 0.2 | 3.6 | 23.7 | 30.9 | 21.8 | 15.3 | 4.5 | 100.0% | 3,305 |
| Oropharynx | 0.0 | 0.2 | 2.7 | 14.9 | 27.6 | 24.3 | 20.4 | 10.0 | 100.0% | 3,418 |
| Hypopharynx | 0.0 | 0.3 | 1.4 | 13.9 | 29.4 | 27.4 | 21.7 | 5.9 | 100.0% | 1,568 |
| Other oral cavity & pharynx | 0.0 | 0.1 | 2.2 | 13.8 | 25.9 | 26.9 | 22.4 | 8.8 | 100.0% | 8,044 |
| Digestive System: | 0.1 | 0.5 | 2.3 | 9.5 | 18.4 | 23.7 | 28.9 | 16.6 | 100.0% | 676,276 |
| Esophagus | 0.0 | 0.3 | 2.0 | 10.9 | 23.8 | 27.4 | 25.7 | 9.9 | 100.0% | 67,513 |
| Stomach | 0.0 | 1.2 | 3.9 | 10.0 | 16.4 | 22.5 | 28.8 | 17.2 | 100.0% | 57,458 |
| Small intestine | 0.0 | 0.8 | 3.6 | 10.3 | 18.8 | 23.3 | 28.4 | 14.8 | 100.0% | 5,598 |
| Colon & Rectum: | 0.0 | 0.6 | 2.4 | 8.4 | 16.0 | 22.0 | 29.8 | 20.7 | 100.0% | 265,857 |
| Colon & Rectum (Male) | 0.0 | 0.7 | 2.6 | 9.3 | 18.7 | 25.0 | 29.1 | 14.6 | 100.0% | 134,400 |
| Colon & Rectum (Female) | 0.0 | 0.6 | 2.3 | 7.4 | 13.3 | 19.0 | 30.5 | 26.9 | 100.0% | 131,457 |
| Anus, anal canal & anorectum | 0.0 | 0.6 | 6.4 | 19.5 | 23.3 | 19.7 | 19.7 | 10.9 | 100.0% | 3,157 |
| Liver & intrahep. bile duct: | 0.3 | 0.7 | 2.2 | 14.7 | 23.2 | 23.1 | 25.4 | 10.5 | 100.0% | 83,280 |
| Liver | 0.4 | 0.7 | 2.1 | 16.3 | 24.3 | 22.6 | 24.1 | 9.6 | 100.0% | 64,674 |
| Intrahepatic bile duct | 0.0 | 0.6 | 2.6 | 9.3 | 19.2 | 24.9 | 29.9 | 13.4 | 100.0% | 18,606 |
| Gallbladder | 0.0 | 0.3 | 1.7 | 7.0 | 16.7 | 25.1 | 32.1 | 17.2 | 100.0% | 9,810 |
| Other biliary | 0.0 | 0.2 | 1.5 | 5.7 | 14.2 | 22.2 | 33.5 | 22.7 | 100.0% | 7,165 |
| Pancreas | 0.0 | 0.2 | 1.6 | 8.2 | 18.7 | 25.5 | 30.5 | 15.3 | 100.0% | 167,335 |
| Retroperitoneum | 0.6 | 2.1 | 3.4 | 11.0 | 18.0 | 24.3 | 27.2 | 13.3 | 100.0% | 1,077 |
| Peritoneum, omentum & mesentery | 0.1 | 0.7 | 2.1 | 7.7 | 18.8 | 29.7 | 30.4 | 10.4 | 100.0% | 3,919 |
| Other digestive system | 0.1 | 0.4 | 1.9 | 7.3 | 15.8 | 21.2 | 30.1 | 23.2 | 100.0% | 4,107 |
| Respiratory System: | 0.0 | 0.1 | 1.3 | 8.0 | 19.8 | 30.4 | 30.4 | 10.0 | 100.0% | 816,325 |
| Nose, nasal cavity & middle ear | 0.2 | 1.5 | 6.1 | 12.8 | 19.5 | 20.9 | 24.4 | 14.7 | 100.0% | 2,359 |
| Larynx | 0.0 | 0.1 | 1.5 | 11.7 | 25.3 | 28.9 | 23.7 | 8.7 | 100.0% | 18,679 |
| Lung & bronchus | 0.0 | 0.1 | 1.3 | 7.9 | 19.6 | 30.5 | 30.6 | 10.0 | 100.0% | 793,097 |
| Lung & bronchus (Male) | 0.0 | 0.1 | 1.2 | 8.0 | 20.7 | 31.3 | 30.0 | 8.7 | 100.0% | 445,827 |
| Lung & bronchus (Female) | 0.0 | 0.1 | 1.4 | 7.7 | 18.3 | 29.4 | 31.3 | 11.7 | 100.0% | 347,270 |
| Pleura | 0.0 | 0.3 | 1.8 | 3.6 | 15.4 | 26.4 | 38.7 | 13.9 | 100.0% | 1,145 |
| Trachea & other respiratory organs | 0.9 | 4.2 | 3.6 | 12.7 | 16.4 | 22.0 | 27.3 | 12.9 | 100.0% | 1,045 |
| Bones & joints | 13.8 | 14.5 | 6.5 | 10.3 | 12.3 | 14.3 | 17.3 | 11.1 | 100.0% | 6,751 |
| Soft tissue (including heart) | 3.9 | 6.3 | 7.1 | 13.4 | 18.3 | 19.1 | 21.5 | 10.5 | 100.0% | 19,658 |
| Skin (excl. basal & squamous): | 0.1 | 2.1 | 4.9 | 12.4 | 18.9 | 20.7 | 25.2 | 15.7 | 100.0% | 54,967 |
| Melanoma of the skin | 0.1 | 2.7 | 5.8 | 13.9 | 19.7 | 21.2 | 24.3 | 12.3 | 100.0% | 41,822 |
| Other non-epithelial skin | 0.0 | 0.3 | 1.8 | 7.8 | 16.3 | 19.1 | 28.1 | 26.5 | 100.0% | 13,145 |
| Breast (Female) | 0.0 | 0.9 | 5.7 | 14.9 | 21.1 | 19.8 | 22.3 | 15.3 | 100.0% | 204,077 |

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
 Percents may not sum to 100 due to rounding.

Table 1.12 - continued

Age Distribution (%) of Deaths by Site, 2004-2008

All Races, Both Sexes

| Site | Age at Death | | | | | | | | All Ages | Deaths |
|------------------------------|--------------|-------|-------|-------|-------|-------|-------|------|----------|---------|
| | <20 | 20-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75-84 | 85+ | | |
| Female Genital System: | 0.0 | 1.2 | 4.5 | 11.9 | 20.1 | 23.4 | 25.2 | 13.7 | 100.0% | 137,708 |
| Cervix uteri | 0.0 | 4.8 | 15.8 | 23.6 | 21.3 | 15.4 | 12.5 | 6.6 | 100.0% | 19,779 |
| Corpus uteri | 0.0 | 0.3 | 2.0 | 7.2 | 21.1 | 27.8 | 27.2 | 14.5 | 100.0% | 16,793 |
| Uterus, NOS | 0.0 | 0.4 | 2.2 | 8.6 | 20.3 | 25.0 | 26.8 | 16.5 | 100.0% | 19,808 |
| Ovary | 0.1 | 0.7 | 2.7 | 11.0 | 20.2 | 24.6 | 27.3 | 13.5 | 100.0% | 73,343 |
| Vagina | 0.0 | 0.7 | 3.6 | 7.5 | 14.8 | 18.3 | 29.0 | 26.2 | 100.0% | 1,987 |
| Vulva | 0.0 | 0.6 | 2.5 | 7.2 | 11.6 | 17.5 | 31.2 | 29.5 | 100.0% | 4,263 |
| Other female genital system | 0.1 | 1.8 | 4.0 | 11.9 | 17.6 | 25.2 | 24.6 | 14.9 | 100.0% | 1,735 |
| Male Genital System: | 0.0 | 0.4 | 0.4 | 1.8 | 7.9 | 19.6 | 38.9 | 30.9 | 100.0% | 146,986 |
| Prostate | 0.0 | 0.0 | 0.1 | 1.5 | 7.7 | 19.8 | 39.5 | 31.4 | 100.0% | 143,843 |
| Testis | 2.8 | 33.1 | 24.2 | 17.4 | 10.3 | 5.1 | 4.2 | 3.0 | 100.0% | 1,758 |
| Penis | 0.0 | 0.8 | 4.5 | 10.8 | 19.7 | 24.2 | 25.7 | 14.3 | 100.0% | 1,185 |
| Other male genital system | 0.5 | 3.0 | 2.5 | 9.5 | 14.0 | 21.0 | 27.0 | 22.5 | 100.0% | 200 |
| Urinary System: | 0.2 | 0.3 | 1.4 | 6.9 | 15.6 | 22.8 | 32.3 | 20.6 | 100.0% | 133,982 |
| Urinary bladder | 0.0 | 0.1 | 0.8 | 4.1 | 11.3 | 20.9 | 36.3 | 26.6 | 100.0% | 67,636 |
| Kidney & renal pelvis | 0.4 | 0.5 | 2.1 | 10.0 | 20.3 | 25.0 | 27.7 | 14.1 | 100.0% | 62,807 |
| Ureter | 0.0 | 0.1 | 0.3 | 3.3 | 9.6 | 22.6 | 40.4 | 23.6 | 100.0% | 1,736 |
| Other urinary system | 0.0 | 0.3 | 1.2 | 6.4 | 13.3 | 21.3 | 36.6 | 20.9 | 100.0% | 1,803 |
| Eye & Orbit | 3.1 | 1.8 | 4.6 | 11.5 | 18.6 | 23.4 | 24.1 | 12.9 | 100.0% | 1,190 |
| Brain & Nervous System: | 4.1 | 3.7 | 6.7 | 14.8 | 22.3 | 22.4 | 19.5 | 6.6 | 100.0% | 65,825 |
| Endocrine System: | 7.4 | 2.5 | 4.2 | 9.8 | 16.8 | 21.5 | 24.7 | 13.0 | 100.0% | 12,073 |
| Thyroid | 0.1 | 0.8 | 2.2 | 7.8 | 16.9 | 24.4 | 30.4 | 17.3 | 100.0% | 7,600 |
| Other endocrine & thymus | 19.9 | 5.4 | 7.4 | 13.3 | 16.6 | 16.5 | 15.0 | 5.9 | 100.0% | 4,473 |
| Lymphoma: | 0.5 | 2.2 | 3.0 | 7.1 | 14.1 | 21.8 | 33.1 | 18.2 | 100.0% | 109,617 |
| Hodgkin lymphoma | 1.6 | 13.2 | 10.2 | 12.3 | 14.7 | 16.6 | 21.9 | 9.4 | 100.0% | 6,317 |
| Non-Hodgkin lymphoma | 0.5 | 1.5 | 2.5 | 6.8 | 14.0 | 22.1 | 33.7 | 18.8 | 100.0% | 103,300 |
| Myeloma | 0.0 | 0.1 | 1.2 | 6.3 | 16.3 | 26.2 | 34.4 | 15.5 | 100.0% | 53,526 |
| Leukemia: | 2.9 | 3.1 | 3.1 | 6.4 | 12.8 | 21.6 | 31.3 | 18.8 | 100.0% | 109,563 |
| Lymphocytic: | 4.5 | 3.8 | 2.4 | 5.0 | 11.2 | 19.0 | 30.0 | 24.1 | 100.0% | 31,074 |
| Acute lymphocytic | 19.5 | 15.8 | 8.6 | 11.2 | 12.9 | 13.0 | 12.9 | 6.2 | 100.0% | 7,066 |
| Chronic lymphocytic | 0.0 | 0.1 | 0.5 | 3.0 | 10.6 | 20.9 | 35.3 | 29.6 | 100.0% | 22,097 |
| Other lymphocytic | 1.4 | 1.7 | 1.7 | 5.8 | 11.2 | 18.9 | 32.1 | 27.3 | 100.0% | 1,911 |
| Myeloid & Monocytic: | 2.2 | 3.2 | 3.8 | 8.1 | 15.0 | 23.7 | 30.5 | 13.5 | 100.0% | 51,610 |
| Acute myeloid | 2.3 | 3.3 | 3.8 | 8.2 | 15.6 | 24.6 | 30.2 | 12.0 | 100.0% | 42,550 |
| Chronic myeloid | 1.1 | 3.5 | 5.2 | 9.0 | 12.9 | 18.9 | 28.3 | 21.2 | 100.0% | 5,292 |
| Acute monocytic | 2.1 | 2.3 | 1.9 | 4.4 | 10.2 | 20.5 | 37.0 | 21.7 | 100.0% | 479 |
| Other myeloid & monocytic | 1.7 | 1.5 | 2.0 | 5.0 | 11.8 | 21.4 | 37.6 | 18.9 | 100.0% | 3,289 |
| Other leukemia: | 2.4 | 2.3 | 2.6 | 4.9 | 10.5 | 20.4 | 34.2 | 22.7 | 100.0% | 26,879 |
| Other acute leukemia | 1.3 | 2.7 | 2.8 | 5.1 | 10.4 | 21.3 | 35.3 | 21.1 | 100.0% | 10,886 |
| Aleukemic, subleukemic & NOS | 3.1 | 2.1 | 2.4 | 4.8 | 10.5 | 19.7 | 33.5 | 23.8 | 100.0% | 15,993 |
| Ill-defined & unspecified | 0.2 | 0.7 | 2.2 | 8.2 | 17.0 | 23.3 | 30.3 | 18.0 | 100.0% | 211,527 |

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
 Percents may not sum to 100 due to rounding.