

Table 2.5  
All Cancer Sites (Invasive)

Age-adjusted SEER Incidence<sup>a</sup> Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975	400.40	466.79	365.83	402.10	468.57	369.54	426.49	525.38	356.65
1976	407.39	481.12	367.40	409.71	481.84	373.14	427.36	553.02	340.40
1977	407.74	486.48	363.73	409.30	487.09	367.91	442.59	577.58	352.01
1978	407.33	488.40	362.23	409.84	489.81	367.97	436.49	570.67	344.07
1979	412.52	496.47	365.54	414.10	497.56	369.71	451.57	589.90	359.00
1980	418.04	505.78	368.10	419.48	505.19	373.36	467.50	615.13	366.56
1981	425.25	510.84	377.09	427.65	510.81	383.86	472.73	633.91	362.95
1982	424.46	511.06	376.17	427.73	512.45	383.38	465.38	621.86	359.81
1983	431.17	519.97	380.78	434.41	520.99	387.98	487.03	645.92	382.08
1984	439.96	525.41	392.45	443.61	527.62	399.56	488.07	643.78	386.00
1985	448.81	531.18	403.13	453.40	533.42	411.49	488.92	643.41	388.12
1986	451.30	538.16	402.21	455.87	542.77	408.98	493.84	644.13	396.37
1987	467.95	563.06	413.23	474.81	568.66	423.29	500.91	665.97	394.02
1988	463.74	558.31	409.10	470.19	565.01	417.72	503.65	657.04	404.27
1989	467.62	566.83	408.24	475.00	573.29	418.50	492.41	660.99	384.44
1990	481.76	592.50	415.30	488.67	599.04	424.63	520.97	699.05	407.92
1991	502.88	637.24	418.80	510.26	644.24	428.30	548.84	764.18	412.68
1992	510.39	657.83	414.72	516.60	662.99	423.39	568.00	807.44	413.04
1993	493.18	621.76	408.17	496.74	619.85	417.01	567.79	817.55	403.65
1994	483.19	589.28	413.28	487.10	587.06	422.86	556.99	781.77	408.52
1995	476.48	571.01	414.53	481.71	569.59	425.77	541.21	746.20	403.99
1996	478.72	571.96	417.72	484.35	573.20	427.77	538.99	724.71	416.02
1997	485.98	575.04	428.05	493.06	575.84	440.72	539.04	737.56	409.52
1998	487.56	571.96	433.27	496.12	575.62	446.34	533.29	718.58	414.14
1999	489.83	579.87	429.93	497.78	581.68	442.99	537.77	718.85	418.90
2000	485.15	578.55	421.39	495.27	582.03	437.07	524.66	708.94	401.44
2001	488.50	579.10	426.17	499.31	584.95	441.28	517.24	694.26	398.18
2002	485.90	572.30	426.86	495.39	578.15	439.33	531.58	695.68	422.64
2003	472.95	556.91	415.48	482.76	561.01	429.95	514.31	675.92	407.28
2004	472.97	553.88	417.31	482.07	558.47	429.86	516.53	675.68	412.26
2005	467.30	542.31	415.81	477.45	549.06	428.93	499.12	631.86	411.89
2006	469.28	546.22	415.10	478.25	550.41	427.96	498.61	632.29	407.55
2007	473.41	554.76	415.39	480.85	557.11	426.98	507.35	655.44	406.34
2008	463.37	530.84	416.33	469.34	531.03	427.11	503.18	644.98	407.41
1975-2008	463.17	553.28	405.99	468.54	555.54	415.14	507.72	675.17	396.87

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).  
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.