


Table 2.18  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	181.32	0.11		
High Five States				
Kentucky	213.58 <sup>b</sup>	0.99	(01)	17.79 <sup>c</sup>
Louisiana	208.41 <sup>b</sup>	0.98	(02)	14.94 <sup>c</sup>
West Virginia	207.83 <sup>b</sup>	1.37	(03)	14.62 <sup>c</sup>
Mississippi	206.82 <sup>b</sup>	1.19	(04)	14.06 <sup>c</sup>
Tennessee	202.83 <sup>b</sup>	0.80	(05)	11.86 <sup>c</sup>
Low Five States				
New Mexico	160.84 <sup>b</sup>	1.28	(47)	-11.29 <sup>c</sup>
Arizona	156.15 <sup>b</sup>	0.70	(48)	-13.88 <sup>c</sup>
Colorado	156.09 <sup>b</sup>	0.88	(49)	-13.91 <sup>c</sup>
Hawaii	149.15 <sup>b</sup>	1.44	(50)	-17.74 <sup>c</sup>
Utah	131.78 <sup>b</sup>	1.18	(51)	-27.32 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	199.86 <sup>b</sup>	0.90	(07)	10.23 <sup>c</sup>	Montana	175.70	1.81	(36)	-3.10
Alaska	181.15	3.11	(30)	-0.09	Nebraska	175.43 <sup>b</sup>	1.36	(37)	-3.25
Arizona	156.15 <sup>b</sup>	0.70	(48)	-13.88 <sup>c</sup>	Nevada	186.14 <sup>b</sup>	1.30	(21)	2.66
Arkansas	201.70 <sup>b</sup>	1.14	(06)	11.24 <sup>c</sup>	New Hampshire	184.17	1.64	(24)	1.57
California	165.13 <sup>b</sup>	0.32	(46)	-8.93 <sup>c</sup>	New Jersey	182.57	0.63	(28)	0.69
Colorado	156.09 <sup>b</sup>	0.88	(49)	-13.91 <sup>c</sup>	New Mexico	160.84 <sup>b</sup>	1.28	(47)	-11.29 <sup>c</sup>
Connecticut	176.91 <sup>b</sup>	0.96	(35)	-2.43	New York	169.64 <sup>b</sup>	0.40	(44)	-6.44 <sup>c</sup>
Delaware	196.62 <sup>b</sup>	2.06	(11)	8.44 <sup>c</sup>	North Carolina	189.55 <sup>b</sup>	0.65	(17)	4.54
D.C.	198.33 <sup>b</sup>	2.63	(08)	9.38 <sup>c</sup>	North Dakota	172.97 <sup>b</sup>	2.17	(39)	-4.60
Florida	172.49 <sup>b</sup>	0.39	(40)	-4.87	Ohio	197.23 <sup>b</sup>	0.56	(09)	8.77 <sup>c</sup>
Georgia	183.14	0.69	(25)	1.00	Oklahoma	195.87 <sup>b</sup>	1.01	(13)	8.03 <sup>c</sup>
Hawaii	149.15 <sup>b</sup>	1.44	(50)	-17.74 <sup>c</sup>	Oregon	183.02	0.96	(26)	0.94
Idaho	167.84 <sup>b</sup>	1.55	(45)	-7.43 <sup>c</sup>	Pennsylvania	190.25 <sup>b</sup>	0.50	(16)	4.92
Illinois	189.29 <sup>b</sup>	0.55	(19)	4.40	Rhode Island	184.95	1.75	(23)	2.00
Indiana	197.18 <sup>b</sup>	0.78	(10)	8.75 <sup>c</sup>	South Carolina	191.08 <sup>b</sup>	0.92	(15)	5.38 <sup>c</sup>
Iowa	180.52	1.02	(32)	-0.44	South Dakota	172.37 <sup>b</sup>	1.96	(41)	-4.93
Kansas	180.75	1.11	(31)	-0.32	Tennessee	202.83 <sup>b</sup>	0.80	(05)	11.86 <sup>c</sup>
Kentucky	213.58 <sup>b</sup>	0.99	(01)	17.79 <sup>c</sup>	Texas	174.75 <sup>b</sup>	0.42	(38)	-3.62
Louisiana	208.41 <sup>b</sup>	0.98	(02)	14.94 <sup>c</sup>	Utah	131.78 <sup>b</sup>	1.18	(51)	-27.32 <sup>c</sup>
Maine	195.96 <sup>b</sup>	1.58	(12)	8.08 <sup>c</sup>	Vermont	178.44	2.28	(34)	-1.59
Maryland	186.82 <sup>b</sup>	0.83	(20)	3.03	Virginia	185.65 <sup>b</sup>	0.71	(22)	2.39
Massachusetts	183.01	0.72	(27)	0.93	Washington	178.59 <sup>b</sup>	0.76	(33)	-1.51
Michigan	189.30 <sup>b</sup>	0.60	(18)	4.40	West Virginia	207.83 <sup>b</sup>	1.37	(03)	14.62 <sup>c</sup>
Minnesota	171.47 <sup>b</sup>	0.81	(42)	-5.43 <sup>c</sup>	Wisconsin	181.48	0.78	(29)	0.09
Mississippi	206.82 <sup>b</sup>	1.19	(04)	14.06 <sup>c</sup>	Wyoming	171.02 <sup>b</sup>	2.57	(43)	-5.68 <sup>c</sup>
Missouri	194.46 <sup>b</sup>	0.78	(14)	7.25 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.