


Table 2.19  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	223.04	0.19		
High Five States				
Mississippi	276.12 <sup>b</sup>	2.18	(01)	23.80 <sup>c</sup>
Kentucky	271.17 <sup>b</sup>	1.75	(02)	21.58 <sup>c</sup>
Louisiana	268.14 <sup>b</sup>	1.76	(03)	20.22 <sup>c</sup>
Alabama	261.97 <sup>b</sup>	1.63	(04)	17.45 <sup>c</sup>
Tennessee	261.08 <sup>b</sup>	1.44	(05)	17.05 <sup>c</sup>
Low Five States				
New Mexico	193.02 <sup>b</sup>	2.13	(47)	-13.46 <sup>c</sup>
Colorado	187.26 <sup>b</sup>	1.51	(48)	-16.04 <sup>c</sup>
Arizona	186.67 <sup>b</sup>	1.15	(49)	-16.31 <sup>c</sup>
Hawaii	186.22 <sup>b</sup>	2.42	(50)	-16.51 <sup>c</sup>
Utah	158.27 <sup>b</sup>	1.97	(51)	-29.04 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	261.97 <sup>b</sup>	1.63	(04)	17.45 <sup>c</sup>	Montana	208.07 <sup>b</sup>	2.98	(42)	-6.71 <sup>c</sup>
Alaska	212.35	5.20	(38)	-4.79	Nebraska	217.13	2.33	(32)	-2.65
Arizona	186.67 <sup>b</sup>	1.15	(49)	-16.31 <sup>c</sup>	Nevada	214.66 <sup>b</sup>	2.08	(34)	-3.76
Arkansas	254.87 <sup>b</sup>	1.97	(08)	14.27 <sup>c</sup>	New Hampshire	223.41	2.82	(27)	0.17
California	197.45 <sup>b</sup>	0.54	(46)	-11.48 <sup>c</sup>	New Jersey	218.49 <sup>b</sup>	1.08	(29)	-2.04
Colorado	187.26 <sup>b</sup>	1.51	(48)	-16.04 <sup>c</sup>	New Mexico	193.02 <sup>b</sup>	2.13	(47)	-13.46 <sup>c</sup>
Connecticut	216.40 <sup>b</sup>	1.66	(33)	-2.98	New York	204.58 <sup>b</sup>	0.70	(43)	-8.28 <sup>c</sup>
Delaware	238.46 <sup>b</sup>	3.51	(16)	6.91 <sup>c</sup>	North Carolina	241.37 <sup>b</sup>	1.17	(15)	8.22 <sup>c</sup>
D.C.	260.44 <sup>b</sup>	4.89	(06)	16.77 <sup>c</sup>	North Dakota	212.84	3.63	(37)	-4.58
Florida	209.42 <sup>b</sup>	0.64	(40)	-6.11 <sup>c</sup>	Ohio	246.49 <sup>b</sup>	0.99	(10)	10.51 <sup>c</sup>
Georgia	237.08 <sup>b</sup>	1.28	(17)	6.29 <sup>c</sup>	Oklahoma	245.45 <sup>b</sup>	1.75	(12)	10.04 <sup>c</sup>
Hawaii	186.22 <sup>b</sup>	2.42	(50)	-16.51 <sup>c</sup>	Oregon	217.68	1.60	(31)	-2.41
Idaho	197.85 <sup>b</sup>	2.54	(45)	-11.29 <sup>c</sup>	Pennsylvania	235.61 <sup>b</sup>	0.87	(18)	5.63 <sup>c</sup>
Illinois	233.25 <sup>b</sup>	0.96	(20)	4.58	Rhode Island	234.39 <sup>b</sup>	3.10	(19)	5.09 <sup>c</sup>
Indiana	247.29 <sup>b</sup>	1.38	(09)	10.87 <sup>c</sup>	South Carolina	245.71 <sup>b</sup>	1.64	(11)	10.16 <sup>c</sup>
Iowa	224.69	1.75	(25)	0.74	South Dakota	214.17	3.30	(36)	-3.98
Kansas	224.66	1.91	(26)	0.72	Tennessee	261.08 <sup>b</sup>	1.44	(05)	17.05 <sup>c</sup>
Kentucky	271.17 <sup>b</sup>	1.75	(02)	21.58 <sup>c</sup>	Texas	217.82 <sup>b</sup>	0.74	(30)	-2.34
Louisiana	268.14 <sup>b</sup>	1.76	(03)	20.22 <sup>c</sup>	Utah	158.27 <sup>b</sup>	1.97	(51)	-29.04 <sup>c</sup>
Maine	243.44 <sup>b</sup>	2.74	(13)	9.15 <sup>c</sup>	Vermont	214.22	3.89	(35)	-3.95
Maryland	229.69 <sup>b</sup>	1.47	(23)	2.98	Virginia	232.72 <sup>b</sup>	1.27	(21)	4.34
Massachusetts	227.26	1.26	(24)	1.89	Washington	211.94 <sup>b</sup>	1.27	(39)	-4.98
Michigan	231.08 <sup>b</sup>	1.04	(22)	3.60	West Virginia	257.12 <sup>b</sup>	2.37	(07)	15.28 <sup>c</sup>
Minnesota	208.79 <sup>b</sup>	1.38	(41)	-6.39 <sup>c</sup>	Wisconsin	222.78	1.34	(28)	-0.12
Mississippi	276.12 <sup>b</sup>	2.18	(01)	23.80 <sup>c</sup>	Wyoming	199.36 <sup>b</sup>	4.22	(44)	-10.62 <sup>c</sup>
Missouri	242.03 <sup>b</sup>	1.36	(14)	8.51 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.