

Table 5.4
Cancer of the Cervix Uteri (Invasive)

Delay-adjusted SEER Incidence^a Rates by Year, Race and Age

Year of Diagnosis:	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
1975	14.79	9.66	28.22	13.25	8.85	24.77	33.10	17.55	73.82
1976	14.26	9.50	26.74	12.87	8.72	23.74	29.25	15.84	64.38
1977	13.03	8.44	25.05	11.70	7.56	22.53	28.30	15.90	60.78
1978	12.54	8.31	23.63	11.36	7.73	20.87	24.14	12.35	55.00
1979	12.76	8.68	23.44	10.97	7.55	19.94	28.89	16.10	62.38
1980	12.23	8.11	23.03	10.98	7.30	20.60	23.43	13.50	49.44
1981	10.76	7.34	19.70	9.54	6.59	17.25	24.00	13.61	51.22
1982	10.62	7.12	19.78	9.40	6.68	16.51	21.82	10.15	52.38
1983	10.51	7.31	18.88	9.58	6.73	17.05	18.84	11.89	37.04
1984	11.04	8.00	19.00	10.00	7.40	16.80	21.09	12.60	43.34
1985	10.22	7.06	18.48	9.15	6.74	15.46	19.00	9.52	43.82
1986	10.83	7.87	18.57	9.73	7.32	16.04	18.40	10.73	38.49
1987	10.01	7.11	17.63	8.99	6.87	14.55	18.76	8.71	45.07
1988	10.60	7.88	17.71	9.54	7.46	14.99	19.11	10.46	41.77
1989	10.71	7.92	18.00	10.08	7.90	15.77	15.71	7.97	35.97
1990	10.65	8.05	17.48	9.96	7.74	15.77	16.75	9.67	35.29
1991	10.06	7.06	17.92	9.27	6.81	15.71	16.36	9.23	35.02
1992	9.95	7.47	16.43	9.43	7.21	15.25	14.20	10.06	25.04
1993	9.65	7.05	16.46	9.19	7.03	14.85	13.77	8.89	26.55
1994	9.47	6.90	16.21	8.74	6.82	13.77	14.06	6.96	32.67
1995	8.91	6.83	14.35	8.01	6.57	11.76	13.63	7.91	28.60
1996	9.65	7.40	15.55	8.78	7.17	12.99	14.49	8.29	30.73
1997	9.24	7.12	14.80	8.40	6.91	12.30	13.61	8.05	28.18
1998	9.12	7.09	14.42	8.44	6.86	12.58	13.15	9.00	24.00
1999	8.28	6.55	12.81	7.84	6.62	11.04	13.43	8.06	27.47
2000	7.74	5.89	12.60	7.47	5.96	11.44	10.27	6.43	20.32
2001	7.99	5.90	13.47	7.42	5.82	11.63	11.33	5.04	27.79
2002	7.45	5.70	12.05	7.04	5.71	10.53	10.50	6.63	20.62
2003	7.34	5.75	11.48	6.88	5.66	10.05	10.82	6.11	23.16
2004	7.16	5.51	11.49	6.88	5.65	10.11	10.63	6.63	21.11
2005	6.83	5.44	10.47	6.58	5.48	9.47	8.70	5.25	17.75
2006	6.99	5.59	10.67	6.85	5.72	9.84	8.84	5.11	18.59
2007	6.68	5.22	10.51	6.49	5.53	8.99	8.66	4.48	19.61
2008	6.78	5.39	10.44	6.48	5.36	9.41	9.51	5.83	19.16

Additional information on the model used to delay-adjust SEER Incidence rates can be found at [\(http://surveillance.cancer.gov/delay/\)](http://surveillance.cancer.gov/delay/).

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.