

Table 5.7
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2004-2008</u>						
Age at Diagnosis/Death:						
All ages	8.1	8.0	10.0	2.4	2.2	4.3
Under 65	7.5	7.5	8.2	1.9	1.7	2.9
65 and over	12.3	11.0	22.2	6.2	5.4	14.0
All ages (IARC world std) ^c	6.4	6.3	7.6	1.8	1.6	3.1
<u>Age-Specific Rates, 2004-2008</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	0.1	-	-	-	-	-
20-24	1.5	1.5	2.0	0.1	0.1	0.2
25-29	5.0	5.5	4.3	0.5	0.5	0.8
30-34	10.8	11.7	9.5	1.3	1.3	1.9
35-39	14.1	15.1	11.4	2.4	2.3	3.2
40-44	15.0	15.0	16.5	3.4	3.1	5.5
45-49	14.0	13.6	15.5	4.0	3.7	6.4
50-54	13.4	12.8	16.6	4.5	4.2	7.1
55-59	12.9	12.1	18.6	5.0	4.7	7.7
60-64	13.2	11.9	19.6	5.3	4.8	10.0
65-69	14.4	13.1	23.3	5.9	5.2	12.5
70-74	13.1	11.9	19.9	5.8	5.1	11.7
75-79	11.5	9.9	23.0	5.9	5.1	13.4
80-84	10.5	9.0	20.5	6.9	6.1	15.1
85+	9.7	8.4	25.2	7.5	6.3	22.2

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.