


Table 5.14
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.42	0.02		
High Five States				
Mississippi	4.03 ^b	0.23	(01)	66.95 ^c
Louisiana	3.35 ^b	0.17	(02)	38.55 ^c
Arkansas	3.24 ^b	0.21	(03)	34.34 ^c
D.C.	3.19	0.45	(04)	32.00 ^c
Alabama	3.07 ^b	0.15	(05)	27.06 ^c
Low Five States				
Rhode Island	1.61	0.23	(47)	-33.41 ^c
Minnesota	1.57 ^b	0.11	(48)	-34.83 ^c
Vermont	1.48	0.28	(49)	-38.65 ^c
Massachusetts	1.39 ^b	0.09	(50)	-42.60 ^c
Utah	1.31 ^b	0.16	(51)	-45.77 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.07 ^b	0.15	(05)	27.06 ^c	Montana	1.87	0.27	(39)	-22.41 ^c
Alaska	2.21	0.41	(27)	-8.35	Nebraska	1.80	0.20	(43)	-25.37 ^c
Arizona	1.98 ^b	0.11	(38)	-18.04 ^c	Nevada	2.71	0.21	(12)	12.16
Arkansas	3.24 ^b	0.21	(03)	34.34 ^c	New Hampshire	2.20	0.24	(28)	-8.96
California	2.36	0.05	(23)	-2.23	New Jersey	2.47	0.10	(19)	2.14
Colorado	1.84 ^b	0.12	(40)	-23.80 ^c	New Mexico	2.64	0.23	(15)	9.17
Connecticut	1.67 ^b	0.13	(45)	-30.66 ^c	New York	2.43	0.07	(21)	0.68
Delaware	2.24	0.31	(26)	-7.24	North Carolina	2.37	0.10	(22)	-1.86
D.C.	3.19	0.45	(04)	32.00 ^c	North Dakota	1.61	0.31	(46)	-33.30 ^c
Florida	2.66	0.07	(14)	10.25	Ohio	2.49	0.09	(18)	3.23
Georgia	2.73	0.11	(10)	12.87	Oklahoma	2.89	0.18	(07)	19.48 ^c
Hawaii	1.99	0.24	(36)	-17.44 ^c	Oregon	1.83 ^b	0.14	(41)	-24.34 ^c
Idaho	2.04	0.24	(33)	-15.35 ^c	Pennsylvania	2.26	0.08	(25)	-6.50
Illinois	2.70	0.09	(13)	11.74	Rhode Island	1.61	0.23	(47)	-33.41 ^c
Indiana	2.46	0.12	(20)	1.91	South Carolina	2.72	0.15	(11)	12.42
Iowa	2.11	0.16	(31)	-12.53	South Dakota	1.70	0.29	(44)	-29.42 ^c
Kansas	2.14	0.18	(30)	-11.43	Tennessee	2.86	0.13	(09)	18.32 ^c
Kentucky	2.64	0.15	(16)	9.12	Texas	3.05 ^b	0.07	(06)	26.43 ^c
Louisiana	3.35 ^b	0.17	(02)	38.55 ^c	Utah	1.31 ^b	0.16	(51)	-45.77 ^c
Maine	1.98	0.22	(37)	-18.01 ^c	Vermont	1.48	0.28	(49)	-38.65 ^c
Maryland	2.31	0.12	(24)	-4.30	Virginia	2.14	0.10	(29)	-11.21
Massachusetts	1.39 ^b	0.09	(50)	-42.60 ^c	Washington	2.00 ^b	0.11	(35)	-17.19 ^c
Michigan	2.03 ^b	0.09	(34)	-15.96 ^c	West Virginia	2.88	0.24	(08)	19.08 ^c
Minnesota	1.57 ^b	0.11	(48)	-34.83 ^c	Wisconsin	1.82 ^b	0.11	(42)	-24.60 ^c
Mississippi	4.03 ^b	0.23	(01)	66.95 ^c	Wyoming	2.11	0.41	(32)	-12.73
Missouri	2.62	0.13	(17)	8.51					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.