

Table 5.15  
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2008  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <33 <sup>e</sup>	>=33 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	38,079	36,825	35,079	30,973	24,009	20,597	128,098	196,451	47,433	243,884
White <sup>b</sup>	30,023	29,234	28,677	25,738	19,903	16,898	103,167	159,918	39,949	199,867
Black <sup>b</sup>	5,465	4,995	3,914	3,481	2,981	2,717	16,157	24,517	+	+
Asian/Pacific Islander <sup>c</sup>	1,988	1,723	1,677	+	+	+	6,128	+	+	+
Hispanic <sup>d</sup>	7,993	7,279	5,946	+	+	+	24,497	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2008, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0794%	-	0.0002%	0.0089%	0.0806%	0.1584%	0.1673%	0.1445%	0.1285%	0.0774%	0.0772%
White <sup>c</sup>	0.0807%	-	0.0003%	0.0099%	0.0919%	0.1703%	0.1636%	0.1335%	0.1153%	0.0613%	0.0784%
Black <sup>c</sup>	0.0696%	-	-	0.0075%	0.0626%	0.1262%	0.1675%	0.1652%	0.1486%	0.1470%	0.0742%
Asian/Pacific Islander <sup>c</sup>	0.0764%	-	-	0.0022%	0.0341%	0.1098%	0.1790%	0.1912%	0.1883%	0.1530%	0.0724%
Hispanic <sup>d</sup>	-	-	-	-	-	-	-	-	-	-	-

<sup>a</sup> US 2008 cancer prevalence counts are based on 2008 cancer prevalence proportions from the SEER registries and 1/1/2008 US population estimates based on the average of 2007 and 2008 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 33 years for 1975-2008 SEER 9 data; 18 years for 1990-2008 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 33 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <33 and >=33. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.