

Table 20.20
Cancer of the Oral Cavity and Pharynx by Subsite (Invasive)

SEER Incidence
Age-Adjusted Rates and Trends^a by Race and Sex

	Site									
	Oral Cavity & Pharynx	Lip	Tongue	Floor of Mouth	Gum & Other Mouth	Oropharynx & Tonsil	Hypophar.	Salivary Gland	Nasophar.	Other Oral Cavity & Pharynx
<u>2004-2008, Rate per 100,000 persons^b</u>										
All Races										
Male and Female	10.6	0.7	3.0	0.6	1.5	2.0	0.7	1.3	0.7	0.2
Male	15.7	1.2	4.5	0.9	1.8	3.3	1.1	1.7	1.0	0.3
Female	6.2	0.3	1.7	0.3	1.2	0.8	0.2	1.0	0.4	0.1
White										
Male and Female	10.8	0.8	3.2	0.6	1.5	2.1	0.6	1.3	0.4	0.2
Male	16.1	1.4	4.7	0.9	1.8	3.5	1.1	1.8	0.6	0.3
Female	6.2	0.4	1.8	0.3	1.2	0.8	0.3	1.0	0.3	0.1
Black										
Male and Female	9.9	0.1	2.3	0.6	1.5	2.3	1.1	1.0	0.7	0.3
Male	15.6	0.1	3.7	1.0	1.9	4.2	2.1	1.0	1.2	0.5
Female	5.6	-	1.2	0.3	1.3	0.8	0.3	1.0	0.3	0.2
<u>Trend 1999-2008^c, APC(%)</u>										
All Races										
Male and Female	-0.4	-6.3*	1.7*	-4.2*	-1.9*	2.5*	-3.4*	0.7	0.5	-5.3*
Male	-0.4	-6.8*	2.1*	-4.6*	-2.3*	2.8*	-3.0*	0.6	0.3	-5.0*
Female	-0.6	-4.6*	1.0	-3.7*	-1.4	0.6	-4.5*	0.6	0.7	-5.9*
White										
Male and Female	-0.1	-6.0*	2.3*	-3.6*	-1.9*	3.1*	-3.0*	0.5	1.1	-5.6*
Male	-0.1	-6.6*	2.4*	-4.1*	-2.0*	3.5*	-2.8*	0.6	0.6	-4.6*
Female	-0.5	-4.2*	2.0*	-3.3*	-1.8*	0.8	-4.0	0.0	2.0	-
Black										
Male and Female	-2.8*	-	-3.0*	-8.7*	-4.1	0.2	-2.5	-0.7	-1.6	-
Male	-3.3*	-	-2.9*	-10.3*	-6.7*	0.2	-1.5	-4.3*	-1.8	-
Female	-1.4	-	-2.8	-	-0.7	0.0	-	2.3	-	-

The APC is the Annual Percent Change over the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^c Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

* The APC is significantly different from zero ($p < .05$).