

Table 21.7  
Cancer of the Ovary (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2004-2008</u>						
Age at Diagnosis/Death:						
All ages	12.8	13.5	10.0	8.4	8.8	7.0
Under 65	7.9	8.2	5.7	3.4	3.5	2.9
65 and over	47.1	49.8	40.1	43.3	44.9	35.6
All ages (IARC world std) <sup>c</sup>	9.1	9.6	7.0	5.3	5.6	4.5
<u>Age-Specific Rates, 2004-2008</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	0.3	0.2	-	-	-	-
10-14	0.8	0.9	-	-	-	-
15-19	1.3	1.3	1.0	0.1	0.0	-
20-24	1.7	1.7	1.1	0.2	0.2	-
25-29	2.1	2.1	1.5	0.3	0.3	0.4
30-34	3.2	3.2	2.2	0.6	0.6	0.6
35-39	4.7	4.7	4.3	1.1	1.2	1.0
40-44	8.8	9.2	5.8	2.6	2.6	2.3
45-49	14.8	15.2	10.6	5.3	5.5	4.5
50-54	22.0	22.8	15.1	9.7	10.0	8.3
55-59	27.4	29.0	18.6	15.0	15.7	12.1
60-64	34.8	36.9	26.8	22.3	23.3	18.3
65-69	39.1	41.9	32.7	31.1	32.3	27.5
70-74	46.6	50.1	37.1	39.4	41.2	32.9
75-79	49.7	52.8	40.9	47.4	49.2	38.2
80-84	55.5	57.8	47.3	56.0	58.1	41.7
85+	51.1	52.4	52.9	56.3	57.6	47.7

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473. SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.
- <sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.
- <sup>c</sup> Rates are per 100,000 and are age-adjusted to the IARC world standard population.
- Statistic not shown. Rate based on less than 16 cases for the time interval.