

Table 25.5
Cancer of the Testis (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

Year of Diagnosis:	All Races, Males			White Males			Black Males		
	All	<50	50+	All	<50	50+	All	<50	50+
1975	3.73	4.23	2.41	4.16	4.74	2.65	-	-	-
1976	3.44	4.18	1.51	3.77	4.56	1.72	-	-	-
1977	4.30	5.09	2.24	4.88	5.82	2.41	-	-	-
1978	3.58	4.25	1.84	3.96	4.66	2.11	-	-	-
1979	3.88	4.54	2.15	4.33	5.10	2.31	-	-	-
1980	4.35	5.28	1.89	4.91	5.98	2.11	-	-	-
1981	4.22	5.12	1.85	4.77	5.79	2.07	-	-	-
1982	4.40	5.22	2.25	5.00	5.98	2.44	-	-	-
1983	4.59	5.50	2.21	5.23	6.30	2.43	-	-	-
1984	4.39	5.36	1.86	5.05	6.18	2.08	-	-	-
1985	4.47	5.47	1.83	5.14	6.32	2.05	-	-	-
1986	4.82	5.83	2.17	5.57	6.79	2.39	-	-	-
1987	5.05	6.26	1.86	5.68	7.11	1.94	-	-	-
1988	4.63	5.83	1.46	5.45	6.88	1.72	-	-	-
1989	5.48	6.71	2.24	6.32	7.79	2.46	-	-	-
1990	5.09	6.27	2.02	5.99	7.40	2.30	-	-	-
1991	5.10	6.24	2.12	5.75	7.06	2.31	-	-	-
1992	5.15	6.44	1.76	6.08	7.65	1.97	-	-	-
1993	5.07	6.38	1.65	5.98	7.53	1.93	-	-	-
1994	5.47	6.75	2.12	6.28	7.79	2.30	1.23	1.54	-
1995	4.56	5.58	1.88	5.25	6.45	2.11	1.46	1.87	-
1996	5.25	6.60	1.73	6.20	7.85	1.89	-	-	-
1997	5.39	6.52	2.43	6.33	7.74	2.64	1.27	-	-
1998	5.60	7.12	1.62	6.59	8.42	1.79	1.52	1.97	-
1999	5.45	6.83	1.81	6.33	7.97	2.02	1.63	2.02	-
2000	5.74	7.17	1.99	6.74	8.43	2.33	1.52	1.86	-
2001	5.49	6.81	2.04	6.48	8.10	2.24	1.67	1.81	-
2002	5.73	7.18	1.96	6.86	8.60	2.31	1.37	1.78	-
2003	5.28	6.61	1.81	6.37	8.06	1.94	1.13	-	-
2004	5.87	7.24	2.27	7.20	8.95	2.62	1.03	-	-
2005	5.91	7.36	2.13	7.09	8.93	2.29	0.99	1.28	-
2006	5.52	6.83	2.08	6.58	8.18	2.38	1.57	1.64	-
2007	5.86	7.23	2.28	6.97	8.65	2.57	1.37	1.42	-
2008	5.89	7.12	2.68	7.05	8.64	2.90	1.35	1.60	-
1975-2008	5.07	6.23	2.02	5.87	7.25	2.25	1.13	1.31	0.65

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.