

Table 25.7
Cancer of the Testis (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2004-2008</u>						
Age at Diagnosis/Death:						
All ages	5.5	6.5	1.3	0.2	0.3	0.1
Under 65	6.1	7.2	1.4	0.2	0.3	0.1
65 and over	1.1	1.2	-	0.3	0.3	-
All ages (IARC world std) ^c	5.2	6.1	1.2	0.2	0.2	0.1
<u>Age-Specific Rates, 2004-2008</u>						
Age at Diagnosis/Death:						
<1	1.0	0.9	-	-	-	-
1-4	0.4	0.4	-	-	-	-
5-9	-	-	-	-	-	-
10-14	0.3	0.3	-	-	-	-
15-19	3.7	4.5	-	0.1	0.1	-
20-24	10.1	12.1	1.4	0.3	0.4	0.2
25-29	13.5	16.2	2.2	0.4	0.4	0.2
30-34	13.1	15.7	3.4	0.4	0.4	-
35-39	10.9	12.9	2.2	0.4	0.4	0.3
40-44	9.0	10.5	2.4	0.4	0.4	0.2
45-49	6.2	7.2	1.8	0.3	0.3	-
50-54	4.6	5.2	1.7	0.3	0.3	-
55-59	2.8	3.2	-	0.3	0.3	-
60-64	1.7	1.8	-	0.2	0.2	-
65-69	1.4	1.6	-	0.2	0.2	-
70-74	1.0	1.1	-	0.2	0.2	-
75-79	0.9	1.0	-	0.2	0.2	-
80-84	1.0	1.1	-	0.4	0.4	-
85+	1.3	1.4	-	0.7	0.7	-

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.