

Table 2.9  
All Cancer Sites (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	69.9	71.4	73.9	75.6	77.2	77.8	77.2	77.0	77.1	77.5	78.2	78.5	79.1	79.5	80.0	80.2	79.9	80.5	80.7	81.3	81.6	81.8
2-year	60.1	61.6	64.8	67.1	69.0	70.2	69.7	69.3	69.6	70.4	71.1	71.7	72.5	73.0	73.7	74.1	73.6	74.2	74.4	75.4	75.7	
3-year	55.0	56.4	60.0	62.7	64.8	66.4	65.8	65.5	65.9	66.8	67.6	68.2	69.1	69.7	70.4	71.0	70.4	71.0	71.2	72.2		
4-year	51.6	53.0	56.9	59.8	62.2	63.9	63.3	62.9	63.4	64.4	65.2	65.9	66.9	67.6	68.4	68.9	68.3	68.8	69.2			
5-year	49.1	50.4	54.6	57.9	60.2	62.0	61.4	61.1	61.7	62.6	63.5	64.1	65.3	66.1	66.9	67.4	66.8	67.3				
6-year	47.2	48.3	52.8	56.2	58.6	60.5	60.1	59.6	60.3	61.3	62.1	62.8	64.0	64.9	65.8	66.2	65.5					
7-year	45.5	46.6	51.2	54.7	57.2	59.1	58.7	58.4	59.1	60.2	61.1	61.8	63.0	63.8	64.7	65.2						
8-year	44.2	45.2	49.9	53.5	56.0	58.0	57.6	57.4	58.2	59.3	60.2	60.9	62.0	62.8	63.9							
9-year	43.0	44.0	48.8	52.4	55.0	57.1	56.6	56.5	57.3	58.4	59.3	60.0	61.3	62.0								
10-year	41.9	43.0	47.8	51.4	54.2	56.2	55.7	55.6	56.5	57.5	58.6	59.2	60.6									
11-year	41.0	42.0	46.9	50.4	53.4	55.4	54.8	54.8	55.7	56.9	57.8	58.5										
12-year	40.1	41.2	46.0	49.7	52.7	54.7	54.1	54.0	54.9	56.2	57.2											
13-year	39.5	40.4	45.2	49.0	52.0	54.1	53.4	53.4	54.1	55.5												
14-year	38.8	39.7	44.5	48.3	51.4	53.5	52.8	52.6	53.4													
15-year	38.1	39.1	43.9	47.7	50.6	52.9	52.0	52.1														
16-year	37.5	38.3	43.3	47.1	50.0	52.3	51.4															
17-year	37.0	37.7	42.7	46.4	49.4	51.6																
18-year	36.5	37.2	42.2	45.8	48.8																	
19-year	36.1	36.7	41.8	45.3																		
20-year	35.6	36.2	41.3																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 2.10  
All Cancer Sites (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	64.5	67.4	70.6	73.7	76.3	77.5	76.6	76.1	75.9	76.8	77.4	77.3	78.5	79.1	79.6	79.7	79.4	80.2	80.3	81.2	81.7	81.5
2-year	53.8	56.8	60.8	64.6	68.0	69.9	69.1	68.1	68.3	69.7	70.4	70.6	71.7	72.7	73.3	73.7	73.2	74.0	74.2	75.5	75.9	
3-year	48.4	51.5	55.7	60.2	64.1	66.5	65.5	64.6	64.8	66.3	67.1	67.3	68.6	69.6	70.5	71.0	70.3	71.2	71.2	72.5		
4-year	44.9	47.9	52.4	57.3	61.4	64.2	63.3	62.1	62.5	64.0	65.0	65.1	66.7	67.7	68.6	69.2	68.5	69.2	69.5			
5-year	42.4	45.2	50.1	55.3	59.5	62.5	61.5	60.5	61.0	62.4	63.3	63.6	65.3	66.4	67.5	68.1	67.3	68.1				
6-year	40.4	43.2	48.2	53.6	57.9	61.1	60.3	59.1	59.6	61.2	62.0	62.4	64.2	65.3	66.5	67.1	66.1					
7-year	38.7	41.4	46.5	51.9	56.5	59.8	59.0	57.9	58.6	60.2	61.2	61.4	63.3	64.3	65.6	66.4						
8-year	37.2	40.0	45.2	50.7	55.2	58.5	58.1	56.8	57.7	59.4	60.4	60.6	62.5	63.4	65.0							
9-year	35.9	38.8	44.0	49.6	54.3	57.6	57.0	55.9	56.9	58.6	59.6	59.8	61.7	62.8								
10-year	34.7	37.7	42.9	48.5	53.4	56.7	56.1	55.1	56.1	57.7	59.0	59.1	61.2									
11-year	33.6	36.7	42.0	47.5	52.5	56.0	55.2	54.3	55.4	57.0	58.3	58.4										
12-year	32.7	35.8	41.0	46.7	51.9	55.3	54.5	53.5	54.5	56.3	57.7											
13-year	32.0	35.0	40.2	46.1	51.2	54.6	53.7	52.9	53.7	55.8												
14-year	31.2	34.2	39.4	45.4	50.6	54.0	53.0	52.0	52.8													
15-year	30.5	33.5	38.7	44.6	49.8	53.3	52.2	51.5														
16-year	29.8	32.7	38.0	43.9	49.1	52.7	51.8															
17-year	29.2	32.1	37.5	43.3	48.5	52.1																
18-year	28.6	31.5	36.9	42.7	48.1																	
19-year	28.1	31.0	36.4	42.3																		
20-year	27.7	30.4	36.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 2.11  
All Cancer Sites (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	75.2	75.5	77.3	77.7	78.3	78.2	77.9	77.9	78.3	78.3	79.0	79.7	79.8	79.9	80.4	80.8	80.5	80.8	81.1	81.3	81.6	82.1
2-year	66.4	66.5	68.9	69.8	70.2	70.5	70.4	70.6	71.0	71.2	71.9	72.8	73.2	73.2	74.1	74.6	74.1	74.4	74.7	75.2	75.4	
3-year	61.5	61.4	64.4	65.4	65.8	66.3	66.2	66.6	67.0	67.4	68.1	69.1	69.6	69.8	70.3	71.0	70.5	70.9	71.2	71.8		
4-year	58.1	58.1	61.5	62.6	63.1	63.5	63.4	63.8	64.4	64.9	65.5	66.7	67.2	67.4	68.1	68.5	68.1	68.3	68.9			
5-year	55.7	55.5	59.2	60.7	61.0	61.3	61.4	61.9	62.5	62.9	63.7	64.7	65.3	65.8	66.3	66.7	66.2	66.5				
6-year	53.8	53.5	57.4	59.0	59.4	59.8	59.8	60.3	61.0	61.4	62.2	63.3	63.8	64.5	65.0	65.3	64.9					
7-year	52.2	51.8	55.9	57.6	58.1	58.3	58.3	58.9	59.7	60.4	61.0	62.2	62.7	63.2	63.7	64.0						
8-year	50.9	50.4	54.7	56.5	56.9	57.4	57.1	58.0	58.7	59.2	60.1	61.2	61.6	62.1	62.7							
9-year	49.8	49.3	53.6	55.4	55.9	56.6	56.1	57.2	57.8	58.2	59.1	60.3	60.8	61.2								
10-year	48.8	48.2	52.6	54.4	55.0	55.7	55.3	56.2	57.0	57.4	58.2	59.3	60.0									
11-year	47.9	47.2	51.7	53.6	54.4	54.8	54.4	55.3	56.1	56.8	57.4	58.5										
12-year	47.1	46.4	50.9	52.9	53.6	54.1	53.7	54.6	55.5	56.1	56.8											
13-year	46.5	45.7	50.2	52.1	52.9	53.4	53.2	54.0	54.6	55.3												
14-year	45.8	45.0	49.5	51.4	52.2	53.0	52.5	53.3	54.0													
15-year	45.2	44.4	48.9	50.8	51.6	52.3	51.8	52.8														
16-year	44.7	43.7	48.4	50.3	51.0	51.8	51.0															
17-year	44.2	43.1	47.8	49.7	50.4	51.1																
18-year	43.7	42.6	47.4	49.0	49.7																	
19-year	43.3	42.0	47.0	48.4																		
20-year	42.8	41.6	46.4																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 3.9  
Cancer of the Brain and Other Nervous System (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	47.5	48.6	52.2	54.1	55.0	52.9	53.2	54.4	53.5	54.5	52.5	55.1	55.9	56.5	58.2	60.0	58.2	58.1	59.0	62.3	62.6	63.0
2-year	32.4	34.3	37.1	37.3	40.5	39.2	38.7	40.4	39.5	40.7	38.2	40.8	40.3	42.0	42.5	46.2	44.0	43.3	45.9	47.0	47.2	
3-year	28.0	29.5	32.8	32.3	36.7	34.6	35.0	36.4	36.1	36.7	34.1	35.5	35.8	38.1	38.8	40.0	37.9	38.0	39.5	39.9		
4-year	25.2	26.6	29.9	30.1	34.3	31.5	32.9	33.8	34.4	34.3	31.8	32.7	33.4	35.9	36.5	37.4	35.7	35.7	36.7			
5-year	23.1	24.6	28.0	28.6	32.2	29.9	31.4	32.0	33.4	32.5	30.1	30.8	31.9	34.6	34.5	35.2	33.9	34.7				
6-year	21.6	23.2	26.6	27.6	30.8	27.9	30.4	31.1	32.2	31.0	28.1	29.5	30.4	33.0	33.4	33.4	32.4					
7-year	20.5	21.9	25.4	26.3	29.8	26.7	29.0	29.4	31.2	30.1	27.3	28.3	29.7	31.4	32.1	32.6						
8-year	19.4	20.9	24.1	25.6	28.8	25.6	28.7	28.2	30.3	28.6	26.5	27.4	28.9	30.7	31.2							
9-year	18.4	19.8	23.3	24.8	28.4	24.4	27.9	27.5	29.8	27.5	25.4	26.8	27.8	29.6								
10-year	17.8	18.9	22.5	24.0	27.3	23.6	27.1	27.1	28.9	26.9	24.4	26.0	27.3									
11-year	17.1	18.1	21.9	23.4	26.9	23.0	26.6	26.0	27.9	26.0	23.5	25.5										
12-year	16.6	17.6	21.3	22.9	25.9	22.6	25.7	25.5	27.6	24.9	22.6											
13-year	16.2	17.0	20.8	22.4	25.2	22.2	25.3	25.2	26.5	24.5												
14-year	15.8	16.5	20.1	22.1	24.5	21.6	24.4	24.4	25.9													
15-year	15.3	16.0	19.5	21.5	23.8	21.2	23.6	23.7														
16-year	15.0	15.5	19.1	20.9	23.1	21.0	23.4															
17-year	14.6	15.2	18.7	20.4	22.8	20.1																
18-year	14.4	14.9	18.1	20.2	22.4																	
19-year	14.0	14.3	17.7	19.9																		
20-year	13.7	13.9	17.4																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 3.10  
Cancer of the Brain and Other Nervous System (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	47.3	49.5	52.5	57.0	56.7	54.9	54.7	55.9	55.5	56.0	55.4	55.3	55.4	55.2	56.0	59.9	58.0	59.9	58.3	64.5	62.6	63.0
2-year	31.4	33.9	36.4	38.9	40.2	39.7	40.3	40.7	40.9	39.7	39.6	39.9	38.6	39.8	38.6	46.1	42.8	44.7	44.7	46.6	46.0	
3-year	26.5	29.4	31.7	32.8	36.4	35.2	35.9	36.9	37.3	35.8	34.9	35.1	34.1	35.5	35.1	38.9	36.0	39.0	38.6	38.7		
4-year	24.0	26.0	28.5	30.5	34.2	32.4	33.7	34.7	36.0	33.4	32.7	31.6	31.2	32.9	32.0	35.7	34.3	36.4	36.1			
5-year	21.5	23.6	26.5	29.0	31.6	30.8	31.3	32.6	34.8	31.2	30.4	29.8	29.5	31.8	30.5	32.9	32.8	35.3				
6-year	20.1	22.2	25.0	28.5	30.8	28.7	30.1	31.9	33.3	29.6	28.2	27.9	28.1	30.3	29.4	30.7	31.4					
7-year	19.1	21.2	23.7	26.7	29.2	27.2	29.1	30.4	32.1	28.6	27.1	26.7	27.5	28.1	28.1	29.6						
8-year	17.9	20.2	22.6	26.4	28.1	26.2	28.8	29.4	30.8	26.9	26.5	25.9	26.6	27.4	27.1							
9-year	16.9	19.0	21.9	25.7	27.7	24.7	28.2	28.4	30.0	25.8	25.7	25.0	24.9	26.4								
10-year	16.3	18.1	21.1	24.8	26.8	23.9	27.0	27.9	29.0	25.3	24.7	24.3	24.0									
11-year	15.5	17.5	20.5	24.3	26.5	23.2	26.5	26.2	28.1	24.5	24.1	23.5										
12-year	15.0	17.0	20.0	23.9	25.6	22.9	25.5	25.9	27.9	23.4	22.8											
13-year	14.6	16.4	19.6	23.4	24.7	22.6	24.9	25.6	26.4	23.0												
14-year	14.4	15.9	19.0	23.2	23.9	22.2	23.9	24.6	25.4													
15-year	13.9	15.4	18.4	22.3	22.9	21.5	23.3	24.0														
16-year	13.8	14.8	18.0	21.9	22.3	21.2	23.3															
17-year	13.5	14.4	17.7	21.5	21.9	20.4																
18-year	13.2	14.1	17.1	21.2	21.7																	
19-year	12.8	13.7	16.6	21.0																		
20-year	12.6	13.4	16.4																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 3.11  
Cancer of the Brain and Other Nervous System (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	47.8	47.5	51.9	50.8	52.8	50.2	51.3	52.5	50.7	52.5	48.8	54.8	56.5	58.1	61.2	60.0	58.6	56.0	59.9	59.4	62.6	63.0
2-year	33.6	34.7	38.1	35.4	40.9	38.5	36.7	40.0	37.7	41.9	36.4	41.9	42.5	44.8	47.4	46.4	45.3	41.6	47.3	47.4	48.8	
3-year	29.9	29.6	34.1	31.7	37.0	33.7	33.8	35.7	34.6	37.8	33.1	35.9	38.1	41.4	43.6	41.5	40.1	36.8	40.6	41.4		
4-year	26.8	27.3	31.7	29.7	34.5	30.2	32.0	32.7	32.1	35.5	30.8	34.0	36.2	39.6	42.2	39.7	37.4	35.0	37.4			
5-year	25.1	26.0	30.0	28.0	32.9	28.6	31.4	31.4	31.6	34.1	29.6	31.9	35.0	38.1	39.5	38.1	35.1	33.9				
6-year	23.5	24.3	28.5	26.5	30.9	26.8	30.7	30.1	30.6	32.9	28.0	31.2	33.4	36.5	38.6	36.8	33.7					
7-year	22.3	22.9	27.5	25.8	30.6	25.9	28.9	28.1	30.1	32.0	27.7	30.3	32.5	35.5	37.3	36.3						
8-year	21.3	21.7	26.1	24.5	29.8	24.9	28.5	26.6	29.7	30.8	26.5	29.3	31.9	34.8	36.4							
9-year	20.3	20.9	25.1	23.7	29.2	24.0	27.5	26.4	29.5	29.7	25.0	28.9	31.6	33.8								
10-year	19.7	19.9	24.3	23.2	27.9	23.2	27.2	26.1	28.7	29.0	23.9	27.9	31.5									
11-year	19.1	18.9	23.6	22.2	27.5	22.7	26.7	25.7	27.5	27.9	22.8	27.7										
12-year	18.6	18.4	23.0	21.8	26.2	22.1	26.0	25.0	27.1	27.0	22.4											
13-year	18.2	17.7	22.3	21.2	25.8	21.6	25.6	24.5	26.5	26.5												
14-year	17.6	17.4	21.6	20.8	25.2	20.8	25.0	24.1	26.4													
15-year	17.1	16.9	20.9	20.6	24.9	20.7	24.0	23.4														
16-year	16.6	16.4	20.6	19.9	24.1	20.6	23.5															
17-year	16.1	16.1	20.0	19.1	24.0	19.6																
18-year	15.9	15.8	19.4	18.9	23.3																	
19-year	15.6	15.1	19.0	18.6																		
20-year	15.1	14.6	18.7																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 4.15  
Cancer of the Female Breast (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	94.7	95.4	96.6	97.0	97.3	97.4	97.2	97.5	97.3	97.2	97.5	97.7	97.6	97.7	97.5	97.7	98.0	97.9	97.9	98.0	97.9	97.8
2-year	89.0	90.3	92.6	93.4	93.8	94.4	94.1	94.3	94.1	93.9	94.5	95.1	95.3	95.6	95.3	95.7	95.7	95.8	95.9	95.8	96.2	
3-year	83.4	84.8	88.5	89.8	90.2	91.1	91.0	91.1	91.5	91.1	92.2	93.1	93.3	93.7	93.0	93.7	93.5	93.4	93.9	93.9		
4-year	78.7	80.4	85.3	87.0	87.7	88.2	88.3	88.7	89.0	89.1	90.1	91.2	91.5	91.8	91.1	91.9	91.7	91.5	92.0			
5-year	74.8	76.6	82.4	84.8	85.4	86.0	85.9	86.7	86.9	86.8	88.6	89.6	89.7	90.4	89.6	90.4	89.9	89.9				
6-year	71.6	73.6	80.1	82.7	83.5	84.2	84.0	85.1	85.5	85.2	87.2	88.5	88.2	89.2	88.4	88.9	88.7					
7-year	68.8	70.8	78.0	80.8	81.8	82.6	82.3	83.7	83.9	84.1	86.1	87.3	87.1	87.9	87.0	87.7						
8-year	66.6	68.5	76.5	79.5	80.2	81.8	80.7	82.9	82.8	82.6	85.2	86.0	86.0	86.6	86.1							
9-year	64.4	66.5	74.8	78.2	78.9	80.8	79.7	81.8	81.6	81.6	84.0	84.8	85.1	85.4								
10-year	62.6	64.6	73.5	77.2	77.9	79.9	78.8	80.6	80.7	80.5	82.9	83.6	84.3									
11-year	61.0	62.8	72.2	76.2	77.2	78.9	77.5	79.2	79.7	79.7	82.0	82.7										
12-year	59.4	61.6	71.0	75.3	76.1	77.8	76.9	78.1	78.7	78.7	81.2											
13-year	58.2	60.5	70.0	74.4	75.3	76.8	76.0	77.4	77.6	77.6												
14-year	57.2	59.2	69.1	73.2	74.5	76.1	75.0	76.5	76.9													
15-year	56.1	58.4	68.2	72.0	73.8	75.2	74.1	75.9														
16-year	55.2	57.3	67.4	70.9	73.1	74.3	73.0															
17-year	54.3	56.4	66.6	70.0	72.0	73.4																
18-year	53.5	55.6	65.9	69.1	71.1																	
19-year	52.6	55.0	65.2	68.4																		
20-year	51.9	54.2	64.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 4.16  
Cancer of the Female Breast (Ages <50, Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	96.8	97.6	98.0	97.6	98.5	98.4	98.0	98.5	98.0	97.8	98.1	98.4	98.7	98.5	98.4	98.6	98.9	98.5	98.9	98.9	99.0	98.7
2-year	91.0	91.5	92.8	93.1	94.0	94.7	94.3	93.7	93.8	93.8	94.8	95.2	95.2	95.7	94.9	95.9	96.1	96.0	96.0	96.7	97.1	
3-year	84.7	85.7	87.5	88.5	90.2	90.9	90.3	89.4	90.0	90.7	91.8	92.3	92.3	93.1	91.7	93.4	93.7	93.3	93.7	94.7		
4-year	80.1	81.0	83.5	85.1	86.9	86.5	87.2	86.3	86.7	88.2	88.8	89.5	90.1	90.6	89.3	90.7	91.8	91.2	92.0			
5-year	76.2	77.2	80.3	82.5	84.0	83.9	84.8	84.1	84.2	85.5	86.9	87.4	88.1	89.1	87.3	88.7	89.8	89.3				
6-year	73.4	74.3	77.9	80.2	81.8	81.8	83.0	82.1	82.3	83.3	84.8	85.5	86.1	87.4	85.2	86.7	88.0					
7-year	70.9	71.7	75.8	78.6	79.6	80.3	81.3	80.4	80.3	81.7	83.2	83.7	84.8	85.8	83.5	85.1						
8-year	69.0	69.5	74.2	76.7	78.2	78.7	79.7	79.1	78.7	80.0	81.5	82.4	83.6	83.8	82.0							
9-year	67.1	67.8	72.6	75.5	77.3	77.6	78.5	77.6	77.6	79.0	80.5	80.8	82.4	82.5								
10-year	65.6	66.3	71.4	74.4	76.4	76.8	77.3	76.6	76.3	77.6	79.1	79.7	81.5									
11-year	64.2	65.0	70.4	73.4	75.4	75.7	76.7	75.7	75.4	76.6	78.1	78.7										
12-year	62.8	64.0	69.3	72.5	74.0	75.1	76.0	74.3	74.5	75.9	77.1											
13-year	61.7	63.0	68.3	71.4	73.2	74.0	75.3	73.5	73.7	74.5												
14-year	61.0	61.9	67.6	70.8	72.5	73.4	74.2	72.6	72.8													
15-year	60.2	61.1	66.8	70.1	71.8	72.6	73.4	72.3														
16-year	59.5	60.4	66.1	69.7	70.8	72.2	72.2															
17-year	58.9	59.7	65.5	69.1	70.1	71.6																
18-year	58.3	59.1	64.9	68.6	69.4																	
19-year	57.7	58.7	64.3	67.8																		
20-year	57.0	58.1	63.8																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).



Table 4.17  
Cancer of the Female Breast (Ages 50+, Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	94.0	94.8	96.1	96.8	96.9	97.1	96.9	97.2	97.1	96.9	97.4	97.4	97.3	97.4	97.3	97.4	97.6	97.7	97.6	97.6	97.5	97.6
2-year	88.3	89.9	92.5	93.5	93.8	94.3	94.0	94.6	94.3	94.0	94.5	95.0	95.4	95.5	95.4	95.6	95.5	95.7	95.8	95.4	95.9	
3-year	83.0	84.6	88.9	90.3	90.2	91.2	91.2	91.7	92.0	91.2	92.3	93.3	93.6	93.9	93.4	93.8	93.4	93.5	93.9	93.6		
4-year	78.2	80.2	85.9	87.7	88.0	88.9	88.7	89.5	89.8	89.4	90.5	91.8	92.0	92.3	91.8	92.4	91.7	91.6	92.0			
5-year	74.3	76.4	83.2	85.6	85.9	86.7	86.4	87.7	87.9	87.3	89.2	90.5	90.3	90.9	90.4	91.1	90.0	90.2				
6-year	71.1	73.4	80.8	83.6	84.2	85.2	84.3	86.2	86.7	85.9	88.1	89.6	88.9	89.8	89.6	89.7	89.0					
7-year	68.1	70.5	78.8	81.6	82.6	83.5	82.7	84.9	85.3	85.0	87.1	88.6	88.0	88.7	88.3	88.6						
8-year	65.8	68.2	77.3	80.6	81.0	82.9	81.1	84.4	84.3	83.6	86.6	87.4	86.9	87.6	87.6							
9-year	63.5	66.1	75.6	79.3	79.5	82.0	80.2	83.4	83.1	82.6	85.3	86.4	86.1	86.6								
10-year	61.6	64.0	74.2	78.3	78.4	81.0	79.3	82.1	82.5	81.7	84.3	85.1	85.3									
11-year	59.8	62.0	72.8	77.2	77.9	80.0	77.7	80.5	81.3	80.9	83.5	84.3										
12-year	58.1	60.7	71.5	76.4	77.0	78.9	77.2	79.6	80.3	79.8	82.8											
13-year	56.9	59.6	70.5	75.5	76.1	77.9	76.3	78.8	79.1	78.8												
14-year	55.8	58.2	69.6	74.1	75.3	77.2	75.3	77.9	78.4													
15-year	54.5	57.3	68.6	72.7	74.7	76.1	74.4	77.2														
16-year	53.5	56.1	67.8	71.1	74.1	75.1	73.3															
17-year	52.3	55.0	66.9	69.9	72.7	73.9																
18-year	51.4	54.2	66.1	68.8	71.7																	
19-year	50.5	53.4	65.4	68.2																		
20-year	49.7	52.6	64.7																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 5.9  
Cancer of the Cervix Uteri (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	87.1	87.6	87.5	88.3	88.9	87.4	89.3	87.9	89.9	89.2	89.4	88.6	89.9	88.7	86.1	88.1	87.0	86.7	87.6	86.4	89.0	87.5
2-year	78.2	77.5	78.1	79.9	79.9	78.4	80.0	80.5	82.4	82.3	80.7	81.5	83.1	81.5	79.4	79.7	79.4	77.5	77.5	78.6	78.8	
3-year	73.3	72.8	73.3	75.6	74.4	74.1	75.8	76.4	78.9	78.0	76.0	76.8	78.5	76.5	75.6	74.7	75.8	73.1	72.1	75.4		
4-year	70.4	69.7	70.6	73.3	70.9	70.4	73.0	73.8	76.2	74.8	73.1	74.0	76.2	74.2	72.9	72.1	72.4	70.0	69.8			
5-year	68.4	67.7	68.6	72.1	68.7	67.8	71.0	72.0	74.9	73.7	71.8	72.5	74.5	72.2	70.4	69.4	70.2	67.1				
6-year	67.2	65.8	67.3	70.7	67.0	66.5	70.5	70.3	73.6	72.9	70.1	71.0	73.4	71.2	69.2	68.5	68.4					
7-year	66.1	64.1	66.2	69.9	66.2	65.4	69.7	68.9	72.3	71.8	69.7	69.7	72.9	70.1	67.7	68.0						
8-year	65.1	62.8	65.1	68.9	64.9	64.7	69.1	68.0	71.6	70.7	69.1	69.3	72.5	69.8	67.3							
9-year	64.3	62.1	64.5	67.0	64.2	64.1	68.4	67.5	70.6	69.4	67.6	68.6	72.3	68.8								
10-year	63.2	61.4	63.7	65.6	63.9	63.7	68.0	66.7	69.8	68.7	66.8	67.7	70.8									
11-year	62.4	60.8	63.1	64.9	63.7	62.6	67.8	65.8	69.5	68.1	65.8	66.9										
12-year	61.7	60.0	62.3	64.6	63.3	61.5	67.0	65.2	68.3	66.8	65.6											
13-year	61.1	59.4	61.8	64.1	62.5	61.3	66.6	64.4	66.8	66.0												
14-year	60.5	58.6	61.2	63.3	61.9	60.7	65.8	64.0	66.0													
15-year	59.8	57.8	60.4	62.9	61.3	59.9	65.2	63.5														
16-year	59.2	57.2	59.7	62.9	60.0	59.3	64.4															
17-year	58.6	56.3	58.9	62.4	60.0	58.6																
18-year	58.1	55.4	58.7	61.4	58.8																	
19-year	57.3	54.8	58.2	60.4																		
20-year	56.6	54.2	57.1																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 6.15  
Cancer of the Colon and Rectum (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	74.7	77.8	80.6	81.7	82.4	81.5	80.6	81.4	80.0	81.8	80.8	82.5	82.7	82.5	83.5	83.7	82.9	83.9	84.3	84.0	84.4	85.0
2-year	63.5	67.3	71.5	73.1	73.6	72.9	72.2	72.6	71.7	73.7	72.8	74.9	74.9	74.8	76.1	77.2	76.1	77.0	77.1	77.4	77.2	
3-year	57.2	60.8	66.0	67.9	67.9	67.2	67.0	67.5	66.0	68.8	67.6	69.7	70.2	70.5	71.4	72.4	71.1	71.8	72.7	72.1		
4-year	53.2	56.7	62.1	63.5	64.3	63.5	62.9	63.4	62.2	65.2	64.6	66.6	67.1	67.2	68.5	69.2	67.9	68.5	68.9			
5-year	50.6	53.9	59.5	61.1	61.9	61.3	60.1	60.4	60.0	62.8	61.8	63.5	64.8	65.1	66.3	66.5	65.5	65.9				
6-year	48.8	51.9	57.8	59.6	59.9	59.7	58.4	58.7	57.5	61.4	60.0	61.7	63.4	63.3	64.4	64.6	63.6					
7-year	47.5	50.4	56.2	57.9	58.4	58.1	56.7	57.3	55.8	60.3	59.1	60.3	62.3	61.8	62.4	63.1						
8-year	46.6	49.4	55.2	56.4	57.1	56.6	55.7	56.4	55.2	59.6	58.2	59.6	60.9	60.4	61.3							
9-year	46.0	48.7	54.3	55.7	55.9	55.7	54.8	56.0	54.9	58.8	57.3	58.3	60.1	59.6								
10-year	45.3	47.9	53.5	55.0	55.3	55.0	54.1	54.9	54.2	58.2	56.6	57.5	59.4									
11-year	44.8	47.3	53.1	54.4	55.1	54.3	53.2	54.5	53.3	57.5	55.8	56.6										
12-year	44.3	46.9	52.6	53.9	54.9	54.2	52.3	53.8	52.6	57.4	55.4											
13-year	43.9	46.6	51.9	53.4	54.5	53.5	51.8	53.7	52.0	57.0												
14-year	43.4	46.1	51.6	52.9	54.0	53.1	51.3	53.1	51.5													
15-year	43.0	45.8	51.1	52.2	53.9	53.1	50.4	52.4														
16-year	42.6	45.3	50.8	52.1	53.9	53.1	50.4															
17-year	42.2	44.9	50.2	51.9	53.1	52.6																
18-year	41.8	44.6	49.9	51.2	52.2																	
19-year	41.8	44.1	49.5	51.1																		
20-year	41.6	43.8	49.1																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 6.16  
Cancer of the Colon and Rectum (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	74.6	78.7	81.6	82.9	83.1	82.6	81.3	82.4	81.5	83.1	81.7	82.6	83.8	84.1	85.5	83.6	83.6	84.9	85.1	85.1	85.8	86.6
2-year	63.5	67.7	72.8	74.0	74.6	73.8	72.7	73.4	73.0	75.3	74.1	74.9	75.5	76.7	77.8	76.9	76.5	78.3	78.1	78.5	78.6	
3-year	56.6	60.9	67.1	69.0	68.7	67.8	67.4	68.5	67.2	70.0	68.5	69.1	70.8	72.3	73.0	72.2	71.4	72.9	73.4	73.1		
4-year	52.4	56.5	62.7	64.2	64.5	63.6	62.9	63.5	63.3	65.5	65.5	65.5	67.7	69.1	70.1	68.9	68.1	69.3	69.0			
5-year	49.6	53.5	60.0	61.1	61.6	61.0	59.7	59.9	60.2	62.6	62.6	62.6	65.1	66.4	67.5	65.9	65.6	66.4				
6-year	47.4	51.3	58.0	59.3	58.9	59.4	58.2	58.0	57.5	61.1	60.6	60.4	63.9	64.1	65.2	63.7	63.2					
7-year	46.1	49.7	56.4	57.1	57.0	57.8	56.5	56.5	55.9	59.8	59.9	58.9	62.7	62.2	63.2	61.9						
8-year	45.0	48.6	55.4	55.6	55.6	56.0	55.6	54.9	55.3	59.5	58.5	57.6	61.4	60.3	62.4							
9-year	44.3	47.7	54.4	55.1	54.6	55.1	54.2	54.3	54.8	59.0	57.5	56.4	60.2	59.8								
10-year	43.5	46.9	53.8	54.4	53.8	54.5	53.2	53.0	54.2	58.1	56.9	55.7	59.4									
11-year	43.0	46.2	53.3	53.9	53.8	53.8	51.9	52.9	53.3	57.3	56.0	54.8										
12-year	42.4	45.7	52.7	53.4	53.7	53.8	51.4	51.9	52.4	57.1	55.8											
13-year	41.9	45.5	52.1	52.9	53.3	53.1	50.3	51.5	52.0	56.7												
14-year	41.2	44.8	52.0	52.4	52.2	52.5	49.9	51.1	51.1													
15-year	41.0	44.6	51.4	51.7	51.8	52.4	49.0	50.6														
16-year	40.5	44.5	51.2	51.7	51.8	52.4	49.0															
17-year	40.1	43.8	50.7	51.6	50.8	52.4																
18-year	39.7	43.5	50.4	50.8	50.0																	
19-year	39.6	43.2	50.1	50.8																		
20-year	39.4	42.7	49.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 6.17  
Cancer of the Colon and Rectum (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	74.7	77.0	79.6	80.5	81.6	80.4	79.9	80.4	78.5	80.4	79.9	82.4	81.5	80.7	81.5	83.8	82.1	82.9	83.5	82.9	82.9	83.3
2-year	63.5	66.9	70.2	72.2	72.6	72.0	71.6	71.8	70.4	72.1	71.5	74.9	74.2	72.9	74.3	77.6	75.7	75.6	76.1	76.2	75.7	
3-year	57.7	60.6	64.9	66.6	67.0	66.6	66.5	66.6	64.9	67.6	66.7	70.3	69.7	68.7	69.6	72.6	70.7	70.7	71.8	71.2		
4-year	54.0	56.8	61.5	62.8	64.0	63.4	62.9	63.3	61.1	64.8	63.5	67.6	66.5	65.2	66.8	69.5	67.7	67.6	68.9			
5-year	51.6	54.3	59.1	61.0	62.1	61.6	60.4	60.9	59.8	62.9	61.0	64.5	64.5	63.8	65.1	67.1	65.5	65.4				
6-year	50.0	52.5	57.5	59.9	60.8	60.0	58.7	59.5	57.5	61.7	59.4	63.0	62.8	62.5	63.5	65.6	64.1					
7-year	48.9	51.2	56.0	58.6	59.8	58.3	57.0	58.1	55.7	60.8	58.1	61.7	61.8	61.5	61.5	64.4						
8-year	48.0	50.3	54.8	57.3	58.7	57.1	55.8	57.7	55.0	59.6	57.8	61.3	60.4	60.6	60.2							
9-year	47.5	49.5	54.1	56.3	57.3	56.4	55.4	57.7	54.8	58.7	57.2	60.2	59.9	59.3								
10-year	46.9	48.9	53.3	55.6	56.8	55.4	55.1	56.7	54.2	58.1	56.3	59.3	59.5									
11-year	46.4	48.3	52.8	54.8	56.2	54.8	54.6	56.1	53.3	57.7	55.5	58.4										
12-year	46.1	47.9	52.5	54.5	56.1	54.4	53.1	55.8	52.8	57.7	54.7											
13-year	45.8	47.6	51.7	53.9	55.6	53.8	53.1	55.8	52.0	57.2												
14-year	45.3	47.3	51.2	53.4	55.6	53.6	52.6	55.2	51.8													
15-year	44.9	46.9	50.7	52.7	55.6	53.6	51.8	54.3														
16-year	44.5	46.1	50.3	52.5	55.6	53.5	51.2															
17-year	44.2	45.9	49.6	52.0	55.5	52.5																
18-year	43.7	45.5	49.3	51.2	54.5																	
19-year	43.7	45.1	49.0	50.9																		
20-year	43.4	44.9	48.5																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 7.9  
Cancer of the Corpus And Uterus, NOS (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	94.0	91.7	92.5	92.7	92.1	93.1	92.5	91.9	93.2	91.8	93.2	93.7	94.2	93.9	92.4	92.3	93.2	93.0	92.9	93.1	93.3	93.4
2-year	90.3	86.5	87.7	88.0	88.1	89.6	88.5	87.2	88.6	88.2	89.3	89.7	89.7	89.6	88.3	88.2	88.7	88.5	89.2	89.6	89.5	
3-year	87.9	83.6	85.0	85.1	85.6	87.1	85.6	84.8	86.2	85.8	86.3	86.6	87.5	87.5	86.0	85.8	86.2	85.5	85.8	87.3		
4-year	86.4	82.1	83.5	83.5	83.8	85.6	84.2	83.2	84.8	84.7	85.3	84.6	85.9	86.2	84.2	83.8	84.8	83.8	84.1			
5-year	85.9	81.0	82.5	82.6	83.2	85.2	83.1	82.3	84.2	83.3	84.4	83.4	84.5	85.4	83.4	82.9	83.5	82.5				
6-year	85.2	80.4	81.8	81.3	82.0	84.3	82.3	81.2	83.5	82.4	83.4	82.6	84.0	84.6	82.4	82.5	82.5					
7-year	84.7	79.9	81.4	81.3	81.8	84.1	82.1	80.3	83.0	81.5	82.7	82.3	82.9	84.3	81.7	81.2						
8-year	84.6	79.2	80.9	80.8	81.0	83.6	80.9	79.7	82.7	80.6	81.8	81.9	82.2	83.5	81.0							
9-year	84.6	78.9	80.6	80.7	80.8	83.6	80.6	79.7	81.8	80.1	81.2	81.4	82.2	83.4								
10-year	84.4	78.6	80.4	79.9	80.6	83.1	79.8	79.6	81.4	79.8	81.2	80.9	81.7									
11-year	84.1	78.2	80.1	79.9	80.6	83.1	79.0	79.6	81.3	79.8	81.2	80.9										
12-year	84.0	78.0	79.7	79.5	80.1	82.8	78.1	79.6	81.1	79.8	81.2											
13-year	84.0	77.6	79.4	78.8	79.7	82.8	78.0	79.6	80.8	79.3												
14-year	83.8	77.2	79.1	78.7	78.8	82.8	77.2	78.9	80.3													
15-year	83.8	77.0	78.8	78.7	77.7	82.8	77.1	78.9														
16-year	83.5	76.3	78.4	78.7	76.9	82.8	76.4															
17-year	83.5	76.1	78.2	78.7	76.8	82.8																
18-year	83.5	75.7	78.1	78.6	76.8																	
19-year	83.5	75.0	78.0	77.9																		
20-year	83.5	75.0	77.3																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 8.9  
Cancer of the Esophagus (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	27.5	31.3	36.2	39.2	41.5	41.8	38.2	38.6	41.5	43.6	41.6	41.7	44.9	47.6	47.6	42.9	48.1	49.0	47.1	49.2	51.1	48.8
2-year	11.6	14.5	19.6	20.3	21.1	24.5	22.1	23.0	23.7	24.4	24.0	24.3	27.1	28.8	29.7	26.1	30.9	30.3	31.5	30.8	33.6	
3-year	7.7	10.1	13.6	14.5	15.0	19.4	15.9	17.9	17.3	18.7	17.6	18.6	21.0	23.9	23.2	20.8	24.2	25.8	24.5	25.2		
4-year	5.9	8.3	10.9	11.5	12.8	16.7	13.5	14.8	13.4	15.9	14.3	14.3	18.5	20.6	19.2	18.2	20.2	22.5	21.6			
5-year	4.9	6.9	9.6	10.0	11.8	14.8	12.0	12.9	11.6	15.0	12.5	12.6	17.5	19.0	17.8	16.8	17.4	20.4				
6-year	4.3	6.1	8.5	9.5	10.3	13.9	10.7	11.6	10.8	13.8	11.3	11.7	16.0	18.0	15.8	15.3	14.6					
7-year	3.8	5.6	7.7	8.5	9.8	12.4	9.9	10.6	10.2	12.9	10.8	11.3	14.9	16.7	14.7	14.8						
8-year	3.4	5.0	7.1	8.0	8.6	11.8	9.5	9.8	10.0	12.3	9.8	9.8	14.4	15.8	13.7							
9-year	3.2	4.5	6.5	7.3	8.5	11.1	8.6	9.6	9.1	11.6	9.4	9.6	13.6	15.6								
10-year	2.9	4.3	6.0	6.9	8.2	10.0	8.1	8.2	8.6	10.8	9.1	9.5	13.0									
11-year	2.7	3.9	5.5	6.8	8.2	8.7	7.2	7.9	7.7	9.7	9.0	9.1										
12-year	2.5	3.8	4.9	5.1	7.5	7.8	7.1	7.8	7.3	9.4	7.4											
13-year	2.4	3.5	4.7	5.0	7.5	6.9	7.1	7.8	6.9	9.4												
14-year	2.4	3.4	4.3	5.0	7.1	6.4	6.7	7.6	5.9													
15-year	2.4	3.4	4.2	4.8	7.0	5.8	6.7	6.5														
16-year	2.3	3.2	3.7	4.8	7.0	5.5	6.0															
17-year	2.3	3.0	3.3	4.7	6.9	5.1																
18-year	2.2	2.7	3.1	4.4	6.9																	
19-year	2.1	2.5	3.0	3.9																		
20-year	2.1	2.4	2.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 8.10  
Cancer of the Esophagus (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	25.3	30.5	35.1	41.2	41.1	41.2	39.4	38.0	41.0	44.9	42.0	41.9	44.5	48.8	48.9	44.3	47.7	48.2	48.4	50.0	52.9	49.9
2-year	10.3	13.9	19.1	21.1	20.7	24.4	21.8	21.6	24.0	24.9	24.7	24.8	26.8	29.1	30.2	26.8	30.2	30.1	31.4	31.4	34.1	
3-year	6.8	9.3	13.2	14.4	15.1	19.4	16.2	17.8	17.4	18.4	18.5	19.0	20.9	23.7	23.2	21.3	23.6	25.5	24.6	26.0		
4-year	5.1	7.5	10.4	11.3	12.4	15.9	13.8	14.5	13.2	15.1	15.2	14.3	18.3	19.5	19.4	18.5	19.7	22.2	21.1			
5-year	4.2	6.0	9.4	9.6	11.7	14.1	12.1	13.0	11.9	13.9	13.2	12.1	17.4	17.4	18.3	16.8	16.9	19.8				
6-year	3.6	5.2	8.4	9.1	10.0	13.1	10.9	11.6	11.0	13.0	11.8	11.5	16.3	16.5	16.2	15.6	14.1					
7-year	3.3	4.5	7.6	8.0	10.0	11.7	10.6	10.7	10.7	11.9	11.5	11.2	15.2	15.5	15.4	15.4						
8-year	3.0	4.0	7.0	7.3	8.5	11.1	10.1	9.8	10.3	11.5	10.3	9.4	15.0	14.6	14.5							
9-year	2.8	3.7	6.5	6.6	8.3	10.6	9.0	9.3	9.8	10.6	10.0	9.3	14.3	14.3								
10-year	2.6	3.4	6.2	6.1	7.8	9.3	8.1	8.3	9.3	10.1	9.7	9.3	13.4									
11-year	2.5	3.3	5.8	6.0	7.8	7.7	7.4	8.3	8.4	9.0	9.5	9.1										
12-year	2.3	3.2	5.1	4.6	6.9	6.9	7.4	8.1	7.8	8.7	8.3											
13-year	2.2	3.1	4.6	4.6	6.8	6.9	7.4	8.1	7.2	8.7												
14-year	2.2	2.9	4.2	4.6	6.8	6.5	6.9	8.1	6.4													
15-year	2.2	2.9	4.0	4.4	6.8	5.8	6.7	7.1														
16-year	2.1	2.9	3.4	4.4	6.8	5.7	6.5															
17-year	2.1	2.8	2.9	4.1	6.7	5.0																
18-year	2.1	2.5	2.8	3.7	6.6																	
19-year	2.1	2.3	2.5	3.6																		
20-year	2.0	2.2	2.3																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).



Table 8.11  
Cancer of the Esophagus (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	33.2	33.2	39.1	32.9	42.4	43.8	35.1	40.3	42.9	39.6	40.6	40.9	46.4	43.8	42.9	38.4	49.5	51.8	42.3	46.2	44.7	44.2
2-year	15.1	15.8	21.1	17.8	22.0	25.0	22.9	27.0	22.7	22.9	22.0	22.5	28.3	27.9	28.0	24.0	33.4	31.2	32.0	29.1	31.7	
3-year	10.1	12.1	14.6	14.6	14.9	19.3	14.9	18.3	17.0	19.8	15.1	17.6	21.0	24.7	23.2	19.2	26.3	26.8	24.4	22.7		
4-year	8.0	10.5	12.3	12.0	13.6	18.8	12.8	15.3	13.7	18.1	11.8	14.4	19.1	23.9	18.7	17.3	22.3	23.3	23.4			
5-year	6.5	9.4	10.1	11.1	12.1	17.2	11.6	12.6	10.8	18.1	10.6	14.1	17.5	23.8	16.1	16.6	19.1	22.1				
6-year	6.0	8.6	8.5	10.7	10.9	16.6	10.0	11.7	10.0	16.3	9.8	12.4	14.8	22.7	14.2	14.4	16.3					
7-year	5.2	8.2	8.1	9.8	9.2	14.5	8.1	10.2	9.0	15.7	8.6	11.8	13.6	20.7	12.2	13.1						
8-year	4.4	7.3	7.3	9.8	8.9	14.2	7.8	9.7	9.0	14.7	8.1	10.9	11.6	19.6	10.5							
9-year	4.1	6.5	6.4	9.2	8.9	12.5	7.6	9.7	7.1	14.1	7.4	10.5	10.5	19.1								
10-year	3.5	6.4	5.4	8.9	8.9	12.2	7.6	7.9	6.6	12.8	7.4	9.6	10.5									
11-year	3.3	5.4	4.8	8.9	8.9	11.9	6.6	6.8	5.7	11.8	7.3	9.3										
12-year	3.1	5.1	4.6	6.6	8.9	10.8	6.0	6.8	5.7	11.4	4.6											
13-year	2.8	4.7	4.6	5.9	8.9	6.9	6.0	6.5	5.7	11.4												
14-year	2.8	4.5	4.6	5.9	7.2	6.3	6.0	6.2	4.0													
15-year	2.8	4.5	4.5	5.9	7.2	5.6	6.0	5.0														
16-year	2.6	3.8	4.2	5.9	7.2	4.9	4.2															
17-year	2.4	3.1	3.9	5.9	7.2	4.9																
18-year	2.2	3.0	3.8	5.9	7.2																	
19-year	2.2	2.8	3.8	4.7																		
20-year	2.2	2.7	3.2																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 9.9  
Hodgkin Lymphoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	87.2	89.7	91.1	90.8	93.0	92.6	91.6	93.3	92.9	91.7	93.6	94.5	91.1	92.6	93.9	93.4	93.4	91.8	93.5	92.7	93.5	94.7
2-year	80.9	83.8	86.3	86.6	87.1	89.0	86.5	89.7	89.0	88.4	91.0	90.4	87.9	90.5	90.8	90.1	92.2	89.8	91.0	90.1	92.4	
3-year	76.7	80.2	83.1	84.3	84.6	87.4	83.6	86.2	86.7	86.7	89.3	88.3	85.8	88.7	88.4	88.5	90.2	87.6	90.3	89.2		
4-year	73.8	77.5	80.6	82.2	82.3	85.4	81.7	84.3	83.7	84.5	88.4	86.6	84.3	87.8	86.5	86.2	88.1	86.2	89.5			
5-year	71.5	74.8	79.1	81.1	80.2	84.1	79.7	83.0	82.3	83.2	87.3	84.9	83.4	86.4	85.1	85.1	87.4	85.7				
6-year	69.0	72.3	77.5	79.4	78.9	82.3	78.0	82.1	80.1	81.6	85.9	83.8	82.1	85.6	84.3	84.7	87.0					
7-year	67.2	70.7	75.1	77.0	77.6	81.4	76.1	81.3	79.0	80.0	85.3	82.9	80.8	84.5	84.2	83.2						
8-year	65.0	69.2	74.0	75.7	76.2	81.0	75.1	80.9	78.5	79.2	84.7	82.4	79.9	83.5	83.9							
9-year	63.0	67.8	73.1	74.5	75.7	79.9	74.2	79.5	77.2	78.1	84.4	81.4	79.1	82.6								
10-year	61.7	66.7	72.5	73.6	74.9	78.9	74.1	79.2	76.7	76.7	83.9	80.1	78.2									
11-year	60.4	65.7	71.6	73.1	73.8	78.0	73.6	79.1	75.7	75.7	83.3	79.5										
12-year	59.5	64.6	70.7	71.9	73.6	77.2	72.6	78.0	75.5	74.9	82.7											
13-year	58.7	63.7	69.5	71.7	72.7	76.4	71.6	77.1	75.4	74.2												
14-year	57.4	62.5	68.6	71.5	72.1	75.9	71.5	76.8	75.4													
15-year	56.4	61.7	67.5	70.9	71.4	75.3	71.2	75.4														
16-year	55.6	60.7	66.6	70.9	70.4	75.3	70.9															
17-year	54.7	59.3	66.1	69.9	70.0	74.5																
18-year	53.8	58.4	65.4	68.9	68.8																	
19-year	53.1	57.8	64.7	67.8																		
20-year	52.0	56.8	64.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 9.10  
Hodgkin Lymphoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	86.1	89.6	90.0	90.7	93.0	90.9	89.3	93.6	92.2	92.9	92.1	94.2	89.3	90.8	92.9	91.0	93.0	89.7	93.6	91.9	92.6	93.5
2-year	80.4	83.8	84.8	84.7	86.4	86.4	83.5	88.8	88.6	89.1	89.5	90.9	86.5	88.4	89.2	88.5	92.2	88.4	91.1	89.1	91.6	
3-year	75.5	80.4	81.5	82.6	82.8	84.6	80.2	85.6	85.4	87.1	87.5	88.8	83.8	85.4	86.4	86.9	90.1	86.7	90.2	88.8		
4-year	72.0	77.4	78.5	80.7	79.7	82.4	77.9	83.1	81.3	85.1	87.0	86.4	82.8	83.7	85.0	84.8	86.8	85.9	88.9			
5-year	69.4	74.1	76.6	80.2	76.8	80.8	75.3	81.9	80.5	83.0	86.0	84.4	82.5	81.9	83.5	83.7	86.0	85.6				
6-year	66.6	71.5	74.8	78.6	75.7	78.7	72.9	80.8	78.8	81.8	83.6	83.5	80.4	80.9	82.0	82.9	86.0					
7-year	64.5	69.8	71.9	75.7	73.5	78.6	71.7	80.5	77.6	79.7	82.9	82.1	78.6	80.0	81.6	81.0						
8-year	62.5	68.2	70.4	74.7	71.3	77.5	70.3	80.3	76.5	78.6	82.3	81.6	77.5	78.8	81.2							
9-year	60.0	67.0	69.2	73.5	71.0	76.9	69.1	78.5	75.4	77.5	82.2	80.2	76.2	78.6								
10-year	58.9	65.6	68.5	72.5	70.5	75.7	69.1	78.4	75.4	75.5	81.3	78.1	75.1									
11-year	57.5	64.7	67.1	72.1	69.7	74.5	68.7	78.3	74.3	74.3	80.7	77.5										
12-year	56.4	63.3	66.0	70.7	69.2	73.5	67.7	77.6	74.3	72.7	80.3											
13-year	55.8	62.5	64.3	70.4	67.8	72.7	66.0	77.2	74.3	72.1												
14-year	54.4	61.2	63.4	70.4	67.5	72.7	65.7	77.0	74.3													
15-year	53.2	60.4	62.1	69.6	67.0	72.2	65.0	74.9														
16-year	52.4	59.4	61.2	69.4	66.0	72.2	64.6															
17-year	51.2	58.1	60.7	67.9	66.0	71.3																
18-year	50.2	56.9	60.0	66.7	65.3																	
19-year	49.5	56.5	59.4	66.1																		
20-year	48.5	55.2	58.5																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 9.11  
Hodgkin Lymphoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	88.7	89.9	92.5	90.9	93.1	94.6	94.4	92.9	93.7	90.0	95.1	94.9	93.2	95.1	95.3	95.9	93.8	93.9	93.4	93.6	94.6	96.1
2-year	81.6	83.7	88.3	89.0	87.9	91.8	90.0	90.7	89.7	87.4	92.6	89.9	89.4	93.5	92.8	91.8	92.2	91.1	90.9	91.3	93.2	
3-year	78.3	79.8	85.2	86.3	86.8	90.5	87.7	87.0	88.1	86.1	91.2	87.7	88.0	93.0	90.9	90.2	90.2	88.6	90.4	89.6		
4-year	76.3	77.7	83.3	83.9	85.7	88.7	86.0	85.6	86.7	83.8	89.7	86.8	85.7	93.0	88.5	87.7	89.7	86.3	90.1			
5-year	74.4	75.7	82.2	82.2	84.4	87.5	84.9	84.3	84.7	83.4	88.6	85.4	84.3	92.5	87.1	86.6	89.0	85.8				
6-year	72.3	73.5	80.9	80.4	83.1	86.2	83.8	83.5	81.8	81.3	88.5	84.0	83.9	92.1	86.9	86.6	88.0					
7-year	70.8	71.8	79.2	78.6	82.6	84.4	81.3	82.0	80.6	80.3	87.7	83.8	83.1	90.8	86.9	85.6						
8-year	68.5	70.6	78.4	76.9	82.3	84.4	80.9	81.6	80.6	80.0	87.2	83.2	82.6	90.0	86.9							
9-year	67.0	69.0	78.1	75.6	81.6	83.2	80.1	80.7	79.4	78.7	86.7	82.7	82.1	88.1								
10-year	65.4	68.2	77.5	74.8	80.1	82.3	79.9	80.0	78.1	78.2	86.5	82.3	81.2									
11-year	64.3	67.1	77.2	74.2	79.0	81.7	79.2	79.7	77.4	77.4	86.1	81.7										
12-year	63.5	66.3	76.6	73.2	79.0	81.0	78.2	78.3	76.8	77.2	85.0											
13-year	62.6	65.3	75.9	73.0	79.0	80.4	77.9	77.1	76.0	76.9												
14-year	61.5	64.3	75.2	72.6	78.0	79.3	77.9	76.7	76.0													
15-year	60.7	63.3	74.1	72.5	76.9	78.5	77.9	75.8														
16-year	59.9	62.3	73.4	72.5	75.9	78.2	77.9															
17-year	59.3	60.9	72.7	72.2	74.5	77.9																
18-year	58.5	60.2	72.2	71.8	73.3																	
19-year	57.9	59.5	71.4	69.8																		
20-year	56.7	58.9	70.9																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 10.7  
Kaposi Sarcoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	95.6	66.9	60.3	61.9	62.8	63.2	59.2	55.9	62.3	69.7	74.3	74.0	76.6	74.3	80.3	78.0	75.0	82.5	81.7	81.0	82.5	79.4
2-year	90.1	44.9	34.0	33.5	30.5	28.6	30.4	27.8	47.0	60.5	65.4	65.2	65.0	67.8	73.0	71.6	72.2	73.9	78.3	77.5	80.0	
3-year	89.0	36.0	20.2	20.1	15.6	16.5	19.7	21.1	42.8	54.6	62.3	60.7	61.3	61.3	67.5	67.1	66.6	70.7	74.3	74.9		
4-year	84.2	31.7	13.9	14.7	11.1	11.1	15.6	18.5	38.7	51.6	59.0	58.5	56.5	57.2	64.6	66.0	64.6	69.2	73.5			
5-year	84.2	28.0	10.4	10.1	8.8	9.2	14.1	17.4	35.6	48.3	56.9	58.1	55.7	54.2	63.4	59.4	64.5	68.0				
6-year	78.8	24.4	8.8	8.0	7.8	8.6	13.8	16.2	35.0	45.7	54.7	55.2	53.1	52.7	62.9	57.6	64.2					
7-year	76.9	23.3	7.6	7.6	7.6	7.9	13.2	15.1	33.4	44.8	50.5	52.7	52.6	52.7	62.1	57.6						
8-year	76.9	20.5	6.7	7.0	7.3	7.7	12.7	14.7	32.1	44.6	47.8	50.6	52.1	52.0	60.8							
9-year	76.9	19.1	6.0	6.9	6.9	7.6	12.5	14.0	31.0	42.7	44.3	49.4	52.1	50.3								
10-year	76.9	16.9	5.7	6.8	6.8	7.4	12.4	13.7	29.8	42.6	43.6	49.4	51.2									
11-year	74.9	15.6	5.5	6.7	6.7	7.4	11.8	13.4	28.6	42.1	42.4	48.7										
12-year	74.9	15.3	5.3	6.7	6.7	7.2	11.7	13.3	27.8	41.7	40.6											
13-year	74.9	14.7	5.0	6.7	6.5	7.2	11.7	13.3	27.3	41.7												
14-year	74.9	14.7	4.8	6.3	6.5	6.8	11.7	13.1	26.7													
15-year	74.9	14.7	4.7	5.8	6.5	6.6	11.6	12.9														
16-year	74.9	14.7	4.6	5.7	6.3	6.4	11.4															
17-year	74.9	14.4	4.6	5.6	6.3	6.2																
18-year	74.9	13.1	4.6	5.3	6.3																	
19-year	74.9	13.1	4.4	5.3																		
20-year	74.9	13.1	4.3																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 10.8  
Kaposi Sarcoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	95.8	65.2	59.9	61.5	62.4	63.0	59.0	55.3	61.4	69.5	73.9	73.3	76.7	73.6	79.5	76.8	75.3	81.3	82.7	80.3	82.6	80.8
2-year	89.6	42.2	33.4	33.0	29.7	28.1	29.4	27.2	45.7	59.9	64.4	63.6	64.5	66.0	73.3	70.5	72.1	72.6	78.9	76.6	80.4	
3-year	87.3	33.5	19.3	19.4	14.7	16.1	19.0	20.5	41.8	54.3	60.9	59.2	60.8	59.8	68.6	66.2	66.7	69.0	75.2	74.2		
4-year	84.3	29.2	13.1	14.0	10.1	10.8	14.8	17.9	37.6	51.1	57.1	56.5	55.5	56.4	65.1	65.7	64.6	67.1	74.4			
5-year	84.2	25.7	9.5	9.4	7.8	8.8	13.3	16.8	35.0	48.0	56.0	55.7	54.4	53.8	63.4	59.1	64.2	65.6				
6-year	75.6	22.0	7.9	7.3	6.9	8.3	12.9	15.6	34.3	45.5	53.5	53.5	51.5	52.7	62.9	57.3	63.8					
7-year	74.3	20.6	6.7	6.8	6.6	7.6	12.5	14.5	32.5	44.5	49.9	51.9	50.7	52.7	62.9	57.3						
8-year	74.3	18.3	5.9	6.3	6.3	7.5	12.0	14.1	31.3	44.3	46.9	50.9	50.6	51.9	61.6							
9-year	74.3	17.1	5.2	6.1	5.8	7.4	11.8	13.4	30.3	42.3	43.6	49.7	50.6	50.1								
10-year	74.3	15.1	5.0	6.1	5.7	7.0	11.6	13.2	29.2	42.1	42.8	49.7	50.1									
11-year	69.1	13.8	4.8	6.1	5.5	7.0	11.0	12.8	28.1	41.7	41.4	48.8										
12-year	69.1	13.6	4.7	6.1	5.4	6.9	10.9	12.7	27.2	41.2	40.0											
13-year	69.1	13.0	4.4	6.1	5.2	6.9	10.9	12.6	26.7	41.2												
14-year	69.1	13.0	4.3	5.6	5.2	6.5	10.9	12.3	26.2													
15-year	69.1	12.9	4.2	5.3	5.1	6.3	10.8	12.1														
16-year	69.1	12.7	4.1	5.3	5.0	6.1	10.6															
17-year	69.1	12.7	4.1	5.1	4.9	5.9																
18-year	69.1	12.4	4.1	4.8	4.8																	
19-year	69.1	12.4	3.9	4.8																		
20-year	69.1	12.4	3.8																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 10.9  
Kaposi Sarcoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	93.9	91.7	83.7	79.9	90.2	74.9	71.5	82.0	87.6	78.6	79.8	85.5	75.7	81.8	86.7	89.4	69.4	100.0	64.6	89.8	80.3	39.9
2-year	90.9	85.7	79.0	56.3	79.9	53.9	71.5	56.2	86.4	78.6	75.1	85.5	69.8	81.8	67.5	81.1	69.4	97.0	64.6	89.8	72.5	
3-year	90.9	74.9	74.8	56.3	75.4	44.5	52.9	50.0	74.9	64.6	75.1	85.4	65.9	77.7	52.1	71.3	62.2	97.0	57.3	81.1		
4-year	83.6	71.3	74.3	52.0	75.4	33.1	52.9	46.0	70.5	64.6	75.1	85.4	65.9	67.5	52.1	63.1	62.2	97.0	57.3			
5-year	81.5	64.4	74.1	49.9	73.2	33.1	52.9	46.0	50.5	55.2	69.1	85.4	65.9	57.6	52.1	59.0	62.2	97.0				
6-year	81.5	62.6	74.1	45.0	69.4	28.2	52.9	45.6	50.5	49.7	69.1	81.2	65.9	47.5	52.1	56.8	62.2					
7-year	81.5	62.6	74.1	45.0	69.4	28.2	47.8	40.9	50.5	49.7	58.1	54.7	65.9	47.5	50.9	56.8						
8-year	81.5	55.2	66.7	45.0	69.4	24.4	47.8	40.9	50.5	49.7	58.1	19.6	65.9	47.5	46.2							
9-year	81.5	52.0	66.7	45.0	69.4	24.4	44.3	38.9	47.7	49.7	52.0	19.6	65.9	47.5								
10-year	81.5	46.3	60.4	45.0	69.4	24.4	44.3	38.9	44.9	49.7	52.0	19.6	63.5									
11-year	81.5	43.6	58.3	40.8	69.4	24.4	44.3	38.9	37.9	49.7	52.0	19.6										
12-year	81.5	43.1	49.5	40.8	69.4	24.4	44.3	38.9	37.9	49.7	46.8											
13-year	81.5	42.2	44.7	40.8	69.4	24.4	44.3	38.9	37.9	49.7												
14-year	81.5	42.2	32.9	40.8	69.4	24.4	44.3	38.9	36.7													
15-year	81.5	42.2	28.0	19.3	69.4	24.4	44.3	38.9														
16-year	81.5	42.2	28.0	19.3	69.4	24.4	44.3															
17-year	81.5	30.2	28.0	19.3	69.4	24.4																
18-year	80.9	17.1	28.0	19.3	69.4																	
19-year	80.9	17.1	28.0	19.3																		
20-year	80.9	17.1	28.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 11.9  
Cancer of the Kidney and Renal Pelvis (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																						
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
1-year	69.6	71.2	73.5	76.2	77.1	77.4	76.6	77.0	77.3	77.6	76.9	78.3	78.3	80.2	81.6	80.0	82.1	85.4	84.2	85.3	86.4	85.3	
2-year	60.7	62.0	65.3	68.3	69.0	69.0	70.8	69.7	69.8	70.7	70.1	71.3	72.0	74.2	74.2	75.0	76.4	79.8	80.2	81.7	82.4		
3-year	56.6	57.2	60.9	65.4	64.5	66.1	66.7	67.0	65.5	67.5	66.1	67.8	67.7	69.5	71.0	71.5	73.4	76.6	77.7	78.2			
4-year	53.1	54.1	57.9	62.8	62.1	62.9	63.1	64.9	63.4	64.4	63.5	64.8	64.6	67.6	68.4	69.3	71.2	73.6	76.0				
5-year	51.0	52.2	55.9	59.9	60.4	60.5	61.1	62.1	61.5	62.6	61.8	63.4	62.5	65.7	66.6	67.6	69.2	71.3					
6-year	49.5	49.9	54.0	57.8	58.4	59.0	59.3	60.8	59.4	60.9	59.8	61.6	60.6	64.2	64.9	65.5	67.9						
7-year	47.5	48.7	52.4	55.5	57.4	56.4	58.6	59.1	58.0	59.2	58.2	59.7	59.1	62.4	63.2	63.5							
8-year	46.3	47.5	51.2	54.7	56.3	55.4	56.9	57.5	56.8	58.6	55.9	58.1	57.4	60.4	61.8								
9-year	45.1	46.2	49.8	53.6	54.7	54.1	55.9	56.9	55.4	58.1	54.2	56.9	55.7	59.1									
10-year	44.0	45.0	48.1	52.8	52.8	53.2	53.4	54.7	54.4	56.1	53.1	55.8	53.8										
11-year	43.2	43.7	46.6	51.7	51.3	51.0	51.9	53.1	52.4	55.1	51.9	55.2											
12-year	42.3	42.9	45.5	50.1	50.3	49.8	51.5	51.6	50.7	53.7	51.2												
13-year	41.6	42.1	44.5	49.3	49.6	48.4	50.5	49.9	49.4	51.9													
14-year	40.6	41.4	43.2	47.8	48.1	47.7	49.1	48.6	48.2														
15-year	39.7	41.0	42.3	46.1	48.1	46.8	48.5	47.9															
16-year	39.3	40.4	41.3	45.8	47.1	46.1	48.2																
17-year	38.8	39.9	40.4	45.1	46.4	45.5																	
18-year	37.8	39.3	39.5	44.1	45.3																		
19-year	36.8	38.3	38.6	42.9																			
20-year	36.1	37.5	37.9																				

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).



Table 11.10  
Cancer of the Kidney and Renal Pelvis (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	70.3	71.3	74.0	77.8	78.7	78.6	77.8	76.8	79.6	78.5	78.5	78.9	77.8	79.5	81.0	80.5	80.9	85.8	84.3	84.9	86.4	85.3
2-year	61.0	62.2	65.7	69.4	69.5	69.3	71.3	69.0	71.3	72.1	71.1	72.2	71.0	73.2	73.8	75.3	76.1	80.7	80.6	81.5	82.2	
3-year	56.6	57.6	61.6	66.5	65.8	66.4	67.3	65.9	67.1	68.3	66.8	68.3	66.8	68.0	70.4	71.1	73.6	77.4	77.9	77.7		
4-year	53.0	54.6	58.4	63.4	64.0	62.4	63.6	64.0	64.3	65.0	64.1	65.3	63.5	65.9	67.5	68.6	70.8	74.9	75.7			
5-year	51.1	52.6	56.4	60.2	62.1	59.4	61.3	61.4	62.4	63.2	61.9	64.0	61.5	63.1	65.6	67.3	69.3	72.2				
6-year	49.6	50.3	54.0	57.4	59.4	58.3	59.2	60.4	59.4	62.2	59.4	61.7	59.0	61.9	63.5	65.2	68.3					
7-year	47.4	49.1	52.3	55.1	58.4	55.7	58.8	58.8	58.4	60.4	57.8	59.6	57.5	59.8	61.1	63.4						
8-year	45.9	47.9	50.9	54.1	56.9	54.8	57.3	57.1	57.0	59.4	54.6	57.5	55.6	57.8	59.4							
9-year	44.9	46.5	49.1	53.0	55.8	53.8	56.3	56.7	56.0	59.0	52.7	56.6	53.7	56.1								
10-year	43.5	45.3	47.7	52.0	53.9	52.7	53.7	54.9	54.7	56.5	52.0	55.5	51.3									
11-year	42.5	44.1	45.7	50.3	52.2	50.7	52.8	53.4	53.3	55.7	50.4	54.9										
12-year	41.4	43.3	44.2	48.6	51.5	49.7	52.7	51.7	50.9	54.1	49.4											
13-year	40.7	42.7	43.0	47.8	51.0	48.1	51.3	49.9	49.5	52.5												
14-year	39.6	41.5	41.8	46.8	49.1	47.4	50.0	48.9	49.2													
15-year	38.5	41.3	41.0	45.2	48.9	46.5	49.3	48.6														
16-year	38.0	40.5	39.8	45.2	47.5	46.2	48.8															
17-year	37.6	40.1	38.9	45.2	46.2	45.2																
18-year	36.6	39.6	38.1	44.7	44.7																	
19-year	35.6	38.6	37.0	43.6																		
20-year	34.8	37.7	36.4																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 11.11  
Cancer of the Kidney and Renal Pelvis (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	68.5	71.0	72.6	73.7	74.5	75.6	74.9	77.1	73.6	76.1	74.3	77.3	79.1	81.2	82.5	79.2	84.0	84.8	84.2	85.8	86.3	85.3
2-year	60.0	61.7	64.6	66.6	68.0	68.5	70.0	70.8	67.3	68.6	68.6	69.8	73.4	75.8	74.9	74.6	76.8	78.4	79.4	82.1	82.6	
3-year	56.5	56.5	59.8	63.5	62.4	65.6	65.7	68.4	62.9	66.2	64.9	66.9	69.1	72.0	72.0	72.0	73.1	75.3	77.3	79.1		
4-year	53.2	53.3	57.2	61.8	59.1	63.5	62.5	66.3	62.0	63.4	62.6	64.0	66.3	70.4	69.6	70.4	71.8	71.5	76.2			
5-year	50.7	51.5	55.0	59.4	57.7	62.1	60.8	63.1	60.0	61.8	61.4	62.6	64.1	69.9	68.2	68.1	68.8	69.9				
6-year	49.2	49.3	53.8	58.4	56.6	60.1	59.2	61.3	59.1	59.0	60.3	61.4	63.2	67.9	67.2	65.8	67.3					
7-year	47.7	48.1	52.5	56.0	55.9	57.4	58.4	59.6	57.5	57.3	58.7	59.8	61.2	66.6	66.5	63.5						
8-year	46.9	46.9	51.6	55.5	55.1	56.2	56.3	58.0	56.3	57.2	57.8	58.9	60.1	64.4	65.6							
9-year	45.4	45.6	50.8	54.5	53.0	54.4	55.4	56.8	54.3	56.4	56.3	57.3	58.9	63.9								
10-year	45.0	44.5	48.7	53.7	50.9	53.8	52.6	54.4	53.9	55.4	54.6	56.1	57.5									
11-year	44.4	43.0	47.9	53.7	49.8	51.5	50.6	52.7	51.0	54.2	53.8	55.5										
12-year	43.7	42.1	47.3	52.4	48.4	50.0	49.7	51.2	50.4	53.0	53.8											
13-year	43.1	41.2	46.7	51.4	47.3	48.8	49.0	50.0	49.1	50.8												
14-year	42.1	40.8	45.3	48.8	46.6	48.1	47.6	48.0	46.5													
15-year	41.5	40.4	44.1	47.3	46.4	47.0	47.1	46.9														
16-year	41.2	40.3	43.6	45.6	46.1	46.1	46.7															
17-year	40.6	39.6	42.5	43.6	46.1	45.6																
18-year	39.7	38.7	41.7	43.1	46.0																	
19-year	38.6	37.6	40.9	42.0																		
20-year	38.3	37.1	40.1																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 12.9  
Cancer of the Larynx (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	88.6	89.0	88.9	87.7	90.6	88.3	87.6	87.2	89.4	88.7	86.4	86.9	86.0	86.0	85.7	84.8	84.8	87.1	87.1	86.8	88.4	84.8
2-year	79.2	80.0	79.3	77.7	80.7	78.0	79.0	77.0	80.1	79.6	78.1	76.4	75.1	77.0	75.6	75.0	74.8	75.8	78.3	74.6	76.1	
3-year	73.3	74.4	73.7	71.1	74.0	73.5	72.3	69.6	75.1	74.4	72.3	70.9	71.0	71.2	70.6	70.1	67.2	69.0	72.7	68.8		
4-year	68.9	70.2	68.9	66.5	69.9	69.2	66.5	65.3	69.5	69.8	68.8	67.7	68.8	67.0	66.8	67.6	62.7	64.4	70.2			
5-year	65.7	66.9	66.1	64.8	67.4	65.8	62.5	62.5	66.1	66.1	65.5	63.4	64.2	63.7	63.2	62.9	60.3	62.2				
6-year	63.3	64.0	62.8	60.5	63.5	62.7	60.3	58.2	63.6	64.0	62.4	59.1	60.7	60.8	60.7	60.1	56.0					
7-year	60.9	61.4	60.2	58.7	62.4	60.0	58.1	56.1	60.3	61.6	60.1	55.5	57.4	56.7	57.2	56.4						
8-year	58.9	59.7	57.3	56.6	58.7	57.3	56.0	54.6	57.7	58.4	57.3	52.5	54.5	53.7	54.8							
9-year	56.5	57.5	54.6	54.3	56.2	55.1	53.0	50.9	54.9	54.5	55.9	50.2	52.3	51.0								
10-year	54.2	55.6	52.4	51.7	54.0	51.9	50.3	49.4	51.6	52.2	53.8	48.2	51.2									
11-year	52.7	53.1	50.2	49.8	52.3	49.3	48.0	47.8	48.1	49.5	51.1	47.0										
12-year	51.0	50.4	48.2	47.4	51.1	46.7	46.1	47.0	46.3	47.2	46.4											
13-year	49.7	48.1	47.0	44.0	49.5	43.0	44.3	44.8	44.4	45.0												
14-year	48.2	46.6	44.4	42.3	47.5	40.9	42.5	43.1	41.6													
15-year	45.9	44.8	41.9	40.7	46.3	39.0	40.5	41.8														
16-year	43.8	42.3	40.4	38.3	44.7	36.7	37.9															
17-year	41.5	40.7	38.8	36.4	42.0	35.3																
18-year	40.3	38.8	37.4	32.9	40.1																	
19-year	38.6	37.3	34.7	30.2																		
20-year	36.6	35.6	33.5																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 12.10  
Cancer of the Larynx (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	88.9	89.1	89.2	88.2	90.8	88.6	88.0	89.4	91.3	88.3	89.0	87.9	87.4	86.1	86.2	85.3	84.7	87.7	87.0	87.1	89.3	84.8
2-year	79.3	80.2	79.7	78.4	81.4	79.0	79.7	79.0	81.9	79.7	80.9	77.2	76.3	77.6	75.8	75.4	75.5	76.0	77.3	75.7	76.3	
3-year	73.7	74.8	74.3	71.4	74.4	73.7	73.9	71.4	76.7	74.5	75.1	71.6	71.9	71.9	71.3	71.2	68.6	69.0	73.0	69.7		
4-year	69.2	70.7	69.7	67.3	70.4	69.8	68.3	66.6	71.8	69.9	71.4	69.1	70.0	67.7	66.9	69.0	63.4	63.8	70.8			
5-year	66.1	67.4	66.9	65.4	68.4	66.5	64.5	63.8	68.2	66.6	68.2	65.5	65.9	64.5	63.4	63.8	60.5	61.8				
6-year	63.7	64.5	63.5	61.3	64.5	63.2	62.3	59.8	65.8	64.2	65.3	62.3	62.3	61.8	61.0	61.3	56.3					
7-year	61.4	61.9	61.1	59.7	63.4	60.3	60.4	57.7	62.4	61.9	63.0	58.8	58.8	58.8	58.9	57.3						
8-year	59.3	60.4	58.4	57.6	59.9	58.2	58.1	56.7	59.6	58.5	60.2	55.4	56.1	56.6	56.5							
9-year	56.8	58.1	56.2	55.7	57.4	56.0	54.6	52.8	57.1	54.3	58.3	52.9	53.6	54.3								
10-year	54.8	56.4	54.1	53.2	54.8	52.7	51.9	50.9	54.0	51.9	56.4	50.8	52.4									
11-year	53.6	54.0	52.1	51.1	53.5	50.3	49.6	49.4	50.4	49.3	54.3	50.1										
12-year	52.1	51.2	50.3	49.0	52.2	47.5	47.6	48.3	49.3	46.7	49.4											
13-year	50.9	48.9	49.4	45.6	51.3	44.4	45.7	46.3	47.5	44.5												
14-year	49.3	47.3	47.0	44.3	48.7	42.0	44.2	44.4	45.1													
15-year	47.1	45.2	44.4	42.6	47.2	40.3	41.9	42.7														
16-year	44.9	43.0	42.9	40.2	46.2	38.3	39.2															
17-year	42.6	41.9	41.4	38.0	43.8	37.0																
18-year	41.4	39.8	39.7	34.8	41.4																	
19-year	39.5	38.2	36.7	31.7																		
20-year	37.7	36.5	35.7																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 12.11  
Cancer of the Larynx (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	87.2	88.4	87.5	85.9	89.5	87.1	85.8	78.5	81.4	90.0	76.7	83.6	81.1	85.4	83.3	82.3	85.4	84.8	87.6	85.3	84.9	84.7
2-year	78.5	78.9	77.7	74.6	78.1	73.8	75.9	69.0	72.7	79.0	67.7	73.9	70.7	74.5	75.0	73.5	71.7	75.0	82.0	70.0	75.1	
3-year	71.5	72.4	71.4	69.9	72.5	72.4	65.0	62.2	68.2	74.0	62.0	68.3	67.6	68.2	67.4	65.9	61.3	68.8	71.5	64.8		
4-year	67.0	67.6	65.7	63.4	67.8	66.7	58.4	59.9	60.2	69.4	59.2	63.0	64.5	64.1	66.0	61.6	59.5	66.4	68.3			
5-year	63.4	64.4	62.9	61.9	62.9	62.9	53.9	57.5	57.4	63.6	55.4	57.1	58.4	60.3	62.2	59.2	58.3	63.6				
6-year	61.1	61.6	59.7	57.7	59.7	60.3	51.5	51.8	54.4	62.8	51.8	49.5	55.1	55.9	58.9	55.4	54.9					
7-year	58.2	59.0	56.2	54.9	57.6	58.7	48.6	49.8	51.5	60.0	49.0	45.3	52.4	48.3	49.5	52.5						
8-year	56.9	56.6	52.7	52.5	54.1	53.2	47.0	46.4	49.7	58.0	46.6	43.8	49.1	41.7	46.6							
9-year	54.3	54.4	48.3	48.9	51.6	51.0	46.0	43.4	45.9	55.2	46.6	42.0	47.6	37.8								
10-year	51.0	51.8	45.6	45.9	50.9	48.9	43.5	43.0	41.7	53.0	44.0	40.1	46.9									
11-year	48.5	49.3	42.5	44.8	48.0	45.0	41.3	41.4	38.9	50.0	39.5	36.9										
12-year	45.6	46.9	40.0	41.3	45.6	42.8	39.5	41.2	34.2	48.4	35.7											
13-year	44.0	44.5	38.0	37.9	42.9	37.5	38.3	38.6	32.5	45.7												
14-year	42.6	43.5	34.5	34.4	42.5	35.8	35.5	38.1	28.2													
15-year	40.2	42.7	32.7	33.5	42.1	33.5	33.8	37.7														
16-year	38.4	39.2	30.9	31.2	39.3	30.1	32.3															
17-year	35.9	35.8	29.5	30.1	35.3	28.4																
18-year	35.1	34.7	29.0	25.8	34.6																	
19-year	34.0	33.2	27.4	24.4																		
20-year	31.5	32.1	25.3																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 13.17  
LeukemiaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	61.3	63.4	65.3	67.3	66.5	67.7	67.9	67.1	68.4	68.7	68.3	66.7	65.8	66.7	70.4	71.9	73.3	74.6	73.6	75.0	75.8	75.0
2-year	50.5	52.8	55.6	56.5	57.0	58.4	58.4	58.1	58.9	59.8	59.8	57.8	58.3	58.3	62.6	64.1	65.6	66.9	67.2	68.4	68.6	
3-year	44.0	46.8	49.8	51.5	51.3	53.2	54.4	53.5	54.0	55.0	54.9	53.6	54.3	54.3	58.3	61.0	61.3	63.3	63.1	64.7		
4-year	39.1	41.9	45.7	47.9	46.9	49.4	50.8	49.9	50.3	51.5	50.8	49.7	51.1	51.0	55.7	58.5	58.4	60.5	60.8			
5-year	34.9	37.9	42.5	45.3	44.3	46.3	47.3	47.1	47.6	48.1	47.9	47.6	47.8	48.7	53.2	56.5	56.1	58.0				
6-year	31.6	35.2	39.9	41.5	42.1	43.6	45.0	44.5	45.3	46.0	45.5	45.3	45.4	47.5	51.1	54.6	54.1					
7-year	29.3	32.6	37.3	38.8	39.5	40.7	42.2	42.3	43.9	44.7	43.4	43.8	43.6	46.1	49.7	52.7						
8-year	27.1	30.5	35.0	36.6	37.0	38.5	41.0	39.6	41.3	43.3	42.8	42.2	42.0	44.4	48.1							
9-year	25.2	28.7	33.0	34.9	35.5	36.7	39.2	38.1	40.0	41.5	41.4	41.3	41.0	43.5								
10-year	23.7	27.1	31.6	33.3	34.2	35.5	38.1	37.3	38.4	40.1	40.3	40.1	39.5									
11-year	22.4	25.8	29.9	31.7	32.8	33.9	37.2	36.0	37.1	39.2	38.4	38.6										
12-year	21.3	24.7	28.8	29.9	31.9	32.9	36.3	34.9	36.5	38.0	37.1											
13-year	20.5	23.6	27.9	28.5	31.4	32.3	35.1	33.5	36.0	37.6												
14-year	19.8	23.1	27.2	27.8	30.7	31.2	34.6	32.5	34.7													
15-year	19.3	22.3	26.7	27.6	30.1	30.6	33.7	32.0														
16-year	18.5	21.5	26.0	27.1	29.3	29.5	33.1															
17-year	18.1	20.7	25.5	26.4	28.5	28.9																
18-year	17.7	20.2	25.2	25.7	27.6																	
19-year	17.2	19.5	24.7	24.9																		
20-year	16.6	19.2	24.3																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 13.18  
LeukemiaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	61.0	63.7	66.1	68.0	67.7	68.1	69.1	69.1	69.8	70.3	69.1	66.9	66.8	68.3	72.6	72.9	74.3	76.2	75.6	75.7	76.8	75.6
2-year	50.0	52.8	56.4	56.4	58.1	58.5	60.0	59.4	60.2	61.6	60.9	58.4	59.1	60.1	64.5	64.3	66.3	67.6	69.2	68.6	69.8	
3-year	43.0	46.4	50.1	50.8	52.0	52.9	56.1	54.1	55.3	57.0	55.1	54.3	54.9	56.5	60.2	61.3	61.6	64.2	65.2	64.7		
4-year	38.1	41.4	46.1	46.9	46.6	49.3	52.6	51.3	51.3	53.3	50.3	50.2	51.5	52.7	57.0	58.5	58.9	61.1	63.3			
5-year	33.7	37.4	43.2	43.3	44.2	46.6	48.7	48.3	47.8	49.1	47.4	48.0	47.5	51.0	54.1	55.9	56.4	58.4				
6-year	30.5	34.6	40.5	39.2	42.3	43.5	46.5	45.2	44.8	46.7	45.3	45.6	45.3	49.8	51.5	53.7	54.0					
7-year	27.9	32.0	37.6	36.7	39.5	40.3	43.4	43.0	43.5	45.4	43.2	43.9	43.2	48.6	49.9	51.2						
8-year	25.5	29.6	35.4	34.0	37.1	38.3	42.1	39.9	41.4	44.2	42.5	42.5	41.4	46.4	48.3							
9-year	23.4	27.8	33.3	32.6	35.7	36.7	40.4	37.9	39.9	41.9	41.5	41.3	40.4	44.9								
10-year	21.9	26.3	31.6	30.5	34.2	35.8	39.3	37.3	37.8	40.2	40.7	40.2	38.9									
11-year	20.5	25.1	30.1	29.1	32.7	34.7	38.4	36.0	37.1	39.3	38.6	38.5										
12-year	19.4	24.4	29.0	27.4	31.6	33.7	37.2	35.0	36.4	38.5	37.2											
13-year	18.8	23.5	28.3	26.4	31.0	32.9	36.0	33.2	36.0	38.5												
14-year	18.0	23.0	27.7	26.0	29.8	31.5	35.5	31.7	34.9													
15-year	17.5	22.3	26.9	25.7	29.0	31.3	34.0	31.4														
16-year	16.8	21.4	26.3	25.4	28.4	30.3	33.3															
17-year	16.4	20.6	25.9	24.5	27.6	29.4																
18-year	16.0	20.0	25.6	23.5	26.9																	
19-year	15.5	19.4	25.1	22.8																		
20-year	14.6	18.9	24.8																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 13.19  
LeukemiaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	61.7	63.0	64.3	66.4	64.9	67.1	66.4	64.5	66.4	66.7	67.2	66.5	64.5	64.6	67.5	70.6	71.9	72.5	70.7	74.2	74.5	74.2
2-year	51.3	52.8	54.5	56.5	55.5	58.1	56.4	56.3	57.0	57.4	58.3	57.0	57.1	56.0	59.9	64.0	64.7	65.8	64.5	68.3	67.1	
3-year	45.2	47.4	49.3	52.4	50.4	53.6	52.2	52.7	52.2	52.4	54.6	52.6	53.3	51.4	55.7	60.6	60.8	62.0	60.3	64.7		
4-year	40.4	42.7	45.1	49.2	47.2	49.5	48.4	48.2	49.1	49.2	51.4	49.0	50.4	48.7	54.0	58.7	57.7	59.8	57.2			
5-year	36.4	38.5	41.6	48.0	44.4	45.8	45.6	45.5	47.4	46.7	48.7	47.0	48.1	45.6	51.9	57.1	55.7	57.5				
6-year	33.0	36.0	39.2	44.5	41.9	43.7	43.0	43.6	45.9	45.0	45.8	44.8	45.4	44.6	50.5	55.7	54.2					
7-year	31.1	33.4	36.9	41.8	39.4	41.2	40.6	41.3	44.3	43.8	43.6	43.6	44.2	42.8	49.3	54.7						
8-year	29.1	31.6	34.5	40.0	36.8	38.8	39.5	39.2	41.1	42.2	43.2	41.8	42.8	41.7	47.7							
9-year	27.5	29.9	32.6	38.0	35.3	36.6	37.7	38.4	40.1	41.0	41.0	41.1	41.9	41.6								
10-year	26.0	28.1	31.5	36.9	34.2	35.2	36.5	37.4	39.1	39.8	39.5	39.7	40.3									
11-year	24.8	26.8	29.6	35.0	32.9	32.8	35.6	36.0	37.0	39.1	38.2	38.8										
12-year	23.6	25.0	28.6	33.1	32.3	31.7	35.0	34.6	36.5	37.3	37.0											
13-year	22.7	23.8	27.5	31.3	32.0	31.4	33.9	33.8	35.8	36.4												
14-year	22.1	23.2	26.6	30.1	31.7	30.7	33.5	33.3	34.5													
15-year	21.4	22.4	26.3	30.0	31.5	29.5	33.4	32.5														
16-year	20.7	21.6	25.6	29.4	30.4	28.4	32.7															
17-year	20.1	20.9	25.0	28.8	29.7	28.2																
18-year	19.8	20.5	24.6	28.6	28.6																	
19-year	19.3	19.7	24.0	27.5																		
20-year	19.0	19.4	23.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).



Table 14.9  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	14.3	15.5	18.6	20.6	22.7	20.5	21.1	25.4	24.3	25.4	28.4	27.6	28.6	29.4	31.8	33.6	36.6	38.8	39.8	40.3	42.5	45.7
2-year	6.9	8.5	9.8	11.2	13.0	12.6	11.1	14.8	14.1	15.0	17.7	17.2	17.8	20.2	21.7	22.3	24.6	26.4	27.1	29.0	29.8	
3-year	4.8	6.0	7.4	9.0	9.5	8.4	7.9	9.2	9.5	10.7	12.6	13.1	14.1	15.2	16.1	17.4	20.1	20.9	21.0	23.3		
4-year	4.1	4.6	6.4	6.4	7.5	6.8	6.1	7.7	7.0	9.1	10.2	10.8	11.9	12.3	13.6	15.4	16.7	17.7	18.1			
5-year	3.4	3.8	5.3	5.1	5.9	5.6	4.5	6.5	5.7	8.0	8.1	9.4	10.0	11.4	11.9	14.0	15.1	14.5				
6-year	3.0	3.5	5.0	4.6	5.6	5.1	3.8	5.7	4.9	6.9	7.5	8.5	8.8	10.7	10.9	11.8	14.4					
7-year	2.8	3.1	4.8	4.3	4.9	4.6	3.5	5.2	4.4	6.5	6.8	7.1	8.2	9.9	9.8	11.2						
8-year	2.7	2.9	4.3	4.3	4.5	4.4	3.4	5.0	4.2	6.3	5.9	6.5	7.6	9.6	8.7							
9-year	2.6	2.8	4.2	4.3	4.4	4.4	3.3	4.7	4.1	5.8	5.6	5.5	7.3	9.2								
10-year	2.6	2.8	4.0	3.7	4.3	4.4	3.2	4.6	3.6	5.5	5.3	5.5	7.2									
11-year	2.6	2.6	3.9	3.4	3.9	4.4	3.2	4.5	3.2	5.3	5.3	4.7										
12-year	2.5	2.6	3.8	3.3	3.9	4.1	3.1	4.5	3.1	5.0	5.3											
13-year	2.3	2.4	3.8	3.3	3.9	4.0	3.1	4.0	3.1	4.8												
14-year	2.3	2.4	3.7	3.2	3.8	3.9	3.1	4.0	3.1													
15-year	2.2	2.4	3.7	3.2	3.7	3.6	3.1	4.0														
16-year	2.2	2.3	3.7	3.0	3.5	3.6	3.1															
17-year	2.1	2.3	3.7	3.0	3.5	3.6																
18-year	2.0	2.2	3.7	2.8	3.4																	
19-year	2.0	2.1	3.7	2.8																		
20-year	2.0	2.1	3.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 14.10  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	11.9	13.4	15.6	18.1	21.4	20.3	19.1	24.6	22.6	25.3	27.4	26.5	29.7	28.9	31.3	33.0	35.9	38.2	40.4	40.3	41.5	46.2
2-year	5.1	6.6	7.4	9.0	13.0	12.6	9.8	13.6	12.9	15.8	16.7	16.3	17.8	19.4	20.6	21.4	24.9	26.0	27.9	29.7	29.2	
3-year	3.2	4.9	5.4	7.6	9.6	8.0	6.8	8.4	9.0	11.1	12.0	12.1	13.5	15.3	15.4	17.8	20.0	20.3	21.4	23.6		
4-year	2.7	3.5	4.5	5.3	7.3	6.1	5.4	7.6	6.2	9.1	9.8	10.6	11.3	12.2	12.6	15.6	16.8	17.7	18.3			
5-year	2.0	2.9	3.6	4.0	5.2	5.3	4.3	6.5	5.0	7.7	7.9	8.9	9.8	11.1	10.9	14.3	15.6	14.5				
6-year	1.7	2.6	3.3	3.3	4.8	4.9	4.0	6.0	4.3	6.7	7.5	7.8	8.7	10.4	10.1	11.8	14.8					
7-year	1.4	2.2	3.3	3.3	4.1	4.3	3.7	5.5	3.7	6.3	6.7	6.4	8.2	9.8	9.1	11.1						
8-year	1.4	1.9	3.0	3.3	3.7	4.2	3.5	5.1	3.6	6.3	5.5	6.1	7.5	9.7	7.7							
9-year	1.3	1.9	3.0	3.3	3.5	4.2	3.4	4.6	3.4	5.7	5.4	5.3	7.3	9.5								
10-year	1.3	1.9	2.9	3.0	3.3	4.2	3.2	4.4	2.8	5.2	5.4	5.3	7.1									
11-year	1.3	1.8	2.9	2.7	3.1	4.2	3.2	4.3	2.3	4.9	5.4	4.4										
12-year	1.3	1.7	2.7	2.5	3.1	4.2	3.2	4.3	2.1	4.6	5.4											
13-year	1.1	1.4	2.7	2.5	3.1	4.0	3.2	3.7	2.1	4.5												
14-year	1.1	1.4	2.6	2.5	3.1	4.0	3.2	3.7	2.1													
15-year	1.0	1.4	2.6	2.3	3.1	4.0	3.2	3.7														
16-year	1.0	1.4	2.6	1.9	3.1	4.0	3.2															
17-year	1.0	1.4	2.6	1.9	3.1	4.0																
18-year	1.0	1.4	2.6	1.9	2.9																	
19-year	1.0	1.4	2.6	1.9																		
20-year	1.0	1.4	2.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 14.11  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	19.0	19.5	24.5	25.8	25.3	20.9	25.2	27.1	27.6	25.5	30.4	30.0	26.1	30.5	33.1	34.9	38.6	40.3	38.1	40.4	45.2	44.6
2-year	10.3	12.2	14.6	15.7	13.2	12.6	13.9	17.3	16.5	13.2	19.7	19.0	17.9	21.8	24.4	24.6	23.9	27.8	25.4	27.2	31.7	
3-year	7.8	8.2	11.3	11.8	9.4	9.2	10.3	10.8	10.4	9.9	13.7	15.2	15.3	15.0	17.7	16.6	20.4	22.5	20.2	22.4		
4-year	6.6	6.7	9.8	8.7	8.0	8.2	7.6	8.0	8.3	9.0	11.1	11.1	13.4	12.3	15.9	14.8	16.6	17.6	17.8			
5-year	6.1	5.6	8.8	7.4	7.3	6.2	4.9	6.4	7.0	8.5	8.5	10.3	10.3	12.0	14.0	13.3	13.9	14.2				
6-year	5.4	5.2	8.3	7.0	7.1	5.4	3.4	5.2	6.0	7.3	7.4	9.9	9.2	11.3	12.7	11.7	13.4					
7-year	5.4	4.7	7.6	6.1	6.3	5.1	3.0	4.8	5.7	6.7	7.0	8.7	8.1	10.0	11.3	11.5						
8-year	5.2	4.7	6.9	6.1	5.9	4.8	3.0	4.8	5.4	6.5	6.3	7.4	7.9	9.2	10.7							
9-year	4.9	4.5	6.5	6.1	5.9	4.8	3.0	4.8	5.4	5.9	5.9	6.1	7.3	8.2								
10-year	4.9	4.4	6.1	5.2	5.9	4.8	3.0	4.8	5.1	5.9	5.0	6.1	7.3									
11-year	4.9	4.3	5.9	4.7	5.3	4.3	3.0	4.8	4.8	5.9	5.0	5.5										
12-year	4.8	4.3	5.9	4.7	5.3	3.9	3.0	4.8	4.8	5.7	5.0											
13-year	4.4	4.2	5.7	4.7	5.3	3.9	3.0	4.5	4.8	5.4												
14-year	4.4	4.2	5.7	4.6	4.9	3.6	3.0	4.5	4.8													
15-year	4.4	4.2	5.7	4.6	4.5	2.7	3.0	4.5														
16-year	4.3	4.1	5.7	4.6	4.5	2.7	3.0															
17-year	4.2	3.9	5.7	4.6	4.5	2.7																
18-year	3.8	3.7	5.6	4.2	4.5																	
19-year	3.8	3.3	5.6	4.2																		
20-year	3.8	3.3	5.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 15.15  
Cancer of the Lung and Bronchus (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	36.6	38.5	39.8	40.2	41.0	40.9	40.6	40.8	41.4	40.9	41.5	41.1	42.6	42.2	41.4	41.8	42.9	43.4	44.5	45.1	45.2	46.0
2-year	21.5	22.3	22.9	23.4	23.7	24.5	24.3	24.3	25.2	25.1	25.2	25.6	26.2	26.3	25.9	26.8	26.8	28.0	28.5	29.7	30.0	
3-year	16.4	17.1	17.3	17.7	18.4	18.7	18.6	18.8	19.3	19.5	19.5	19.9	19.9	20.5	20.0	20.9	20.8	22.4	22.6	23.5		
4-year	14.0	14.5	14.7	14.9	15.6	15.8	15.8	15.9	16.6	16.3	16.7	17.1	17.0	17.7	16.9	17.6	17.9	18.9	19.4			
5-year	12.6	13.0	13.1	13.3	13.7	13.9	14.2	14.3	14.6	14.6	14.9	14.7	15.2	15.7	15.0	15.6	16.0	16.8				
6-year	11.5	11.8	11.9	12.2	12.6	12.7	13.0	13.0	13.2	13.2	13.4	13.3	13.8	14.2	13.5	14.2	14.5					
7-year	10.8	10.9	10.9	11.2	11.5	11.6	12.0	11.9	12.4	12.1	12.3	12.2	12.7	12.9	12.3	12.9						
8-year	10.1	10.0	10.1	10.3	10.6	10.7	11.0	11.1	11.5	11.2	11.3	11.2	11.9	11.9	11.4							
9-year	9.4	9.3	9.4	9.4	10.0	9.9	10.3	10.3	10.7	10.4	10.4	10.5	11.1	11.2								
10-year	8.8	8.6	8.8	8.7	9.4	9.3	9.5	9.7	10.0	9.5	9.8	10.0	10.5									
11-year	8.3	8.2	8.2	8.1	8.7	8.6	8.8	9.1	9.5	8.9	9.2	9.4										
12-year	7.9	7.7	7.7	7.5	8.2	8.2	8.2	8.3	8.9	8.3	8.4											
13-year	7.5	7.2	7.2	7.2	7.5	7.7	7.6	7.9	8.2	8.0												
14-year	7.1	6.8	6.8	6.8	6.8	7.4	7.3	7.4	7.7													
15-year	6.6	6.5	6.4	6.5	6.4	6.9	6.7	7.0														
16-year	6.4	6.0	6.1	6.1	5.9	6.5	6.4															
17-year	6.1	5.7	5.7	5.7	5.5	6.2																
18-year	5.7	5.4	5.3	5.5	5.2																	
19-year	5.4	5.0	4.9	5.2																		
20-year	5.2	4.6	4.7																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 15.16  
Cancer of the Lung and Bronchus (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	34.7	36.6	37.8	38.0	38.9	38.9	38.1	38.5	39.3	39.1	39.2	38.3	39.5	39.9	38.2	38.2	40.5	40.5	41.0	42.2	41.9	42.9
2-year	19.9	20.6	21.3	21.2	21.8	22.9	22.1	21.8	23.4	23.4	23.4	23.1	23.2	24.1	22.6	23.3	24.0	25.1	25.0	27.2	26.3	
3-year	14.9	15.4	15.8	15.8	16.8	17.3	16.3	16.4	17.5	18.0	17.9	17.5	16.8	18.4	17.4	18.1	18.4	19.7	19.5	20.9		
4-year	12.6	13.0	13.2	13.1	13.8	14.4	13.7	13.7	15.0	14.8	15.0	14.8	14.1	15.8	14.4	15.1	15.6	16.7	16.5			
5-year	11.2	11.5	11.7	11.5	12.1	12.7	12.2	12.3	13.2	13.2	13.4	12.5	12.6	14.1	12.8	13.5	14.1	14.9				
6-year	10.2	10.4	10.7	10.5	11.2	11.4	11.1	11.0	11.9	11.8	11.9	11.0	11.3	12.6	11.6	12.1	12.6					
7-year	9.5	9.5	9.8	9.7	10.0	10.5	10.2	10.0	11.3	10.7	10.8	10.1	10.4	11.3	10.4	11.1						
8-year	8.9	8.7	9.0	8.8	9.1	9.4	9.3	9.3	10.3	9.9	10.0	9.1	9.6	10.7	9.6							
9-year	8.3	8.0	8.5	7.9	8.6	8.7	8.7	8.7	9.6	9.2	9.2	8.5	8.8	9.9								
10-year	7.7	7.4	7.9	7.2	8.2	8.0	8.1	8.2	8.8	8.3	8.6	8.1	8.4									
11-year	7.2	7.1	7.4	6.6	7.5	7.4	7.3	7.7	8.4	7.5	8.0	7.5										
12-year	6.8	6.6	6.8	6.1	7.1	7.0	6.7	6.9	8.0	6.9	7.2											
13-year	6.4	6.2	6.4	5.8	6.4	6.5	6.1	6.7	7.2	6.7												
14-year	6.1	5.8	6.0	5.5	5.8	6.3	5.7	6.1	6.9													
15-year	5.6	5.5	5.7	5.1	5.3	6.0	5.3	5.8														
16-year	5.4	5.1	5.3	4.7	4.9	5.5	5.1															
17-year	5.1	4.7	4.9	4.5	4.7	5.3																
18-year	4.8	4.5	4.5	4.3	4.4																	
19-year	4.5	4.2	4.2	4.2																		
20-year	4.3	3.8	3.9																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 15.17  
Cancer of the Lung and Bronchus (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	41.7	42.6	43.6	43.6	44.4	44.0	44.4	44.0	44.5	43.3	44.5	44.8	46.3	45.1	45.4	46.1	45.9	46.7	48.4	48.3	48.8	49.4
2-year	25.7	26.0	25.8	26.8	26.6	26.8	27.6	27.7	27.7	27.3	27.4	28.9	29.9	29.0	30.2	30.8	30.1	31.3	32.4	32.7	34.0	
3-year	20.4	20.7	19.8	20.7	20.9	20.8	21.9	21.9	21.8	21.4	21.5	23.0	23.7	23.1	23.1	24.4	23.8	25.6	25.9	26.5		
4-year	17.9	17.8	17.3	17.7	18.3	18.0	18.8	19.0	18.9	18.4	18.7	20.0	20.6	20.0	19.9	20.6	20.6	21.5	22.5			
5-year	16.2	16.0	15.4	16.1	16.1	15.7	17.2	17.0	16.5	16.6	16.6	17.5	18.4	17.7	17.8	18.1	18.2	18.8				
6-year	15.0	14.7	14.2	14.7	14.8	14.6	15.8	15.8	15.0	15.1	15.2	16.1	16.9	16.1	15.8	16.5	16.7					
7-year	14.0	13.7	12.9	13.4	13.7	13.4	14.5	14.5	13.9	14.1	14.1	14.9	15.6	14.9	14.6	15.1						
8-year	13.2	12.8	12.0	12.7	12.8	12.6	13.6	13.7	13.1	13.0	13.0	13.9	14.7	13.5	13.7							
9-year	12.4	11.9	11.1	11.8	12.0	11.8	12.7	12.6	12.3	11.9	12.0	13.0	13.9	12.7								
10-year	11.6	11.1	10.5	11.0	11.3	11.1	11.5	11.7	11.6	11.0	11.3	12.3	13.1									
11-year	11.1	10.6	9.8	10.3	10.6	10.4	11.0	10.9	11.0	10.6	10.6	11.7										
12-year	10.7	9.9	9.2	9.5	9.9	9.9	10.2	10.1	10.2	9.9	9.9											
13-year	10.3	9.3	8.8	9.2	9.1	9.3	9.9	9.5	9.5	9.5												
14-year	9.7	8.9	8.2	8.7	8.4	8.9	9.5	9.1	8.7													
15-year	9.2	8.4	7.8	8.5	7.9	8.2	8.7	8.6														
16-year	8.9	8.0	7.4	8.1	7.3	8.0	8.3															
17-year	8.4	7.7	7.0	7.5	6.8	7.5																
18-year	8.1	7.2	6.7	7.1	6.3																	
19-year	7.7	6.6	6.2	6.6																		
20-year	7.4	6.3	6.1																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 16.9  
Melanoma of the Skin (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	94.6	95.9	96.5	96.7	97.2	96.3	96.9	97.1	97.5	97.7	96.9	96.8	97.7	98.0	97.5	98.0	97.9	97.8	98.1	97.8	97.9	98.0
2-year	90.0	91.8	93.4	93.8	94.2	93.8	94.5	94.0	94.5	95.7	94.5	94.6	95.7	96.2	95.7	95.7	96.2	96.3	96.3	96.3	96.3	96.3
3-year	86.2	88.1	90.8	91.8	92.1	91.7	92.6	91.7	92.8	94.0	92.3	92.8	94.1	94.7	94.2	94.4	95.0	94.8	95.0	94.8		
4-year	83.5	85.5	89.2	89.9	90.8	89.7	90.9	90.0	91.4	93.2	91.3	91.5	93.2	93.2	92.9	93.3	93.7	93.6	94.2			
5-year	82.0	83.6	87.8	89.3	89.8	88.8	90.2	88.9	90.2	92.3	90.0	91.0	92.4	92.2	92.2	92.9	93.3	93.2				
6-year	80.3	82.4	86.7	88.2	88.6	88.2	89.7	87.9	90.1	91.5	89.4	90.4	91.7	91.7	91.4	92.5	92.5					
7-year	79.0	81.7	85.9	87.7	87.8	87.9	88.6	87.4	90.1	91.2	88.8	90.4	91.6	91.5	91.1	92.0						
8-year	78.0	80.9	85.3	86.9	87.0	87.4	88.2	86.4	90.0	90.9	87.9	90.0	91.1	91.4	91.0							
9-year	77.2	80.2	85.0	86.8	86.8	87.1	88.0	86.2	89.6	90.4	87.7	89.9	90.9	91.4								
10-year	76.7	79.6	84.9	86.6	86.8	86.8	87.6	86.0	89.5	90.0	87.7	89.3	90.9									
11-year	76.1	79.3	84.4	86.3	86.5	86.5	87.4	86.0	89.2	89.6	87.7	89.3										
12-year	75.6	79.1	84.2	86.3	86.1	86.4	87.4	85.6	89.2	89.4	87.2											
13-year	75.3	78.6	84.1	86.3	85.9	86.2	87.4	85.4	88.8	89.4												
14-year	75.1	78.4	83.9	86.2	85.3	86.2	87.3	85.4	88.7													
15-year	74.9	78.2	83.9	85.8	85.0	86.2	87.3	85.4														
16-year	74.6	78.1	83.8	85.8	84.7	86.2	87.3															
17-year	74.6	77.8	83.8	85.8	84.7	86.2																
18-year	74.5	77.8	83.8	85.8	84.7																	
19-year	74.5	77.8	83.8	85.8																		
20-year	74.5	77.8	83.8																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 16.10  
Melanoma of the Skin (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	92.8	94.4	95.4	95.9	96.2	96.0	96.3	96.9	96.5	96.8	96.4	96.4	97.3	97.5	96.6	97.1	97.1	97.7	97.8	97.4	97.7	97.3
2-year	86.8	88.9	91.5	91.9	93.2	92.9	93.5	92.5	92.3	94.4	93.5	94.0	94.4	95.5	93.9	94.3	94.9	95.8	95.9	95.7	95.6	
3-year	82.4	84.4	88.4	89.6	90.8	90.5	91.0	89.7	90.0	92.3	90.8	92.0	92.6	93.7	91.6	92.7	93.5	93.1	94.1	93.8		
4-year	79.1	81.4	86.3	87.5	89.4	87.7	89.4	87.8	88.1	91.3	89.8	90.5	91.4	92.1	90.0	91.1	91.9	91.9	92.8			
5-year	77.7	78.6	84.5	86.3	88.4	86.4	88.7	86.8	86.6	90.5	88.1	90.1	90.5	91.3	89.2	90.4	91.8	91.1				
6-year	75.6	77.4	83.2	84.9	87.2	85.5	87.7	86.0	86.3	89.6	87.2	89.6	89.1	90.4	88.2	90.2	90.8					
7-year	73.9	76.8	82.2	84.6	86.3	85.2	85.8	85.1	86.3	89.5	87.2	89.6	89.1	90.4	88.0	89.8						
8-year	72.7	76.1	81.6	83.9	85.5	85.0	85.4	83.5	86.0	89.0	86.3	89.4	88.7	90.3	88.0							
9-year	71.9	75.2	81.2	83.9	84.9	84.1	85.0	83.2	85.6	88.5	85.9	88.8	88.5	90.3								
10-year	71.3	74.7	81.2	83.8	84.9	83.8	84.2	83.1	85.5	88.1	85.6	88.4	88.5									
11-year	70.4	74.3	81.1	83.5	84.9	83.4	83.6	82.9	85.5	87.8	85.6	88.4										
12-year	69.9	74.1	80.9	83.5	84.9	83.2	83.4	82.3	85.5	87.8	84.9											
13-year	69.6	73.8	80.8	83.5	84.9	83.0	83.4	82.1	85.5	87.8												
14-year	69.6	73.5	80.4	82.8	84.9	82.9	83.1	82.1	85.5													
15-year	69.5	73.5	80.3	82.2	84.9	82.9	83.1	82.1														
16-year	69.2	73.5	80.0	82.2	84.9	82.9	83.1															
17-year	69.1	73.4	80.0	82.1	84.9	82.9																
18-year	69.1	73.4	80.0	82.1	84.9																	
19-year	69.1	73.4	80.0	82.1																		
20-year	69.1	73.4	80.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).



Table 16.11  
Melanoma of the Skin (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	96.4	97.5	97.7	97.6	98.4	96.7	97.7	97.4	98.6	98.8	97.4	97.3	98.2	98.5	98.5	99.1	98.9	98.0	98.3	98.2	98.2	98.9
2-year	93.2	94.9	95.4	95.9	95.4	94.8	95.7	95.8	97.0	97.4	95.5	95.4	97.0	97.0	97.7	97.2	97.8	97.0	96.8	97.0	97.2	
3-year	89.9	92.0	93.5	94.2	93.5	93.2	94.4	94.2	96.0	96.1	94.1	93.7	95.8	95.8	97.2	96.4	96.7	96.7	96.0	95.8		
4-year	87.9	89.8	92.4	92.8	92.3	92.0	92.5	92.7	95.1	95.3	93.0	92.7	95.4	94.6	96.2	95.8	95.7	95.6	95.9			
5-year	86.4	88.6	91.4	92.5	91.3	91.4	92.1	91.5	94.3	94.5	92.2	91.9	94.7	93.4	95.7	95.5	95.1	95.4				
6-year	84.9	87.5	90.4	92.0	90.1	91.4	91.8	90.4	94.3	93.7	91.9	91.4	94.5	93.3	95.2	95.0	94.4					
7-year	84.0	86.6	89.9	91.1	89.5	90.9	91.8	90.2	94.3	93.3	90.7	91.1	94.4	92.7	94.7	94.6						
8-year	83.1	85.9	89.3	90.2	88.8	90.2	91.5	90.1	94.3	93.2	89.9	90.6	93.8	92.7	94.2							
9-year	82.4	85.4	89.0	89.9	88.6	90.2	91.3	89.8	94.1	92.7	89.7	90.6	93.7	92.7								
10-year	81.9	84.7	88.9	89.6	88.6	90.0	91.3	89.6	94.0	92.3	89.7	90.4	93.6									
11-year	81.6	84.3	88.0	89.2	87.6	89.9	91.3	89.6	93.1	91.8	89.7	90.4										
12-year	81.1	84.1	87.8	89.2	86.8	89.9	91.3	89.6	93.1	91.0	89.7											
13-year	80.9	83.6	87.5	89.2	86.5	89.5	91.3	89.1	92.2	91.0												
14-year	80.3	83.5	87.5	89.2	84.8	89.5	91.3	89.1	92.0													
15-year	80.1	82.9	87.5	89.2	84.6	89.5	91.3	89.1														
16-year	79.9	82.6	87.5	89.2	84.5	89.5	91.3															
17-year	79.9	82.3	87.5	89.2	84.5	89.5																
18-year	79.8	82.1	87.5	89.2	84.5																	
19-year	79.8	82.1	87.5	89.2																		
20-year	79.8	82.1	87.5																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 17.5  
MesotheliomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																						
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
1-year	37.3	38.8	37.8	40.6	35.4	40.3	33.0	38.9	36.8	42.9	36.8	39.2	33.6	41.5	42.7	33.0	48.5	39.8	44.7	45.9	43.9	43.7	
2-year	16.5	19.2	17.0	15.8	13.7	19.6	13.7	17.7	13.8	26.1	16.9	22.1	18.0	15.1	24.7	16.9	22.3	16.7	21.1	28.7	20.0		
3-year	10.9	11.9	10.9	10.5	11.5	12.4	8.7	12.3	11.8	17.6	10.3	15.3	12.0	10.5	14.5	6.9	16.4	9.9	13.7	18.7			
4-year	9.4	8.4	8.1	5.9	8.6	10.1	6.1	9.6	9.0	11.9	7.8	12.6	9.7	7.0	10.9	4.7	13.0	8.1	10.2				
5-year	7.9	7.0	6.6	5.9	8.3	8.8	4.0	8.2	7.2	10.2	6.2	12.6	7.9	3.1	10.5	4.3	8.4	5.4					
6-year	6.7	5.7	5.2	4.1	7.4	6.4	4.0	7.9	6.1	10.0	4.7	11.4	7.0	1.3	9.4	3.8	7.4						
7-year	6.2	5.1	4.5	3.0	7.0	6.0	4.0	7.4	5.1	8.6	3.0	11.0	6.5	1.3	7.2	3.2							
8-year	5.6	4.9	4.0	2.5	6.3	6.0	3.0	6.6	5.1	6.6	3.0	8.5	6.5	0.7	7.2								
9-year	5.5	4.6	3.8	2.5	4.7	5.7	3.0	6.6	3.7	5.0	2.5	8.5	5.1	0.7									
10-year	5.2	4.3	3.4	2.5	4.7	4.7	2.5	6.4	3.1	4.4	2.5	7.7	5.1										
11-year	5.0	4.3	3.4	2.5	4.4	4.3	2.5	6.4	3.1	4.0	1.9	7.4											
12-year	4.9	4.1	3.4	2.5	4.4	3.7	2.5	5.5	2.5	3.5	1.9												
13-year	4.7	4.0	3.3	2.5	4.4	3.7	2.0	5.0	2.5	3.0													
14-year	4.7	4.0	3.3	2.5	4.4	3.7	2.0	5.0	2.5														
15-year	4.5	3.5	3.3	2.5	3.0	3.7	2.0	5.0															
16-year	4.1	3.5	3.3	2.5	3.0	3.7	2.0																
17-year	4.0	3.5	3.2	2.3	3.0	3.7																	
18-year	4.0	3.5	3.2	2.3	1.7																		
19-year	4.0	3.5	3.2	2.3																			
20-year	4.0	3.5	3.1																				

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 17.6  
MesotheliomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	35.8	37.3	36.0	37.0	31.2	36.0	35.1	34.5	36.1	46.3	34.0	35.1	31.7	41.9	38.8	32.2	47.9	38.1	44.4	42.2	42.4	40.9
2-year	11.9	16.3	14.6	12.6	8.5	14.9	13.6	11.3	11.1	25.8	14.3	18.2	13.7	14.2	21.2	16.1	19.3	15.4	17.4	22.9	17.5	
3-year	6.6	9.5	8.5	7.6	7.8	7.7	7.7	6.8	9.0	16.2	8.7	12.4	6.8	9.0	11.6	6.1	14.0	8.9	9.0	12.2		
4-year	5.5	6.7	6.0	4.3	4.2	6.0	5.0	3.8	5.9	11.3	6.2	10.2	4.4	6.7	8.2	3.1	12.8	6.6	6.4			
5-year	4.0	5.1	4.7	4.3	4.2	4.3	2.2	3.3	4.2	10.3	4.2	10.2	3.8	4.0	7.6	3.1	7.2	3.9				
6-year	3.3	4.3	3.2	3.3	4.2	2.6	2.2	3.3	3.5	10.3	2.2	8.2	3.8	1.7	6.8	2.4	6.5					
7-year	2.9	3.4	2.5	1.7	4.2	2.1	2.2	2.7	2.2	9.6	1.5	8.2	3.8	1.7	4.4	1.7						
8-year	2.3	3.3	2.5	0.9	4.2	2.1	0.8	2.1	2.2	6.8	1.5	4.8	3.8	0.9	4.4							
9-year	2.1	3.1	2.3	0.9	2.8	2.1	0.8	2.1	0.9	5.3	0.8	4.8	2.9	0.9								
10-year	2.1	3.1	1.9	0.9	2.8	1.5	0.8	2.1	0.0	4.5	0.8	4.1	2.9									
11-year	2.1	3.1	1.9	0.9	2.2	1.5	0.8	2.1	0.0	3.9	0.0	4.1										
12-year	2.0	3.0	1.9	0.9	2.2	1.5	0.8	1.4	0.0	3.2	0.0											
13-year	1.7	2.8	1.9	0.9	2.2	1.5	0.8	1.4	0.0	2.4												
14-year	1.7	2.8	1.7	0.9	2.2	1.5	0.8	1.4	0.0													
15-year	1.1	2.5	1.6	0.9	2.2	1.5	0.8	1.4														
16-year	1.1	2.5	1.6	0.9	2.2	1.5	0.8															
17-year	1.1	2.4	1.6	0.0	0.0	1.5																
18-year	1.1	2.4	1.6	0.0	0.0																	
19-year	1.1	2.4	1.6	0.0																		
20-year	1.1	2.4	1.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 17.7  
MesotheliomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	42.4	43.7	43.9	53.0	50.6	59.9	25.6	57.1	38.9	32.3	48.7	54.6	41.7	39.6	54.2	35.7	50.1	45.1	45.8	56.3	47.9	55.4
2-year	31.1	29.4	25.0	27.0	32.7	40.7	14.2	44.0	23.2	26.8	28.0	37.1	37.0	19.1	34.9	19.4	31.3	20.8	33.3	45.0	26.7	
3-year	24.4	20.0	18.9	20.6	23.9	33.1	12.0	34.9	21.2	21.3	17.2	26.0	34.9	16.8	22.9	9.8	23.0	12.6	29.2	36.6		
4-year	21.5	14.4	15.4	11.6	23.9	28.4	9.8	33.1	19.4	13.7	14.4	21.7	32.9	8.6	18.9	9.8	14.0	12.6	22.8			
5-year	20.1	13.3	13.4	11.6	22.5	28.4	9.8	28.6	17.7	9.9	14.4	21.7	25.6	0.0	18.9	7.9	11.9	10.2				
6-year	17.1	10.6	11.9	7.1	17.9	23.4	9.8	26.7	15.2	7.9	14.4	21.7	20.3	0.0	17.0	7.9	9.8					
7-year	16.4	10.6	10.9	7.1	15.6	23.4	9.8	26.7	15.2	6.0	8.9	20.6	17.6	0.0	15.1	7.9						
8-year	15.8	10.4	8.7	7.1	13.2	23.4	9.8	25.0	15.2	6.0	8.9	20.6	17.6	0.0	15.1							
9-year	15.8	9.5	8.7	7.1	11.3	21.5	9.8	25.0	12.9	4.0	8.9	20.6	15.0	0.0								
10-year	14.5	7.7	8.4	7.1	11.3	19.0	8.6	23.8	12.9	4.0	8.9	20.2	15.0									
11-year	13.8	7.7	8.4	7.1	11.3	16.3	8.6	23.8	12.9	4.0	8.9	18.1										
12-year	13.8	7.7	8.2	7.1	11.3	13.1	8.6	22.1	10.6	4.0	8.9											
13-year	13.8	7.7	8.2	7.1	11.3	13.1	6.0	19.5	10.6	4.0												
14-year	13.8	7.7	8.2	7.1	11.3	13.1	6.0	19.5	10.6													
15-year	13.8	6.8	8.2	7.1	7.0	13.1	6.0	19.5														
16-year	12.9	6.8	8.2	7.1	7.0	13.1	6.0															
17-year	12.3	6.8	8.2	7.1	7.0	13.1																
18-year	12.3	6.8	7.9	7.1	4.0																	
19-year	12.3	6.8	7.9	7.1																		
20-year	12.3	6.5	7.4																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 18.9  
MyelomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	68.8	70.7	73.0	71.4	75.5	69.5	72.8	73.4	74.7	71.1	74.8	75.3	74.2	73.0	73.2	74.9	77.6	74.7	76.1	73.9	76.5	76.5
2-year	52.6	55.0	58.5	56.5	60.0	56.6	58.8	59.1	59.2	58.3	61.1	62.6	60.2	60.2	60.0	63.2	65.2	66.5	64.1	65.7	65.6	
3-year	40.3	43.2	45.9	44.0	48.0	45.7	46.6	48.3	48.1	45.8	49.0	51.5	48.2	49.9	49.6	53.4	55.8	58.2	56.3	57.1		
4-year	31.7	34.5	36.2	37.8	38.2	35.1	37.8	37.6	40.5	38.2	38.9	39.8	38.6	41.0	43.9	46.6	47.6	50.4	49.2			
5-year	25.5	27.1	27.7	30.2	31.0	27.5	31.1	31.3	33.6	31.8	31.8	34.0	31.1	34.1	37.1	39.8	41.7	44.5				
6-year	20.2	21.3	22.3	24.1	24.8	21.3	26.8	26.2	28.3	27.1	26.4	29.5	26.7	28.8	32.3	34.7	37.1					
7-year	16.4	16.9	17.7	18.8	21.1	18.7	22.2	21.0	23.6	23.8	22.5	23.9	22.9	24.7	28.7	31.2						
8-year	13.6	13.5	15.1	17.0	17.5	15.8	17.9	18.3	20.4	21.4	19.8	21.7	19.4	22.0	25.2							
9-year	11.5	11.4	12.0	15.3	16.1	13.8	16.0	15.5	18.2	19.9	17.1	19.0	17.7	19.6								
10-year	10.1	9.7	10.5	13.6	14.4	11.9	14.5	13.7	17.0	17.3	15.4	17.1	15.6									
11-year	8.8	8.7	9.2	12.4	12.8	10.8	13.4	13.1	15.8	15.7	14.1	16.0										
12-year	7.9	7.7	8.4	11.5	11.1	9.0	11.9	11.5	14.2	14.9	12.9											
13-year	6.7	6.8	7.6	10.8	10.3	8.2	10.6	11.1	13.4	13.3												
14-year	6.1	6.1	7.0	9.7	9.5	7.7	9.9	10.4	12.2													
15-year	5.4	5.7	6.3	9.1	9.1	7.6	9.0	9.9														
16-year	4.9	5.2	5.7	8.7	8.3	7.4	8.2															
17-year	4.6	4.7	5.4	8.7	8.0	6.6																
18-year	4.0	4.4	5.1	8.0	7.4																	
19-year	3.8	3.8	5.0	7.3																		
20-year	3.3	3.5	4.7																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 18.10  
MyelomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	68.7	70.8	73.9	71.0	76.6	73.2	70.5	74.5	75.4	71.9	74.5	76.4	76.1	74.4	74.8	75.6	77.5	76.7	76.2	73.6	77.0	75.6
2-year	52.8	55.4	60.0	56.8	62.0	60.1	57.6	59.7	59.9	59.7	61.4	63.8	61.7	62.2	62.1	62.1	65.9	68.1	63.7	66.4	66.8	
3-year	39.6	43.1	46.6	44.9	49.2	48.9	44.9	47.4	49.6	46.0	50.8	54.0	51.4	52.5	51.3	52.2	56.8	60.2	55.7	57.5		
4-year	31.2	34.1	37.2	37.2	39.6	36.5	36.1	37.3	41.8	39.8	41.0	43.6	43.0	43.2	46.4	46.7	48.1	51.3	49.0			
5-year	24.7	26.0	28.6	30.4	32.8	29.1	29.9	30.5	36.4	33.8	32.5	38.1	34.8	35.9	39.8	41.4	42.3	45.5				
6-year	19.4	21.0	23.7	22.9	26.9	22.4	26.8	27.8	31.6	29.6	26.6	33.7	29.9	29.9	34.9	35.8	37.8					
7-year	15.6	17.1	19.4	18.5	22.6	19.5	22.1	23.2	26.1	25.8	22.4	27.6	25.9	25.6	31.2	32.6						
8-year	13.1	13.8	16.3	17.7	19.6	17.6	17.5	20.2	22.2	24.1	20.3	25.1	21.5	23.0	27.3							
9-year	11.1	11.9	13.0	16.1	18.0	15.9	16.0	16.8	20.0	21.4	18.3	22.8	19.1	19.2								
10-year	9.5	9.9	11.4	14.6	16.0	13.5	14.9	14.8	19.0	18.5	16.4	21.1	17.3									
11-year	7.7	9.0	10.0	12.5	14.1	13.0	13.4	13.9	17.8	17.2	14.9	20.3										
12-year	7.1	7.7	9.4	12.3	12.4	11.6	11.4	11.4	15.5	16.6	14.3											
13-year	5.9	6.9	8.6	11.6	11.8	10.7	10.0	10.9	14.9	14.4												
14-year	5.4	6.1	8.0	10.5	11.1	10.4	9.6	9.9	13.2													
15-year	4.7	5.7	7.3	10.5	10.5	10.2	8.4	9.2														
16-year	4.5	5.2	6.8	10.2	10.0	10.2	7.9															
17-year	4.1	4.9	6.5	9.9	9.6	9.1																
18-year	3.6	4.5	6.2	8.8	8.7																	
19-year	3.4	4.2	6.2	7.8																		
20-year	2.7	3.9	6.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 18.11  
MyelomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	69.0	70.5	72.2	71.8	74.2	65.6	75.7	72.5	74.1	70.2	75.0	74.1	72.0	71.4	71.4	74.0	77.7	72.4	76.1	74.3	75.8	77.4
2-year	52.4	54.7	56.9	56.1	57.6	52.9	60.2	58.5	58.4	56.9	60.8	61.3	58.5	57.8	57.5	64.6	64.4	64.7	64.5	65.1	63.9	
3-year	41.0	43.4	45.1	43.0	46.7	42.4	48.7	49.1	46.4	45.5	47.1	48.7	44.5	46.8	47.6	54.8	54.7	55.9	57.1	56.6		
4-year	32.2	35.0	35.2	38.4	36.6	33.5	39.8	37.9	39.1	36.5	36.7	35.5	33.6	38.3	41.1	46.5	46.9	49.3	49.5			
5-year	26.3	28.3	26.8	29.9	28.9	25.9	32.5	31.9	30.6	29.8	31.1	29.5	26.9	31.9	34.1	37.9	41.1	43.3				
6-year	21.1	21.5	20.9	25.4	22.4	20.2	26.8	24.7	24.6	24.4	26.1	24.7	23.0	27.6	29.4	33.4	36.2					
7-year	17.2	16.7	16.1	19.1	19.3	17.9	22.4	19.1	20.9	21.6	22.5	20.0	19.6	23.7	25.8	29.4						
8-year	14.2	13.1	13.8	16.2	14.9	14.0	18.5	16.4	18.3	18.6	19.2	18.0	17.1	20.7	22.9							
9-year	12.0	10.9	11.0	14.4	13.9	11.7	16.0	14.3	16.1	18.1	15.7	14.8	16.0	20.2								
10-year	10.7	9.4	9.6	12.6	12.5	10.3	13.9	12.7	14.9	16.0	14.4	12.8	13.8									
11-year	9.9	8.5	8.4	12.3	11.1	8.5	13.4	12.3	13.4	14.2	13.2	11.5										
12-year	8.8	7.6	7.3	10.7	9.5	6.3	12.5	11.5	12.5	13.1	11.1											
13-year	7.5	6.7	6.6	9.9	8.5	5.7	11.2	11.1	11.6	12.0												
14-year	6.9	6.0	6.0	8.9	7.5	5.0	10.2	10.9	11.0													
15-year	6.0	5.7	5.3	7.7	7.3	5.0	9.7	10.6														
16-year	5.3	5.1	4.6	7.2	6.3	4.2	8.6															
17-year	5.1	4.6	4.4	7.2	6.0	4.0																
18-year	4.5	4.2	4.0	7.2	5.9																	
19-year	4.2	3.4	3.7	6.6																		
20-year	3.8	3.2	3.4																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 19.9  
Non-Hodgkin Lymphoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	71.1	73.8	72.0	70.7	70.3	71.4	71.8	71.2	70.7	73.7	76.4	77.2	77.9	77.5	79.2	81.4	80.6	81.4	81.7	81.3	81.7	84.4
2-year	60.6	64.2	63.1	61.4	61.3	62.5	63.7	63.3	62.2	66.0	68.6	70.2	71.3	71.5	73.5	76.1	75.8	76.0	77.1	76.9	77.3	
3-year	54.9	58.7	58.4	56.7	56.8	58.1	58.9	58.8	57.6	62.2	65.0	66.4	67.0	67.8	70.3	73.2	73.2	73.6	74.5	74.4		
4-year	50.8	54.7	54.5	52.5	54.0	54.7	55.9	55.4	54.8	58.8	62.1	63.8	64.6	65.5	68.2	71.2	71.8	71.4	72.5			
5-year	47.0	51.0	51.2	49.9	51.2	51.5	53.1	53.0	52.1	56.1	59.6	61.2	62.3	63.7	66.2	69.4	70.1	70.0				
6-year	43.7	47.7	48.5	47.4	48.4	49.0	50.9	51.0	49.9	54.2	57.3	58.4	59.5	62.1	64.8	68.1	68.4					
7-year	41.0	45.2	46.0	44.5	45.5	46.6	49.2	49.1	48.1	52.3	55.2	56.5	57.7	60.3	63.1	66.2						
8-year	38.6	43.0	43.6	42.4	43.1	45.0	47.2	47.3	46.3	50.4	53.0	55.3	56.2	58.7	61.5							
9-year	36.6	40.9	41.6	40.9	41.5	43.9	45.7	45.6	45.1	49.3	51.7	53.5	54.6	57.3								
10-year	35.2	39.0	39.9	39.8	40.3	42.3	44.3	43.9	44.4	48.1	50.4	52.3	53.4									
11-year	33.6	37.5	38.5	38.0	38.9	41.0	42.9	42.4	43.3	46.5	48.8	51.1										
12-year	32.5	36.0	37.0	36.5	38.0	40.0	41.8	41.4	42.1	45.6	48.2											
13-year	31.5	34.8	35.7	35.3	37.4	38.9	40.2	39.9	41.3	44.6												
14-year	30.3	33.6	34.7	34.3	36.2	37.6	39.0	38.6	40.4													
15-year	29.0	32.3	33.6	33.9	34.9	36.9	38.6	37.8														
16-year	28.0	31.2	32.8	32.7	34.1	36.2	37.4															
17-year	27.4	30.4	32.0	32.3	33.1	35.6																
18-year	26.8	29.6	31.3	31.8	32.2																	
19-year	26.1	28.8	30.5	31.1																		
20-year	25.3	28.0	30.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).



Table 19.10  
Non-Hodgkin Lymphoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	70.3	73.5	69.7	66.6	66.8	68.4	68.7	68.0	67.2	71.7	75.6	75.4	76.6	76.6	78.2	81.1	80.1	80.5	81.1	81.4	81.5	83.8
2-year	59.8	64.2	60.7	57.6	57.2	58.2	59.7	60.0	58.8	63.7	67.3	68.4	69.6	70.4	71.5	75.4	74.9	75.5	76.0	76.4	77.7	
3-year	53.7	58.4	55.9	52.4	52.6	54.3	54.9	55.1	54.5	60.0	63.9	64.7	64.4	66.6	68.3	72.3	72.7	72.8	72.8	73.4		
4-year	49.6	54.1	51.7	48.5	49.9	50.6	51.8	51.7	51.7	56.4	61.0	62.2	61.7	63.7	65.8	70.2	71.0	70.1	70.8			
5-year	45.6	50.4	48.4	46.1	47.7	46.6	48.9	49.1	48.8	53.7	58.4	59.4	59.3	61.6	63.8	68.1	69.0	68.8				
6-year	42.6	46.9	45.6	44.2	45.0	44.8	46.9	47.5	46.8	51.2	56.2	56.6	56.3	60.2	62.1	66.8	67.0					
7-year	40.0	44.3	43.3	41.4	42.2	42.6	46.1	45.9	45.1	49.2	54.3	54.7	54.6	58.4	60.2	65.4						
8-year	37.9	42.2	41.1	39.7	39.8	41.2	44.3	43.9	43.2	47.4	52.0	53.5	53.6	56.5	58.5							
9-year	35.7	40.2	39.2	38.2	38.2	40.2	43.0	42.0	42.2	46.4	50.5	51.5	52.2	54.9								
10-year	34.2	38.1	37.7	37.3	37.2	38.6	42.2	40.6	41.7	45.5	49.1	50.6	51.4									
11-year	32.9	36.4	36.4	36.0	36.2	38.1	40.6	39.5	40.8	43.8	47.6	49.6										
12-year	32.0	35.2	35.1	34.5	35.5	37.4	40.0	38.7	39.4	42.7	47.2											
13-year	30.8	34.2	34.2	33.3	34.9	36.2	38.6	37.4	38.8	41.9												
14-year	29.5	33.1	33.2	32.4	33.9	34.4	37.3	36.5	37.6													
15-year	28.1	32.0	32.3	32.2	32.4	33.7	37.1	35.8														
16-year	27.1	30.8	31.6	30.6	31.7	33.1	35.8															
17-year	26.5	30.0	30.8	30.3	31.0	32.2																
18-year	26.1	29.3	30.2	29.9	30.4																	
19-year	25.6	28.8	29.3	29.1																		
20-year	25.1	27.9	29.1																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 19.11  
Non-Hodgkin Lymphoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	71.9	74.2	74.7	75.8	74.9	75.1	75.8	75.4	75.3	76.3	77.3	79.2	79.5	78.5	80.4	81.7	81.3	82.4	82.4	81.1	82.0	85.1
2-year	61.5	64.2	65.9	66.2	66.7	67.9	68.8	67.7	66.6	68.9	70.1	72.1	73.3	72.8	75.8	76.9	76.7	76.6	78.5	77.4	76.8	
3-year	56.1	59.1	61.4	61.9	62.3	62.9	64.0	63.6	61.6	65.1	66.3	68.3	70.2	69.2	72.7	74.2	73.9	74.6	76.5	75.5		
4-year	52.1	55.4	57.9	57.5	59.5	59.9	61.1	60.2	58.9	61.9	63.5	65.6	68.2	67.5	71.1	72.4	72.6	72.8	74.4			
5-year	48.5	51.7	54.6	54.6	55.8	57.5	58.4	57.9	56.6	59.1	61.1	63.1	65.8	66.4	69.1	70.9	71.4	71.4				
6-year	44.9	48.6	51.8	51.3	53.0	54.2	56.0	55.5	53.9	58.0	58.5	60.4	63.4	64.5	68.0	69.5	69.8					
7-year	42.0	46.1	49.3	48.4	49.7	51.6	53.1	53.1	52.0	56.2	56.4	58.6	61.3	62.5	66.5	67.1						
8-year	39.4	43.8	46.5	45.8	47.4	49.7	50.9	51.7	50.3	54.1	54.2	57.2	59.4	61.3	65.0							
9-year	37.6	41.7	44.4	44.3	45.8	48.5	49.1	50.3	49.0	52.9	53.1	55.8	57.6	60.3								
10-year	36.3	40.0	42.4	42.9	44.2	47.0	47.0	48.1	47.9	51.2	52.0	54.2	55.8									
11-year	34.4	38.6	41.0	40.5	42.4	44.6	45.8	46.0	46.7	49.9	50.2	52.8										
12-year	33.0	36.9	39.2	39.1	41.2	43.2	44.1	44.9	45.8	49.1	49.4											
13-year	32.3	35.4	37.6	37.8	40.5	42.2	42.3	43.2	44.8	47.9												
14-year	31.1	34.2	36.5	36.6	39.0	41.6	41.0	41.1	43.8													
15-year	30.0	32.6	35.2	35.8	38.0	40.9	40.5	40.4														
16-year	28.9	31.7	34.2	35.3	37.3	39.9	39.3															
17-year	28.3	30.8	33.6	34.8	35.8	39.6																
18-year	27.5	29.8	32.6	34.1	34.6																	
19-year	26.6	28.7	31.9	33.6																		
20-year	25.5	27.9	31.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 20.12  
Cancer of the Oral Cavity and Pharynx (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	80.3	80.0	81.3	82.1	81.8	80.3	82.0	83.7	83.4	82.2	84.2	81.5	84.2	85.3	83.3	84.2	84.5	84.8	83.7	85.7	85.1	86.6
2-year	66.3	66.3	68.0	69.9	69.5	69.0	68.9	72.1	71.9	70.3	71.4	70.4	72.6	73.7	73.3	73.2	75.1	75.6	74.2	75.5	76.2	
3-year	60.0	59.8	61.3	63.2	62.9	61.7	62.8	66.4	65.6	63.9	66.2	64.5	65.7	68.7	67.3	67.7	70.2	70.7	68.5	70.5		
4-year	56.0	55.7	57.4	59.9	58.2	57.0	58.7	62.0	61.8	60.1	62.2	60.2	62.1	64.9	62.7	64.3	67.4	66.7	65.7			
5-year	53.0	52.5	54.2	57.1	54.8	54.7	55.8	59.1	58.7	57.9	58.6	57.1	59.0	61.9	60.1	62.1	65.1	64.8				
6-year	50.6	49.8	51.7	54.9	52.2	52.2	53.4	56.3	56.1	55.5	56.1	55.8	56.2	60.0	58.2	59.9	63.8					
7-year	48.5	47.6	49.1	52.7	50.1	49.4	51.0	55.0	53.5	53.7	54.4	53.9	53.8	58.0	56.3	58.3						
8-year	46.3	46.0	47.0	51.0	48.7	47.2	48.8	51.8	51.3	51.3	53.1	52.8	52.0	55.5	53.9							
9-year	44.5	44.1	45.1	48.3	47.0	45.2	46.5	49.2	49.3	49.8	51.7	52.2	50.2	53.6								
10-year	42.4	42.4	43.5	46.5	45.3	43.7	45.1	47.0	47.8	47.7	49.9	50.9	48.9									
11-year	40.6	40.9	42.2	45.4	43.1	41.5	43.3	45.2	45.9	46.4	47.8	48.7										
12-year	39.2	39.4	40.6	44.4	40.9	40.3	42.5	44.0	44.0	45.4	45.9											
13-year	37.9	37.6	39.1	42.4	40.0	38.8	41.1	42.2	42.6	43.1												
14-year	36.8	36.7	37.8	41.1	39.0	37.3	40.1	40.3	39.8													
15-year	35.4	35.0	36.7	40.4	38.1	36.0	39.1	39.1														
16-year	34.1	33.9	35.6	40.0	36.2	35.3	38.4															
17-year	33.2	32.9	34.7	39.1	34.5	34.7																
18-year	32.1	31.9	33.3	38.1	34.0																	
19-year	31.3	30.3	32.0	37.0																		
20-year	30.1	29.8	31.4																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 20.13  
Cancer of the Oral Cavity and Pharynx For Selected Subsites (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
<u>Cancer of the Tongue</u>																						
1-year	75.3	74.3	77.7	80.3	82.2	77.1	81.9	82.0	80.1	80.0	82.4	80.2	82.1	85.4	82.2	82.1	85.2	87.4	85.0	84.9	86.1	87.0
2-year	55.3	57.8	60.7	64.3	67.2	60.2	65.5	68.5	65.6	67.2	71.1	66.3	68.4	71.8	71.1	69.5	71.7	75.9	72.4	74.0	76.2	
3-year	47.7	51.0	54.7	58.1	59.1	50.6	57.1	62.9	58.7	61.2	65.1	59.4	60.3	66.4	64.6	63.9	67.1	71.1	65.2	68.9		
4-year	43.7	47.5	50.2	55.4	53.7	46.6	54.1	58.4	56.5	57.3	61.9	55.7	57.9	63.5	61.2	61.6	64.9	66.5	61.6			
5-year	40.3	44.8	47.5	52.4	52.8	43.7	51.7	54.4	54.2	55.5	57.9	53.0	54.9	62.0	58.5	59.9	62.0	64.8				
6-year	38.4	41.9	45.3	50.3	49.3	42.2	49.7	52.0	50.6	53.0	56.2	52.5	52.9	60.2	57.2	57.4	61.6					
7-year	37.2	39.3	42.5	47.7	47.5	40.3	46.8	51.2	48.6	51.7	55.3	49.6	50.6	58.2	54.5	56.2						
8-year	35.7	37.5	40.6	45.6	46.7	38.8	44.5	47.5	46.5	49.9	53.8	48.0	50.6	55.9	50.7							
9-year	34.4	34.8	38.9	42.6	45.1	35.0	42.0	45.4	44.8	48.6	52.8	47.4	48.7	54.6								
10-year	32.6	33.4	37.7	40.7	42.3	32.6	41.5	43.8	43.5	46.5	52.3	45.6	48.1									
<u>Cancer of the Lip</u>																						
1-year	99.4	98.1	99.6	98.9	100.0	100.0	99.3	100.0	99.9	98.5	100.0	99.5	97.6	98.0	95.0	99.6	99.4	99.6	98.1	97.1	95.9	99.9
2-year	97.6	96.2	98.2	97.7	98.3	98.3	98.4	100.0	99.7	94.3	97.3	98.3	95.4	96.0	94.0	99.6	99.4	99.6	95.5	94.6	95.9	
3-year	97.3	93.8	97.3	96.2	98.3	96.4	98.2	100.0	99.1	92.7	95.4	95.7	93.9	95.2	94.0	97.7	97.7	98.5	95.1	92.6		
4-year	96.0	92.4	97.1	96.2	97.7	94.5	98.2	98.8	96.4	91.3	93.5	93.8	92.1	92.5	91.4	97.7	97.7	95.5	95.1			
5-year	94.6	90.1	96.3	94.9	95.4	93.8	96.1	98.4	95.4	88.6	90.9	91.3	90.6	90.0	90.5	95.2	97.7	94.0				
6-year	92.7	88.0	95.4	92.9	95.4	90.3	95.6	98.1	95.4	87.5	89.3	89.8	89.5	90.0	87.6	93.5	97.5					
7-year	91.5	86.6	93.6	92.2	95.4	87.2	94.0	98.0	95.4	86.5	88.3	89.6	89.0	89.9	84.8	93.5						
8-year	90.1	86.2	92.2	92.2	95.4	84.5	91.2	96.7	91.5	83.1	88.3	86.9	89.0	89.9	82.3							
9-year	88.1	85.3	91.5	90.2	95.1	83.1	89.9	94.6	86.7	82.7	88.3	86.8	88.4	86.9								
10-year	86.0	83.9	91.5	89.6	93.3	81.0	87.7	91.0	82.1	82.7	85.8	86.2	88.4									
<u>Cancer of the Oropharynx and Tonsil</u>																						
1-year	71.3	72.3	73.9	76.8	76.6	71.4	77.5	74.4	81.0	80.7	81.8	81.5	87.4	82.8	84.0	85.7	82.7	84.3	84.3	85.6	86.5	86.5
2-year	54.0	53.1	56.6	60.4	62.5	58.3	64.5	63.2	70.0	67.0	67.0	66.8	75.4	67.5	72.2	76.5	75.8	74.8	74.3	75.7	77.9	
3-year	44.5	44.7	46.1	51.5	56.9	49.5	55.3	57.0	62.6	60.6	61.8	59.9	67.6	62.2	66.9	71.7	71.3	71.1	70.4	70.7		
4-year	39.1	39.5	41.7	47.6	52.2	44.6	49.3	52.5	58.3	54.1	56.2	54.7	63.2	57.3	60.6	67.5	68.4	67.2	67.0			
5-year	35.8	35.4	38.0	42.9	45.5	41.0	45.7	49.1	54.0	52.8	51.5	52.8	60.5	53.5	59.0	64.4	64.9	64.8				
6-year	32.3	32.2	34.8	42.6	40.5	39.8	43.6	45.8	50.2	48.1	48.2	51.2	57.4	52.3	54.8	63.5	63.6					
7-year	30.3	29.6	32.1	40.5	38.8	38.3	42.5	45.0	46.5	43.8	47.2	49.6	55.3	48.6	52.9	62.6						
8-year	26.6	28.0	30.4	39.1	36.0	37.0	40.4	41.5	45.4	42.4	46.2	48.8	51.6	46.6	52.9							
9-year	23.9	25.7	28.4	35.4	33.9	35.2	37.6	40.0	44.0	41.4	44.2	48.6	51.5	44.5								
10-year	22.6	23.2	26.3	35.2	32.0	35.2	34.2	39.0	42.8	40.2	43.1	46.0	49.6									

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 20.14  
Cancer of the Oral Cavity and Pharynx (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	79.7	79.5	80.6	81.2	81.4	79.6	80.9	83.8	83.3	82.1	83.4	81.1	84.6	85.1	83.9	84.4	84.8	85.4	84.2	85.6	86.3	86.9
2-year	65.5	65.4	66.5	67.9	68.5	67.7	67.3	72.3	71.1	69.8	69.9	70.3	71.7	73.2	73.4	72.0	74.9	75.5	74.6	74.9	76.9	
3-year	59.2	58.4	59.2	61.4	61.2	59.7	60.5	65.2	65.1	62.8	64.8	63.9	64.2	67.6	67.8	65.6	69.8	69.9	68.3	69.3		
4-year	54.8	54.3	55.1	57.8	55.2	55.5	56.7	60.5	60.8	58.8	60.0	59.3	60.4	63.9	62.5	62.4	66.9	65.5	65.6			
5-year	51.9	50.9	51.6	54.9	51.7	53.2	53.8	57.9	57.8	56.7	56.4	56.1	57.5	60.9	59.8	60.6	65.0	63.8				
6-year	49.5	48.2	49.4	52.7	49.2	50.9	51.2	55.4	55.0	53.7	53.9	54.8	54.5	58.6	57.6	58.4	63.9					
7-year	47.3	46.0	47.1	50.4	46.9	47.8	48.3	54.3	52.0	51.4	52.6	52.7	52.6	56.3	55.5	57.4						
8-year	45.3	44.4	44.8	49.1	45.6	45.6	46.9	51.4	49.7	49.2	51.5	51.3	51.0	53.7	52.8							
9-year	43.4	42.7	42.9	46.5	44.2	43.6	44.5	48.6	47.6	48.1	49.8	50.4	49.4	51.6								
10-year	41.3	40.7	41.0	45.5	42.1	42.1	43.3	46.4	45.8	45.9	48.3	49.2	48.5									
11-year	39.4	39.2	39.7	44.3	40.2	40.3	41.6	44.9	43.8	44.1	46.4	47.0										
12-year	38.1	37.9	38.4	43.5	37.7	39.0	40.6	43.7	42.0	43.3	44.3											
13-year	36.9	36.4	37.0	41.4	37.1	37.5	38.8	42.0	40.4	41.0												
14-year	35.9	35.4	35.7	39.9	36.5	36.0	37.9	39.7	37.1													
15-year	34.6	34.0	34.7	39.2	35.6	34.6	37.2	38.3														
16-year	33.2	32.8	33.8	38.4	33.2	33.9	36.6															
17-year	32.3	31.8	32.8	37.6	31.9	33.1																
18-year	31.3	30.7	31.6	36.4	31.4																	
19-year	30.6	28.9	30.0	35.5																		
20-year	29.6	28.3	29.4																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 20.15  
Cancer of the Oral Cavity and Pharynx (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	81.6	81.2	82.7	83.8	82.7	81.8	84.3	83.3	83.4	82.5	86.1	82.3	83.3	85.6	82.2	83.7	83.7	83.4	82.6	85.9	82.5	85.6
2-year	68.2	68.1	70.9	74.3	71.3	71.8	72.2	71.6	73.4	71.4	74.6	70.6	74.5	74.7	72.9	75.6	75.6	75.8	73.4	77.0	74.4	
3-year	61.7	62.7	65.5	67.3	66.3	65.7	67.7	68.8	66.6	66.0	69.5	65.6	68.7	71.2	66.5	72.1	71.1	72.5	69.1	73.4		
4-year	58.8	58.7	62.1	64.4	64.1	60.2	62.8	65.2	63.6	62.7	66.9	62.0	65.8	66.9	63.1	68.4	68.5	69.5	66.1			
5-year	55.8	56.0	59.6	61.7	61.0	58.0	59.7	61.7	60.4	60.5	63.5	59.2	62.1	64.2	60.7	65.3	65.4	66.9				
6-year	53.2	53.2	56.4	59.7	58.0	54.8	58.3	58.1	58.3	59.0	60.9	57.7	59.6	63.2	59.5	62.9	63.5					
7-year	51.2	50.9	53.2	57.5	56.4	52.8	56.6	56.5	56.3	58.3	58.4	56.4	56.4	61.6	57.9	60.2						
8-year	48.7	49.3	51.6	55.1	54.6	50.7	52.8	52.8	54.4	55.5	56.8	55.8	53.9	59.6	56.2							
9-year	47.0	47.1	49.7	52.4	52.7	48.5	50.8	50.5	52.4	53.3	55.8	55.8	51.8	58.1								
10-year	45.0	45.7	48.5	49.0	51.5	47.0	48.9	48.3	51.0	51.4	53.5	54.3	49.7									
11-year	43.1	44.3	47.1	47.8	48.7	43.9	46.9	45.6	49.5	50.9	51.0	52.2										
12-year	41.6	42.6	45.2	46.4	47.3	42.6	46.5	44.8	47.6	49.7	49.3											
13-year	40.1	40.3	43.5	44.6	45.7	41.4	46.0	42.7	46.6	47.3												
14-year	38.8	39.3	42.1	43.5	44.1	39.9	44.7	41.5	44.8													
15-year	37.1	37.3	40.8	43.0	43.0	38.9	43.2	40.6														
16-year	36.1	36.2	39.4	43.0	41.9	38.2	42.3															
17-year	35.1	35.2	38.6	42.2	39.7	37.9																
18-year	33.9	34.3	36.8	41.7	38.9																	
19-year	32.8	33.2	35.9	39.8																		
20-year	31.4	32.7	35.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 21.9  
Cancer of the Ovary (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	65.0	70.4	70.5	72.0	72.6	74.5	72.3	71.9	73.7	74.4	77.0	75.5	74.2	77.3	76.4	76.2	76.2	75.8	75.8	75.6	75.7	75.6
2-year	49.3	53.5	54.4	57.3	58.4	60.0	58.4	60.7	60.4	61.8	65.6	64.4	61.9	64.6	65.4	63.9	66.2	64.8	64.4	64.1	63.9	
3-year	42.4	46.1	46.6	48.4	48.8	50.9	48.8	53.2	51.2	53.0	56.8	55.3	53.1	54.9	56.0	54.4	58.4	56.1	56.0	55.8		
4-year	38.8	41.6	41.6	43.8	43.3	46.1	42.7	46.9	46.2	46.3	49.0	50.2	47.8	47.5	50.3	48.4	49.9	49.1	49.5			
5-year	36.6	39.0	38.2	40.3	39.8	41.8	38.6	44.1	42.4	42.3	44.2	45.1	42.8	43.2	45.6	42.9	44.2	43.5				
6-year	35.2	37.0	36.1	38.3	37.3	39.2	36.4	40.5	38.8	38.9	40.6	42.3	40.2	39.8	42.8	39.4	40.8					
7-year	33.9	36.1	34.7	36.9	35.4	37.4	35.4	38.6	36.9	37.3	38.1	40.4	38.1	37.3	40.2	37.5						
8-year	33.0	34.7	33.7	35.5	34.0	36.4	34.0	37.8	35.1	36.1	36.6	38.3	36.5	35.6	37.7							
9-year	32.4	33.7	32.9	34.4	33.1	35.7	32.8	35.8	33.5	35.3	35.2	36.6	34.8	34.1								
10-year	32.0	32.9	32.3	34.0	32.3	35.3	31.7	34.8	32.7	34.6	34.2	35.6	33.9									
11-year	31.6	32.5	32.0	33.3	31.5	34.7	30.7	33.8	31.9	33.7	32.9	34.7										
12-year	31.0	31.8	31.3	33.0	30.7	33.8	30.1	33.5	31.8	33.2	31.9											
13-year	30.7	31.2	30.8	32.4	30.5	33.4	30.0	32.9	31.2	32.5												
14-year	30.2	30.8	30.3	32.1	30.3	32.9	29.9	32.6	31.2													
15-year	29.6	30.6	30.2	31.5	29.6	32.2	29.8	32.5														
16-year	29.2	30.4	30.2	31.2	29.5	31.9	29.8															
17-year	28.9	29.9	30.0	30.7	29.5	31.1																
18-year	28.8	29.8	29.8	30.5	29.5																	
19-year	28.5	29.6	29.8	29.7																		
20-year	28.4	29.3	29.5																			

<sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 22.9  
Cancer of the Pancreas (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	15.1	17.0	18.2	19.3	20.0	20.3	18.2	19.6	18.5	19.9	21.4	20.8	23.5	23.4	24.9	26.2	23.9	24.1	27.9	26.9	28.7	30.6
2-year	6.1	6.5	7.0	8.2	9.2	9.0	8.4	9.1	8.4	9.6	10.0	8.9	10.6	11.0	11.7	12.1	11.4	12.0	12.8	13.3	13.7	
3-year	3.6	4.1	4.8	5.3	6.4	5.8	5.2	6.5	5.3	6.4	7.1	5.8	6.7	7.3	7.4	8.5	7.5	8.4	8.8	9.3		
4-year	2.9	3.3	3.8	4.1	5.2	5.1	4.3	5.2	4.0	4.8	5.6	4.5	5.6	6.1	6.0	6.9	5.7	6.4	6.9			
5-year	2.5	2.9	3.3	3.7	4.6	4.6	3.5	4.8	3.5	4.2	5.0	3.8	5.0	5.0	5.0	6.2	4.8	5.4				
6-year	2.3	2.6	3.0	3.5	4.2	4.6	3.2	4.2	3.2	3.8	4.5	3.7	4.5	4.6	4.6	5.7	4.3					
7-year	2.1	2.3	2.7	3.1	3.9	4.0	3.0	3.8	3.1	3.5	4.3	3.4	4.3	4.2	4.1	5.0						
8-year	1.9	2.1	2.5	3.1	3.6	3.8	3.0	3.3	2.7	3.4	4.1	3.1	4.0	3.7	3.8							
9-year	1.8	2.0	2.2	2.8	3.6	3.5	2.8	3.2	2.6	3.3	4.0	2.7	3.7	3.6								
10-year	1.7	1.9	2.2	2.5	3.3	3.4	2.5	3.0	2.5	3.2	3.9	2.7	3.5									
11-year	1.7	1.8	2.1	2.4	3.0	3.1	2.4	2.8	2.4	2.9	3.7	2.5										
12-year	1.6	1.7	2.1	2.4	2.9	3.0	2.2	2.6	2.3	2.8	3.5											
13-year	1.6	1.6	2.0	2.3	2.9	2.8	2.2	2.5	2.0	2.7												
14-year	1.5	1.6	1.9	2.1	2.7	2.8	2.0	2.5	1.7													
15-year	1.5	1.5	1.8	2.0	2.3	2.8	2.0	2.4														
16-year	1.5	1.4	1.7	2.0	2.1	2.5	1.7															
17-year	1.4	1.4	1.7	1.9	2.1	2.4																
18-year	1.3	1.3	1.7	1.8	2.0																	
19-year	1.2	1.3	1.6	1.8																		
20-year	1.1	1.3	1.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).



Table 22.10  
Cancer of the Pancreas (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	15.0	16.0	17.9	19.1	19.6	17.9	16.9	20.0	17.8	21.3	22.9	20.3	24.5	23.4	24.1	26.0	25.4	26.2	26.3	28.0	29.0	32.4
2-year	5.9	6.1	6.5	8.4	8.8	8.7	7.5	9.5	7.4	11.1	10.0	9.1	9.9	11.4	12.4	11.3	11.5	12.6	11.6	13.9	13.8	
3-year	3.5	3.8	4.4	5.2	6.4	5.5	4.6	7.4	4.2	7.3	6.9	5.7	6.3	7.0	7.9	8.2	7.3	8.2	7.6	9.8		
4-year	2.9	3.0	3.3	3.7	5.3	4.9	3.4	5.7	3.5	5.8	5.6	4.5	5.5	5.9	6.3	6.8	5.5	5.7	5.3			
5-year	2.5	2.5	2.9	3.2	4.7	4.2	2.6	5.3	2.9	5.1	4.7	3.9	4.9	5.1	5.3	6.0	4.2	4.7				
6-year	2.3	2.3	2.5	3.2	4.3	4.2	2.3	4.7	2.6	4.7	3.9	3.7	4.1	4.8	4.9	5.2	3.8					
7-year	2.2	1.9	2.3	2.9	3.8	3.5	2.0	4.4	2.6	4.2	3.6	3.4	3.9	4.4	4.4	4.7						
8-year	1.9	1.7	2.1	2.9	3.5	3.1	1.9	3.7	2.4	4.2	3.4	3.2	3.9	3.7	3.9							
9-year	1.8	1.6	1.8	2.6	3.5	3.0	1.8	3.5	2.2	4.0	3.4	2.8	3.6	3.6								
10-year	1.7	1.5	1.7	2.3	3.2	3.0	1.5	3.0	2.1	3.8	3.2	2.8	3.5									
11-year	1.7	1.5	1.6	2.2	3.1	3.0	1.3	3.0	2.0	3.2	2.9	2.8										
12-year	1.5	1.5	1.6	2.2	3.1	3.0	1.1	2.8	1.8	3.0	2.8											
13-year	1.5	1.4	1.5	2.2	3.1	3.0	1.1	2.5	1.7	3.0												
14-year	1.5	1.4	1.5	2.0	3.0	3.0	0.9	2.4	1.2													
15-year	1.5	1.3	1.3	2.0	2.9	3.0	0.9	2.3														
16-year	1.5	1.3	1.3	2.0	2.7	2.4	0.4															
17-year	1.4	1.3	1.3	2.0	2.5	2.4																
18-year	1.2	1.3	1.2	2.0	2.5																	
19-year	1.1	1.3	1.2	2.0																		
20-year	1.0	1.2	1.2																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 22.11  
Cancer of the Pancreas (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	15.2	18.1	18.4	19.5	20.5	22.6	19.3	19.1	19.2	18.5	20.1	21.3	22.6	23.3	25.7	26.3	22.5	22.0	29.6	25.8	28.4	28.8
2-year	6.3	6.9	7.5	8.0	9.5	9.3	9.2	8.7	9.3	8.3	10.1	8.7	11.2	10.5	11.1	12.8	11.2	11.4	14.1	12.8	13.6	
3-year	3.9	4.4	5.2	5.5	6.3	6.2	5.8	5.6	6.3	5.6	7.3	5.9	7.1	7.7	6.9	8.8	7.6	8.5	10.0	8.8		
4-year	3.0	3.6	4.3	4.4	5.2	5.3	5.2	4.7	4.4	4.0	5.6	4.4	5.7	6.3	5.8	7.0	5.9	7.2	8.6			
5-year	2.4	3.2	3.7	4.1	4.5	4.9	4.3	4.4	4.1	3.4	5.3	3.7	5.2	5.0	4.6	6.3	5.4	6.0				
6-year	2.3	2.9	3.4	3.7	4.2	4.9	4.1	3.6	3.7	3.0	5.0	3.7	5.0	4.3	4.3	6.1	4.8					
7-year	2.0	2.8	3.1	3.2	4.0	4.3	3.9	3.2	3.4	2.8	5.0	3.4	4.6	3.9	3.7	5.3						
8-year	1.8	2.5	2.8	3.2	3.7	4.3	3.9	3.0	2.9	2.6	4.8	2.9	4.1	3.6	3.5							
9-year	1.8	2.3	2.7	2.9	3.6	3.9	3.6	2.8	2.9	2.5	4.5	2.6	3.8	3.6								
10-year	1.7	2.2	2.6	2.6	3.4	3.5	3.5	2.8	2.8	2.5	4.5	2.5	3.6									
11-year	1.7	2.0	2.6	2.6	2.8	3.0	3.3	2.6	2.7	2.5	4.5	2.3										
12-year	1.7	1.9	2.5	2.5	2.7	2.7	3.1	2.4	2.7	2.5	4.2											
13-year	1.6	1.8	2.4	2.2	2.7	2.7	3.1	2.4	2.4	2.3												
14-year	1.5	1.7	2.4	2.1	2.4	2.7	3.0	2.4	2.1													
15-year	1.4	1.6	2.3	1.9	1.6	2.5	2.9	2.4														
16-year	1.4	1.5	2.0	1.8	1.6	2.5	2.7															
17-year	1.4	1.5	2.0	1.8	1.6	2.4																
18-year	1.4	1.4	2.0	1.6	1.5																	
19-year	1.4	1.4	2.0	1.4																		
20-year	1.3	1.4	2.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 23.9  
Cancer of the Prostate (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	91.9	94.0	95.9	97.7	98.8	99.6	99.4	99.2	98.9	99.3	99.5	99.6	99.7	99.9	100.0	99.9	99.9	100.0	99.9	99.9	99.9	99.9
2-year	85.1	87.9	91.5	94.8	97.4	98.4	98.2	98.1	97.8	98.2	98.9	99.1	99.7	99.7	100.0	99.9	99.8	99.9	99.8	99.9	99.9	99.9
3-year	78.8	82.0	87.2	92.4	95.9	97.8	97.0	96.8	97.2	97.6	98.6	98.6	99.7	99.4	100.0	99.9	99.6	99.9	99.7	99.9		
4-year	73.6	76.8	83.8	90.4	94.4	97.2	96.5	95.7	96.5	97.1	98.1	98.4	99.4	99.1	100.0	99.9	99.6	99.9	99.7			
5-year	69.2	72.4	80.9	88.9	93.0	96.4	95.5	94.9	95.9	96.7	97.7	98.4	99.4	99.1	100.0	99.9	99.4	99.9				
6-year	65.6	68.8	78.2	87.4	91.6	95.3	94.7	94.1	95.5	96.3	97.5	98.3	99.4	98.9	100.0	99.9	99.4					
7-year	62.1	65.3	75.7	85.3	90.8	94.3	93.6	93.1	95.1	95.7	97.5	98.0	99.4	98.8	100.0	99.9						
8-year	58.9	62.3	73.7	83.8	89.6	93.0	93.0	92.5	94.9	95.2	97.4	98.0	99.1	98.7	100.0							
9-year	56.1	60.0	71.7	82.5	88.6	92.2	91.8	91.8	94.5	94.8	97.1	97.4	98.9	98.7								
10-year	53.5	58.1	69.9	80.8	87.6	91.2	90.9	91.3	93.9	94.6	96.9	97.1	98.6									
11-year	51.0	55.9	68.3	79.3	86.3	90.7	90.3	90.2	93.2	94.3	96.7	96.8										
12-year	48.7	53.9	66.4	78.4	85.6	89.9	89.0	89.4	92.4	93.9	96.6											
13-year	46.9	52.1	64.8	77.7	84.6	89.1	88.5	88.7	91.4	93.5												
14-year	45.2	50.5	63.3	76.6	84.2	88.5	87.8	87.5	90.6													
15-year	44.0	49.2	61.8	75.6	83.3	87.7	86.6	87.2														
16-year	42.2	47.3	60.2	73.9	82.1	86.7	85.8															
17-year	40.5	46.2	59.7	72.8	81.6	85.9																
18-year	39.1	44.8	58.6	72.1	81.2																	
19-year	38.3	43.4	58.0	71.6																		
20-year	37.4	42.0	57.1																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 24.9  
Cancer of the Stomach (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	39.6	42.6	45.0	46.3	44.5	47.1	45.0	46.0	46.9	46.6	45.9	49.0	47.5	48.4	49.6	50.6	51.3	53.7	51.4	52.9	52.5	56.1
2-year	25.2	27.0	29.7	32.2	30.2	32.2	30.5	33.2	31.8	32.2	33.1	33.9	34.2	33.0	33.7	35.6	38.8	38.3	36.4	39.9	39.4	
3-year	19.8	21.3	23.6	25.0	24.8	26.0	24.0	26.6	26.6	27.2	26.3	28.3	27.7	27.2	29.0	30.5	33.6	32.9	31.7	34.1		
4-year	17.3	18.4	20.7	22.2	22.1	23.5	20.9	23.3	23.9	23.5	25.3	24.8	24.8	27.6	27.0	30.4	29.7	28.1				
5-year	15.7	16.7	19.4	20.0	20.7	22.1	19.4	22.0	21.9	22.5	22.0	23.0	23.2	23.0	26.3	25.9	27.7	27.9				
6-year	14.7	15.9	18.5	18.3	19.1	21.0	18.4	20.1	20.7	20.6	20.8	21.8	21.2	22.2	25.1	24.2	26.5					
7-year	14.2	15.2	17.6	17.7	18.6	19.6	17.3	19.1	19.5	20.1	20.0	21.1	20.1	21.3	24.4	23.6						
8-year	13.5	14.5	17.1	17.3	18.0	18.4	16.2	17.9	19.5	19.8	19.4	20.5	18.9	20.6	22.6							
9-year	13.2	14.1	16.6	16.1	17.2	17.5	15.4	17.9	19.0	19.4	19.3	19.6	18.5	19.8								
10-year	12.8	13.9	15.9	15.4	16.7	16.9	14.5	16.9	18.1	18.9	18.4	18.7	17.7									
11-year	12.3	13.4	15.4	15.1	16.7	16.1	14.1	16.6	17.8	17.7	17.8	18.5										
12-year	12.1	12.9	15.1	14.7	16.7	15.6	13.8	16.3	17.2	17.0	17.8											
13-year	11.8	12.9	14.9	14.7	16.2	15.5	13.3	15.6	16.8	15.9												
14-year	11.3	12.9	14.6	14.7	16.0	15.3	12.9	15.3	16.5													
15-year	10.9	12.4	14.3	14.6	15.6	14.7	12.6	15.1														
16-year	10.6	11.9	14.1	14.0	14.4	14.3	11.9															
17-year	10.3	11.7	13.9	13.8	14.2	14.3																
18-year	9.8	11.5	13.8	12.8	13.4																	
19-year	9.6	11.3	13.3	12.4																		
20-year	9.2	10.9	13.1																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 24.10  
Cancer of the Stomach (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	39.1	42.1	44.1	46.0	42.6	45.4	43.5	46.2	47.8	44.6	46.2	50.5	47.3	47.6	50.4	48.9	50.9	54.2	52.9	52.7	51.9	54.6
2-year	24.2	26.1	28.0	31.1	26.8	30.1	27.9	32.6	31.9	29.0	31.7	36.0	33.9	33.3	33.2	33.0	36.6	37.6	37.1	37.7	37.2	
3-year	18.6	20.4	21.6	23.7	21.3	24.2	22.3	25.8	25.9	24.3	24.9	29.6	27.4	27.4	28.5	27.1	31.6	30.9	31.7	32.7		
4-year	16.1	17.3	18.3	20.3	18.3	21.8	19.6	23.5	22.5	21.7	21.8	25.8	24.3	24.6	26.6	24.8	27.8	27.2	27.6			
5-year	14.5	15.5	16.8	17.7	16.2	20.5	18.0	21.5	20.8	20.4	20.0	23.4	22.3	22.4	25.5	23.3	24.9	25.9				
6-year	13.5	14.6	16.0	16.5	14.4	19.1	16.9	19.3	19.7	19.1	18.7	22.4	20.6	21.2	23.7	21.8	23.7					
7-year	12.8	13.9	15.1	15.8	13.5	17.9	15.4	18.2	18.7	18.8	17.8	21.4	18.8	20.2	23.1	21.2						
8-year	12.4	13.3	14.5	15.0	12.7	16.7	14.2	17.1	18.7	18.8	17.7	20.8	17.7	19.3	21.3							
9-year	12.2	13.1	14.0	14.2	12.2	15.8	13.4	16.8	18.6	17.9	17.5	20.1	16.9	17.8								
10-year	11.8	12.9	13.4	13.4	11.7	15.1	12.2	15.7	17.3	17.5	16.7	19.1	16.4									
11-year	11.5	12.5	12.9	12.5	11.7	14.6	12.1	15.5	17.1	15.6	15.9	19.1										
12-year	11.3	11.9	12.7	12.2	11.7	14.2	11.6	15.1	16.6	14.8	15.7											
13-year	11.2	11.8	12.7	12.2	10.8	14.1	11.6	14.0	16.0	14.6												
14-year	10.5	11.8	12.3	11.8	10.3	13.6	10.9	13.4	15.4													
15-year	10.0	11.3	12.0	11.3	9.4	12.5	10.4	13.0														
16-year	9.8	10.8	11.8	10.7	9.1	12.5	9.8															
17-year	9.4	10.6	11.5	10.6	9.1	12.5																
18-year	8.8	10.4	11.4	10.3	9.1																	
19-year	8.7	10.3	11.0	9.5																		
20-year	8.6	9.9	10.9																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 24.11  
Cancer of the Stomach (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	40.5	43.5	46.7	46.7	47.6	50.0	47.7	45.7	45.4	49.8	45.4	46.6	47.7	49.7	48.1	53.3	52.0	52.8	48.9	53.3	53.6	58.8
2-year	26.9	28.5	32.5	34.0	35.6	35.7	35.1	34.2	31.5	37.4	35.2	30.8	34.6	32.6	34.6	39.6	42.2	39.5	35.0	43.3	43.1	
3-year	21.8	22.9	26.9	27.0	30.3	28.8	26.9	28.0	27.7	31.9	28.5	26.5	28.2	26.9	29.7	35.4	36.8	36.1	31.8	36.3		
4-year	19.3	20.3	24.6	25.3	28.0	26.2	23.0	24.8	24.6	27.6	26.2	24.4	25.7	25.1	29.1	30.2	34.3	33.8	29.1			
5-year	17.5	18.8	23.8	23.7	27.6	24.6	21.6	23.0	23.6	25.8	24.9	22.3	24.3	23.9	27.7	29.6	32.0	31.2				
6-year	16.7	18.0	22.6	21.1	26.4	24.2	20.9	21.3	22.3	23.1	23.8	20.8	22.2	23.8	27.3	27.6	30.8					
7-year	16.3	17.2	22.0	20.5	26.1	22.3	20.2	20.5	21.0	21.9	23.2	20.6	21.9	23.0	26.7	27.2						
8-year	15.3	16.3	21.4	20.5	26.1	21.0	19.4	19.1	20.6	21.2	22.1	20.1	20.6	22.7	24.8							
9-year	14.6	15.9	21.0	18.8	24.9	20.4	18.5	19.1	19.6	21.2	22.1	18.9	20.6	22.7								
10-year	14.4	15.5	19.9	18.8	24.2	19.9	18.2	18.8	19.4	21.2	21.1	17.7	19.6									
11-year	13.8	14.9	19.6	18.6	24.2	18.3	17.4	18.3	18.9	21.0	20.6	17.4										
12-year	13.3	14.7	19.1	18.6	24.2	17.7	17.4	18.1	18.2	20.5	20.6											
13-year	12.7	14.6	18.7	18.6	24.2	17.5	16.2	18.1	18.0	17.8												
14-year	12.5	14.4	18.5	18.6	24.2	17.5	16.1	18.1	18.0													
15-year	12.2	13.9	18.1	18.6	24.2	17.5	16.1	18.1														
16-year	11.7	13.7	17.9	18.6	22.6	17.3	15.4															
17-year	11.5	13.5	17.7	18.6	21.4	17.0																
18-year	11.4	13.3	17.7	16.6	19.8																	
19-year	10.9	12.9	17.3	16.6																		
20-year	10.2	12.5	16.7																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 25.9  
Cancer of the Testis (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	93.4	96.1	98.1	98.2	98.5	98.0	96.6	97.7	98.3	97.9	97.4	98.2	96.9	97.9	98.5	98.2	98.5	97.9	98.3	98.9	98.9	98.8
2-year	88.6	92.7	96.2	96.7	97.0	96.7	95.7	96.5	97.4	96.4	95.6	96.6	96.3	96.7	97.4	97.1	96.8	96.4	97.3	98.5	97.7	
3-year	86.5	92.1	95.5	96.2	96.1	95.5	95.0	96.1	96.4	95.8	95.1	96.2	95.7	95.9	97.1	96.8	96.5	95.7	97.1	98.4		
4-year	85.6	91.5	95.0	96.1	95.7	95.4	94.9	96.1	96.3	95.8	94.9	96.1	95.6	95.9	96.8	96.7	96.3	95.3	96.8			
5-year	85.0	91.2	94.7	96.1	95.7	95.3	94.8	96.0	96.3	95.8	94.9	95.8	95.6	95.9	96.5	96.2	96.0	95.2				
6-year	84.7	91.0	94.5	95.8	95.5	94.9	94.8	96.0	96.3	95.8	94.7	95.6	95.5	95.9	96.1	96.0	96.0					
7-year	84.6	90.7	94.4	95.7	95.4	94.6	94.8	95.8	96.3	95.8	94.7	95.4	95.5	95.9	96.1	96.0						
8-year	84.5	90.5	94.1	95.6	95.2	94.0	94.8	95.1	96.3	95.8	94.7	95.4	95.4	95.9	96.1							
9-year	84.5	90.1	93.9	95.5	95.0	94.0	94.6	95.1	96.2	95.8	94.7	95.4	95.1	95.9								
10-year	83.6	89.9	93.8	95.4	94.7	93.9	94.6	95.1	96.1	95.8	94.7	94.6	95.1									
11-year	83.1	89.7	93.6	95.3	94.7	93.8	94.6	95.1	96.1	95.8	94.7	94.1										
12-year	82.9	89.7	93.5	94.9	94.5	93.8	94.6	95.0	96.1	95.8	94.7											
13-year	82.5	89.2	93.3	94.9	94.5	93.8	94.6	94.9	96.0	95.8												
14-year	82.0	88.8	93.1	94.9	94.5	93.6	94.6	94.8	95.8													
15-year	81.8	88.5	93.1	94.9	94.4	93.5	94.6	94.8														
16-year	81.3	87.9	92.9	94.9	94.4	92.9	94.6															
17-year	80.7	87.6	92.6	94.7	94.4	92.9																
18-year	80.1	87.4	92.6	94.7	93.8																	
19-year	79.5	87.2	92.4	94.7																		
20-year	79.4	86.9	92.2																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 26.9  
Cancer of the Thyroid (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
1-year	95.0	95.2	96.0	96.6	96.6	95.6	95.2	97.5	96.7	96.9	96.5	97.8	97.4	97.7	97.6	97.2	98.0	98.4	97.8	98.5	98.6	98.3
2-year	93.6	94.5	95.3	96.2	95.9	94.8	94.7	97.0	96.2	96.5	96.2	97.1	97.1	97.1	97.4	97.1	97.9	97.9	97.3	98.4	98.2	
3-year	92.9	93.7	94.8	95.5	95.6	94.2	94.7	96.4	96.0	96.0	96.2	96.9	97.1	97.1	96.9	97.1	97.6	97.9	97.3	98.3		
4-year	92.5	93.4	94.3	95.5	95.1	93.8	94.3	96.0	95.9	95.4	95.7	96.4	97.0	97.0	96.9	97.1	97.4	97.8	97.3			
5-year	92.1	93.3	94.1	95.5	94.4	93.0	94.3	95.7	95.8	94.9	95.4	96.3	96.8	96.4	96.2	97.1	97.3	97.4				
6-year	91.7	93.2	93.8	95.0	94.1	92.8	94.1	95.3	95.8	94.9	95.2	95.9	96.3	96.4	96.1	96.8	96.9					
7-year	91.5	92.7	93.5	94.7	93.7	92.7	93.9	95.2	95.6	94.5	95.2	95.9	96.2	96.2	96.1	96.5						
8-year	91.1	92.6	93.3	94.7	93.7	92.7	93.9	94.5	95.5	94.3	95.2	95.9	96.0	96.0	95.9							
9-year	90.7	92.3	93.1	94.7	93.6	92.7	93.7	94.2	95.5	93.8	95.1	95.9	95.6	96.0								
10-year	90.6	92.0	92.9	94.0	93.3	92.7	93.1	94.0	95.1	93.6	95.1	95.9	95.3									
11-year	90.3	91.7	92.7	93.5	93.1	92.4	92.8	94.0	95.0	93.3	95.1	95.9										
12-year	90.3	91.5	92.6	93.4	92.7	92.1	92.7	93.6	95.0	93.0	95.1											
13-year	90.1	91.2	92.5	93.0	92.7	92.1	92.7	93.6	95.0	92.9												
14-year	90.0	91.1	92.4	92.5	92.4	92.1	92.3	93.6	95.0													
15-year	89.8	90.7	92.3	92.4	91.7	91.8	92.0	93.0														
16-year	89.7	90.6	92.3	92.2	91.2	91.8	91.9															
17-year	89.6	90.6	92.2	92.2	91.2	91.6																
18-year	89.3	90.4	92.2	92.2	91.2																	
19-year	89.2	90.4	91.9	92.2																		
20-year	89.2	90.4	91.9																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).



Table 26.10  
Cancer of the Thyroid (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	94.6	92.5	95.0	93.9	95.3	93.9	93.3	96.7	93.4	93.8	92.4	97.3	97.0	96.4	96.2	95.2	96.2	96.7	97.0	96.4	97.4	96.9
2-year	92.6	92.0	93.8	92.5	92.5	92.9	92.5	95.5	91.9	93.5	91.0	95.4	96.2	95.0	95.5	94.9	95.5	95.1	95.5	96.2	96.6	
3-year	91.6	90.8	93.6	91.5	91.3	92.0	91.8	94.8	91.7	92.3	91.0	94.7	96.2	95.0	94.2	94.9	94.2	94.8	95.5	96.1		
4-year	90.5	90.4	92.2	91.5	91.1	91.3	90.2	94.2	91.4	91.7	89.4	93.7	96.2	95.0	93.6	94.6	93.6	94.4	95.5			
5-year	89.9	90.3	91.6	91.1	89.4	90.7	90.0	93.4	91.3	89.9	88.5	93.7	96.2	94.0	92.3	94.3	93.3	94.2				
6-year	89.6	89.9	91.3	90.4	88.4	90.7	89.8	93.0	90.4	89.9	88.0	93.5	95.8	94.0	92.0	93.9	91.7					
7-year	89.6	89.4	90.1	90.4	87.5	90.7	89.6	92.3	88.8	89.6	88.0	93.3	95.4	93.6	92.0	92.3						
8-year	89.3	89.3	89.7	90.4	87.1	90.7	89.2	91.3	88.8	88.9	88.0	93.3	95.3	93.5	90.9							
9-year	88.8	88.7	88.6	90.4	87.1	90.7	88.9	90.9	88.8	86.9	87.3	93.3	93.1	93.5								
10-year	88.6	87.9	88.5	90.3	87.1	90.7	87.3	90.9	87.5	86.6	87.3	93.3	92.8									
11-year	87.5	87.4	88.5	90.2	86.9	90.5	87.3	90.9	86.3	85.2	87.0	93.3										
12-year	87.5	86.5	88.5	89.0	86.8	89.1	87.3	89.8	86.1	84.3	87.0											
13-year	87.4	86.1	87.9	86.9	86.8	89.1	87.3	89.4	86.1	83.5												
14-year	87.3	86.1	87.8	86.7	86.5	89.0	86.7	89.4	86.1													
15-year	86.7	85.3	87.8	83.7	85.1	88.9	85.8	89.4														
16-year	86.5	85.3	87.8	83.4	85.1	88.4	85.5															
17-year	86.5	84.9	87.8	83.4	85.1	87.3																
18-year	86.5	84.4	87.8	83.4	84.4																	
19-year	86.5	84.4	87.8	83.4																		
20-year	86.5	84.4	87.8																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 26.11  
Cancer of the Thyroid (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	95.1	96.0	96.4	97.4	97.1	96.3	95.8	97.8	97.8	97.9	97.7	97.9	97.5	98.0	98.0	97.7	98.5	98.9	98.0	99.0	98.9	98.7
2-year	93.9	95.4	95.7	97.3	96.9	95.4	95.6	97.4	97.4	97.5	97.7	97.6	97.3	97.7	98.0	97.7	98.5	98.7	97.9	99.0	98.6	
3-year	93.3	94.7	95.2	96.8	96.9	95.0	95.6	96.8	97.3	97.1	97.7	97.6	97.3	97.5	97.7	97.7	98.5	98.7	97.9	99.0		
4-year	93.2	94.5	95.0	96.8	96.3	94.7	95.6	96.6	97.3	96.5	97.5	97.2	97.1	97.5	97.7	97.7	98.5	98.7	97.9			
5-year	92.8	94.3	94.9	96.8	96.0	93.8	95.6	96.4	97.2	96.4	97.5	97.0	96.8	97.1	97.3	97.7	98.4	98.3				
6-year	92.4	94.2	94.7	96.4	95.9	93.6	95.6	95.9	97.2	96.3	97.3	96.5	96.5	97.0	97.3	97.6	98.3					
7-year	92.1	93.8	94.6	95.8	95.6	93.4	95.4	95.9	97.2	95.8	97.3	96.5	96.2	97.0	97.3	97.6						
8-year	91.7	93.7	94.5	95.8	95.6	93.4	95.4	95.5	97.2	95.8	97.3	96.5	96.1	96.6	97.3							
9-year	91.4	93.6	94.5	95.8	95.6	93.4	95.4	95.2	97.2	95.8	97.3	96.5	96.1	96.6								
10-year	91.2	93.4	94.3	95.1	95.2	93.4	95.3	94.8	97.2	95.7	97.3	96.5	95.9									
11-year	91.2	93.1	94.0	94.5	95.0	93.0	94.8	94.8	97.2	95.7	97.3	96.5										
12-year	91.2	93.1	94.0	94.5	94.5	93.0	94.7	94.8	97.2	95.6	97.3											
13-year	91.0	92.9	94.0	94.5	94.4	93.0	94.6	94.7	97.2	95.6												
14-year	90.8	92.8	93.8	94.2	94.2	93.0	94.4	94.7	97.2													
15-year	90.7	92.5	93.8	94.2	93.6	92.7	94.3	93.9														
16-year	90.7	92.3	93.6	94.2	93.0	92.7	94.1															
17-year	90.5	92.3	93.5	94.2	93.0	92.7																
18-year	89.9	92.3	93.4	94.2	93.0																	
19-year	89.7	92.3	93.2	94.2																		
20-year	89.7	92.3	93.2																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 27.9  
Cancer of the Urinary Bladder (Invasive and *In Situ*)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	87.4	89.2	90.4	90.0	90.1	90.3	90.8	91.0	91.0	90.1	91.7	91.3	90.5	91.4	91.4	90.4	90.9	91.2	91.0	91.5	90.5	89.5
2-year	81.1	83.8	85.4	85.8	85.3	85.0	85.9	86.2	87.0	85.9	86.5	86.7	85.3	86.8	86.7	86.1	86.0	86.4	85.7	86.8	85.2	
3-year	77.9	80.7	82.4	82.7	82.9	82.3	83.2	84.2	84.3	83.7	83.3	83.7	82.6	83.9	84.8	83.9	83.8	83.9	82.7	84.0		
4-year	75.1	78.2	80.2	81.0	81.3	81.3	81.5	81.9	82.0	81.4	81.0	81.8	80.6	81.9	82.8	81.4	81.8	81.7	81.4			
5-year	73.2	76.4	78.3	79.9	79.6	79.3	79.8	80.4	81.4	79.8	78.9	79.6	78.5	80.5	81.9	80.2	80.0	80.4				
6-year	71.3	74.7	76.7	78.3	78.5	77.8	78.1	78.6	79.5	78.5	77.1	77.9	77.1	79.6	80.4	78.7	78.7					
7-year	69.2	73.2	75.2	77.1	76.1	75.8	76.3	76.9	77.6	77.7	75.9	76.8	75.9	78.2	78.6	77.6						
8-year	67.7	72.0	73.6	75.1	74.5	74.4	75.4	76.2	76.6	76.1	74.9	75.1	74.8	76.6	77.4							
9-year	66.1	70.3	72.4	73.0	73.3	73.6	73.7	75.1	75.0	74.6	73.5	74.1	73.6	75.7								
10-year	64.1	69.0	71.1	70.9	72.0	72.2	72.9	74.3	74.1	73.2	72.2	72.3	73.0									
11-year	62.3	67.7	70.0	69.2	71.0	71.1	71.5	72.9	73.4	72.1	71.4	71.3										
12-year	61.1	66.4	68.5	67.5	70.2	70.4	70.7	71.8	72.2	70.5	70.3											
13-year	60.0	65.3	67.2	66.9	69.0	69.6	69.3	71.6	69.8	69.4												
14-year	58.7	63.8	65.7	65.5	69.0	68.4	68.4	70.2	68.1													
15-year	57.4	62.4	64.4	64.5	66.0	66.3	66.8	69.6														
16-year	56.3	60.8	62.9	62.6	65.3	65.0	65.7															
17-year	55.2	59.8	62.0	60.9	64.1	64.0																
18-year	54.4	58.8	61.0	59.5	63.2																	
19-year	53.9	57.9	60.3	59.2																		
20-year	52.9	57.3	59.7																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 27.10  
Cancer of the Urinary Bladder (Invasive and *In Situ*)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	89.4	90.4	92.1	91.9	91.5	91.9	92.4	92.7	92.4	91.7	93.4	93.0	92.2	92.7	92.0	92.5	92.6	92.4	92.6	92.8	92.0	91.1
2-year	83.1	85.4	87.3	87.9	87.4	87.3	88.2	87.8	88.4	87.5	88.5	88.5	87.0	88.2	87.2	88.8	87.6	88.1	87.4	88.3	86.7	
3-year	79.7	82.3	84.4	85.1	85.3	84.9	85.5	85.9	85.9	85.2	85.4	85.1	84.4	85.0	85.5	86.5	85.2	85.6	84.4	85.5		
4-year	76.8	79.7	82.1	83.3	83.8	84.2	83.5	83.5	83.4	82.7	82.8	83.0	82.1	82.7	83.5	83.7	83.2	83.5	83.2			
5-year	74.5	77.8	80.3	82.3	81.8	82.1	81.7	82.1	83.0	81.1	80.3	80.6	79.7	81.2	82.7	82.4	81.2	82.3				
6-year	72.6	76.0	78.6	80.4	80.4	80.9	80.4	80.2	81.1	79.9	78.2	79.2	78.1	80.5	81.2	80.9	79.8					
7-year	70.3	74.7	76.9	78.9	77.6	78.7	78.4	78.2	79.3	78.7	76.9	78.1	76.9	78.9	78.9	80.0						
8-year	68.5	73.4	75.2	76.8	75.9	77.0	77.7	77.5	78.0	77.2	76.0	76.2	75.7	77.0	78.0							
9-year	66.7	71.7	74.0	74.3	74.5	75.6	76.5	76.6	75.8	76.0	74.5	75.4	74.9	75.8								
10-year	64.6	70.4	72.6	72.7	72.8	74.4	75.6	76.0	75.0	74.3	73.3	73.5	74.7									
11-year	62.5	69.1	71.7	70.8	71.9	73.2	74.5	74.4	74.3	73.1	72.4	72.3										
12-year	61.2	67.6	70.0	69.3	71.4	72.2	73.7	73.0	72.9	71.4	71.0											
13-year	59.9	66.6	68.6	68.9	70.3	71.5	71.9	73.0	71.2	70.3												
14-year	58.6	64.8	67.0	67.4	70.3	71.4	70.8	71.9	69.3													
15-year	57.5	63.4	65.7	65.8	67.9	69.8	69.9	70.9														
16-year	55.9	61.5	64.2	63.8	67.5	67.6	69.1															
17-year	54.9	60.5	63.4	62.1	66.7	66.5																
18-year	53.9	59.5	62.2	60.9	66.7																	
19-year	53.1	58.8	61.4	60.2																		
20-year	52.2	58.4	61.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 27.11  
Cancer of the Urinary Bladder (Invasive and *In Situ*)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	81.8	85.6	85.4	84.2	85.8	85.5	86.2	86.0	86.9	85.7	86.8	86.3	85.8	87.5	89.6	84.5	86.1	87.5	86.4	87.5	85.4	84.5
2-year	75.5	79.3	80.0	79.4	79.2	78.5	79.2	81.3	82.8	81.2	80.7	81.5	80.4	82.3	85.2	78.4	81.5	81.5	80.8	82.0	80.2	
3-year	72.7	76.1	76.8	75.6	75.7	74.6	76.4	79.2	79.9	79.1	77.4	79.5	77.5	80.6	82.6	76.4	79.6	78.8	77.6	79.3		
4-year	70.3	73.9	74.7	74.1	73.8	73.1	75.7	77.1	78.2	77.3	75.8	77.9	76.0	79.2	80.6	74.8	77.7	76.5	76.2			
5-year	69.3	72.4	72.6	72.7	73.1	71.0	74.3	75.4	76.5	76.0	74.7	76.5	75.1	78.3	79.6	73.9	76.5	74.1				
6-year	67.7	70.7	71.1	71.8	72.6	68.7	71.4	74.2	75.0	74.5	73.9	74.1	74.2	76.5	78.1	72.3	75.6					
7-year	66.1	69.0	70.2	71.6	71.5	67.4	70.3	72.9	73.0	74.3	72.9	72.7	73.1	76.1	77.6	70.2						
8-year	65.3	68.0	68.8	70.0	70.4	66.9	68.4	72.2	72.4	73.1	71.6	71.8	72.2	75.2	75.4							
9-year	64.3	66.3	67.9	69.1	69.9	66.9	65.8	70.4	72.1	70.5	70.5	70.1	69.6	75.2								
10-year	62.7	64.9	66.5	65.5	69.5	65.8	64.9	69.4	71.3	70.0	68.8	68.6	68.3									
11-year	61.5	63.6	65.2	63.9	68.1	64.6	63.0	68.7	70.3	69.2	68.0	68.3										
12-year	60.4	62.7	64.1	61.4	66.7	64.5	62.1	67.9	69.9	67.8	68.0											
13-year	59.9	61.6	63.1	60.7	65.1	63.7	61.8	67.1	65.7	66.8												
14-year	58.2	60.7	61.6	59.7	64.8	59.3	61.3	65.1	64.8													
15-year	56.9	59.6	60.4	59.7	60.6	56.7	58.0	65.1														
16-year	56.7	58.7	59.2	58.8	58.5	56.7	56.2															
17-year	55.6	57.6	57.8	57.1	56.3	56.7																
18-year	55.3	56.4	57.4	55.5	53.2																	
19-year	55.3	54.9	56.7	55.5																		
20-year	54.3	54.1	55.6																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 28.10  
All Cancer Sites (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Ages 0-14

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	81.3	85.0	87.3	89.5	91.0	91.1	88.6	90.6	90.5	92.9	91.2	91.0	91.3	92.5	92.1	91.8	94.1	92.9	92.5	94.1	92.2	93.0
2-year	71.1	75.8	78.4	81.8	82.7	84.1	82.3	84.7	84.8	86.4	84.8	84.4	85.3	85.9	87.1	87.5	89.3	88.4	88.5	89.3	88.1	
3-year	65.4	70.8	74.3	77.7	79.4	81.1	79.0	81.7	81.3	83.7	82.4	81.7	82.8	83.5	83.9	83.9	86.0	85.1	85.1	88.0		
4-year	62.1	67.6	71.8	75.5	76.7	78.7	78.2	79.0	78.9	80.7	80.5	78.9	81.1	81.6	82.3	82.1	85.1	83.6	83.5			
5-year	59.9	65.9	70.7	74.4	75.6	77.6	76.6	78.4	77.4	80.3	79.2	78.0	79.2	80.7	81.9	81.0	83.9	82.7				
6-year	58.2	64.8	69.7	73.6	74.9	76.3	76.1	78.1	76.4	79.6	78.4	76.5	78.7	79.5	80.9	80.2	83.5					
7-year	57.1	63.9	68.7	72.7	74.3	75.9	75.6	77.4	75.9	79.0	77.9	76.1	78.0	79.0	79.8	79.7						
8-year	56.4	63.2	68.1	72.3	73.6	75.8	75.0	77.1	75.0	78.5	76.7	75.6	77.7	78.3	79.8							
9-year	56.0	62.6	67.7	71.8	73.2	75.6	74.7	76.2	74.6	78.1	76.5	75.0	77.1	78.1								
10-year	55.8	62.1	67.3	71.4	73.1	75.5	74.7	76.0	74.5	77.9	75.9	74.7	76.7									
11-year	55.4	61.7	67.0	70.9	72.8	75.4	74.5	75.4	74.3	77.7	75.6	74.2										
12-year	55.1	61.2	66.7	70.8	72.6	75.3	74.5	74.7	74.0	77.6	75.2											
13-year	54.9	60.9	66.5	70.2	72.4	75.1	74.3	74.5	74.0	77.5												
14-year	54.5	60.6	66.1	70.0	72.2	74.7	73.8	74.4	74.0													
15-year	54.3	60.3	66.0	69.7	71.9	74.4	73.8	74.3														
16-year	53.9	60.0	65.7	69.2	71.8	74.2	73.4															
17-year	53.5	59.8	65.5	68.7	71.6	73.9																
18-year	53.3	59.5	65.3	68.3	71.5																	
19-year	53.0	59.3	65.2	68.2																		
20-year	52.9	59.1	64.9																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 28.11  
All Cancer Sites (Invasive)

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Ages 0-19

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	83.1	86.3	88.8	90.6	91.4	90.5	89.8	90.7	90.3	93.2	91.7	92.1	91.2	91.3	92.3	92.5	94.1	92.5	92.8	94.3	93.2	94.2
2-year	73.2	77.5	80.4	83.1	83.7	83.5	82.9	84.3	84.9	87.4	85.1	85.4	85.8	84.3	87.3	87.6	89.2	88.1	89.0	89.0	88.6	
3-year	68.1	72.5	76.5	78.7	80.1	80.5	79.8	80.9	81.6	85.1	82.0	82.6	83.0	82.0	84.1	84.1	86.2	84.5	86.2	87.4		
4-year	64.9	69.6	73.9	76.5	78.0	78.3	78.4	78.9	79.9	82.2	80.4	80.0	81.1	80.3	82.3	82.1	85.1	83.3	84.6			
5-year	62.9	67.9	72.9	75.2	76.7	77.0	76.8	78.2	78.5	81.6	78.7	78.8	79.5	79.2	81.7	80.9	84.0	82.2				
6-year	61.2	66.5	71.9	74.2	76.0	75.7	76.3	77.7	77.4	81.0	77.8	77.6	79.1	78.3	81.0	79.9	83.5					
7-year	60.3	65.7	71.0	73.3	75.2	75.2	75.9	77.1	76.9	80.6	77.3	77.3	78.6	77.7	79.9	79.5						
8-year	59.5	64.9	70.4	72.7	74.6	75.1	75.2	76.8	76.0	80.0	76.6	76.9	78.4	77.1	79.8							
9-year	58.9	64.2	70.0	72.2	74.3	74.9	74.9	76.1	75.7	79.6	76.3	76.4	78.0	76.9								
10-year	58.6	63.8	69.6	72.0	73.9	74.9	74.8	75.7	75.5	79.4	75.8	76.1	77.6									
11-year	58.3	63.4	69.3	71.3	73.5	74.8	74.6	75.2	75.4	79.1	75.4	75.6										
12-year	57.9	63.0	69.0	71.2	73.1	74.7	74.6	74.8	75.1	79.0	75.2											
13-year	57.8	62.7	68.7	70.5	73.0	74.4	74.3	74.6	75.1	78.9												
14-year	57.4	62.3	68.2	70.3	72.8	74.1	73.8	74.4	75.0													
15-year	57.2	62.0	68.0	70.0	72.6	73.8	73.8	74.4														
16-year	56.9	61.6	67.7	69.5	72.5	73.6	73.4															
17-year	56.5	61.4	67.5	68.7	72.4	73.5																
18-year	56.3	61.1	67.3	68.5	72.2																	
19-year	56.0	60.9	67.0	68.4																		
20-year	55.7	60.6	66.7																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).