

Table 5.7
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2005-2009</u>						
Age at Diagnosis/Death:						
All ages	8.1	8.0	9.8	2.4	2.2	4.3
Under 65	7.5	7.6	8.1	1.9	1.7	2.9
65 and over	12.1	10.7	21.5	6.1	5.3	13.6
All ages (IARC world std) ^c	6.4	6.3	7.4	1.8	1.6	3.1
<u>Age-Specific Rates, 2005-2009</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	0.2	0.2	-	-	-	-
20-24	1.5	1.5	1.7	0.1	0.1	0.2
25-29	5.1	5.6	4.2	0.5	0.4	0.7
30-34	10.5	11.4	8.8	1.3	1.2	1.9
35-39	14.6	15.6	12.1	2.4	2.4	3.2
40-44	15.1	15.3	16.2	3.3	3.0	5.3
45-49	14.0	13.7	14.6	4.1	3.8	6.6
50-54	13.2	12.6	16.9	4.6	4.2	7.2
55-59	12.5	11.8	17.1	5.0	4.6	7.8
60-64	13.3	11.9	20.3	5.2	4.7	9.6
65-69	14.1	13.0	22.1	5.8	5.0	12.4
70-74	13.0	11.7	18.8	5.7	5.1	11.3
75-79	10.9	9.5	21.0	5.9	5.1	13.8
80-84	10.4	8.8	22.2	6.7	6.0	14.8
85+	9.8	8.2	25.9	7.1	6.2	19.4

^a SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.